

Self-directed treatment for lower limb wounds in persons with diabetes: a short report

Patrick Harnarayan
Shamir O Cawich
Shariful Islam
Shivaa Ramsewak
Vijay Naraynsingh

Department of Clinical Surgical
Sciences, University of the West
Indies, St Augustine Campus,
Trinidad and Tobago

Aim: There has been little focus on self-directed treatment for lower limb wounds, although it is a common practice among persons with diabetes across the Caribbean. We sought to document this practice in a Caribbean nation.

Methods: We prospectively interviewed all consecutive patients with diabetes who were admitted with lower limb wounds at the San Fernando General Hospital in Trinidad and Tobago over a period of 18 months. A questionnaire was used to collect data on patient demographics, use of self-directed treatment, and details of these treatments.

Results: Of 839 persons with diabetes who were admitted with infected lower limb wounds, 344 (41%) admitted to self-directed treatment before seeking medical attention. These patients were predominantly male (59.9%) at a mean age of 56.4 ± 12.4 years. The practice was most common in persons of Afro-Caribbean descent (45.9%) and those with type 2 diabetes (93.9%). In this group, 255 (74.4%) patients were previously admitted to hospital for lower limb infections. And of those, 32 (12.6%) had a prior amputation and 108 (42.4%) had at least one operative debridement specifically for foot infections.

Conclusion: Caribbean cultural practices may be an important contributor to negative outcomes when treating lower limb wounds in persons with diabetes. Despite being acutely aware of the potentially devastating consequences of inadequate treatment, 41% of our patients with diabetes still opted to use self-directed treatment for lower limb wounds. This deserves further study in order to give a more tailored approach in care delivery.

Keywords: diabetes, foot infection, alternative, home, remedy, amputation

Background

The treatment of infected lower limb wounds in persons with diabetes required an annual national expenditure of US\$85 million (0.4% of the gross domestic product) in Trinidad and Tobago between June 2012 and June 2013.¹ This has generated significant interest from policy makers and researchers across the region, with a heavy focus on the epidemiology,²⁻⁶ prevention,⁷⁻⁹ and therapeutic protocols¹⁰⁻¹³ used to treat infected wounds on the lower limbs of persons with diabetes.

However, there has been little focus on these patients' practices of self-directed treatment despite it being a common practice across the Caribbean. This study sought to document the practice of self-directed treatment of infected lower limb wounds in persons with diabetes in a small Caribbean country.

Methods

This study was carried out at the San Fernando General Hospital in Trinidad and Tobago. This is one of three tertiary referral hospitals on the island and serves a catchment of approximately 400,000 persons. In order to ensure that there is no financial

Correspondence: Shamir O Cawich
Department of Clinical Surgical
Sciences, University of the West
Indies, St Augustine Campus,
Trinidad and Tobago
Tel +1 868 798 9627
Email socawich@hotmail.com

barrier to health care access, the Government of Trinidad and Tobago prohibits user fees from being generated at this facility for all legal residents.

Patients who present to the San Fernando General Hospital with infected lower limb wounds are evaluated in the casualty department. These patients are admitted to hospital when there are systemic signs of infection, evidence of deep tissue infection, leukocytosis, metabolic derangements and/or ischemia present.

Once patients are admitted to hospital, the attending surgeon is responsible for care. They routinely evaluate the patients for limb ischemia, neuropathy, deep-seated collections/abscesses, ulcers, and generalized sepsis. The PEDIS (Perfusion, Extent/size, Depth/tissue loss, Infection, and Sensation) classification system developed by the International Working Group on the Diabetic Foot¹⁴ was used to classify diabetic foot ulcers at this facility. This is a standardized grading system that assesses foot ulcers based on the presence of five parameters: limb ischemia, wound size, wound depth, infection, and neuropathy.

Permission to interview patients was granted by the University of the West Indies' Ethics Committee. Admission registers were used to identify all consecutive patients with diabetes admitted to the San Fernando General Hospital with a diagnosis of infected lower limb wounds over 18 months from January 1, 2012 to June 30, 2013. All consecutive patients admitted were interviewed within 24 hours of admission. Patients were excluded if they were younger than 16 years of age, declined to participate, or did not admit to the use of self-directed therapy. A questionnaire was used to collect data on patient demographics, social circumstances, lifestyle habits, medical therapy, self-directed treatment, and details of these methods.

The data were analyzed using the Statistical Package for the Social Sciences (SPSS) version 14 (SPSS Inc., Chicago, IL, USA). This statistical software is widely used for data management and statistical analysis on data sets. Descriptive analyses (cross tabulations, frequencies, and descriptive ratio statistics) were generated using the populated data spreadsheets.

Results

There were 839 patients with diabetes admitted with infected lower limb wounds over the study period and 344 (41%) admitted for deliberate use of self-directed treatments before seeking professional medical care. These patients were predominantly male (59.9% versus 40.1%) at an average age of 56.4 years (standard deviation [SD] ± 12.4 ; range 24–93).

Table 1 Baseline characteristics of 344 patients who used self-directed treatment instead of presenting for medical advice and treatment

Characteristic	N (%)
Duration of diabetes (mean \pm SD)	12.9 \pm 7.9 years
Current smokers	98 (28.5%)
Regular alcohol use	120 (34.9%)
Male sex	206 (59.9%)
Female sex	138 (40.1%)
Afro-Caribbean descent	158 (45.9%)
East Indian descent	141 (41.0%)
Mixed descent	41 (11.9%)
Chinese descent	4 (1.2%)
Prior counseling on diabetes management	344 (100%)
Prior counseling on foot complications	344 (100%)

Abbreviation: SD, standard deviation.

The majority used oral hypoglycemics (177, 51.5%) to control their diabetes. Others used combinations of insulin and oral hypoglycemics (103, 29.9%), used insulin only (46, 13.4%), or used only self-directed treatment 18 (5.2%) for glycemic control.

The baseline characteristics of the patients who used self-directed treatment are outlined in Table 1. The practice was most common in persons of Afro-Caribbean descent (45.9%) and persons with type 2 (formerly known as non-insulin dependent or adult onset) diabetes (93.9%). They had lived with diabetes for a mean duration of 12.9 years (SD ± 7.9 ; median 12; mode 10) and all patients reported that they were previously counseled on the importance of glycemic control, foot care, and early presentation of lower limb wounds.

Nevertheless, the patients had poor glycemic control in the weeks prior to hospital presentation as evidenced by mean glycosylated hemoglobin (HbA_{1c}) levels of 7.82% (SD ± 1.56 ; median 7.9; mode 8.78). Additionally, 255 (74.4%) of these patients had previous hospital admissions for diabetic foot complications. And of those, 32 (12.6%) had a prior amputation and 108 (42.4%) had at least one operative debridement specifically for foot infections.

The patients attempted trials of self-directed treatment that lasted for a mean duration of 8.8 days (SD ± 5.4 ; range 1–60). Multiple agents were used as outlined in Table 2. Topical agents were more commonly employed than oral treatments (85.1% versus 9.6%, respectively). Unspecified oral therapies were used in 33 cases; 9 (2.6%) patients had “pills” purchased from informal vendors, 12 (3.5%) had “left-over pills” from uncompleted antibiotic courses previously prescribed for other persons to treat other diseases, and 12 (3.5%) had unspecified herbal concoctions specifically to treat lower limb infections.

Table 2 Self-directed treatment for diabetic foot infections in Trinidad and Tobago

Agent	n=344	Method of use	Dose	Frequency
Oral Therapies	33			
Unspecified herbal concoctions	12 (3.5%)	Premixed herbal mixtures	One cup	Twice daily
"Left-over pills"	12 (3.5%)	"Left-over pills" from uncompleted antibiotic courses that were previously prescribed for other persons to treat other diseases – patients usually followed directions as printed on the dispensed container	Variable	Variable
"Pills" purchased from informal vendors	9 (2.6%)	Red and white "pills" as provided by informal vendors	One pill	Twice daily
Topical Therapies	311			
Soft candle (paraffin wax)	66 (19.2%)	Melt candle with open flame and allow hot wax to drop onto the wound, then cover with a brown paper bag	Unspecified	Variable
Honey	56 (16.3%)	Solution applied directly onto the wound	Generous	Variable
Wonder of world (<i>Kalanchoe Pinnata</i>)	40 (11.6%)	Heat leaves and apply leaf as a plaster onto the wound (\pm paraffin wax)	Unspecified	Once daily
Ichthammol ointment	36 (10.5%)	Apply paste directly onto wound and leave uncovered	Unspecified	Once daily
Aloe vera	36 (10.5%)	Slice aloe leaves lengthwise and apply cut surface directly onto the wound.	Unspecified	Variable
Heated cooking oil, grease or vaseline	36 (10.5%)	Heat over open fire until melted and then poured directly onto wound	Unspecified	Variable
Iodex ointment	23 (6.7%)	Apply paste directly onto wound	Generous	Variable
Epsom salt soak	9 (2.6%)	One cup of Epsom salts dissolved in hot water	15 minute soak	Twice daily
Methylated spirits	5 (1.5%)	Solution poured onto wound and then covered	Generous	Thrice daily
Hydrogen peroxide	2 (0.6%)	Solution poured onto cotton swabs and then applied directly onto wound	Generous	Variable
Green papaya	2 (0.6%)	Minced green papaya paste applied directly onto wound	Generous	Variable

Discussion

Approximately 15% of the general adult population in Trinidad and Tobago has diabetes mellitus.¹ These persons have 0.75% annual risk to develop lower limb infections.¹ When they do, they have a high age-standardized amputation rate, approximately 54 per 100,000 between the ages of 30 and 60 years.¹²

Although much research has been done on diabetic foot complications in the Caribbean,^{1–13} these data are incomplete without the understanding of the existing cultural practices in persons with diabetes, who are prone to developing lower limb infections. This paper demonstrates that up to 41% of persons initially opt for self-directed treatments and delay seeking medical opinions for a mean duration of 8.8 \pm 5.4 days. This is a disappointing result because delayed medical intervention has been previously established as an independent predictor of amputation risks.^{15,16}

Hennis et al¹³ studied a population of Afro-Caribbean patients with diabetic foot infections in Barbados where the amputation rates were among the highest in the world.^{17,18} They concluded that the high amputation risk was due to the use of inadequate footwear, poorly controlled diabetes, undiagnosed peripheral neuropathy, and untreated

peripheral arterial occlusive disease.¹⁵ Teelucksingh et al⁹ also identified the use of traditional footwear as a contributor to poor therapeutic outcomes in a Trinidadian population. However, we believe that the practice of self-directed treatment may also be a significant contributor to the poor outcomes. There has been little focus on this practice in the Caribbean, but our anecdotal experience is that the patients' strong cultural belief in self-directed treatments often supersedes their perceptions of conventional medical treatments.

These outcomes were particularly alarming considering that 74% of the patients were previously hospitalized for diabetic foot complications. We expected that they would be acutely aware of the consequences of this disease, especially since 13% of them were amputees and 42% had operative debridement to avert the threat of limb loss. Further, patients had limited awareness of the disease despite the fact they were able to recall being counseled about the importance of glycemic control and foot care at some point before disease presentation. Most patients were counseled at dedicated diabetes clinics that are scattered throughout the communities in Trinidad and Tobago. Additionally, 69% of persons were poorly controlled in the 12-week period preceding

admission as evidenced by HbA_{1c} levels greater than 7%. This shows that despite all the work dedicated to diabetic foot complications,^{1–18} we have still not properly addressed secondary prevention to this captive audience.

Finally, a disturbing trend uncovered was the use of unknown “pills” by 6.4% of patients, half of which were readily available from unregulated “informal vendors”. This deserves closer attention by the national regulatory bodies.

Study limitations

There were several limitations to this study. Firstly, this was a cohort study so we were not able to compare outcomes with conventional managements to determine whether this method of treatment is beneficial or not. This may be the focus for further work, possibly with a more robust blinded-randomized methodology.

Secondly, an investigator interviewed diabetic patients who developed lower limb infections to collect data for this study, relying wholly on the information supplied by the patients. We acknowledge that this may have introduced some degree of bias because the patients may not have divulged accurate information for the fear that this may alter the care they receive.

Finally, it is evident that these patients made a conscious effort to choose self-directed treatment over conventional medical advice and treatment. Therefore, they may have believed that the medical personnel do not understand these forms of traditional treatments. Hence, they may have not been cooperative with the process of data collection.

Conclusion

Caribbean cultural practices may be an important contributor to poor outcomes from diabetic foot complications. This is an area of research that deserves more attention in order to provide a tailored approach in care delivery.

Despite being acutely aware of the potentially devastating consequences of inadequate treatment, 41% of our patients still opted to use self-directed treatments for up to 8.8 days.

We must reinforce public health education strategies so that patients promptly seek medical care. Policy makers should also focus on removing any barriers including cultural resistance to accessing health care.

Disclosure

There are no existing financial interests or other commitments that may represent potential conflicts of interest for any of the authors. No funding was received for this work.

References

1. Cawich SO, Islam S, Hariharan S, et al. The Economic Impact of Hospitalization for Diabetic Foot infections in a Caribbean Nation. *Perm J*. 2014;18(1):101–104.
2. Samuels TA, Hospedales CJ. From Port-of-Spain Summit to United Nations High Level Meeting CARICOM and the Global Non-communicable Disease Agenda. *West Ind Med J*. 2011;60(4):387–391.
3. Boyne MS. Diabetes in the Caribbean: Trouble in paradise. *Insulin*. 2009;4(2):94–105.
4. Ferguson RS, Tulloch-Reid MK, Wilks RJ. Epidemiology of Diabetes Mellitus in Jamaica and the Caribbean: A Historical Review. *West Ind Med J*. 2010;59(3):259–264.
5. Hennis A, Wu SY, Nemesure B, Li X, Leske MC. Diabetes in a Caribbean Population: Epidemiological Profile and Implications. *Int J Epidemiol*. 2002;31:234–239.
6. Islam S, Harnarayan P, Cawich SO, et al. Epidemiology of Diabetic Foot Infections in an Eastern Caribbean population: A Prospective Study. *Perm J*. 2013;17(2):37–40.
7. Solomon S, Affan AM, Gopie P, et al. Taking the next step in 2005, the year of the diabetic foot. *Primary Care Diabet*. 2008;2:175–180.
8. Islam S, Harnarayan P, Cawich SO, et al. Secondary Prevention of Diabetic Foot Infections in a Caribbean Nation: A Call for Improved Patient Education. *Int J Low Extrem Wounds*. 2013;12(3):232–236.
9. Teelucksingh S, Ramdass MJ, Charan A, Mungalsingh C, Seemungal T, Naraynsingh V. The Slipping Slipper Sign: A Marker of Severe Peripheral Diabetic Neuropathy and Foot Sepsis. *Postgrad Med*. 2009;85:288–291.
10. Teelucksingh S, Naraynsingh V. Injury to diabetic feet by thumb tacks. *Lancet*. 1997;350:7.
11. Islam S, Cawich SO, Budhooram S, et al. Microbial profile of diabetic foot infections in Trinidad and Tobago. *Prim Care Diabetes*. 2013;7(4):303–308.
12. Gulliford MC, Ariyanayagam-Baksh SM, Bickram L, Picou D, Mahabir D. Counting the cost of diabetic hospital admissions from a multi-ethnic population in Trinidad. *Diabet Med*. 1995;12(12):1077–1085.
13. Hennis AJ, Frazer HS, Jonnalagadda R, Fuler J, Chaturvedi N. Explanations for the high risk of diabetes related amputation in a Caribbean population of black African descent and potential for prevention. *Diabet Care*. 2004;27(11):2636–2641.
14. Valk GD, Kriegsman DM, Assendelft WJ. Patient education for preventing diabetic foot ulceration: a systematic review. *Endocrinol Metab Clin North Am*. 2002;31:633–658.
15. Hennis A, Wu SY, Nemesure B, Li X, Leske MC. Diabetes in a Caribbean population: epidemiological profile and implications. *Int J Epidemiol*. 2002;31:234–239.
16. Lipsky BA, Berendt AR, Cornia PB, et al. Infectious Diseases Society of America. 2012 Infectious Diseases Society of America Clinical Practice Guidelines for the Diagnosis and Treatment of Diabetic Foot Infections. *Clin Infect Dis*. 2012;54(12):132–173.
17. Walrond ER. The Caribbean experience with the diabetic foot management of the diabetic foot. *West Ind Med J*. 2001;50(S7):24–26.
18. Hambleton IR, Jonnalagadda R, Davis CR, Fraser HS, Chaturvedi N, Hennis AJ. All-cause mortality after diabetes-related amputation in Barbados: a prospective case-control study. *Diabetes Care*. 2009;32(2):306–307.

Patient Preference and Adherence**Dovepress****Publish your work in this journal**

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focuses on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize

clinical outcomes for existing disease states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/patient-preference-and-adherence-journal>