#### **Open Access Full Text Article**

#### ORIGINAL RESEARCH

Application of carbon nanoparticles in lymph node dissection and parathyroid protection during thyroid cancer surgeries: a systematic review and meta-analysis

Lun Wang Dong Yang Jun-Yuan Lv Dan Yu Shi-Jie Xin

Department of Vascular and Thyroid Surgery, The First Affiliated Hospital, China Medical University, Shenyang, People's Republic of China

Correspondence: Shi-Jie Xin Department of Vascular and Thyroid Surgery, The First Affiliated Hospital, China Medical University, No 155 of Nanjing Street, Shenyang 110001, People's Republic of China Tel +86 24 8328 3288 Fax +86 24 2270 3576 Email sjxin@cmu.edu.cn



**Purpose:** To investigate whether carbon nanoparticles (CNs) are helpful in identifying lymph nodes and metastatic lymph nodes and in parathyroid protection during thyroid cancer surgery. **Methods:** English and Chinese literature in PubMed, Cochrane Database of Systematic Reviews, EMBASE, ClinicalTrials.gov, China Biology Medicine Database, China National Knowledge Infrastructure, China Master's and Doctoral Theses Full-Text Database, Wanfang database, and Cqvip database were searched (till March 22, 2016). Randomized controlled trials (RCTs) that compared the use of CNs with a blank control in patients undergoing thyroid cancer surgery were included. Quality assessment and data extraction were performed, and a meta-analysis was conducted using RevMan 5.1 software. The primary outcomes were the number of retrieved central lymph nodes and metastatic lymph nodes, and the rate of accidental parathyroid removal.

**Results:** We obtained 149 relevant studies, and only 47 RCTs with 4,605 patients (CN group: n=2,197; blank control group: n=2,408) met the inclusion criteria. Compared with the control group, the CN group was associated with more retrieved lymph nodes/patient (weighted mean difference [WMD]: 3.39, 95% confidence interval [CI]: 2.73–4.05), more retrieved metastatic lymph nodes (WMD: 0.98, 95% CI: 0.61–1.35), lower rate of accidental parathyroid removal, and lower rates of hypoparathyroidism and hypocalcemia. However, the total metastatic rate of the retrieved lymph nodes did not differ between the groups (odds ratio: 1.13, 95% CI: 0.87–1.47, P=0.35).

**Conclusion:** CNs can improve the extent of neck dissection and protect the parathyroid glands during thyroid cancer surgery. And the number of identified metastatic lymph nodes can be simultaneously increased.

Keywords: carbon nanoparticles, lymph node tracer, thyroid cancer, parathyroid, meta-analysis

### Introduction

Thyroid cancer is a common malignant tumor, and its occurrence has been increasing appreciably over the last few decades.<sup>1,2</sup> Papillary thyroid carcinoma, the most common pathological type of thyroid carcinoma, is associated with an excellent prognosis if surgery is performed successfully.<sup>3,4</sup> However, complications and tumor metastasis often occur postoperatively. A multicenter revisit study of 25,634 patients with a history of ambulatory surgery revealed that the incidence of hypocalcemia was as high as 20.8%.<sup>5</sup> Moreover, the rate of postoperative cervical lymph node metastasis has been reported to be as high as 20%–90%.<sup>6,7</sup> Therefore, an effective method is urgently

OncoTargets and Therapy 2017:10 1247-1260

Control of the second sec

required to help identify and remove additional lymph nodes and decrease the risk of parathyroid injury.

Nanobiotechnology, a new field defined as biomedical applications of nanosized systems, which involves nanostructure and nanomaterials, has emerged as a key player among various disciplines of biomedical science. Nanomaterials, which measure 1–1,000 nm, have unique physical and chemical properties such as small-size effect, large surface area, high reactivity, and quantum effects. In addition, they have been certified as breaking a new ground in disease detection,<sup>8,9</sup> imaging,<sup>10,11</sup> diagnosis,<sup>12,13</sup> and treatment.<sup>12,14–16</sup> Carbon-based nanoparticles are an important part of nanomaterials; they include carbon nanotubes, fullerene, and graphene and its derivatives.<sup>17</sup> Due to their unique physical and chemical properties, carbon-based nanoparticles have broad applications in the biomedical field. Above all, structure or surface modifications of carbon-based nanoparticles result in different effects.

In recent years, with the development of nanotechnology, nanomaterials or nanosized products have been used in surgeries. Carbon nanoparticles (CNs) suspension (China Food and Drug Administration approval H20041829; Lai Mei Pharmaceutical Co, Chongqing, People's Republic of China), which comprises nanosized polymeric carbon granules with an average diameter of 150 nm (Figure 1),<sup>18</sup> ensures that these CNs pass through the lymphatic vessels (diameter: 120–500 nm) rather than blood capillaries (diameter: 20–50 nm) due to their molecular size. Hence, the lymph nodes and thyroid glands can be stained with CNs but not the parathyroid glands. Therefore, as revealed in Figure 2,<sup>19</sup> CNs have been used as lymph node tracers during thyroid



**Figure I** Transmission electron microscopic image of CNs. **Notes:** The main active component of CNs is nanosized polymeric carbon granules with an average diameter of 150 nm. Republished with permission of SAGE Publications, Inc., from Liu X, Chang S, Jiang X, Huang P, Yuan Z. Identifying parathyroid glands with carbon nanoparticle suspension does not help protect parathyroid function in thyroid surgery: a prospective, randomized control clinical study. *Surg Innov.* 2016;23(4):381–389. © The Author(s) 2016; permission conveyed through Copyright Clearance Center, Inc.<sup>18</sup> **Abbreviation:** CN, carbon nanoparticle.



Figure 2 Intraoperative view of the black-stained clusters of lymph nodes by CN injection.

**Notes:** Reproduced with permission of John Wiley and Sons, from Zhu Y, Chen X, Zhang H, et al. Carbon nanoparticle-guided central lymph node dissection in clinically node-negative patients with papillary thyroid carcinoma. *Head Neck*. 2016;38(6):840–845. © 2015 Wiley Periodicals, Inc.<sup>19</sup> **Abbreviation:** CN, carbon nanoparticle.

surgeries in the People's Republic of China in recent years, but there is no report about their usage in other countries. Previous studies have demonstrated that CNs could help visualize the lymph nodes and preserve the parathyroid. In addition, a meta-analysis has also been published to support this viewpoint.<sup>20</sup> However, controversy still exists about the usefulness of CNs. Liu et al reported that CNs are not beneficial for parathyroid protection during thyroid cancer surgery and that the usage of CNs results in a significantly prolonged operation time.<sup>18</sup> Before 2014, all articles published on the usage of CNs in thyroid cancer surgeries were in Chinese, and there were only limited randomized controlled trials (RCTs) reported. Accordingly, a considerable amount of non-RCTs were included in the previous meta-analysis.20 However, in recent years, more RCTs have been published. Hence, another meta-analysis is needed to determine whether CNs are helpful in thyroidectomy.

With this in mind, the present systematic review and meta-analysis was designed to confirm whether CNs are indeed helpful in thyroid cancer surgery, that is, whether CNs can really improve the extent of thyroidectomy and neck dissection and help identify metastatic lymph nodes while preserving the parathyroid glands, as compared with the performance of blank controls.

## **Methods** Search strategy

The following English and Chinese databases were searched systematically by 2 investigators independently (till March 22, 2016): PubMed, Cochrane Database of Systematic Reviews,

EMBASE, <u>ClinicalTrials.gov</u>, China Biology Medicine Database, China National Knowledge Infrastructure, China Master's and Doctoral Theses Full-Text Database, WANFANG database, and Cqvip database. RCTs on initial thyroid cancer surgeries that compared the use of CNs with a blank control were included. Our search terms included (nano-carbon) or (carbon particle) or (carbon nanoparticle) or (carbon nanoparticles) or (lymph node tracer) or (lymphatic tracer) or (lymphography) and (thyroid or thyroidea). To resolve any disagreement between the 2 investigators, a third reviewer was invited to assess any discrepant items.

## Inclusion criteria

The studies selected were RCTs on thyroid cancer surgeries that included: 1) patients who underwent initial surgery and with a confirmed pathology diagnosis and 2) a control group not injected with anything before thyroidectomy and a CN group injected with CNs.

## Exclusion criteria

Non-independent clinical controlled trials, non-RCTs, studies with a patient number < 10, or studies with incomplete data were excluded.

### Observation indexes

The primary outcomes were the number of retrieved central lymph nodes and metastatic lymph nodes per patient, and the rate of accidental parathyroid removal. Other outcomes extracted from the identified RCTs included the staining rate of lymph nodes, the number of metastatic lymph nodes in all retrieved lymph nodes, and the rate of postoperative transient or permanent hypoparathyroidism and hypocalcemia. All included articles reported at least 1 of the outcomes.

## Quality assessment

To evaluate the quality of these studies, the Jadad scoring system was applied in the RCTs. The scoring system included 3 items: descriptions of the dropouts and withdrawals (0 or 1 point), blinding (0–2 points), and randomization (0–2 points). The maximum score was 5 points. RCTs that scored 3–5 points were considered to be of high quality, whereas a score of 0-2 was considered to indicate low quality.

## Statistical analysis

RevMan version 5.1 software was used for the statistical analyses. We measured the heterogeneity of the studies using the  $I^2$  and  $\chi^2$  tests. Statistical heterogeneity of the studies was defined as an  $I^2$  value <50% or *P*-value <0.10. The random-effects model was applied if heterogeneity existed among

the studies; otherwise, the fixed-effects model was adopted for the analyses. Weighted mean differences (WMDs) with 95% confidence intervals (CIs) were used for the continuous outcome variables, whereas risk differences (RDs) and odds ratios (ORs) with 95% CIs were calculated for the dichotomous outcome variables. To investigate possible bias, funnel plots were created. For all analyses, statistical differences were considered to exist between the 2 groups when *P*-value was <0.05.

# **Results and discussion**

Considering the complicated anatomic structure and lymphatic drainage, the cervical lymph node metastasis rate is amazingly high after thyroid cancer surgery.<sup>6,7</sup> Especially, in the central region of the neck and other places such as the central neck compartment and deep surface of the recurrent laryngeal nerve, which cannot be easily dissected, postoperative cervical lymph node metastasis is common. As a result, reoperation and surgical trauma are common in these patients. According to a previous report,<sup>21</sup> while the incidence of permanent hypoparathyroidism is 3%-10% after the first surgery for thyroid disease, it is as high as 9%-35% after reoperation. Kurmann et al reported that the incidence of permanent recurrent laryngeal nerve palsy was significantly higher in patients undergoing reoperation on the ipsilateral lobe compared to patients undergoing initial operation (3.8% vs 1.1%; P=0.03).<sup>22</sup> In addition, parathyroid injury has been considered inevitable for a long time, mostly due to its unique anatomy. The appearance of the parathyroid is similar to that of the cervical lymph nodes, and the location of the gland is close to the backside of the thyroid gland and varies greatly; for example, the parathyroid may hide within the thyroid lobes, thymus, or carotid sheath. Xu and Gu reported that 6.9%-46% of parathyroid glands were damaged during thyroid surgery.<sup>23</sup> Such damage may cause permanent or transient hypocalcemia and hypoparathyroidism, and will consequently affect the quality of life of the patients. Therefore, some technical methods are urgently needed to help us better visualize the lymph nodes and metastatic lymph nodes, and distinguish the parathyroid glands and preserve them.

CNs, which have a mean diameter of 150 nm and a lymphatic tendency, had been used as lymph node tracers clinically in other cancer surgeries before. Upon injection, CNs are rapidly devoured by macrophages, resulting in the lymph nodes, but not the capillaries, initially becoming black-stained. Subsequently, the thyroid tissue stains black, as do the surrounding lymph nodes, whereas any tissues without lymph vessel connections remain unstained. Some previous studies have demonstrated that CNs are beneficial for visualizing the lymph nodes and for distinguishing and preserving the parathyroid glands. On the contrary, other studies found no advantage of CNs.<sup>18,24</sup> In addition, some indexes, such as postoperative hypocalcemia, are easily influenced by confounding factors such as the postoperative therapeutic selection, calcium supplements, and individual differences. To date, no large-scale meta-analysis has been performed to clarify these diverging results. Therefore, our analysis was designed to resolve the problems.

### Literature search and study description

According to the search strategy, 149 references were obtained, and 47 RCTs met the inclusion criteria.<sup>18,19,23–67</sup> The flowchart of the literature search is shown in Figure 3. A total of 4,605 patients were included in the analysis, including 2,197 patients in the CN group and 2,408 patients in the blank control group. All patients had confirmed thyroid cancer by postoperative pathologic diagnosis and had been divided into the CN and blank control groups randomly. All of the CNs used in these studies were the same product.

The characteristics of all 47 RCTs are presented in Table 1. The Jadad scale system was used to assess the

quality of the included studies (Table 2). Most investigators preferred multipoint injections (2–4 points) prior to the thyroidectomy, with a total dose of approximately 0.3–0.8 mL. Besides, according to our experience and the studies analyzed herein, no adverse reactions to CNs have been reported.

### Intervention effects

Number of retrieved lymph nodes and metastatic lymph nodes, and metastatic rate of retrieved lymph nodes in the CN and blank control groups

Compared with the blank control groups, the use of CNs resulted in an increased number of retrieved lymph nodes, approximately 3.39 per patient (WMD=3.39, 95% CI=2.73–4.05, P<0.00001; Figure 4A). The number of retrieved metastatic lymph nodes per patient in the CN group was significantly higher than in the blank control group (WMD=0.98, 95% CI=0.61–1.35, P<0.00001; Figure 4B). However, interestingly, the total metastatic rate of the lymph nodes, metastatic rate of the stained lymph nodes, and metastatic rate of the unstained lymph nodes were not significantly different between the CN and blank control groups



Figure 3 Flowchart of the literature search.

Table I Characteristic	s of the	47 RCT	s included	I in the meta-ana	ılysis					
Study	c	Male/fer	nale	Age, years, mé	san (SD)	Injection site	Dose	Waiting	Staining	Indices <sup>a</sup>
		CNs	BL	CNs	BL		(mL)	time	rate (%)	
Chen et al (2016) <sup>25</sup>	173	15/72	13/73	33.46 (4.58)	34.32 (4.36)	Upper, middle, and lower points of the thyroid	0.3	5-10 min	AN	l, 4–6
Zhang et al (2016) <sup>26</sup>	37	3/14	3/17	41.1 (8.7)	43 (7.2)	In the bilateral thyroid	0.2–0.4	5-20 min	95.5	4
Liu et al (2016) <sup>18</sup>	156	16/62	12/61	AA	NA	Upper, middle, and lower parts of the lobes	0.4-0.8	NA	NA	4, 6
Li (2015) <sup>27</sup>	40	4/16	8/12	47.9 (11.2)	49.5 (12.4)	Top of the tumor	0.4-0.8	3 min	NA	4
Feng and He (2015) <sup>28</sup>	60	0/30	0/30	NA	NA	4 points around the tumor	0.8	AN	AN	4-6
Guo et al (2015) <sup>29</sup>	59	9/21	8/2 I	35 (9.5)	32 (8.2)	Upper, middle, and lower points of the bilateral thyroid	9.6	20 min	AN	6
Duan et al (2015) <sup>30</sup>	80	14/26	15/25	40.4 (7.18)	42.5 (7.65)	3–5 points around the thyroid	0.3-0.5	30 min	95.7	I, 2, 4, 5
Chen and Wu (2015) <sup>31</sup>	96	11/37	16/32	46.1 (10.9)	45.8 (12.6)	NA	NA	NA	AN	I, 5, 6
Du et al (2015) <sup>32</sup>	811	19/41	18/40	42.7 (10.6)	43.5 (10.8)	Upper and middle points of the thyroid	0.4	5 min	AA	4-6
Wang et al (2015) <sup>33</sup>	120	AA	AN	NA	NA	Upper and middle points of the thyroid	0.2	I 5 min	AN	9-1
Liu and Qing (2015) <sup>34</sup>	66	12/21	11/22	46.11 (2.09)	45.63 (2.7)	Upper, middle, and lower points of the thyroid	0.3	20 min	AN	4-6
Shao et al (2015) <sup>35</sup>	60	AA	AA	NA	NA	Multipoint injection of the tumor	0.4–1.2	10 min	AA	I, 5, 6
Fu (2015) <sup>43</sup>	250	AA	AN	NA	NA	4-6 points around the tumor	0.8–1.2	l 5–20 min	AN	5
Wu et al (2015) <sup>36</sup>	245	62/11	33/122	NA	NA	Top and middle points of the tumor	0.2–0.6	5-10 min	NA	4-6
Li et al $(2015)^{37}$	52	AA	AA	NA	NA	4 points of the bilateral thyroid	0.1	10 min	AA	4-6
Chu et al (2015) <sup>38</sup>	57	10/18	8/21	46.28 (12.635)	40.39 (1.711)	Upper, middle, and lower points of the ipsilateral thyroid	0.3-0.6	I 0 min	AN	Ι, 4
Wang et al (2015) <sup>39</sup>	88	10/34	9/35	36.6 (11.2)	36.8 (11.4)	Upper, middle, and lower points of the thyroid	0.3-0.8	NA	AN	_
Wu et al (2015) <sup>40</sup>	86	AA	AN	NA	NA	Upper, middle, and lower points of the thyroid	0.6–1.2	30 min	82.9	3 3
Yin et al (2015) <sup>41</sup>	80	17/23	18/22	AA	NA	Upper, middle, and lower points of the thyroid	0.6	5 min	NA	4-6
Li et al (2015) <sup>42</sup>	72	9/27	10/26	AN	AN	2–4 points around the thyroid	NA	I5 min	84	9-1
Xu and Gu (2016) <sup>23</sup>	4	5/52	4/53	45.37 (10.71)	42.68 (14.43)	Around the tumor	0.5	5–10 min	NA	I, 4–6
Wang et al (2015) <sup>64</sup>	55	1/27	2/25	30.25 (6.04)	29.44 (6.27)	In the thyroid gland	0.1-0.2	2–3 min	85	I, 4–6
Zhu et al (2016) <sup>19</sup>	162	14/67	16/65	46.75 (12.09)	44.31 (10.73)	I–2 points in the thyroid gland	0.1-0.2	Few minutes	92.75	9-1
Gu et al (2015) <sup>65</sup>	001	10/40	6/44	46.98 (9.027)	47.76 (13.912)	Upper and lower points of the thyroid	0.2-0.3	3–5 min	NA	I, 2, <del>4–</del> 6
Liu et al (2014) <sup>44</sup>	47	3/20	5/19	37.79 (11.2)	33.94 (7.76)	Upper, middle, and lower points of the thyroid	9.0	I day	NA	I, 4–6
Chen et al (2014) <sup>45</sup>	72	5/31	8/28	38.23 (10.67)	34.64 (8.75)	Upper, middle, and lower points of the thyroid	0.4/0.6	30 min	95.4	I, 2, 4
Gao and Zhao (2014) <sup>46</sup>	001	12/38	9/41	42 (1.28)	41 (1.36)	3-4 points around the tumor	0.53-1	30 min	95.56	I5
Yang et al (2014) <sup>47</sup>	379	23/155	27/174	42.3 (8.5)	44.2 (7.3)	Upper and lower points of the tumor or the thyroid gland	0.1-0.3	3–5 min	NA	I, 6
Wang and Rang (2014) <sup>48</sup>	70	AN	AA	44.3 (8.8)	45.2 (7.9)	4 points in the contralateral thyroid gland	0.4-0.8	l 5–20 min	NA	I, 2, <del>4–</del> 6
Liu et al (2014) <sup>49</sup>	55	5/21	9/20	36.58 (11.31)	33.58 (7.77)	In the ipsilateral thyroid	14 days	NA	NA	I, 4–6
Du (2014) <sup>50</sup>	40	4/16	8/12	47.9 (10.12)	49.5 (12.4)	Top point of the thyroid gland	0.4-0.8	3 min	NA	I, 4–6
Zhao (2014) <sup>51</sup>	183	23/79	19/62	NA	NA	Upper and lower points of the thyroid	0.1-0.2	5 min	73.5	Ι, 2
Chun (2014) <sup>52</sup>	67	6/27	11/23	41.06 (12.84)	44.65 (12.84)	Upper, middle, and lower points of the thyroid	0.1-0.3	3–5 min	86.8	Ι, 5
Liu (2014) <sup>53</sup>	184	23/53	30/78	46.5 (12.8)	47.2 (13.5)	Multipoint injection of the tumor	0.8	NA	AA	5,6
Zhang et al (2014) <sup>54</sup>	72	12/24	10/26	AA	NA	Upper and lower points of the thyroid	0.2–0.4	5 min	NA	4-6
Shen et al (2014) <sup>55</sup>	109	AN	AA	AA	NA	Around the tumor	0.2	10 min	90.5	I, 2, 4
Shao et al (2014) <sup>57</sup>	29	AA	AA	NA	NA	Upper, middle, and lower points of the thyroid	0.3-0.6	I5 min	AN	4-6
Long et al (2014) <sup>56</sup>	150	15/60	12/63	41.4 (1.62)	42.I (2.56)	Upper, middle, and lower points of the ipsilateral thyroid	0.3	20 min	NA	I, 2, 6
Tian et al (2014) <sup>66</sup>	001	5/45	11/39	36.4 (2.5)	44.5 (5.8)	Upper, middle, and lower points of the thyroid	9.0	10–15 min	NA	I, 2, <del>4–</del> 6
									•	Continued)

auuy	c	Male/f∈	emale	Age, years, m	ean (SD)	Injection site	Dose	Waiting	Staining	Indices <sup>a</sup>
		CNs	BL	CNs	BL		(mL)	time	rate (%)	
Sun et al (2014) <sup>67</sup>	80	AN	NA	AN	AN	2-4 points around the tumor	0.2-0.8	15 min	69.89	4
Yang et al (2013) <sup>58</sup>	43	1/20	1/21	32.48 (4.69)	32.32 (5.35)	In the ipsilateral thyroid gland	0.1	20 min	AN	I, 2, 4, 5
Yang et al (2013) <sup>59</sup>	68	17/19	12/20	34.5 (9.1)	33.9 (10.3)	I-2 points around the tumor	0.1-0.3	I0 min	7.16	Ι, 5
Wu (2013) <sup>60</sup>	55	5/21	9/20	36.58 (11.31)	33.58 (7.77)	Upper, middle, and lower points of the ipsilateral thyroid	0.6	AN	AN	1, 2, 4–6
Huang et al (2013) <sup>24</sup>	72	12/24	10/26	41.22 (2.53)	40.69 (2.42)	Lower and upper points of the tumor	0.2–0.4	AN	NA	5
Bai et al (2013) <sup>61</sup>	88	9/39	7/33	46.28 (9.2)	45.39 (12.03)	Lower and upper points of the tumor	0.2	AN	AN	<u>9</u>
Zeng et al (2012) <sup>62</sup>	80	0/40	0/40	NA	NA	4–6 points around the tumor	I.0	20 min	AN	5, 6
Wang et al (2009) <sup>63</sup>	36	10/8	7/11	NA	NA	4-6 points around the tumor	0.4-0.6	30 min	95.2	24

**Dove**press

(OR =1.13, 95% CI =0.87-1.47, P=0.35; OR =1.33, 95% CI =0.91-1.94, P=0.14; and OR =0.55, 95% CI =0.23-1.36, P=0.20, respectively; Figure 5A–C).

In the present meta-analysis of these studies, we found that the number of retrieved lymph nodes per patient in the CN group was higher than that of the blank control group. This finding of an increasing number of retrieved lymph nodes corresponds to the improvement in the extent of neck dissection. Further, with increasing removal of dyed lymph nodes, metastatic lymph nodes will also be cleared away simultaneously. Hence, the number of metastatic lymph nodes in the CN group was statistically higher than that in the blank control group. Besides, 4 studies reported that the number of retrieved small lymph nodes (diameter <5 mm) was significantly higher in the CN group.44,60,61,67 These findings suggest that CNs may help identify tiny, suspicious lymph nodes.

However, it should be noted that the total metastatic rate of the retrieved lymph nodes and the metastatic rate of stained or unstained lymph nodes did not significantly differ between the 2 groups, consistent with the findings of previous studies.46,68 In Gao and Zhao study,46 more metastatic lymph nodes were eliminated in the CN group (6±2.37 vs 4 $\pm$ 2.49; P<0.01), but the rate of metastatic lymph nodes did not differ (45.97% vs 47.10%; P>0.05). In the study by Yan et al,<sup>68</sup> no increase in the number of sentinel lymph node metastasis-positive cases was observed with the utilization of CNs, as compared to that in the control group (36.8% vs 63.2%). Concerning the mechanism of CNs, it is considered that the tissue damage and inflammation caused by the tumor alter the lymphatic drainage channels of the thyroid, which in turn will affect the diffusion of CNs and the identification of metastatic lymph nodes. Thus, it is actually quite hard for CNs to distinguish metastatic lymph nodes among normal lymph nodes.

### Anatomic structure and physical function of the parathyroid in the CN and blank control groups

Compared with the blank control group, the use of CNs was associated with a lower rate of accidental parathyroid removal, approximately 22% (OR =0.22, 95% CI =0.16-0.30, P<0.00001; RD =-0.13, 95% CI =-0.15 to -0.11, P < 0.00001; Figure 6). The transient and permanent hypoparathyroidism rates declined by approximately 31% and 24% in the CN group, respectively (transient hypoparathyroidism rate: OR =0.31, 95% CI =0.25–0.39, P<0.00001; RD =-0.15, 95% CI =-0.18 to -0.12, P < 0.00001; Figure 7A; permanent hypoparathyroidism rate: OR =0.24,

**Dove**press

#### Table 2 Quality assessment of the 47 randomized controlled trials included using the Jadad scale system

	-
follow-up (%) a	ssessment
Chen et al (2016) <sup>25</sup> No detailed description Only mentioned randomized Unclear 0 2	)
Zhang et al (2016) <sup>26</sup> No detailed description Only mentioned randomized Unclear 0 2	
Liu et al (2016) <sup>18</sup> Computer-generated permuted Only mentioned randomized Unclear 0 3	1
Li (2015) <sup>27</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Feng and He (2015) <sup>28</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Guo et al (2015) <sup>29</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Duan et al (2015) <sup>30</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Chen and Wu (2015) <sup>31</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Du et al (2015) <sup>32</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Wang et al (2015) <sup>33</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Liu and Oing (2015) <sup>34</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Shao et al (2015) <sup>35</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Fu (2015) <sup>43</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Wu et al (2015) <sup>36</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Li et al (2015) <sup>37</sup> Random-number table Only mentioned randomized Unclear 0 3	}
Chu et al (2015) <sup>38</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Wang et al (2015) <sup>39</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Wu et al (2015) <sup>40</sup> Random-number table Only mentioned randomized Unclear 0 3	}
Yin et al (2015) <sup>41</sup> Random-number table Only mentioned randomized Unclear 0 3	}
Li et al (2015) <sup>42</sup> Random-number table Only mentioned randomized Unclear 0 3	}
Xu and Gu (2016) <sup>23</sup> No detailed description Only mentioned randomized Unclear 0 2	1
Wang et al (2015) <sup>64</sup> Computer-generated Only mentioned randomized Unclear 0 3	-
Thu et al (2016) <sup>19</sup> Computer-generated Only mentioned randomized Unclear 0 3	
random-number tables	
Gu et al (2015) <sup>65</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Liu et al (2014) <sup>44</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Chen et al (2014) <sup>45</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Gao and Zhao (2014) <sup>46</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Yang et al (2014) <sup>47</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Wang and Rang (2014) <sup>48</sup> Random-number table Only mentioned randomized Unclear 0 3	;
Liu et al (2014) <sup>49</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Du (2014) <sup>50</sup> No detailed description Only mentioned randomized Unclear 0 2	1
Zhao (2014) <sup>51</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Chun (2014) <sup>52</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Liu (2014) <sup>53</sup> No detailed description Only mentioned randomized Unclear 0 2	1
Zhang et al (2014) <sup>54</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Shen et al (2014) <sup>55</sup> No detailed description Only mentioned randomized Unclear 0 2	<u>.</u>
Shao et al (2014) <sup>57</sup> No detailed description Only mentioned randomized Unclear 0 2	1
Long et al (2014) <sup>56</sup> Odd or even number Only mentioned randomized Unclear 0 I	
Tian et al (2014) <sup>66</sup> Randomization chart Only mentioned randomized Unclear 0 3	1
Sun et al (2014) <sup>67</sup> Computer-generated permuted Only mentioned randomized Unclear 0 3	}
block sequencing	
Yang et al (2013) <sup>58</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Yang et al (2013) <sup>59</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Wu (2013) <sup>60</sup> No detailed description Only mentioned randomized Unclear 15.4 3	;
Huang et al (2013) <sup>24</sup> Computer-generated permuted Sealed envelopes Single 0 3	}
blinding	
Bai et al (2013) <sup>61</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Zeng et al (2012) <sup>62</sup> No detailed description Only mentioned randomized Unclear 0 2	2
Wang et al (2009) <sup>63</sup> Random-number table Only mentioned randomized Unclear 0 3	}

Notes: The Jadad scale was used to assess the quality of these RCTs. Thirteen studies included had a score of 3 points, which reflected the high quality of the study. The majority of studies had 2 points or lower. These studies can be considered to be of relatively low quality. **Abbreviation:** RCT, randomized controlled trials.

95% CI =0.07–0.85, P=0.03; RD =–0.02, 95% CI =–0.03 to –0.00, P<0.02; Figure 7B). In addition, the rate of post-operative transient hypocalcemia in the blank control group was 30% higher than in the CN group (OR =0.30, 95%

CI =0.25–0.38, P<0.00001; RD =–0.16, 95% CI =–0.18 to –0.13, P<0.00001; Figure 8A). On the other hand, the postoperative permanent hypocalcemia rate did not significantly differ between the CN and blank control groups

Δ

Study or	Expe	riment	al	Cont	rol		Weight	Mean difference IV.	Year		Mean	differen	ce IV.		
subgroup	Mean	SD	Total	Mean	SD	Total	(%)	random, 95% Cl			rando	m, 95%	CI		
Bai et al (2013)61	7.6	4.1	48	3.7	4.9	33	3.9	3.90 (1.87, 5.93)	2013			-			
Wu (2013) <sup>60</sup>	9.96	4.53	26	5.62	2.58	29	4.0	4.34 (2.36, 6.32)	2013						
Yang et al (2013)59	8.99	2.24	36	6.41	1.56	32	5.5	2.58 (1.67, 3.49)	2014			-			
Yang et al (2013)58	9.67	5.26	21	4.95	2.54	22	3.3	4.72 (2.23, 7.21)	2013			-		-	
Chen et al (2014)45	8.67	3.15	36	5.61	3.36	36	4.7	3.06 (1.56, 4.56)	2014						
Wang and Rang (2014)48	11.4	1.88	35	7.6	1.59	35	5.6	3.80 (2.98, 4.62)	2014						
Zhao (2014)51	10.9	4.3	102	8.1	3.7	81	5.2	2.80 (1.64, 3.96)	2014						
Chun (2014)52	5.97	3.92	33	4.44	0.2	34	4.7	1.53 (0.03, 3.03)	2014			-	_		
Gao and Zhao (2014)46	9.45	3.16	50	5.75	2.75	50	5.2	3.70 (2.54, 4.86)	2014						
Shen et al (2014)55	9.7	1.7	45	6.7	1.9	64	5.8	3.00 (2.32, 3.68)	2014						
Liu et al (2014)44	10.43	3.64	23	5.54	1.79	24	4.4	4.89 (3.24, 6.54)	2014						
Liu et al (2014)49	9.96	4.53	26	5.62	2.58	29	4.0	4.34 (2.36, 6.32)	2014			· ·	•		
Du (2014) <sup>50</sup>	7.6	2.52	20	4.7	2.52	20	4.6	2.90 (1.34, 4.46)	2014			_	+		
Zhu et al (2016) <sup>19</sup>	10.07	6.46	81	4.04	2.84	81	4.6	6.03 (4.49, 7.57)	2016					_	
Duan et al (2015)30	10.86	7.39	40	6.88	2.37	40	3.4	3.98 (1.57, 6.39)	2015						
Li et al (2015)42	8.7	1.8	36	5.3	1.4	36	5.7	3.40 (2.66, 4.14)	2015						
Chu et al (2015)38	7.89	0.685	28	3.38	2.111	29	5.6	4.51 (3.70, 5.32)	2015						
Wu et al (2015)40	10.19	4.27	43	6.26	2.98	43	4.6	3.93 (2.37, 5.49)	2015						
Gu et al (2015)65	5.78	4.55	50	6.62	5.07	50	4.1	-0.84 (-2.73, 1.05)	2015		-				
Chen et al (2016) <sup>25</sup>	9.71	4.32	87	5.54	2.54	86	5.3	4.17 (3.12, 5.22)	2016						
Xu and Gu (2016) <sup>23</sup>	0.6	0.98	57	4.58	0.6	57	6.1	1.42 (1.12, 1.72)	2016			-			
Total (95% CI)			923			911	100	3.39 (2.73, 4.05)					•		
Heterogeneity: r <sup>2</sup> =1.86; ;	χ²=163	.75, df	=20 (F	<b>~</b> 0.00	001); <i>I</i>	<sup>2</sup> =88%	0			10				10	_
Test for overall effect: Z=	10.04	( <i>P</i> <0.0	0001)							-10	-5	0	5 _	10	
										Fav (blank)	vors control)	(0	F arbon n	avors anoparticle	s)

В										
Study or subgroup	Expe Mean	riment SD	al Total	Conti Mean	rol SD	Total	Weight (%)	t Mean difference IV, random, 95% CI	Year	Mean difference IV, random, 95% Cl
Yang et al (2013)58	1.23	1.57	21	0.55	0.79	22	10.0	0.68 (-0.07, 1.43)	2013	
Liu et al (2014)44	8.92	3.38	23	4.47	1.91	24	4.1	4.45 (2.87, 6.03)	2014	
Gao and Zhao (2014)46	0.6	2.37	50	0.4	2.49	50	8.0	2.00 (1.05, 2.95)	2014	
Long et al (2014)56	2.86	0.13	75	1.87	0.09	75	16.9	0.99 (0.95, 1.03)	2014	•
Duan et al (2015)30	0.5	4.15	40	2.64	1.52	40	5.1	2.36 (0.99, 3.73)	2015	
Zhu et al (2016)19	2.74	3.66	81	1.68	2.4	81	8.0	1.06 (0.11, 2.01)	2016	
Chu et al (2015)38	2.21	3.348	28	0.86	1.302	29	5.3	1.35 (0.02, 2.68)	2015	
Gu et al (2015)65	1.66	2.353	50	3.06	4.501	50	4.9	-1.40 (-2.81, 0.01)	2015	
Wu et al (2015)40	3.88	0.49	43	3.81	3.43	43	7.3	0.07 (–0.97, 1.11)	2015	
Chen et al (2016)25	1.3	1.59	87	0.57	0.74	86	14.5	0.73 (0.36, 1.10)	2016	
Xu and Gu (2016) <sup>23</sup>	0.96	0.77	57	0.47	0.5	57	15.8	0.49 (0.25, 0.73)	2016	+
Total (95% CI)			555			557	100	0.98 (0.61, 1.35)		•
Heterogeneity: $\tau^2=0.21$ ;	$\chi^2 = 60.0$	04, <i>df=</i>	10 ( <i>P</i> <	<0.000	01); /²=	-83%			_	
Test for overall effect: Z:	=5.18 (/	P<0.00	001)							-4 -2 0 2 4
			,							Favors Favors (blank control) (carbon nanoparticles)

Figure 4 Forest plot showing the association of CNs with retrieved lymph nodes and metastatic lymph nodes. Notes: (A) Number of retrieved lymph nodes per patient in the CN and blank control groups. (B) Number of retrieved metastatic lymph nodes per patient. Abbreviations: CN, carbon nanoparticle; SD, standard deviation; IV, interval variable; CI, confidence interval; random, random effect; df, degrees of freedom.

(OR =0.33, 95% CI =0.04–3.03, *P*=0.33; RD =–0.01, 95% CI =–0.03 to –0.01, *P*=0.35; Figure 8B).

In terms of the protection of the parathyroid, some previous studies showed that the usage of CNs was not beneficial;<sup>18,24</sup> however, our study demonstrated that the rates of accidental parathyroid removal (Figure 6), postoperative transient or permanent hypoparathyroidism (Figure 7), and transient hypocalcemia (Figure 8A) were lower in the CN group. This finding suggests that the usage of CNs will help distinguish and preserve the parathyroid glands. However, there was no significant difference in the rate of permanent hypocalcemia. This might have resulted from the quantitative

restrictions of the RCTs included. In addition, Yang et al reported that the use of CNs resulted in a decreased rate of parathyroid auto-transplantation.<sup>47</sup>

### Publication bias and limitations

A funnel plot analysis of all RCTs was performed as part of the present meta-analysis. The findings indicated that the publication bias was low (Figure 9). But considering the difficulties in publishing studies with negative findings, many studies likely remain unpublished, and these were not available for analysis. In addition, the quality of assessment scores of the 47 RCTs included was relatively low, owing largely to a lack -

A											
Study or subgroup	Exper Event	rimenta s Total	al Cont Even	rol ts Tota	Weig I (%)	ht Odds ratio M–H, Yea random, 95% Cl	r	Od ran	ds ratio M–H dom, 95% C	, I	
Wang et al (2009)63	85	186	80	124	4.9	0.46 (0.29, 0.74) 200	9				
Bai et al (2013)61	76	365	32	148	4.9	0.95 (0.60, 1.52) 201	3				
Yang et al (2013)58	26	203	12	109	4.0	1.19 (0.57, 2.46) 201	3				
Wu (2013) <sup>60</sup>	78	227	45	165	5.0	1.40 (0.90, 2.16) 201	3				
Wang and Rang (2014) <sup>₄</sup>	°81	171	56	114	4.8	0.93 (0.58, 1.50) 201	1				
Long et al (2014) <sup>56</sup>	120	327	56	328	5.2	2.82 (1.95, 4.06) 201	1				
Gao and Zhao (2014)46	228	496	148	310	5.4	0.93 (0.70, 1.24) 201	4		-		
Chen et al (2014) <sup>45</sup>	197	327	120	202	5.2	1.04 (0.72, 1.48) 201	1		-		
Sun et al (2014) <sup>67</sup>	5/	269	35	144	4.8	0.84 (0.52, 1.35) 201	1				
Shen et al (2014)55	349	815	494	899	5.6	0.61 (0.51, 0.74) 201	4				
Tian et al (2014) <sup>66</sup>	118	135	68	116	4.3	4.90 (2.61, 9.19) 201	4				
Znao (2014) <sup>31</sup>	120	1,511	15	659	4.6	3.70 (2.15, 6.39) 201	-				
Li et al $(2015)^{-2}$	89	312	40	189	5.0	1.24 (0.82, 1.88) 201	-				
	167	420	20	90	4.0	1.14 (0.66, 1.96) 201	-		_		
Wang at al (2015) <sup>10</sup>	107	430	104	2/0	5.5	0.40(0.29, 0.54) 201	-		-		
Wang et al $(2015)^{30}$	409	093	102	407	5.5 E 4	1.32 (1.05, 1.00) 201	-		_		
$2015)^{-2}$	197	401	100	212	5.4	1.17 (0.00, 1.59) 201	2				
Zilu et al (2010) <sup>13</sup>	01	020	42	321	5.0	0.44(0.26, 0.07) 201	2				
Chap at al $(2016)^{25}$	222	04Z	125	201	4.9	2.09 (1.00, 4.30) 201	2				
Cheff et al (2010)-	223	092	155	546	5.5	1.02 (0.00, 1.30) 201	5				
Total (95% CI)		9,419	1	5,75	0100	1.13 (0.87, 1.47)			•		
Total events	2,813		1,888								
Heterogeneity: $\tau^2=0.30$	$\gamma^2 = 189$	971 df	=19 (P	<0 000	$(1)^{1} I^{2} =$	-90%					
Test for overall effect: Z	=0.94 (1	P=0.35	)	0.000	.,,.	00,0	0.01	0.1	1	10	100
		2.00	,					Favors		Favors	
								(blank control)	(ca	rbon nanopartic	es)

Study or subgroup	Experim Events	ental Total	Contro Events	l Total	Weight (%)	Odds ratio M–H, random, 95% Cl	Year
Wang et al (2009)63	83	177	80	124	9.6	0.49 (0.30, 0.78)	2009
Bai et al (2013)61	71	141	32	148	9.3	3.68 (2.20, 6.14)	2013
Sun et al (2014)67	39	188	35	144	9.3	0.82 (0.49, 1.37)	2014
Chen et al (2014)45	194	312	120	202	10.2	1.12 (0.78, 1.61)	2014
Zhao (2014)51	120	1,110	15	659	9.1	5.20 (3.01, 8.98)	2014
Shen et al (2014)55	339	738	494	899	10.9	0.70 (0.57, 0.85)	2014
Gao and Zhao (2014)46	218	474	148	310	10.6	0.93 (0.70, 1.24)	2014
Wang et al (2015)33	400	853	182	467	10.8	1.38 (1.10, 1.74)	2015
Li et al (2015)42	72	263	46	189	9.8	1.17 (0.76, 1.80)	2015
Zhu et al (2016) <sup>19</sup>	195	768	42	327	10.2	2.31 (1.61, 3.32)	2016
Total (95% CI)		5,024		3,469	100	1.33 (0.91, 1.94)	
Total events	1,731	10 df	1,194	00004	1). 12-02	0/	
Test for overall effect: Z	, χ==108. =1.48 (P	=0.14)	=9 (P<0 )	.0000	1), 1-=92	70	0.01



# С

В

Study or subgroup	Experi Events	menta Total	l Contro Events	l Total	Weigh (%)	nt Odds ratio M–H, random, 95% Cl	Year	Odds rando	ratio M- m, 95%	-H, Cl	
Wang et al (2009)63	2	9	80	124	8.2	0.16 (0.03, 0.79)	2009		-		
Bai et al (2013)61	5	224	32	148	9.9	0.08 (0.03, 0.22)	2013				
Shen et al (2014)55	10	77	494	899	10.5	0.12 (0.06, 0.24)	2014				
Chen et al (2014)45	3	15	120	202	9.0	0.17 (0.05, 0.62)	2014	<b>.</b>			
Sun et al (2014)67	18	81	35	144	10.5	0.89 (0.47, 1.70)	2014		•		
Zhao (2014)51	15	401	15	659	10.4	1.67 (0.81, 3.45)	2014				
Gao and Zhao (2014)	<sup>46</sup> 10	22	148	310	10.1	0.91 (0.38, 2.17)	2014		•		
Li et al (2015)42	17	49	46	189	10.5	1.65 (0.84, 3.25)	2015		+		
Zhu et al (2016)19	27	60	42	327	10.6	5.55 (3.04, 10.15)	2016				
Wang et al (2015)33	9	40	182	467	10.3	0.45 (0.21, 0.98)	2015		_		
Total (95% CI)		978		3.469	9100	0.55 (0.23, 1.36)					
Total events	116	••••	1,194	•,.•		,					
Heterogeneity: $\tau^2=1.8$	8; $\chi^2 = 11$	3.69, a	lf=9 (P<0	0.0000	)1); /2=9	92%		1			_
Test for overall effect:	Z=1.29 (	P=0.2	0)				0.01	0.1	1	10	100
	· · · · · · · · · · · · · · · · · · ·							Favors (blank control)	(0	Favors carbon nanoparticles)	

Figure 5 Forest plot showing the relationship of CNs and different kinds of retrieved lymph nodes.

Notes: (A) Total metastatic rate of the retrieved lymph nodes. Metastatic rates of the (B) stained and (C) unstained lymph nodes in the CN and blank control groups. Abbreviations: CN, carbon nanoparticle; M–H, Mantel–Haenszel; random, random effect; CI, confidence interval; df, degrees of freedom.

Study or subgroup	Experim Events	ental Total	Control Events	Total	Weight (%)	Odds ratio M–H, fixed, 95% Cl	Year	Odds ratio M–H, fixed, 95% Cl
Wang et al (2009)63	7	18	8	18	2.1	0.80 (0.21, 3.00)	2009	
Yang et al (2013)58	2	21	10	22	3.9	0.13 (0.02, 0.68)	2013	
Wu (2013)60	0	26	7	29	3.1	0.06 (0.00, 1.05)	2013 +	
Tian et al (2014)66	1	50	8	50	3.4	0.11 (0.01, 0.89)	2014	
Chen et al (2014)45	5	36	13	36	4.9	0.29 (0.09, 0.91)	2014	
Zhang et al (2014)54	0	36	7	36	3.2	0.05 (0.00, 0.98)	2014 +	
Sun et al (2014)67	0	40	3	40	1.5	0.13 (0.01, 2.65)	2014 +	
Liu et al (2014)44	0	23	7	24	3.2	0.05 (0.00, 0.93)	2014 🔶	
Shao et al (2014)57	0	9	5	20	1.5	0.15 (0.01, 3.00)	2014 🔶	
Shen et al (2014)55	5	45	11	64	3.5	0.60 (0.19, 1.87)	2014	<b>-</b> _
Liu et al (2014)49	0	26	7	29	3.1	0.06 (0.00, 1.05)	2014 🔶	
Gao and Zhao (2014)46	0	50	4	50	2.0	0.10 (0.01, 1.95)	2014 +	
Du (2014)50	1	20	7	20	2.9	0.10 (0.01, 0.89)	2014 -	
Wang and Rang (2014)48	0	35	3	35	1.5	0.13 (0.01, 2.63)	2014 +	
Liu and Qing (2015)34	2	33	7	33	2.9	0.24 (0.05, 1.25)	2015	
Duan et al (2015)30	0	40	5	40	2.4	0.08 (0.00, 1.49)	2015 +	
Li et al (2015)42	0	36	0	36		Not estimable	2015	
Du et al (2015)32	1	60	3	58	1.3	0.31 (0.03, 3.08)	2015	
Wang et al (2015)64	0	28	5	27	2.4	0.07 (0.00, 1.37)	2015 +	
Zhu et al (2016) <sup>19</sup>	4	81	14	81	5.8	0.25 (0.08, 0.79)	2016	
Wang et al (2015)39	0	44	9	44	4.1	0.04 (0.00, 0.75)	2015 +	•
Li et al (2015)37	1	26	5	26	2.1	0.17 (0.02, 1.55)	2015	
Wu et al (2015)36	4	90	20	155	6.1	0.31 (0.10, 0.95)	2015	
Yin et al (2015)41	0	40	6	40	2.8	0.07 (0.00, 1.21)	2015 +	
Li (2015) <sup>27</sup>	1	20	7	20	2.9	0.10 (0.01, 0.89)	2015 -	
Feng and He (2015)28	0	30	7	30	3.2	0.05 (0.00, 0.95)	2015 +	
Chu et al (2015)38	8	28	2	29	0.6	5.40 (1.03, 28.23)	2015	
Gu et al (2015)65	3	50	13	50	5.4	0.18 (0.05, 0.68)	2015	
Chen et al (2016)25	6	87	23	86	9.4	0.20 (0.08, 0.53)	2016	
Xu and Gu (2016)23	2	57	7	57	3.0	0.26 (0.05, 1.31)	2016	
Liu et al (2016) <sup>18</sup>	0	78	0	78		Not estimable	2016	
Total (95% CI)		1,263		1,363	100	0.22 (0.16, 0.30)		•
Total events	53		233					
Heterogeneity: $\chi^2$ =31.83,	df=28 (P=	0.28); <i>I</i> <sup>2</sup> =	12%				⊢	
Test for overall effect: Z=9	.73 (P<0.0	00001)					0.01	1 0.1 1 10 100
								Favors Favors (blank control) (carbon nanoparticles)

**Figure 6** Accidental parathyroid removal rate in the CN and blank control groups. **Abbreviations:** CN, carbon nanoparticle; M–H, Mantel–Haenszel; CI, confidence interval; *df*, degrees of freedom.

Α									
Study or subgroup	Experin Events	nental Total	Control Events	Total	Weight (%)	Odds ratio M–H, fixed, 95% Cl	Year	Odds ratio fixed, 95%	o M–H, ⊌ CI
Wu (2013) <sup>60</sup>	1	26	1	29	0.3	1.12 (0.07, 18.86)	2013		
Tian et al (2014)66	1	50	7	50	2.3	0.13 (0.01, 1.06)	2014 -		
Shao et al (2014)57	2	9	11	20	1.8	0.23 (0.04, 1.42)	2014	+	-
Long et al (2014)56	26	75	45	75	9.8	0.35 (0.18, 0.69)	2014		
Du (2014)50	3	20	8	20	2.3	0.26 (0.06, 1.21)	2014	+	
Liu et al (2014)49	1	26	1	29	0.3	1.12 (0.07, 18.86)	2014		
Wang and Rang (2014)48	0	35	5	35	1.8	0.08 (0.00, 1.47)	2014 🔶		-
Zhang et al (2014)54	1	36	5	36	1.6	0.18 (0.02, 1.60)	2014		_
Liu et al (2014)44	0	23	2	24	0.8	0.19 (0.01, 4.21)	2014 🔶		
Liu (2014)53	7	76	36	108	9.0	0.20 (0.08, 0.49)	2014	<b>_</b>	
Wu et al (2015) <sup>36</sup>	10	90	48	155	10.4	0.28 (0.13, 0.58)	2015		
Li et al (2015)37	5	26	10	26	2.7	0.38 (0.11, 1.34)	2015		-
Liu and Qing (2015)34	2	33	10	33	3.1	0.15 (0.03, 0.74)	2015		
Guo et al (2015)29	4	30	15	29	4.4	0.14 (0.04, 0.52)	2015		
Shao et al (2015)35	2	30	9	30	2.8	0.17 (0.03, 0.85)	2015		
Gu et al (2015)65	24	50	28	50	4.9	0.73 (0.33, 1.59)	2015		_
Wang et al (2015)33	5	60	12	60	3.7	0.36 (0.12, 1.11)	2015		
Du et al (2015)32	2	60	7	58	2.3	0.25 (0.05, 1.26)	2015		
Wang et al (2015)64	3	28	6	27	1.8	0.42 (0.09, 1.89)	2015		<u> </u>
Li et al (2015)42	1	36	8	36	2.6	0.10 (0.01, 0.85)	2015 —		
Zhu et al (2016)19	8	81	20	81	6.0	0.33 (0.14, 0.81)	2016		
Wu et al (2015)40	4	43	4	43	1.2	1.00 (0.23, 4.29)	2015		
Feng and He (2015) <sup>28</sup>	1	30	8	30	2.6	0.09 (0.01, 0.82)	2015 —		
Yin et al (2015)41	5	40	14	40	4.1	0.27 (0.08, 0.83)	2015		
Chen and Wu (2015)31	7	48	10	48	2.8	0.65 (0.22, 1.88)	2015		_
Chen et al (2016)25	0	87	3	86	1.2	0.14 (0.01, 2.68)	2016 🔶		
Yang et al (2014)47	18	178	48	201	13.5	0.36 (0.20, 0.64)	2014		
Total (95% CI)		1,326		1,459	100	0.31 (0.25, 0.39)		•	
Total events	143		381						
Heterogeneity: $\chi^2$ =19.55,	df=26 (P=0	0.81); <i>I</i> 2	=0%					0.1 1	10 100
Test for overall effect: Z=1	0.23 ( <i>P</i> <0.	.00001)					0.01	Favors (blank control)	Favors (carbon nanoparticles)

Figure 7 (Continued)

В										
Study or subgroup	Experin Events	nental Total	Control Events	Total	Weight (%)	Odds ratio M–H, fixed, 95% Cl	Year	Odds ra fixed, 9	atio M–H, 5% Cl	
Zeng et al (2014)62	0	40	0	40		Not estimable	2014			
Gao and Zhao (2014)46	0	50	0	50		Not estimable	2014			
Wu et al (2015)36	0	90	2	155	15.1	0.34 (0.02, 7.14)	2015 -			
Guo et al (2015)29	1	30	6	29	48.6	0.13 (0.01, 1.18)	2015 —		+	
Liu and Qing (2015)34	0	33	1	33	12.2	0.32 (0.01, 8.23)	2015 —			
Du et al (2015)32	0	60	1	58	12.5	0.32 (0.01, 7.94)	2015 —		<u> </u>	
Yang et al (2014)47	0	178	1	201	11.6	0.37 (0.02, 9.25)	2014 -	•		
Total (95% CI)		481		566	100	0.24 (0.07, 0.85)				
Total events	1		11							
Heterogeneity: $\chi^2=0.47$ , c	df=4 (P=0.9	98); /²=0	%						+ +	
Test for overall effect: Z=	2.21 (P=0.0	03)					0.01	0.1	1 10	100
		,						Favors (blank control)	Favors (carbon nanopartic	les)

Figure 7 (A) Postoperative transient and (B) permanent hypoparathyroidism rates in the CN and blank control groups. Abbreviations: CN, carbon nanoparticle; M–H, Mantel–Haenszel; CI, confidence interval; *df*, degrees of freedom.

Α								
Study or subgroup	Experin Events	nental Total	Contro Events	Total	Weight (%)	Odds ratio M–H, fixed, 95% Cl	Year	Odds ratio M–H, fixed, 95% Cl
Bai et al (2013)61	5	48	7	40	2.0	0.55 (0.16, 1.88)	2013	
Wu (2013)60	3	26	8	29	2.0	0.34 (0.08, 1.46)	2013	
Yang et al (2013)59	4	36	9	32	2.5	0.32 (0.09, 1.17)	2014	
Huang et al (2013) <sup>24</sup>	3	36	10	36	2.7	0.24 (0.06, 0.95)	2013	
Yang et al (2013)58	2	21	0	22	0.1	5.77 (0.26, 127.60)	2013	
Gao and Zhao (2014)46	0	50	12	50	3.7	0.03 (0.00, 0.53)	2014	<b>←</b> • • • • • • • • • • • • • • • • • • •
Liu et al (2014)49	3	26	8	29	2.0	0.34 (0.08, 1.46)	2014	
Tian et al (2014)66	2	50	10	50	2.9	0.17 (0.03, 0.81)	2014	
Liu (2014)53	6	76	30	108	6.8	0.22 (0.09, 0.57)	2014	
Zhang et al (2014)54	2	36	6	36	1.7	0.29 (0.06, 1.57)	2014	
Liu et al (2014)44	3	23	4	24	1.0	0.75 (0.15, 3.79)	2014	
Chun (2014)52	3	33	7	34	1.9	0.39 (0.09, 1.64)	2014	
Du (2014)50	3	20	7	20	1.8	0.33 (0.07, 1.52)	2014	
Shao et al (2014)57	3	9	12	20	1.5	0.33 (0.06, 1.73)	2014	
Wang and Rang (2014)48	1	35	6	35	1.7	0.14 (0.02, 1.25)	2014	
Duan et al (2015)30	0	40	5	40	1.6	0.08 (0.00, 1.49)	2015	•
Chen and Wu (2015)31	6	48	9	48	2.4	0.62 (0.20, 1.90)	2015	
Du et al (2015)32	2	60	10	58	2.9	0.17 (0.03, 0.79)	2015	
Feng and He (2015)28	2	30	10	30	2.8	0.14 (0.03, 0.72)	2015	
Liu and Qing (2015)34	4	33	10	33	2.6	0.32 (0.09, 1.14)	2015	
Wu et al (2015)40	5	43	7	43	1.8	0.68 (0.20, 2.33)	2015	
Wang et al (2015)33	6	60	15	60	4.0	0.33 (0.12, 0.93)	2015	
Fu (2015) <sup>43</sup>	15	50	106	200	8.9	0.38 (0.20, 0.74)	2015	<b>_</b> _
Li et al (2015)37	3	26	7	26	1.8	0.35 (0.08, 1.56)	2015	
Wu et al (2015)36	12	90	53	155	10.1	0.30 (0.15, 0.59)	2015	<b>_</b>
Li et al (2015)42	1	36	8	36	2.3	0.10 (0.01, 0.85)	2015	
Wang et al (2015)64	3	28	8	27	2.2	0.28 (0.07, 1.22)	2015	
Zhu et al (2016)19	6	81	17	81	4.7	0.30 (0.11, 0.81)	2016	
Yin et al (2015)41	5	40	15	40	3.9	0.24 (0.08, 0.74)	2015	
Shao et al (2015)35	1	30	12	30	3.5	0.05 (0.01, 0.43)	2015	<b>←</b>
Gu et al (2015)65	10	50	12	50	2.9	0.79 (0.31, 2.05)	2015	
Xu and Gu (2016)23	3	57	9	57	2.5	0.30 (0.08, 1.16)	2016	
Chen et al (2016)25	4	87	16	86	4.6	0.21 (0.07, 0.66)	2016	
Total (95% CI)		1,414		1,665	100	0.30 (0.25, 0.38)		♦
Total events	131		465					
Heterogeneity: $\chi^2=23.88$ ,	df=32 (F	P=0.85)	; /²=0%					
Test for overall effect: Z=	10.92 (P<	<0.0000	1)				0.	0.01 0.1 1 10 100
								Favors Favors (blank control) (carbon nanoparticles)

### В

Study or subgroup	Experimental Events Total		Control Events	l Total	Weight (%)	Odds ratio M–H, fixed, 95% Cl	Year	Odds ratio M–H, fixed, 95% Cl		
Zeng et al (2012)62	0	40	0	40		Not estimable	2014			
Huang et al (2013)24	0	36	0	36		Not estimable	2013			
Gao and Zhao (2014)46	0	50	0	50		Not estimable	2014			
Du et al (2015)32	0	60	1	58	45.2	0.32 (0.01, 7.94)	2015 —		<u> </u>	
Wu et al (2015)36	0	90	2	155	54.8	0.34 (0.02, 7.14)	2015 -			
Total (95% CI)		276		339	100	0.33 (0.04, 3.03)				
Total events	0		3							
Heterogeneity: $\chi^2=0.00$ ,	df=1 (P=0	0.98); <i>l<sup>2</sup></i>	<sup>2</sup> =0%						+ +	
Test for overall effect: Z=	=0.98 (P=	0.33)					0.01	0.1	1 10	100
		,						Favors (blank control)	Favors (carbon nanoparticle	s)

Figure 8 (A) Postoperative transient and (B) permanent hypocalcemia rates in the CN and blank control groups. Abbreviations: CN, carbon nanoparticle; M–H, Mantel–Haenszel; CI, confidence interval; df, degrees of freedom.



Figure 9 Funnel plots for publication bias.

Notes: Publication bias for (A) the number of retrieved lymph nodes per patient, (B) number of retrieved metastatic lymph nodes per patient, (C) total metastatic rate of the retrieved lymph nodes, (D) metastatic rate of the stained lymph nodes, (E) metastatic rate of the unstained lymph nodes, (F) accidental parathyroid removal rate, (G) postoperative transient hypoparathyroidism rate, (H) postoperative permanent hypoparathyroidism rate, (I) postoperative transient hypocalcemia rate, and (J) postoperative permanent hypocalcemia rate in the CN and blank control groups. Each point represents a separate study for the indicated association. Abbreviations: CN, carbon nanoparticle; SE, standard error; MD, mean difference; OR, odds ratio.

of blinding; however, it is difficult to apply double-blinding during surgery. Thus, further high-quality research is needed to verify the conclusions of the present study.

# Conclusion

This systematic review and meta-analysis demonstrated that the usage of CNs can improve the extent of neck dissection and preserve the normal anatomic structure and physiological function of the parathyroid. At the same time, the number of retrieved metastatic lymph nodes can also be improved during thyroid cancer surgery.

# Disclosure

The authors report no conflicts of interest in this work.

### References

 Chen AY, Jemal A, Ward EM. Increasing incidence of differentiated thyroid cancer in the United States, 1988–2005. *Cancer*. 2009;115(16): 3801–3807.

- La Vecchia C, Malvezzi M, Bosetti C, et al. Thyroid cancer mortality and incidence: a global overview. *Int J Cancer*. 2015;136(9): 2187–2195.
- 3. Brito JP, Hay ID, Morris JC. Low risk papillary thyroid cancer. *BMJ*. 2014;348:g3045.
- 4. Markovina S, Grigsby PW, Schwarz JK, et al. Treatment approach, surveillance, and outcome of well-differentiated thyroid cancer in childhood and adolescence. *Thyroid*. 2014;24(7):1121–1126.
- Orosco RK, Lin HW, Bhattacharyya N. Ambulatory thyroidectomy: a multistate study of revisits and complications. *Otolaryngol Head Neck Surg.* 2015;152(6):1017–1023.
- Podnos YD, Smith D, Wagman LD, Ellenhorn JD. The implication of lymph node metastasis on survival in patients with well-differentiated thyroid cancer. *Am Surg.* 2005;71(9):731–734.
- Zaydfudim V, Feurer ID, Griffin MR, Phay JE. The impact of lymph node involvement on survival in patients with papillary and follicular thyroid carcinoma. *Surgery*. 2008;144(6):1070–1077; discussion 1077–1078.
- Zhu X, Li J, He H, Huang M, Zhang X, Wang S. Application of nanomaterials in the bioanalytical detection of disease-related genes. *Biosens Bioelectron*. 2015;74:113–133.
- Konvalina G, Haick H. Sensors for breath testing: from nanomaterials to comprehensive disease detection. Acc Chem Res. 2014;47(1):66–76.
- Islam T, Harisinghani MG. Overview of nanoparticle use in cancer imaging. *Cancer Biomark*. 2009;5(2):61–67.
- Peng F, Su Y, Zhong Y, Fan C, Lee ST, He Y. Silicon nanomaterials platform for bioimaging, biosensing, and cancer therapy. *Acc Chem Res.* 2014;47(2):612–623.
- Yezhelyev MV, Gao X, Xing Y, Al-Hajj A, Nie S, O'Regan RM. Emerging use of nanoparticles in diagnosis and treatment of breast cancer. *Lancet Oncol.* 2006;7(8):657–667.
- Liu Z, Liu J, Wang R, Du Y, Ren J, Qu X. An efficient nanobased theranostic system for multi-modal imaging-guided photothermal sterilization in gastrointestinal tract. *Biomaterials*. 2015;56: 206–218.
- Gismondi A, Nanni V, Reina G, Orlanducci S, Terranova ML, Canini A. Nanodiamonds coupled with 5,7-dimethoxycoumarin, a plant bioactive metabolite, interfere with the mitotic process in B16F10 cells altering the actin organization. *Int J Nanomedicine*. 2016;11:557–574.
- Gismondi A, Reina G, Orlanducci S, et al. Nanodiamonds coupled with plant bioactive metabolites: a nanotech approach for cancer therapy. *Biomaterials*. 2015;38:22–35.
- Davis ME, Chen ZG, Shin DM. Nanoparticle therapeutics: an emerging treatment modality for cancer. *Nat Rev Drug Discov*. 2008;7(9): 771–782.
- Yanamala N, Kagan VE, Shvedova AA. Molecular modeling in structural nano-toxicology: interactions of nano-particles with nanomachinery of cells. *Adv Drug Deliv Rev.* 2013;65(15):2070–2077.
- Liu X, Chang S, Jiang X, Huang P, Yuan Z. Identifying parathyroid glands with carbon nanoparticle suspension does not help protect parathyroid function in thyroid surgery: a prospective, randomized control clinical study. *Surg Innov.* 2016;23(4):381–389.
- Zhu Y, Chen X, Zhang H, et al. Carbon nanoparticle-guided central lymph node dissection in clinically node-negative patients with papillary thyroid carcinoma. *Head Neck.* 2016;38(6):840–845.
- Li Y, Jian WH, Guo ZM, Li QL, Lin SJ, Huang HY. A meta-analysis of carbon nanoparticles for identifying lymph nodes and protecting parathyroid glands during surgery. *Otolaryngol Head Neck Surg.* 2015; 152(6):1007–1016.
- Youssef T, Gaballah G, Abd-Elaal E, El-Dosoky E. Assessment of risk factors of incidental parathyroidectomy during thyroid surgery: a prospective study. *Int J Surg*. 2010;8(3):207–211.
- Kurmann A, Herden U, Schmid SW, Candinas D, Seiler CA. Morbidity rate of reoperation in thyroid surgery: a different point of view. *Swiss Med Wkly*. 2012;142:w13643.
- Xu XF, Gu J. The application of carbon nanoparticles in the lymph node biopsy of cN0 papillary thyroid carcinoma: a randomized controlled clinical trial. *Asian J Surg*. Epub 2016 Jan 15.

- Huang K, Luo D, Huang M, Long M, Peng X, Li H. Protection of parathyroid function using carbon nanoparticles during thyroid surgery. *Otolaryngol Head Neck Surg.* 2013;149(6):845–850.
- Chen ZL, Chen GY, Fan HL. Clinical research on the identification and protection effects of nano carbon particles on parathyroid gland in thyroid carcinoma surgery. *J Beihua Univ Nat Sci.* 2016;17(1):89–91.
- Zhang J, Zhang YQ, Zhang AM, et al. Protective effect of nanocarbon on parathyroid gland during thyroid. *Acta Acad Med Weifang*. 2016;38(1):70–72.
- Li DJ. Application value of nano carbon for the protection of parthyroid in thyroid carcinoma surgery. *Chin J Mod Drug Appl.* 2015;9(3):50–51.
- Feng LF, He Z. Value of carbon nanoparticles in the surgery of thyroid carcinoma. J Basic Clin Oncol. 2015;28(1):39–40.
- Guo M, Zhang H, Li WH, Gao JL, Yang J. The protection of the parathyroid and recurrent laryngeal nerve by carbon nanoparticles in cN0 phase thyroidectomy. *China Med Eng.* 2015;23(4):169–171.
- Duan XW, Li ZL, Xu J. Application of carbon nanoparticle tracer in radical neck dissection for thyroid cancer. *Chin J Gen Surg.* 2015;24(5): 638–642.
- Chen SH, Wu ZY. The effection of lymph node tracer in central neck dissection and thyroidectomy. *Chin J Gen Surg.* 2015;24(5):760–763.
- Du GN, Xiao YG, Tan YH, et al. Application of nano-carbon parathyroid negative imaging technique to radical thyroidectomy for thyroid cancer. *J Chin Oncol.* 2015;21(6):469–471.
- Wang W, Hang B, Lin P, et al. Research on the feasibility and significance of Delphian lymph node labeling by nanometer carbon in thyroid cancer surgery. *MMJC*. 2015;17(7):7–10.
- Liu JW, Qing XY. The usage of carbon nanopartciles in preserving parathyroid and the value in thyroidectomy. *Guide China Med.* 2015; 13(25):124–125.
- Shao Y, Zhao MR, Bai YX, et al. The application of the nano-carbon tracer in 30 cases of thyroid gland carcinoma operation. *Mod Oncol.* 2015;23(20):2915–2917.
- Wu HS, Xiao Y, Zheng C, Wang XM. Application of negative stained parathyroid in thyroid carcinoma surgery by using carbon nanoparticles. *Shenzhen J Integr Trad Chin West Med.* 2015;25(18):6–9.
- Li XT, Lin Y, Dai P, Ai QB, Zhang GR. Clinical study of avoiding intraoperative parathyroid cut by mistake or function impairment. *China Mod Med.* 2015;22(29):26–28.
- Chu HD, Jiang LX, Zheng HT. Activated carbon nanoparticle in dissection of central lymph nodes of papillary thyroid carcinomas: a control study. *Chin J Oper Proc Gen Surg (Electronic Edition)*. 2015; 9(5):57–59.
- Wang LP, Liu CS, Luo XY. Application value of nano carbon for the protection of parathyroid glands in thyroid carcinoma surgery. *Zhong Guo She Qu Yi Sheng*. 2015;31(32):31–33.
- 40. Wu G, Cai L, Hu J, et al. [Role of carbon nanoparticles in patients with thyroid carcinoma undergoing total thyroidectomy plus bilateral central neck dissection]. *Zhonghua Yi Xue Za Zhi*. 2015;95(12):912–916. Chinese [with English abstract].
- Yin FY, Xin H, Zhang XJ, et al. Protective effects of nano-carbon on parathyroid in thyroid surgery. *Chin J Endocr Surg.* 2015;9(2): 144–146.
- Li HQ, Yin DT, Wang YF, et al. Carbon nanoparticles in central lymph node dissection in treatment of papillary thyroid carcinoma. *Chin J Endocr Surg.* 2015;9(5):398–400.
- Fu K. The Comparison of Different Methods in Protecting Parathyroid Gland and Clinical Significant [dissertation]. Shijiazhuang: Hebei Medical University; 2015.
- 44. Liu XL, Han B, Wu WW, Sun DS, Wei W. Experience of preoperative ultrasound guided carbon nanoparticles thyroid injection in thyroid carcinoma surgery: an analysis of 47 cases. *Zhong Guo Shi Yong Wai Ke Za Zhi*. 2014;34(Suppl 1):7–9.
- 45. Chen W, Lv Y, Xie R, Xu D, Yu J. [Application of lymphatic mapping to recognize and protect parathyroid in thyroid carcinoma surgery by using carbon nanoparticles]. *Lin Chung Er Bi Yan Hou Tou Jing Wai Ke Za Zhi.* 2014;28(24):1918–1920, 1924. Chinese [with English abstract].

- 46. Gao Q, Zhao D. [Clinical application of carbon nanoparticles labeled lymph node in cervical lymph node dissection with papillary thyroid cancer staged preoperatively as N0]. *Lin Chung Er Bi Yan Hou Tou Jing Wai Ke Za Zhi*. 2014;28(24):1938–1940. Chinese [with English abstract].
- Yang JF, Li H, Hu B, et al. The protection of carbon nanoparticles for parathyroid gland in total thyroidectomy and central lymph dissection. J Clin Otorhinolaryngol Head Neck Surg (China). 2014;28(18): 1382–1384.
- Wang JW, Rang CF. Protective effects of carbon nanoparticles on parathyroid in thyroid surgery. *Medical Journal of National Defending Forces in Southwest China*. 2014;24(6): 609–612.
- Liu N, Han B, Wei W. Application of carbon nanoparticles in thyroid cancer surgery. *Guangdong Med J.* 2014;35(12):1918–1920.
- Du J. Clinical Research of Carbon Nano Lymph Tracer on Parathyroid Protective Effect in Operation of Thyroid Carcinoma [dissertation]. Dalian: Dalian Medical University; 2014.
- Zhao B. Carbon Nanoparticles Tracer in the Early Thyroid Papillary Carcinoma Clinical Studies of Surgical Treatment [dissertation]. Kunming: Kunming Medical University; 2014.
- Chun L. The Applied Significance of Carbon Nanoparticles in Surgery of Papillary Thyroid Microcarcinoma [dissertation]. Chongqing: Chongqing Medical University; 2014.
- Liu ZF. Research on Nano-Carbon Protection of Parathyroid Glands in Operation of Thyroid Carcinoma [dissertation]. Qingdao: Shandong University; 2014.
- Zhang Q, Zhang Q, Deng LF. Protective effects of carbon nanoparticles on parathyroid gland during thyroidectomy. *Chin J Prim Med Pharm*. 2014;21(22):3408–3409.
- 55. Shen H, Wei B, Feng S, Zhou Q. [Efficiency of carbon nanoparticles in level VI lymphadenectomy for thyroid carcinoma and prevention of postoperative hypoparathyroidism]. *Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi.* 2014;49(10):817–820. Chinese [with English abstract].
- Long MY, Long HY, Huang MQ, et al. Activated nano carbon in prophylactic central lymph node dissection of T1 papillary thyroid non-microcarcinoma. *Chin J Endocr Surg.* 2014;8(5):422–424.
- Shao Q, Zou RL, Shan HL, et al. Application of carbon nanoparticles during thyroid surgery. J Clin Med Pract. 2014;18(2):34–35.

- Yang XH, Wang Y, Wang P. Application of carbon nanoparticle in endoscopic surgery of thyroid carcinoma. *J Laparosc Surg.* 2013; 18(4):262–265.
- Yang ZF, Yue RX, Zhu Z, et al. The applied significance of carbon nanoparticles in central compartment lymph node dissection in treatment of cN0 papillary thyroid carcinoma. *Chin J Bases Clin Gen Surg.* 2013;20(9): 976–980.
- 60. Wu WW. The Significance of Applying Preoperative Ultrasound Guided Carbon Nanoparticles Injection in Thyroid Cancer Surgery [dissertation]. Shantou: Shantou University; 2013.
- Bai YC, Zhang JM, Su YJ, Diao C, Qian J, Chen RC. Role of lymphatic tracers in lymph node dissection and pathological examination of papillary thyroid carcinoma. *Chin J Clin Oncol.* 2013;40(17):1034–1037.
- 62. Zeng YJ, Qian J, Chen RC. Protective effect of lymphatic tracer on parathyroid glands in lymph node dissection in thyroid carcinoma. *Chongqing Med J.* 2012;41(11):1076–1088.
- Wang XL, Wu YH, Xu ZG, Ni S, Liu J. [Parathyroid glands are differentiated from lymph node by activated-carbon particles]. *Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi*. 2009;44(2):136–140. Chinese [with English abstract].
- Wang B, Qiu N, Zhang W, et al. The role of carbon nanoparticles in identifying lymph nodes and preserving parathyroid in total endoscopic surgery of thyroid carcinoma. *Surg Endosc.* 2015;29(10):2914–2920.
- 65. Gu JL, Wang JF, Nie XL, Wang WD, Shang JB. Potential role for carbon nanoparticles identification and preservation in situ of parathyroid glands during total thyroidectomy and central compartment node dissection. *Int J Clin Exp Med.* 2015;8(6):9640–9648.
- Tian W, Jiang Y, Gao B, et al. Application of nano-carbon in lymph node dissection for thyroid cancer and protection of parathyroid glands. *Med Sci Monit*. 2014;20:1925–1930.
- Sun SP, Zhang Y, Cui ZQ, et al. Clinical application of carbon nanoparticle lymph node tracer in the VI region lymph node dissection of differentiated thyroid cancer. *Genet Mol Res.* 2014;13(2):3432–3437.
- Yan X, Zeng RC, Ma Z, et al. The utility of sentinel lymph node biopsy in papillary thyroid carcinoma with occult lymph nodes. *PLoS One*. 2015;10(6):e129304.

#### **OncoTargets and Therapy**

#### Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: http://www.dovepress.com/oncotargets-and-therapy-journal

#### **Dove**press

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.