Open Access Full Text Article

The smaller the better?

LETTER

Cornelis Slagt

Department of Anaesthesia, Pain and Palliative Medicine, Radboud University Medical Center, Nijmegen, the Netherlands

Dear editor

With great interest, I have read the article written by Khraise et al¹ which retrospectively investigated the incidence and risk factors for postdural puncture headache (PDPH) after cesarean delivery in Jordanian women. Their main conclusions were that repeated puncture attempts and presence of tension headache before the cesarean delivery were risk factors for PDPH.

The presence of PDPH is a potential burden to the mother and the family, with a large, negative impact on the puerperal period. Moreover, the correct diagnosis of PDPH can be difficult. After analyzing 95 women with presumed PDPH, Stella et al found that the majority had tension and migraine headache, and preeclampsia was the cause of the headache.² Khraise et al found an association between tension headache and PDPH which could actually be a diagnostic issue.

The pathophysiology behind PDPH is complex and is suggested to involve leakage of spinal fluids through the punctured hole. Larger needles will cause bigger holes increasing the risk of developing PDPH. Other risk factors which have been identified are younger age, female gender, cutting needle tip, bevel direction (perpendicular to fibers of the ligaments), vaginal delivery and experience of the anesthetist (in training).³

In the study by Khraise et al, two different needles were used: a pencil-point 25-G and a traumatic 27-G Spinostar. A very interesting finding in this study is that the use of the traumatic 27-G Spinostar needle leads to repeated punctures, increasing the risk of developing PDPH. The positioning of fine needles can be difficult, and an introducer needle through which the 27-G spinal needle can be introduced is needed for guidance. The findings of this study pose the following question: although the anesthetists are creatures of habit, is it time to get rid of the 27-G needle?

Disclosure

The author reports no conflicts of interest in this communication.

References

- Khraise WN, Allouh MZ, El-Radaideh KM, Said RS, Al-Rusan AM. Assessment of risk factors for postdural puncture headache in women undergoing cesarean delivery in Jordan: a retrospective analytical study. *Local Reg Anesth.* 2017;10:9–13.
- Stella CL, Jodicke CD, How HY, Harkness UF, Sibai BM. Postpartum headache: is your work-up complete? Am J Obstet Gynecol. 2009;198:318.e1–318.e7.
- Gaiser RR. Postdural puncture headache: a headache for the patient and a headache for the anesthesiologist. Curr Opin Anaesthesiol. 2013;26(3):296–303.

Local and Regional Anesthesia 2017:10 51-52

Construction of the work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress.com/terms. php and incorporate the Creative Commons Artifubution — Non Commercial (unported, v3.0) License (http://reative.commons.org/license/by-nc/3.0/). By accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial use of this work, please see paragraphs 4.2 and 5 of our Terms (https://www.dovepress.com/terms.php).

Correspondence: Cornelis Slagt Department of Anaesthesia, Pain and Palliative Medicine, Radboud University Medical Center, Geert Grooteplein-Zuid 10, 6500 HB Nijmegen, the Netherlands Tel +31 24 361 4406 Email slagtc@gmail.com



Dove Medical Press encourages responsible, free and frank academic debate. The content of the Local and Regional Anesthesia 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Local and Regional Anesthesia editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Local and Regional Anesthesia

Publish your work in this journal

Local and Regional Anesthesia is an international, peer-reviewed, open access journal publishing on the development, pharmacology, delivery and targeting and clinical use of local and regional anesthetics and analgesics. The journal is included in PubMed, and welcomes submitted papers covering original research, basic science, clinical studies,

Submit your manuscript here: https://www.dovepress.com/local-and-regional-anesthesia-journal

52

reviews and evaluations, guidelines, expert opinion and commentary, case reports and extended reports. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit http://www.dovepress.com/ testimonials.php to read real quotes from published authors.