

Nonobese, exercising children diagnosed with dyslipidemia have normal C-reactive protein

Miguel Arturo Salazar Vázquez¹

Beatriz Yadira Salazar Vázquez^{2,3}

M Intaglietta³

Pedro Cabrales⁴

¹Facultad de Ciencias Químicas Gómez Palacio, Universidad Juárez del Estado de Durango, Dgo., México;

²Faculty of Medicine, Universidad Juárez del Estado de Durango, Victoria de Durango Dgo., México;

³Department of Bioengineering, University of California, San Diego, La Jolla, CA, USA; ⁴La Jolla Bioengineering Institute, La Jolla, CA, USA

Abstract: Nonobese children age 10.4 ± 1.1 years diagnosed with dyslipidemia ($n = 51$) were compared to normal children age 10.8 ± 1.1 years ($n = 38$). Affected individuals had increased total cholesterol: 223 ± 23 vs 152 ± 17 mg/dl, $p < 0.001$; and decreased high-density lipoprotein-cholesterol: 41.9 ± 4.1 vs 57.6 ± 5.7 mg/dl, $p < 0.001$ and triglycerides: 90.8 ± 40.5 vs 65.7 ± 25.0 mg/dl, $p < 0.002$. Fasting glucose was also significantly elevated ($p < 0.02$). All other parameters, including blood pressure, were not statistically different between groups. The concentration of C-reactive protein was not statistically different between groups. Analysis of medical records showed that this anomaly may be related to this group (as well as the control group) performing regular, daily exercise. This activity was quantified via a self administered questionnaire, and found to be statistically identical in controls and dyslipidemic individuals. Exercise is associated with the release of antiinflammatory cytokines, therefore our results support the contention that it is a significant factor in promoting health conditions from an early stage in life.

Keywords: dyslipidemia, C-reactive protein, children, exercise, inflammation, diabetes, obesity, metabolic syndrome, hispanics

Introduction

Dyslipidemia is a metabolic condition associated with abdominal obesity, hypertension, insulin resistance (glucose intolerance), prothrombotic conditions and the presence of pro-inflammatory factors, even though this latter condition maybe at a low level. It presents different forms whose common denominator is the alteration of lipid metabolism, manifested by changes in lipids and lipoproteins in blood. A generalized pro-inflammatory condition is presently recognized to be the precursor for atherosclerosis,^{1,2} type 2 diabetes, and associated long range risk factor for vascular and cardiovascular disease.^{3,4}

The incidence of inflammatory conditions and dyslipidemia is relatively high in the adult population. Conversely, its prevalence in children is low, but, increases significantly in overweight children.⁵ However, circumscribing clinical evaluation for identification of proinflammatory conditions solely to obese children and adolescents would limit the opportunities of diagnosis and prevention in lean young individuals.⁶ Recent studies show that children from Hispanic origin (Mexican-American) present a higher risk to develop this syndrome than their Anglo-Saxon counterpart.^{7,8}

The present study was conducted to determine the association between the diagnosis of dyslipidemia in lean children who perform regular exercise and the incidence of plasma C-reactive protein (CRP), to test the hypothesis that dyslipidemia is associated with the presence of inflammatory conditions, that may pose a risk even in the absence of overweight. Results were compared to those from a nondyslipidemic group.

Methods

Nonobese children in the age range of 8 to 12 years residing in the city of Durango were enrolled in this study. They were recruited in the Unidad de Medicina del Deporte de la

Correspondence: Pedro Cabrales
La Jolla Bioengineering Institute, 505
Coast Boulevard South Suite #405,
La Jolla, CA 92037, USA
Tel +1 858 5342315
Email pcabrales@ucsd.edu

Facultad de Medicina Sports Medicine Unit of the Faculty of Medicine) UJED by open invitation. They were entered into the study following individual and parental consent, according to regulations governing the study and treatment of children. Individuals were divided into a group consisting of nonobese children with dyslipidemia, and a control group of nonobese children without dyslipidemia. Exclusion criteria were the use of any medicament taken daily, children with previous diagnosis of renal, renovascular and endocrine disease, and hypertension.

For the purpose of this study we defined a control group selected on the basis of not being dyslipidemic. Anthropometric parameters recorded were height, weight and body mass index.

Fasting glucose (10–12 hours after food ingestion) was measured in blood samples by the glucose oxidase enzymatic method. Hyperglycemia was defined when fasting glucose was equal to or greater than 100 mg/dL. Dyslipidemia was diagnosed following determination of elevated cholesterol (>200 mg/dL) and triglycerides (>150 mg/dL) and reduced high-density lipoprotein-cholesterol (HDL-C) (<50 mg/dL), when 2 out of 3 of these parameters were exceeded. Concentrations were measured from 5 ml blood samples (no anticoagulants), incubated at 37°C for 5 minutes, centrifuged at 3000 RPM for 5 minutes and processed by an autoanalyzer (Data Pro Plus, Thermo Fisher Scientific, Inc. Waltham, MA, USA). C-reactive protein was determined from 3 ml venous blood samples (no anticoagulant) processed using the DPC IMMULITE High Sensitivity CRP instrument (Diagnostic Products Corp., Los Angeles, CA, USA).

Systolic and diastolic pressures were recorded and mean arterial blood pressure (MAP) was determined using the relationship:

$$\text{MAP} = P_{\text{diastolic}} + 1/3(P_{\text{systolic}} - P_{\text{diastolic}})$$

The level of physical activity is increasingly recognized as a factor in promoting the metabolic syndrome therefore it is important to establish the relative level of activity in both

groups. Daily activity was measured by means of a self-report questionnaire, designed following the scheme of Aadahl and Jørgensen⁹ partially modified to reflect the younger age of the tested population and the difference in environment. The test was administered to 20 individuals in each group chosen at random by blind selection.

The data was analyzed to determine statistically significant differences according to the Student's *t* test. Statistical significance is assumed for $p < 0.05$.

Results

Fifty one dyslipidemic nonobese children age 10.4 ± 1.1 years and 38 controls age 10.8 ± 1.1 were entered into the study. Their characteristics are given in Table 1. Table 2 summarizes our results showing that affected individual had triglycerides 90.0 ± 40.5 vs 65.7 ± 25.0 mg/dL, $p < 0.002$; increased total cholesterol: 223 ± 23 vs 152 ± 17 mg/dL, $p < 0.001$; and diminished HDL-C: 41.9 ± 4.1 vs 57.6 ± 5.7 mg/dL, $p < 0.001$. Fasting glucose were also significantly elevated ($p < 0.02$). All other parameters, including blood pressure and the concentration of CRP were not statistically different between groups. Analysis of medical records showed that this anomaly was related to this group (as well as the control group) performing regular, daily exercise for a minimum period of one hour.

Twenty individuals were selected at random for each group to determine the respective levels of physical activity. This was found to be 45.7 ± 3.1 for controls and 45.9 ± 2.9 h MET-time/day (MET: metabolic equivalents or estimation of energy expenditure for specific physical activity) for the sample of study population, where the differences were not statistically significant. As a comparison Aadahl and Jørgensen⁹ report a range of results for young adults including both leisure time and work activity ranging from 41.0 ± 9.9 to 61.7 ± 14.5 24 h MET-time/day. An independently sampled equivalent sampled group of normal children with comparatively low physical measured 38.5 ± 1.2 h MET-time/day using the same

Table 1 Characteristics of the sample populations

	Children with dyslipidemia n = 51	Controls n = 38	Significance (p, Student's <i>t</i> test)
Age, years	10.3 ± 1.1	10.7 ± 1.1	n.s.
Males, n (%)	30 (59%)	16 (42%)	–
Females, n (%)	21 (41%)	22 (58%)	–
History of diabetes in family, n (%)	29 (57%)	19 (50%)	n.a.
History of hypertension in family, n	31 (61%)	17(45%)	n.a.
Weight, kg	37.4 ± 6.7	41.3 ± 9.5	<0.05
BMI, kg/m ²	18.4 ± 2.4	19.1 ± 2.6	n.s.

Abbreviations: BMI, body mass index; n.s., not significant, $p > 0.05$; n.a., not applicable because data was mostly anecdotal.

Table 2 Differences between children with dyslipidemia and controls

	Children with dyslipidemia	Controls	Significance p, student's t test
C-reactive protein, mg/L	2.5 ± 1.3 Range: 0.3–4.5	2.3 ± 1.4 Range: 0.7–4.5	0.40 (n.s.)
Systolic pressure, mmHg	97.7 ± 13.4	96.6 ± 9.5	n.s.
Diastolic pressure, mmHg	56.5 ± 9.7	57.6 ± 10.5	n.s.
Fasting glucose, mg/dl	90.3 ± 7.4	85.4 ± 9.8	<0.02
Triglycerides, mg/dl	90.9 ± 40.5	65.7 ± 25.0	<0.002
Total cholesterol, mg/dl	223 ± 23	152 ± 17	<0.001
HDL-cholesterol, mg/dl	41.9 ± 4.1	57.6 ± 5.7	<0.001

Abbreviations: HDL, high-density lipoprotein; n.s. not significant, $p > 0.05$.

self evaluation questionnaire, the difference being statistically significant from the measurement in control active children ($p < 0.001$).

Discussion

The principal finding of this study is that children diagnosed with dyslipidemia do not present elevated CRP in their plasma when compared with a matched cohort of normal children. Notably the affected cohort was clearly dyslipidemic with diminished HDL-C and increased total cholesterol levels presenting the classic pattern of elevated total cholesterol, and decreased HDL-C, differences that were statistically significant ($p < 0.001$). There was no particular trend in the anthropometric data, with the exception of weight that was somewhat greater for controls ($p < 0.05$).

The result that the concentration of CRP in children diagnosed as dyslipidemic was not statistically significantly different from control is unexpected and required further scrutiny of the population characteristics. As an example a recent study conducted in dyslipidemic children and adolescents with different combinations of low and high HDL-C and triglycerides and average body mass index (BMI) ranging from 21.3 to 24.8 kg/m² (control 20.1 kg/m²) had significantly elevated levels of CRP. Notably the prevalent level of exercise was not evaluated in these populations.¹⁰ Analysis of the clinical questionnaires administered to the population sampled in our study revealed that both groups (diagnosed and controls) routinely carried out a regular regime of daily, one hour or more exercise. Therefore this population may be markedly different from the more sedentary individuals that manifest early signs of overweight or obesity, who encompass a greater number of factors linked to the metabolic syndrome.

A significant factor of cardiovascular risk (particularly in adults) is the presence of pro-inflammatory conditions evidenced by elevated CRP. The significant decrease of this marker in the population studied confirms that the inherent risk factors are reduced in individuals that perform regular exercise,¹¹ an effect noted in previous studies primarily in adults and older people. The present finding shows that exercise is necessary and beneficial at the earliest age. The incidence of increasingly sedentary life styles evidenced also in children should be a factor on the increased incidence of diabetes in younger individuals, and the lowering of the threshold at which these diseases appear in the population.

It has been recently recognized that exercising skeletal muscle is the source of antiinflammatory cytokines (IL-6) which significantly lower the incidence of inflammatory markers, including CRP.¹² This effects has been observed in young and elder individuals¹³ who do consistent daily exercise. This mechanism is now considered to be the basis for the noted beneficial effects resulting from the performance of exercise,^{14–16} as well as the generally improved health condition of exercising individuals compared to sedentary persons. It should be noted however that although reducing CRP concentrations through lifestyle changes or pharmacological interventions is widely held to be of clinical benefit^{17,18} there remains a critical need for long-term studies to determine whether reductions in CRP concentrations translate into a decreased cardiovascular risk factor.¹⁹

Regular exercise has the additional benefit of increasing blood flow in the circulation, with the effect of increasing shears stress and the production of nitric oxide (NO) by the endothelium, increasing NO bioavailability. This effect is generally considered to produce an antiinflammatory environment.^{20,21}

The results and interpretations of the present study are directly linked to physical activity as a determining factor, which is a challenge to evaluate since there is no gold standard to obtain this information. Continuous recording of single axis body acceleration by a transducer designed to measure and record acceleration as a function of time is an option for obtaining reliable first order evaluation of this parameter.²² A doubly labeled water method²³ is also considered to be a valid method for measuring energy expenditure. However, both methods are complex and expensive to implement, and not suitable for comparatively large studies in children populations.

In view of these considerations we adopted the “self-questionnaire” technique developed by Aadahl and Jørgensen⁹ minimally modified to account for the youth of our population and environmental differences, since their approach was developed for adults in the population of Denmark. The significant correlation found by these authors between results of their questionnaire and the combined data from actual measurement and a diary of daily activity suggested that their approach could be effectively used in this study, a perception in part supported by the low within group variability, and the significant difference between active and the low level exercising normal group found in our study.

A tentative conclusion of this study is that children that report similar high levels of physical exercise have low levels of CRP whether they are normal or are diagnosed dyslipidemic. Notably, similar studies in children where exercise was not quantified report a significant difference in CRP. The data on CRP and exercise presents some dispersion; however this was not sufficient for determining a statically significant association between CRP and levels of exercise. The possibility exists that the recorded levels of exercise are sufficient to overcome the inflammatory component manifested by the increase of concentration of CRP, a process that, however, may not be complete since there is a statistically not significant remnant increase in CRP in the dyslipidemic children. Verification of this conjecture rests on the possibility of testing a significant number of nonobese dyslipidemic children that engage in a low level of physical activity, which in the population tested is a small sub-group.

Although exercise is a significant factor in reducing inflammatory and pro-inflammatory conditions, it cannot be excluded that in the absence of obesity, the expression of pro-inflammatory markers such as CRP may be absent. Therefore in a population of active and thin adolescents dyslipidemia may not be necessarily be linked to inflammatory

conditions, and may be due to genetic and familial antecedents or unusual dietary habits.^{24,25}

In conclusion, there is a population of children with a median age of 10 years that presents diminished plasma concentration of HDL-C and increased cholesterol, a condition classified as dyslipidemia, that is prognostic and precursor to the metabolic syndrome, cardiovascular dysfunction and diabetes. However, an elevated level of CRP associated with these syndromes is not present in this population of nonobese children that perform regular exercise. Exercise is regarded as a critical component of the maintenance of “healthy” conditions. This study shows that the effects of exercise may be important in very young individuals. The effects of inflammatory conditions, some of them derived from dietary habits, are most likely cumulative, thus there should be health benefits in engaging in moderate but not inconsequential routine exercise, at an early age.

Acknowledgments

This investigation was supported in part by Laboratorios Nova, Victoria de Durango, Dgo., Mexico. The authors report no conflicts of interest in this work.

References

1. Li S, Chen W, Srinivasan SR, et al. Childhood cardiovascular risk factors and carotid vascular changes in adulthood: the Bogalusa Heart Study. *JAMA*. 2003;290:2271–6.
2. Raitakari OT, Juonala M, Kahonen M, et al. Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood: the Cardiovascular Risk in Young Finns Study. *JAMA*. 2003;290:2277–83.
3. Dandona P, Ajada A, Bandyopadhyay A. Inflammation: the link between insulin resistance, obesity and diabetes. *Trends Immunol*. 2004;25:4–7.
4. Libby P. Inflammation in atherosclerosis. *Nature*. 2002;420:868–74.
5. Ford ES, Ajani UA, Mokdad AH. The metabolic syndrome and concentrations of C-reactive protein among U.S. youth. *Diabetes Care*. 2005;28:878–81.
6. Barlow SE, Dietz WH. Obesity evaluation and treatment: Expert Committee recommendations. The Maternal and Child Health Bureau, Health Resources and Services Administration and the Department of Health and Human Services. *Pediatrics*. 1998;102:E29.
7. Posadas-Sanchez R, Posadas-Romero C, Zamora-Gonzalez J, et al. Lipid and lipoprotein profiles and prevalence of dyslipidemia in Mexican adolescents. *Metabolism*. 2007;56:1666–72.
8. de Ferranti SD, Gauvreau K, Ludwig DS, et al. Prevalence of the metabolic syndrome in American adolescents: findings from the Third National Health and Nutrition Examination Survey. *Circulation*. 2004;110:2494–7.
9. Aadahl M, Jørgensen T. Validation of a new self-report instrument for measuring physical activity. *Med Sci Sports Exerc*. 2003;35:1196–202.
10. Medina-Urrutia A, Juárez-Rojas JG, Martínez-Alvarado R, et al. High-density lipoprotein subclasses distribution and composition in Mexican adolescents with low HDL cholesterol and/or high triglyceride concentrations, and its association with insulin and c-reactive protein. *Atherosclerosis*. 2008;Mar 18 [Epub ahead of print].
11. Petersen AM, Pedersen BK. The anti-inflammatory effect of exercise. *J Appl Physiol*. 2005;98:1154–62.

12. Pedersen BK, Steensberg A, Fischer C, et al. The metabolic role of IL-6 produced during exercise: is IL-6 an exercise factor? *Proc Nutr Soc.* 2004;63:263–7.
13. Abramson JL, Vaccarino V. Relationship between physical activity and inflammation among apparently healthy middle-aged and older US adults. *Arch Intern Med.* 2002;162:1286–92.
14. Geffken DF, Cushman M, Burke GL, et al. Association between physical activity and markers of inflammation in a healthy elderly population. *Am J Epidemiol.* 2001;153:242–50.
15. Hu G, Tuomilehto J, Silventoinen K, et al. The effects of physical activity and body mass index on cardiovascular, cancer and all-cause mortality among 47 212 middle-aged Finnish men and women. *Int J Obes (Lond).* 2005;29:894–902.
16. LaMonte MJ, Blair SN, Church TS. Physical activity and diabetes prevention. *J Appl Physiol.* 2005;99:1205–13.
17. Ridker PM, Cushman M, Stampfer MJ, et al. Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med.* 1997;336(14):973–9.
18. Ridker PM, Hennekens CH, Buring JE, et al. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med.* 2000;342:836–43.
19. Dandona P. Effects of antidiabetic and antihyperlipidemic agents on C-reactive protein. *Mayo Clin Proc.* 2008;83:333–42.
20. Ni W, Egashira K, Kataoka C, et al. Antiinflammatory and anti-arteriosclerotic actions of HMG-CoA reductase inhibitors in a rat model of chronic inhibition of nitric oxide synthesis. *Circ Res.* 2001;89:415–21.
21. Yamawaki H, Lehoux S, Berk BC. Chronic physiological shear stress inhibits tumor necrosis factor-induced proinflammatory responses in rabbit aorta perfused ex vivo. *Circulation.* 2003;108:1619–25.
22. Hendelman D, Miller K, Baggett C, et al. Validity of accelerometry for the assessment of moderate intensity physical activity in the field. *Med Sci Sports Exerc.* 2000;32(9 Suppl):S442–9.
23. Conway JM, Seale JL, Jacobs DR Jr., et al. Comparison of energy expenditure estimates from doubly labeled water, a physical activity questionnaire, and physical activity records. *Am J Clin Nutr.* 2002;75:519–25.
24. Sveger T, Flodmark CE, Nordborg K, et al. Hereditary dyslipidemias and combined risk factors in children with a family history of premature coronary artery disease. *Arch Dis Child.* 2000;82:292–6.
25. Daniels SR, Greer FR. Lipid screening and cardiovascular health in childhood. *Pediatrics.* 2008;122:198–208.

Appendix

Exercise self-report questionnaire

PERSONAL INFORMATION

Name:	Last name:			
Age:			<input type="radio"/> Female	<input type="radio"/> Male

Home address

Street:	
Number:	
Zip code:	
Phone number:	

Occupation or work

Street:	
Number:	
Zip code:	
Phone number	

Weight		Height	
Waist		Hip	
Blood pressure		Heart rate	
Body mass index			
Product $\geq 4\text{kg}$		Product $< 2\text{ kg}$	

FISICAL ACTIVITY

Of the following activities, please mark these that you have done in your free time in the last 12 months and select the circle that better indicates the frequency whereupon the activity was performed:

Activity	¿Qué día las realizó?							5-14	15-30	31-60	1-2 hr	3-4 hr	5-6 hr	> 6 hr	Cada actividad la realiza en forma:		
	L	M	M	J	V	S	D	min sem	min sem	min sem	sem	sem	sem	hr sem	Light	Moderate	Intense
None <input type="radio"/>																	
To walk	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
To run	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Cycling	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Baseball	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Soccer	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Volleyball	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Aerobics	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
To swim	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Bowling	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
To dance	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
Other																	

Notes: **Light:** like the activity that is resembled to walk in calm form. **Moderate:** activity that realizes with effort, causing in you perspiration. **Intense:** activity that realizes like training or preparing themselves for a sport competition

Daily activities	1 Monday 2 Tuesday 3 Wednesday 4 Thursday 5 Friday 6 Saturday 7 Sunday							5 min	10 min	30 min	1 hr	2–3 hr	4–5 hr	6–7 hr sem
	1	2	3	4	5	6	7							
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
								<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>

Additional notes by the interviewer

Some familiar has suffered or suffers	Grandparents to father	Maternal grandparents	Mother	Father	brothers
¿Hypertension?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
NO Yes (¿Who?)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
¿Diabetes mellitus?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
NO Yes (¿Who?)	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
¿myocardial infarction?	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
NO Yes (¿Who?)					
Renal disease?					
NO Yes (¿Who?)					
¿Obesity?					
NO Yes (¿Who?)					
some other disease in its family					

What medications do you take in the present time?

[illegible]

Please check if you ever had done one of the procedures mentioned below or if any doctor has told that you had or have any of the next diseases and the year that you were told for the first time

just check what you have had					
Diabetes					
Blood pressure					
Cholesterol					
Triglycerides					
Renal disease					