

Efficacy of Acupuncture-Related Therapy in the Treatment of Knee Osteoarthritis: A Network Meta-Analysis of Randomized Controlled Trials

Wei Liu^{1,2,*}Yihua Fan^{1,2,*}Yuanhao Wu^{1,2,*}Xu Hou³Bin Xue^{1,2}Peihao Li^{1,2}Shumin Zhang^{1,2}Qingyun Yue^{1,2}

¹First Teaching Hospital of Tianjin University of Traditional Chinese Medicine, Tianjin, 300193, People's Republic of China; ²National Clinical Research Center for Chinese Medicine Acupuncture and Moxibustion, Tianjin, 300381, People's Republic of China; ³Department of Endocrinology and Metabolic Diseases, Shandong Provincial Hospital Affiliated to Shandong First Medical University, Jinan, Shandong Province, 250021, People's Republic of China

*These authors contributed equally to this work

Objective: Knee osteoarthritis (KOA) is prevalent in middle-aged and elderly people. This condition negatively affects the quality of life of patients. Although non-steroidal anti-inflammatory drugs (NSAIDs) are often used to relieve symptoms associated with KOA, it is associated with many side effects. Acupuncture and moxibustion therapies have been applied in the treatment of KOA. However, the efficacy of various acupuncture and moxibustion treatments has not been compared.

Methods: Randomized controlled trials (RCTs) on the application of acupuncture and moxibustion in the treatment of KOA were searched in English databases and Chinese databases. Data were retrieved from establishment of the database to September 2020. Data analysis was performed using Stata14.0 and GeMTC 0.14.3 softwares.

Results: A total of 40 RCTs involving 3215 patients with KOA were retrieved. Network meta-analysis revealed that the fire needle was superior to western medicine, electro-acupuncture, conventional acupuncture, warm needle and sham acupuncture; warm needle was better than conventional acupuncture and western medicine whereas electro-acupuncture was better than conventional acupuncture in improving pain scores in the Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC). Moreover, we found that fire needle and warm needle more effectively improved WOMAC stiffness scores than western medicine and sham moxibustion, whereas electro-acupuncture was superior to western medicine and sham moxibustion in improving WOMAC stiffness scores. Further analysis revealed that fire needle, warm needle and electro-acupuncture were more effective in improving WOMAC joint function scores than conventional acupuncture and western medicine. The fire needle was superior to conventional acupuncture and sham acupuncture, whereas electro-acupuncture was better than western medicine, conventional acupuncture and sham acupuncture in improving visual analogue scale scores.

Conclusion: This study shows that fire needle is superior to warm needle and electro-acupuncture, whereas warm needle and electro-acupuncture are better than conventional acupuncture, western medicine, sham moxibustion and sham acupuncture.

Keywords: knee osteoarthritis, acupuncture, moxibustion, randomized controlled trials, network meta-analysis

Introduction

Knee osteoarthritis (KOA) is a common disease associated with knee joint degeneration among the elderly. The disease has seriously negative effects on the quality of life of patients. Moreover, it is one of the main diseases leading to knee dysfunction and disability among the elderly people.^{1,2} Globally, KOA is the 11th

Correspondence: Wei Liu
No. 88, Chang Ling Road, Li Qi Zhuang
Jie, Xi Qing District, Tianjin, People's
Republic of China
Email fengshiliuwei@163.com

leading cause of disability, affecting about 3.8% of the world's population.³ With the progressively increasing aging population in China, the incidence of KOA has been on the rise, reaching about 85% among those aged over 65 years.⁴ The development of KOA is associated with a variety of factors, including age, sex, aging, trauma, obesity, inflammation, occupation, activity, metabolism, and heredity among others.⁵ Non-steroidal anti-inflammatory drugs (NSAIDs) are recommended for the treatment of early and middle-term KOA.⁶ Although their analgesic effects are very good, patients often develop pains after drug withdrawal. Cases of gastrointestinal discomfort, liver and kidney function damage as well as other adverse reactions have also been reported.⁷ Acupuncture and moxibustion have shown good therapeutic effects on KOA with few adverse reactions,^{8,9} and have been adopted in China's Guidelines for the Diagnosis and Treatment of Osteoarthritis⁷ and the Guidelines of the American Academy of Orthopaedic Surgeons.¹⁰ There are many types of acupuncture and moxibustion treatments, with varying clinical effects. Direct comparisons of the curative effects of different acupuncture and moxibustion therapies have not been done so far. Therefore, we used network meta-analysis to compare the efficacy of different types of acupuncture and moxibustion therapies in KOA patients to provide a basis for selection of optimal acupuncture and moxibustion therapies in the clinical treatment of KOA.

Information and Methodology

Research Registration

The network meta-analysis research protocol was registered on the PROSPERO at <https://www.crd.york.ac.uk/prospero/#recordDetails>; Registration number: CRD42020203602. This network meta-analysis was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses for NMA guidelines. See supplementary materials (Table S1).

Inclusion and Exclusion Criteria

- i. Study type: randomized controlled trials (RCTs), not limited to blinding method, but limited to Chinese and English languages.
- ii. Study participants: KOA patients diagnosed based on definitive diagnostic criteria, gender and age were unlimited.
- iii. Interventions: treatment groups involved different acupuncture therapies, including conventional acupuncture

alone, warm needle, electro-acupuncture, fire needle, blood-letting puncture, moxibustion, auricular acupuncture, auricular point sticking, acupoint catgut embedding and acupoint injection, control groups comprised of treatments such as western medicine, sham acupuncture and sham moxibustion; or a comparison between different acupuncture types.

iv. Outcome indicators: a. pain, stiffness, and joint function scores based on the Western Ontario and McMaster University Osteoarthritis Index (WOMAC); b. Visual Analogue Scale (VAS); c. Adverse events.

v. Exclusion criteria: a. Studies in which participants did not conform to the inclusion criteria, such as patients with other arthritis; b. studies without clear diagnostic criteria; c. Studies without any of the above outcome indicators; d. Studies using acupuncture combination therapy, such as acupuncture combined with moxibustion, acupuncture combined with auricular acupuncture treatment; e. Studies using traditional Chinese medicine treatments in both groups, such as cupping and Chinese medicine compounds; f. For repeated publications, studies with the most complete data were selected; g. Abstracts or articles without specific data on relevant indicators, and which could not be obtained from the corresponding authors.

Search Strategy

Published RCTs on the application of acupuncture and moxibustion in the treatment of KOA were searched in PubMed, Cochrane Library, EMBASE, Web of Science, CNKI, VIP, Wanfang and China Biomedical Literature Databases. Chinese search terms were “zhen ci” (acupuncture), “dian zhen” (electro-acupuncture), “wen zhen jiu” (warm needle), “huo zhen” (fire needle), “ci luo” (blood-letting puncture), “ai jiu” (moxibustion), “er xue tie ya” (auricular acupoint sticking), “er zhen” (auricular acupuncture), “xue wei mai xian” (acupoint catgut embedding), “xue wei zhu she” (acupoint injection), “xi gu guan jie yan” (knee osteoarthritis). English search terms were “acupuncture”, “electro-acupuncture”, “warm needle”, “fire needle”, “blood-letting puncture”, “moxibustion”, “auricular application pressure”, “auricular needle”, “acupoint catgut embedding”, “acupoint injection”, “knee osteoarthritis”, “KOA”. PubMed database retrieval strategies are shown in Table 1.

Studies Screening and Data Extraction

Study screening and data extraction, as well as cross-checking, were independently performed by two

Table I Retrieval Strategy of Studies from the PubMed Database

Number	Search Terms
#1	Acupuncture [MeSH]
#2	Acupuncture [Title/Abstract]
#3	Pharmacopuncture [Title/Abstract]
#4	Electro-acupuncture [Title/Abstract]
#5	Warm needle [Title/Abstract]
#6	Fire needle [Title/Abstract]
#7	Blood-letting puncture [Title/Abstract]
#8	Moxibustion [MeSH]
#9	Moxibustion [Title/Abstract]
#10	Auricular application pressure [Title/Abstract]
#11	Auricular needle [Title/Abstract]
#12	Acupoint catgut embedding [Title/Abstract]
#13	Acupoint injection [Title/Abstract]
#14	#1OR#2OR#3OR#4OR#5OR#6OR#7OR#8OR#9OR#10OR#11OR#12OR#13
#15	Osteoarthritis, knee [MeSH]
#16	Osteoarthritis, knee [Title/Abstract]
#17	Knee osteoarthritis [Title/Abstract]
#18	Knee osteoarthritides [Title/Abstract]
#19	Osteoarthritis of knee [Title/Abstract]
#20	Osteoarthritis of the knee [Title/Abstract]
#21	KOA [Title/Abstract]
#22	#15OR#16OR#17OR#18OR#19OR#20OR#21
#23	#14AND#22

researchers. In case of disagreements, a third researcher was involved to reach a consensus. The following information was obtained: name of first author, publication year, KOA diagnostic criteria, sample size, gender, age, course of disease, study type, intervention, treatment course, and outcome indicators.

Risk Assessment of Bias in the Included Studies

The Cochrane System Evaluation Manual version 5.1.0 RCT bias risk assessment tool was used to evaluate the quality of the included studies. This was done through random sequence generation, allocation concealment, participant and personnel blinding, outcome assessment blinding, incomplete outcome data, selective reporting, and other bias items. Two researchers graded the above contents as “low risk”, “high risk” and “unclear”, and cross-checked the obtained results. A third researcher was consulted if there were any disagreements. Finally, a bias risk diagram was drawn using RevMan5.3 software.

Statistical Analysis

Stata 14.0 software was used to draw an evidence network diagram to show comparisons of the intervention measures

for each outcome indicator. For continuous variables, if the unit or tool of the measurement index was the same, the mean difference (MD) was used for analysis; if the measurement tools or units were inconsistent, the standardized mean difference (SMD) was used for analysis. Chi-square test was used to directly compare heterogeneity between research results, and I^2 was used to determine level of heterogeneity. If results of the included studies showed no statistical heterogeneity ($I^2 < 50\%$, $P > 0.1$), a meta-analysis using the fixed effect model. If heterogeneity was found, the reasons for heterogeneity were further analyzed. If there was no obvious clinical heterogeneity or methodological heterogeneity, the random effect model was used for meta-analysis. Small sample effects or publication bias were detected using comparative corrected funnel plots. The GEMTC 0.14.3 software was used for network meta-analysis based on the Markov Chain Monte Carlo (MCMC) fitting consistent model under the Bayesian framework. Four chains were used for simulation, and the number of iterations was set at 50,000. The potential scale reduction factor (PSRF) was estimated and deduced under the assumption that MCMC reached a stable convergence state. The stability and consistency of results were evaluated using the MCMC fitted inconsistency model.

Results

Study Retrieval Results

A total of 6290 relevant studies were retrieved. After primary screening and re-screening, 40 RCTs^{11–50} involving 3215 patients were finally included in the study. The screening process of the included studies is shown in Figure 1.

Basic Features of the Included Studies

Among the 40 RCTs,^{11–50} 6, 17, 16, 12, 9, 1, 18, 1, 1 and 1 RCTs involved the application of moxibustion, electroacupuncture, warm acupuncture, conventional acupuncture, fire needle, acupoint embedding, western medicine, placebo, sham acupuncture, and sham moxibustion, respectively. Among these studies, there were 2 three-arm trials^{12,38} and 38 double-arm trials;^{11,13–37,39–50} In addition, 23 trials^{13,14,18,20–22,24–28,31–34,36,40–43,46,49,50} reported WOMAC pain scores, 21^{13,14,18,20–22,24–28,31–}

^{34,36,37,42,43,46,49} reported WOMAC stiffness scores, 21–13,14,18,20–22,25–28,31–34,36,37,40–42,46,49 reported WOMAC joint function scores, 25^{11–13,15–17,19,22,23,25,29,30,34–36,38,40,42–49} reported VAS scores while 12–11,20,24,31,32,37,39–41,44,48,49 reported adverse events. Basic characteristics of the included studies are shown in Table 2, whereas the characteristics of the interventions are shown in Table 3.

Risk of Bias Assessment Results of the Included Studies

i. Random sequence generation: Nineteen studies^{12,14,18,20,22,25,26,28,29,32,36,37,39,42–44,46,48,49} used a table of random numbers, seven^{13,16,21,35,40,41,50} used computer-generated random numbers, one¹⁷ used a coin toss for randomization, one¹⁹ used random cards, while the remaining twelve^{11,15,23,24,27,30,31,33,34,38,45,47} only mentioned the word “random”; ii. Allocation concealment: Three studies^{13,28,36} used




Figure 1 Flowchart of study screening.

Table 2 Basic Features of the Included Studies

Included Studies	Diagnostic Criteria	Sample Size (T/C)	Sex (Men/Women)	Age (Year)	Course of Disease (Year)
Zhang 2011 ¹¹	ACR	30/30	22/38	58.2	–
Zhou 2014 ¹²	ACR	39/44/22	T:14/25 C:18/36 C2:5/17	T:67±10 C:180±10 C2:66±12	T:3 C:1.202 C2:1.53
Zhou 2017 ¹³	ACR	30/30	T:14/16 C:13/17	T:59.07±7.89 C:60.60±8.27	T:5.73±2.85 C:5.50±2.54
Ren 2011 ¹⁴	ACR	31/28	T:11/20 C:8/20	T:64.03±7.24 C:62.57±8.12	T:6.82±6.60 C:7.15±7.72
Zhou 2015 ¹⁵	ACR	40/40	T:17/23 C:19/21	T:54.6±5.3 C:53.8±7.6	T:17.2±2.2 (month) C:15.6±3.0 (month)
Zhang 2018 ¹⁶	ACR	39/39	T:13/26 C:15/24	T:61.36±2.24 C:62.08±2.46	T:4.77±0.12 C:4.68±0.28
Liu 2020 ¹⁷	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	41/37	T:2/39 C:3/34	T:61.72±8.05 C:60.98±7.56	T:2.13±0.98 C:2.53±0.83
Guo 2016 ¹⁸	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	45/45	T:19/26 C:17/28	T:56.00±7.25 C:57.17±6.96	T:33.75±14.11 (month) C:32.35±13.72 (month)
Wu 2015 ¹⁹	ACR	47/48	–	T:58.75±1.21 C:60.02±2.17	T:18±2 (month) C:19±3 (month)
Qiu 2006 ²⁰	ACR	30/30	T:5/25 C:4/26	T:56.07 C:55.37	T:8.95 C:9.55
Gang 2016 ²¹	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	43/45	T:19/24 C:22/23	T:54±8 C:54±8	T:1.1±0.6 C:1.2±0.6
Gao 2011 ²²	ACR	34/35	T:13/21 C:15/20	T:57.68±8.67 C:58.57±8.89	T:37.35±10.83 (month) C:38.86±12.12 (month)
Wang 2017 ²³	<i>Diagnostic criteria for blood stasis type of KOA</i>	48/47	T:25/23 C:24/23	T:56.15±7.36 C:56.19±7.33	–
Yin 2017 ²⁴	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	60/60	–	–	–
Ju 2017 ²⁵	ACR	30/30	T:6/24 C:7/23	T:60±10 C:64±6	T:29.89±29.74 (month) C:32.74±31.43 (month)
Wu 2012 ²⁶	ACR	30/30	T:11/19 C:13/17	T:60.63±6.44 C:59.87±6.77	T:3.47±1.27 C:3.23±1.48
Chen 2018 ²⁷	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	45/45	T:31/14 C:28/17	T:65.51±3.26 C:66.36±3.08	T:3.56±1.87 C:3.49±1.76
Liu 2014 ²⁸	ACR	30/30	T:10/20 C:12/18	T:58.07±8.76 C:59.47±7.92	–
Tan 2016 ²⁹	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	40/40	T:18/22 C:17/23	T:52.72±5.83 C:51.93±6.18	T:1–10 C:1–12

(Continued)

Table 2 (Continued).

Included Studies	Diagnostic Criteria	Sample Size (T/C)	Sex (Men/Women)	Age (Year)	Course of Disease (Year)
Ma 2015 ³⁰	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	54/54	T:22/32 C:19/35	T:53.2±7.5 C:56.3±8.6	T:3.2±1.4 C:4.4±1.5
Chen 2012 ³¹	ACR	30/30	T:8/22 C:9/21	T:59±12 C:59±16	T:50.23±47.50 (month) C:62.20±45.67 (month)
Jiang 2013 ³²	ACR	42/44	T:18/24 C:21/23	T:65.7±3.9 C:67.7±4.5	T:3.7±0.9 C:4.2±1.5
Ren 2018 ³³	ACR	54/54	T:23/31 C:25/29	T:67.1±4.6 C:68.7±5.2	T:4.9±1.5 C:4.6±1.7
Zhang 2016 ³⁴	<i>Standard of diagnosis and curative effect of disease and syndrome in TCM</i>	30/30	T:18/12 C:19/11	T:56.0±2.9 C:55.9±2.8	T:5.6±0.5 C:5.7±0.5
Pan 2020 ³⁵	ACR	35/35	T:11/24 C:9/26	T:65.086 C:64.2	T:3.34±1.72 C:3.03±1.33
Ji 2012 ³⁶	<i>Guiding principles of clinical research on new chinese medicine</i>	30/30	T:12/18 C:11/19	T:56.73±10.08 C:58.77±7.98	T:38.17±23.99 (month) C:39.23±25.54 (month)
Shen 2017 ³⁷	<i>Standard of diagnosis and curative effect of disease and syndrome in TCM</i>	100/100	T:54/46 C:51/49	T:66.12±2.09 C:66.51±2.12	T:3.31±0.23 C:3.35±0.12
Miao 2014 ³⁸	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	35/35/35	T:5/30 C1:9/26 C2:7/28	T:57.5±9.7 C1:56.3±8.9 C2:60.4±10.5	T:5.6±3.8 C1:6.4±4.2 C2:6.8±3.3
Zhang 2016 ³⁹	<i>Guidelines for the diagnosis and treatment of arthritis</i>	28/28	T:11/17 C:12/16	T:62.3±5.1 C:61.8±4.7	T:8.7±3.6 C:8.4±3.4
Lin 2018 ⁴⁰	The National Institute for Health and Clinical Excellence Guidelines 2014 Edition criteria	21/21	T:4/17 C:1/20	T:59.5±7.5 C:60.0±7.3	T:60.0±45.9(month) C:63.1±45.4(month)
Zhao 2014 ⁴¹	ACR	55/55	T:16/39 C:21/34	T:65.80±7.45 C:64.55±8.38	–
Zhang 2013 ⁴²	ACR	33/34	T:13/20 C:14/20	T:57±8 C:58±9	T:38±10 (month) C:38±11 (month)
Zheng 2016 ⁴³	ACR	35/35	T:16/19 C:15/20	T:62.39±8.004 C:61.41±8.203	T:135.97±74.068 (month) C:128.03±70.194 (month)
Lu 2011 ⁴⁴	ACR	27/27	T:6/21 C:8/19	T:54.11±9.46 C:60.81±10.09	T:3.83±5.78 C:3.69±2.98
Li 2020 ⁴⁵	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	36/36	T:12/24 C:14/22	T:58.54±8.41 C:56.52±7.91	–
Tang 2017 ⁴⁶	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	30/30	T:8/22 C:9/21	T:59.64 C:60.40	T:2.62 C:2.73
Song 2020 ⁴⁷	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	30/30	T:12/18 C:14/16	T:53.83±5.37 C:53.47±7.34	T:37.90±16.01 (month) C:34.67±19.78 (month)
Lin 2012 ⁴⁸	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	30/30	T:14/16 C:16/14	T:48.47±11.95 C:50.07±9.7	T:4.97±7.1 (month) C:9.83±17.74 (month)

(Continued)

Table 2 (Continued).

Included Studies	Diagnostic Criteria	Sample Size (T/C)	Sex (Men/Women)	Age (Year)	Course of Disease (Year)
He 2018 ⁴⁹	ACR	57/55	T:26/31 C:22/33	T:56±5 C:58±5	T:73.28±29.24 (month) C:71.09±27.96 (month)
Fan 2016 ⁵⁰	<i>Guidelines for the diagnosis and treatment of osteoarthritis</i>	54/54	T:33/21 C:30/24	T:58±6.2 C:56±8.4	T:14.8±8.8 (month) C:12.7±7.3 (month)

Abbreviations: ACR, American College of Rheumatology; T, treatment group; C, control group; -, not mentioned.

sealed opaque envelopes, one study⁴⁰ used identical and ordered drug containers, one study⁴¹ used alphabetic codes, while the remaining thirty-five studies did not mention allocation concealment; iii. Blinding of patients and experimentalists: Four studies^{13,14,40,41} blinded patients and experimentalists; iv. Blinding of outcome evaluators: Five studies^{13,14,36,40,41} blinded outcome evaluators; v. Incomplete result data, selective reporting, other bias: The results of 40 studies^{11–50} were all complete, without selective reporting and other bias. Results of bias risk assessment are shown in Figure 2.

Directly Compared Meta-Analysis Results Visual Analogue Scale

Results of the meta-analysis showed that the VAS score of the moxibustion group was significantly better than that of the western medicine group (4 RCTs; SMD 0.624, 95% CI 1.239 to 0.009; $I^2 = 82.10\%$, $P = 0.001$), whereas the VAS score of the western medicine group was significantly better than that of the electro-acupuncture group (6 RCTs; SMD 1.201, 95% CI 0.169 to 2.223; $I^2 = 95.80\%$, $P < 0.00001$). The VAS score of conventional acupuncture group was significantly better than that of warm needle group (3 RCTs; SMD 2.974, 95% CI 0.798 to 5.150; $I^2 = 97.60\%$, $P < 0.00001$) and fire needle group (3 RCTs; SMD 1.22, 95% CI 0.681 to 1.760; $I^2 = 67.5\%$, $P = 0.046$), and the VAS score of warm needle group was substantially better than that of fire needle group (3 RCTs; SMD 1.432, 95% CI 0.669 to 2.194; $I^2 = 80.90\%$, $P = 0.005$) ($P < 0.05$). Descriptive analysis results showed that VAS score of electro-acupuncture group was significantly better than that of conventional acupuncture group, and VAS score of conventional acupuncture group was significantly better than that of sham acupuncture group. The VAS score of western medicine group was significantly better than that

of fire needle group ($P < 0.05$). There was no significant difference among other interventions in the aforementioned scores ($P > 0.05$). See supplementary materials (Table S2).

WOMAC Pain Score

Meta-analysis results showed that the WOMAC pain score of the western medicine group was significantly better than that of the warm needle group (4 RCTs; SMD 0.664, 95% CI 0.306 to 1.022; $I^2 = 62.10\%$, $P = 0.048$), while the WOMAC pain score of the warm needle group was significantly better than that of the fire needle group (3 RCTs; SMD 0.956, 95% CI 0.139 to 1.774; $I^2 = 88.10\%$, $P < 0.00001$) ($P < 0.05$). Descriptive analysis results showed that WOMAC pain score of electro-acupuncture group was significantly better than that of conventional acupuncture group, and WOMAC pain score of conventional acupuncture group was significantly better than that of warm needle group and that of the fire needle group ($P < 0.05$). There was no differences among other interventions in the aforementioned scores, ($P > 0.05$). See supplementary materials (Table S1).

WOMAC Joint Function Score

Meta-analysis results showed that the WOMAC joint function score of the western medicine group was significantly better than that of the electro-acupuncture group (4 RCTs; SMD 0.419, 95% CI 0.209 to 0.629; $I^2 = 20.90\%$, $P = 0.285$) and that of the warm needle group (4 RCTs; SMD 0.646, 95% CI 0.201 to 1.091; $I^2 = 75.40\%$, $P = 0.007$) ($P < 0.05$). Descriptive analysis results showed that WOMAC joint function score of moxibustion group was significantly better than that of western medicine group, and WOMAC joint function score of western medicine group was significantly better than that of conventional acupuncture group, and WOMAC joint function score of electro-acupuncture group

Table 3 Characteristics of Interventions

Included Studies	Study Type	Interventions		Course of Treatment (Week)	Outcome Indicators
		Treatment Group	Control Group		
Zhang 2011 ¹¹	Double-arm	Moxibustion	celecoxib 200 mg, 1/d	6	d,e
Zhou 2014 ¹²	Three-arm	Moxibustion	C1:electro-acupuncture C2:celecoxib 200 mg, 1/d	4	d
Zhou 2017 ¹³	Double-arm	Moxibustion	Diclofenac sodium gel	4	a,b,c,d
Ren 2011 ¹⁴	Double-arm	Moxibustion	Sham moxibustion	6	a,b,c
Zhou 2015 ¹⁵	Double-arm	Electro-acupuncture	Diclofenac sodium sustained release capsules 50 mg, 1/d	4	d
Zhang 2018 ¹⁶	Double-arm	Electro-acupuncture	Meloxicam dispersible tablets 7.5 mg, 1/d	2	d
Liu 2020 ¹⁷	Double-arm	Electro-acupuncture	Conventional acupuncture	4	d
Guo 2016 ¹⁸	Double-arm	Electro-acupuncture	Diclofenac sodium double release enteric capsules 75 mg, 1/d	3	a,b,c
Wu 2015 ¹⁹	Double-arm	Electro-acupuncture	Votalin emulsion	4	d
Qiu 2006 ²⁰	Double-arm	Electro-acupuncture	Futarin sustained-release tablets 75 mg, 1/d	4	a,b,c,e
Gang 2016 ²¹	Double-arm	Electro-acupuncture	Meloxicam tablets 7.5 mg, 1/d	6	a,b,c
Gao 2011 ²²	Double-arm	Electro-acupuncture	Warm needle	8	a,b,c,d
Wang 2017 ²³	Double-arm	Electro-acupuncture	Warm needle	3	d
Yin 2017 ²⁴	Double-arm	Electro-acupuncture	Glucosamine 240 mg, 1/d	8	a,b,e
Ju 2017 ²⁵	Double-arm	Electro-acupuncture	Celecoxib capsules 200 mg, 1/d	2	a,b,c,d
Wu 2012 ²⁶	Double-arm	Electro-acupuncture	Conventional acupuncture	4	a,b,c
Chen 2018 ²⁷	Double-arm	Warm needle	Glucosamine sulfate capsules 314 mg, 2 capsules/ times, 3 times/ d	4	a,b,c
Liu 2014 ²⁸	Double-arm	Warm needle	Electro-acupuncture	4	a,b,c
Tan 2016 ²⁹	Double-arm	Warm needle	Conventional acupuncture	3	d
Ma 2015 ³⁰	Double-arm	Warm needle	Conventional acupuncture	4	d
Chen 2012 ³¹	Double-arm	Warm needle	Ibuprofen sustained release capsules 300 mg, 2 times/ d	3	a,b,c,e
Jiang 2013 ³²	Double-arm	Warm needle	Glucosamine sulfate tablets 314 mg, 2 tablets/times, 3 times/d	8	a,b,c,e
Ren 2018 ³³	Double-arm	Warm needle	Glucosamine hydrochloride tablets 240 mg, 2 tablets/ times, 3 times/d	20	a,b,c
Zhang 2016 ³⁴	Double-arm	Warm needle	Conventional acupuncture	4	a,b,c,d
Pan 2020 ³⁵	Double-arm	Acupoint catgut embedding	Conventional acupuncture	3	d
Ji 2012 ³⁶	Double-arm	Electro-acupuncture	Warm needle	8	a,b,c,d

(Continued)

Table 3 (Continued).

Included Studies	Study Type	Interventions		Course of Treatment (Week)	Outcome Indicators
		Treatment Group	Control Group		
Shen 2017 ³⁷	Double-arm	Conventional acupuncture	Diclofenac sodium emulsion	4	b,c,e
Miao 2014 ³⁸	Three-arm	Moxibustion	C1: electro-acupuncture C2: celecoxib capsules 200 mg, 1/d	4	d
Zhang 2016 ³⁹	Double-arm	Conventional acupuncture	Warm needle	4	e
Lin 2018 ⁴⁰	Double-arm	Conventional acupuncture	Sham acupuncture	26	a,b,d,e
Zhao 2014 ⁴¹	Double-arm	Moxibustion	Sham moxibustion	6	a,c,e
Zhang 2013 ⁴²	Double-arm	Fire needle	Warm needle	4	a,b,c,d
Guo 2016 ⁴³	Double-arm	Fire needle	Conventional acupuncture	3	a,b,d
Lu 2011 ⁴⁴	Double-arm	Fire needle	Warm needle	4	d,e
Li 2020 ⁴⁵	Double-arm	Fire needle	Conventional acupuncture	24	d
Tang 2017 ⁴⁶	Double-arm	Fire needle	Warm needle	4	a,b,c,d
Song 2020 ⁴⁷	Double-arm	Fire needle	Celecoxib capsules 200 mg, 1/d; diclofenac sodium diethylamine emulsion agent 200 mg, 3 times/d	3	d
Lin 2012 ⁴⁸	Double-arm	Fire needle	Conventional acupuncture	4	d,e
He 2018 ⁴⁹	Double-arm	Electro-acupuncture	Fire needle	4	a,b,c,d,e
Fan 2016 ⁵⁰	Double-arm	Fire needle	Warm needle	4	a

Notes: a, WOMAC pain scores; b, WOMAC stiffness scores; c, WOMAC joint function scores; d, VAS scores; e, Adverse events.

was significantly better than that of conventional acupuncture group, and WOMAC joint function score of conventional acupuncture group was significantly better than that of warm needle group, and the differences were statistically significant ($P<0.05$). There was no statistical significance in the comparison of other interventions ($P>0.05$). See supplementary materials (Table S1).

WOMAC Stiffness Score

Meta-analysis results showed that the WOMAC stiffness score of the western medicine group was significantly better than that of the electro-acupuncture group (6 RCTs; SMD 1.201, 95% CI 0.169 to 2.223; $I^2=95.80\%$, $P<0.00001$), while the WOMAC stiffness score of the electro-acupuncture group was significantly better than that of the warm needle (3 RCTs; SMD 0.671, 95% CI 0.377 to 0.965; $I^2=1.30\%$, $P=0.363$) ($P<0.05$). Descriptive analysis results showed that WOMAC stiffness score of moxibustion group was

significantly better than that of sham moxibustion group, and WOMAC stiffness score of western medicine group and electro-acupuncture group were significantly better than that of conventional acupuncture group, respectively, and WOMAC stiffness score of conventional acupuncture group was significantly better than that of warm needle group and that of fire needle group, respectively ($P<0.05$). There was no significant difference in the aforementioned scores among other interventions ($P>0.05$). See supplementary materials (Table S1).

Heterogeneity Analysis

In the direct comparison meta-analysis, most results were heterogeneous. Through the analysis of original data, we found that the lack of description of blind methods and allocation concealment in included studies may lead to methodological heterogeneity, at the same time, the inclusion of population, KOA stage and other factors may cause clinical

heterogeneity, but since the original study did not specify these details and the number of included studies was small, further subgroup analysis could not be performed to explore the source of heterogeneity. However, the sensitivity analysis of the study results by a one-by-one exclusion method showed that the results were stable after the exclusion of any study. Therefore, we can ignore this heterogeneity and use a random effects model to analyze the results.

Comparison Results of Network Meta-Analysis

Evidence Network Diagram

Twenty-three studies^{13,14,18,20–22,24–28,31–34,36,40–43,46,49,50} reported WOMAC pain scores, forming five closed loops, that is electro-acupuncture-conventional acupuncture-warm needle, western medicine-electro-acupuncture-warm needle, electro-acupuncture-conventional acupuncture-fire needle, fire needle-conventional acupuncture-warm needle and electro-acupuncture-warm needle-fire needle; Twenty-one studies^{13,14,18,20–22,24–28,31–34,36,37,42,43,46,49} reported WOMAC stiffness scores, forming seven closed loops, that is western medicine-electro-acupuncture-conventional acupuncture, electro-acupuncture-conventional acupuncture-fire needle, electro-acupuncture-conventional acupuncture-warm needle, warm needle-fire needle-conventional acupuncture, western medicine-warm needle-conventional acupuncture, western medicine-electro-acupuncture-warm needle, electro-acupuncture-warm needle-fire needle; Twenty-one studies^{13,14,18,20–22,25–28,31–34,36,37,40–42,46,49} reported WOMAC joint function scores, forming five closed loops, that is electro-acupuncture-conventional acupuncture-warm needle, western medicine-electro-acupuncture-conventional acupuncture, western medicine-conventional acupuncture-warm needle, western medicine-electro-acupuncture-warm needle, electro-acupuncture-warm needle-fire needle; Twenty-five studies^{11–13,15–17,19,22,23,25,29,30,34–36,38,40,42–49} reported VAS scores, forming six closed loops, that is moxibustion-western medicine-electro-acupuncture, electro-acupuncture-conventional acupuncture-warm needle, electro-acupuncture-conventional acupuncture-fire needle, fire needle-conventional acupuncture-warm needle, western medicine -electro-acupuncture-fire needle, electro-acupuncture-warm needle-fire needle. The thicker the line between the two measures, the larger the number of pairable studies between the two measures, the larger the node, and the larger the study sample size of

the intervention involved (Figure 3Figure 4Figure 5Figure –6).

Network Meta-Analysis of WOMAC Pain Scores

WOMAC pain scores were reported in 23 studies.^{13,14,18,20–22,24–28,31–34,36,40–43,46,49,50} Convergence assessment revealed that PSRF values tended to 1 and results of the incongruity model were similar to those of the congruity model, indicating that the stability and consistency of the indicators were good, therefore, the MCMC congruity model was used for network meta-analysis of WOMAC pain scores. It was found that the warm needle was superior to conventional acupuncture and western medicine; fire needle was superior to western medicine, electro-acupuncture, conventional acupuncture, warm needle and sham acupuncture while electro-acupuncture was superior to conventional acupuncture. WOMAC pain scores among other different acupuncture treatments showed no significant statistical difference (Table 4). Treatment ranking of WOMAC pain scores were: fire needle > warm needle > electro-acupuncture > western medicine > moxibustion > conventional acupuncture > sham acupuncture > sham moxibustion (Table 5).

Network Meta-Analysis of WOMAC Stiffness Scores

WOMAC stiffness scores were reported in 21 studies.^{13,14,18,20–22,24–28,31–34,36,37,42,43,46,49} Convergence assessment showed that PSRF values tended to 1, and the convergence effect was satisfactory, the results of the incongruity model were similar to those of the congruity model, without significant changes, indicating that the stability and consistency of the indicators were good, therefore, the MCMC congruity model was used for network meta-analysis of WOMAC stiffness scores. It was found that electro-acupuncture was superior to western medicine and sham moxibustion, fire needle and warm needle were superior to western medicine and sham moxibustion, the difference was statistically significant. There was no statistically significant difference in improving WOMAC stiffness scores between other acupuncture treatments (Table 6). In the ranking of conformance model analysis, fire needle > warm needle > electro-acupuncture > conventional acupuncture > western medicine > moxibustion > sham moxibustion (Table 7).

Network Meta-Analysis of WOMAC Joint Function Scores

WOMAC joint function scores were reported in 21 studies.^{13,14,18,20–22,25–28,31–34,36,37,40–42,46,49} According to

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Chen 2012 [31]	?	?	?	?	?	?	?
Chen 2018 [27]	?	?	?	?	?	?	?
Fan 2016 [50]	?	?	?	?	?	?	?
Gang 2016 [21]	?	?	?	?	?	?	?
Gao 2011 [22]	?	?	?	?	?	?	?
Guo 2016 [43]	?	?	?	?	?	?	?
He 2018 [49]	?	?	?	?	?	?	?
Ji 2012 [36]	?	?	?	?	?	?	?
Jiang 2013 [32]	?	?	?	?	?	?	?
Ju 2017 [25]	?	?	?	?	?	?	?
Li 2020 [45]	?	?	?	?	?	?	?
Lin 2012 [48]	?	?	?	?	?	?	?
Lin 2018 [40]	?	?	?	?	?	?	?
Liu 2014 [28]	?	?	?	?	?	?	?
Liu 2020 [17]	?	?	?	?	?	?	?
Lu 2011 [44]	?	?	?	?	?	?	?
Ma 2015 [30]	?	?	?	?	?	?	?
Miao 2014 [38]	?	?	?	?	?	?	?
Pan 2020 [35]	?	?	?	?	?	?	?
Qiu 2006 [20]	?	?	?	?	?	?	?
Ren 2011 [14]	?	?	?	?	?	?	?
Ren 2018 [33]	?	?	?	?	?	?	?
Shen 2017 [37]	?	?	?	?	?	?	?
Song 2020 [47]	?	?	?	?	?	?	?
Tan 2016 [29]	?	?	?	?	?	?	?
Tang 2017 [46]	?	?	?	?	?	?	?
Wang 2017 [23]	?	?	?	?	?	?	?
Wu 2012 [26]	?	?	?	?	?	?	?
Wu 2015 [19]	?	?	?	?	?	?	?
Yin 2017 [24]	?	?	?	?	?	?	?
Zhang 2011 [11]	?	?	?	?	?	?	?
Zhang 2013 [42]	?	?	?	?	?	?	?
Zhang 2016 [34]	?	?	?	?	?	?	?
Zhang 2016 [39]	?	?	?	?	?	?	?
Zhang 2018 [18]	?	?	?	?	?	?	?
Zhao 2014 [41]	?	?	?	?	?	?	?
Zhen 2016 [43]	?	?	?	?	?	?	?
Zhou 2014 [12]	?	?	?	?	?	?	?
Zhou 2015 [15]	?	?	?	?	?	?	?
Zhou 2017 [13]	?	?	?	?	?	?	?

Figure 2 Evaluation results of bias risk.

the Monte Carlo simulation iteration, the PSRF value tended to 1, and the convergence effect was good. Results of the incongruity model were similar to those of




Figure 3 Evidence network diagram of WOMAC pain score for different acupuncture treatments for knee osteoarthritis.

Notes: 1-moxibustion; 2-western medicine; 3-electro-acupuncture; 4-sham moxibustion; 5-conventional acupuncture; 6-warm needle; 7-sham acupuncture; 8-fire needle




Figure 4 Evidence network diagram of a WOMAC stiffness score for different acupuncture treatments for KOA.

Notes: 1-moxibustion; 2-western medicine; 3-electro-acupuncture; 4-sham moxibustion; 5-conventional acupuncture; 6-warm needle; 7-fire needle




Figure 5 Evidence network diagram showing WOMAC joint function score for different acupuncture treatments for KOA.
Notes: 1-moxibustion; 2-western medicine; 3-electro-acupuncture; 4-sham moxibustion; 5-conventional acupuncture; 6-warm needle; 7-sham acupuncture; 8-fire needle




Figure 6 Evidence network diagram of VAS score for different acupuncture treatments for KOA.
Notes: 1-moxibustion; 2-western medicine; 3-electro-acupuncture; 4-conventional acupuncture; 5-warm needle; 6-acupoint catgut embedding; 7-sham acupuncture; 8-fire needle

the congruity model, without significant changes, indicating that stability and consistency of the indicators were good, therefore, the MCMC congruity model was used for

Table 4 Results of Network Meta-Analysis of WOMAC Pain Scores of Different Acupuncture Treatments for KOA

Moxibustion	0.34 (-2.88, 3.42)	-0.91 (-4.37, 2.49)	3.81 (-2.72, 10.17)	1.96 (-1.88, 5.89)	-1.26 (-4.76, 2.20)	2.49 (-2.80, 7.89)	-3.03 (-6.81, 0.61)
Western medicine	1.25 (-0.03, 2.51)	-1.25 (-2.51, 0.03)	3.49 (-3.83, 10.45)	1.64 (-0.57, 3.98)	-1.59 (-2.97, -0.21)	2.16 (-2.10, 6.50)	-3.38 (-5.39, -1.49)
Electro-acupuncture	-3.49 (-10.45, 3.83)	-4.71 (-11.78, 2.68)	4.71 (-2.68, 11.78)	2.89 (0.80, 5.04)	-0.34 (-1.69, 0.97)	3.39 (-0.79, 7.63)	-2.12 (-4.00, -0.43)
Conventional acupuncture	-1.64 (-3.98, 0.57)	-2.89 (-5.04, -0.80)	Sham moxibustion	-1.84 (-9.05, 5.82)	-5.08 (-12.20, 2.33)	-1.33 (-9.39, 7.26)	-6.86 (-14.07, 0.68)
Warm needle	1.59 (0.21, 2.97)	0.34 (-0.97, 1.69)	1.84 (-5.82, 9.05)	Conventional acupuncture	-3.23 (-5.29, -1.23)	0.53 (-3.17, 4.21)	-5.02 (-7.38, -2.87)
Sham acupuncture	-2.16 (-6.50, 2.10)	-3.39 (-7.63, 0.79)	5.08 (-2.33, 12.20)	Sham moxibustion	Warm needle	3.75 (-0.38, 8.07)	-1.78 (-3.40, -0.32)
Fire needle	3.38 (1.49, 5.39)	2.12 (0.43, 4.00)	1.33 (-7.26, 9.39)	Fire needle	1.78 (0.32, 3.40)	5.54 (1.31, 9.93)	-5.54 (-9.93, -1.31)

Notes: The above data represent the confidence interval. The bold font indicates that there was a statistically significant difference between the two treatments.

Table 5 Ranking Probability Table of WOMAC Pain Scores

Intervention	Rank 1	Rank 2	Rank 3	Rank 4	Rank 5	Rank 6	Rank 7	Rank 8
Moxibustion	0.01	0.09	0.13	0.25	0.24	0.11	0.13	0.04
Western medicine	0.01	0.04	0.19	0.45	0.31	0.01	0	0
Electro-acupuncture	0	0	0.01	0.06	0.22	0.5	0.21	0.01
Sham moxibustion	0.58	0.12	0.11	0.05	0.03	0.03	0.04	0.03
Conventional acupuncture	0.11	0.39	0.38	0.09	0.02	0	0	0
Warm needle	0	0	0	0.03	0.11	0.31	0.53	0.01
Sham acupuncture	0.29	0.36	0.18	0.08	0.06	0.02	0.02	0.01
Fire needle	0	0	0	0	0	0.01	0.07	0.91

Notes: The bold font represents the probability of ordering the therapy.

network meta-analysis of WOMAC joint function scores. It was found that fire needle, warm needle and electro-acupuncture yielded significantly better results in improving WOMAC joint function scores compared to conventional acupuncture and western medicine. There were no significant difference in the WOMAC joint function scores among the other acupuncture treatments (Table 8). In the ranking of conformance model analysis, fire needle > electro-acupuncture > warm needle > moxibustion > western medicine > conventional acupuncture > sham acupuncture > sham moxibustion (Table 9).

Network Meta-Analysis of VAS Scores

VAS scores were reported in 25 studies.^{11–13,15–17,19,22,23,25,29,30,34–36,38,40,42–49} Convergence assessment showed that PSRF values tended to 1, and the convergence effect was satisfactory. Results of the incongruity model were similar to those of the congruity model, indicating that the stability and consistency of the indicators were good, therefore, the MCMC congruity model was used for network meta-analysis of VAS scores. The results showed that electro-acupuncture was superior to western medicine, conventional acupuncture and sham acupuncture while fire needle was superior to conventional acupuncture and sham acupuncture (Table 10). The probability ranking conducted with the MCMC method revealed that fire needle > electro-acupuncture > moxibustion > warm needle > western medicine > conventional acupuncture > acupoint catgut embedding > sham acupuncture in improving VAS scores (Table 11).

Small Sample Effect Estimation

A comparative correction funnel plot was used to evaluate the outcome of the WOMAC pain scores (Figure 7). The funnel plot was not completely symmetric, suggesting that

there may be some publication bias or small sample effect in the research network.

Adverse Events

Twelve studies^{11,20,24,31,32,37,39–41,44,48,49} reported the occurrence of adverse events (Table 12). Overall, there were only mild adverse reactions but no serious adverse reactions in the treatment of KOA by different acupuncture and moxibustion treatments.

Discussion

According to the first-line management protocol recommended by Osteoarthritis Research Society International (OARSI), conservative treatment (drug therapy and non-drug therapy) plays an important role in the management of osteoarthritis.⁵¹ However, due to adverse reactions which decrease patient compliance, the benefit-risk ratios of such interventions in KOA patients require urgent investigation.^{52,53} Traditional Chinese medicine classifies KOA as “bi zheng” (arthralgia syndrome). In China, acupuncture and moxibustion therapies have been used in the treatment of arthralgia syndrome for thousands of years, and this in ancient period in the Chinese book *Lingshu*. Acupuncture and moxibustion therapies are widely used in clinical practice and have high efficacy and strong safety.^{54,55} Studies have shown that these therapies can improve the pain threshold by promoting the release of analgesic substances in KOA patients.⁵⁶

Herein, the effects of acupuncture and moxibustion therapy on WOMAC pain, stiffness, joint function scores and VAS scores in KOA patients were investigated. Results showed that warm needle was superior to conventional acupuncture and western medicine, fire needle was superior to western medicine, electro-acupuncture, conventional acupuncture, warm needle and sham acupuncture, while electro-acupuncture was superior to conventional acupuncture in

Table 6 Results of Network Meta-Analysis of WOMAC Stiffness Scores of Different Acupuncture Treatments for KOA

Moxibustion	0.22 (-2.67, 3.20)	-1.06 (-4.21, 2.10)	8.44 (-0.15, 16.98)	0.10 (-3.22, 3.49)	-1.32 (-4.44, 1.89)	-1.70 (-5.12, 1.68)
-0.22 (-3.20, 2.67)	Western medicine	-1.29 (-2.39, -0.16)	8.27 (-0.97, 17.20)	-0.13 (-1.77, 1.52)	-1.54 (-2.71, -0.31)	-1.95 (-3.70, -0.15)
1.06 (-2.10, 4.21)	1.29 (0.16, 2.39)	Electro-acupuncture	9.57 (0.21, 18.64)	1.16 (-0.48, 2.76)	-0.24 (-1.43, 0.91)	-0.66 (-2.33, 1.03)
-8.44 (-16.98, 0.15)	-8.27 (-17.20, 0.97)	-9.57 (-18.64, -0.21)	Sham moxibustion	-8.35 (-17.51, 0.99)	-9.78 (-18.86, -0.43)	-10.21 (-19.34, -0.70)
-0.10 (-3.49, 3.22)	0.13 (-1.52, 1.77)	-1.16 (-2.76, 0.48)	8.35 (-0.99, 17.51)	Conventional acupuncture	-1.39 (-3.07, 0.23)	-1.82 (-3.68, 0.11)
1.32 (-1.89, 4.44)	1.54 (0.31, 2.71)	0.24 (-0.91, 1.43)	9.78 (0.43, 18.86)	1.39 (-0.23, 3.07)	Warm needle	-0.42 (-1.94, 1.17)
1.70 (-1.68, 5.12)	1.95 (0.15, 3.70)	0.66 (-1.03, 2.33)	10.21 (0.70, 19.34)	1.82 (-0.11, 3.68)	0.42 (-1.17, 1.94)	Fire needle

Notes: The above data represent the confidence interval. The bold font indicates that there was a statistically significant difference between the two treatments.

Table 7 Ranking Probability Table of WOMAC Stiffness Scores

Intervention	Rank 1	Rank 2	Rank 3	Rank 4	Rank 5	Rank 6	Rank 7
Moxibustion	0.01	0.36	0.16	0.2	0.08	0.07	0.11
Western medicine	0.02	0.31	0.47	0.18	0.01	0	0
Electro-acupuncture	0	0	0.03	0.17	0.46	0.25	0.09
Sham moxibustion	0.95	0.01	0.01	0.01	0	0.01	0.01
Conventional acupuncture	0.02	0.3	0.31	0.29	0.05	0.02	0.01
Warm needle	0	0	0.01	0.09	0.27	0.44	0.19
Fire needle	0	0.01	0.01	0.05	0.12	0.21	0.59

Notes: The bold font represents the probability of ordering the therapy.

Table 8 Results of Network Meta-Analysis of WOMAC Joint Function Scores of Different Acupuncture Treatments for KOA

Moxibustion	4.62 (-5.64, 15.13)	-1.34 (-12.71, 9.91)	4.77 (-7.97, 16.18)	5.87 (-6.31, 18.18)	-0.73 (-12.02, 10.72)	7.49 (-8.94, 24.30)	-3.96 (-16.47, 8.98)
Western medicine	6.00 (1.98, 10.38)	-6.00 (-10.38, -1.98)	0.13 (-16.45, 15.20)	1.22 (-5.20, 7.31)	-5.38 (-9.77, -1.32)	2.85 (-9.80, 15.53)	-8.54 (-15.74, -1.59)
Electro-acupuncture	-0.13 (-15.20, 16.45)	-6.12 (-21.85, 10.79)	6.12 (-10.79, 21.85)	7.21 (1.00, 13.55)	0.59 (-3.53, 4.88)	8.91 (-3.79, 21.62)	-2.56 (-9.23, 3.89)
Sham moxibustion	-1.22 (-7.31, 5.20)	-7.21 (-13.55, -1.00)	Sham moxibustion	1.04 (-15.13, 18.93)	-5.55 (-21.20, 11.29)	2.71 (-16.74, 23.65)	-8.72 (-25.39, 9.18)
Conventional acupuncture	5.38 (1.32, 9.77)	-0.59 (-4.88, 3.53)	-1.04 (-18.93, 15.13)	Conventional acupuncture	-6.59 (-12.86, -0.30)	1.65 (-9.28, 12.84)	-9.75 (-18.21, -1.32)
Warm needle	-2.85 (-15.53, 9.80)	-8.91 (-21.62, 3.79)	5.55 (-11.29, 21.20)	6.59 (0.30, 12.86)	Warm needle	8.30 (-4.31, 20.98)	-3.14 (-9.22, 2.94)
Sham acupuncture	8.54 (1.59, 15.74)	2.56 (-3.89, 9.23)	-2.71 (-23.65, 16.74)	-1.65 (-12.84, 9.28)	-8.30 (-20.98, 4.31)	Sham acupuncture	-11.48 (-25.41, 2.64)
Fire needle			8.72 (-9.18, 25.39)	9.75 (1.32, 18.21)	3.14 (-2.94, 9.22)	11.48 (-2.64, 25.41)	Fire needle

Notes: The above data represent the confidence interval. The bold font indicates that there was a statistically significant difference between the two treatments.

Table 9 Ranking Probability Table of WOMAC Joint Function Scores

Intervention	Rank 1	Rank 2	Rank 3	Rank 4	Rank 5	Rank 6	Rank 7	Rank 8
Moxibustion	0.02	0.07	0.08	0.14	0.24	0.11	0.17	0.17
Western medicine	0.08	0.21	0.38	0.25	0.08	0	0	0
Electro-acupuncture	0	0	0.01	0.08	0.18	0.31	0.32	0.09
Sham moxibustion	0.3	0.14	0.11	0.14	0.08	0.06	0.08	0.09
Conventional acupuncture	0.15	0.37	0.27	0.13	0.06	0.01	0	0
Warm needle	0	0	0.02	0.13	0.25	0.35	0.21	0.04
Sham acupuncture	0.44	0.2	0.12	0.09	0.06	0.03	0.03	0.03
Fire needle	0	0	0.01	0.03	0.06	0.13	0.19	0.57

Notes: The bold font represents the probability of ordering the therapy.

Table 10 Results of Network Meta-Analysis of VAS Scores of Different Acupuncture Treatments for KOA

Moxibustion	0.69 (-0.28, 1.66)	-0.32 (-1.40, 0.74)	1.16 (-0.28, 2.58)	0.49 (-0.86, 1.86)	0.87 (-1.52, 3.36)	2.10 (-0.49, 4.62)	-0.34 (-1.73, 1.03)
-0.69 (-1.66, 0.28)	Western medicine	-1.01 (-1.76, -0.26)	0.47 (-0.75, 1.70)	-0.19 (-1.35, 0.94)	0.20 (-2.03, 2.52)	1.42 (-1.08, 3.82)	-1.03 (-2.18, 0.12)
0.32 (-0.74, 1.40)	1.01 (0.26, 1.76)	Electro-acupuncture	1.48 (0.46, 2.49)	0.82 (-0.11, 1.72)	1.20 (-0.95, 3.42)	2.43 (0.04, 4.74)	-0.02 (-1.01, 0.95)
-1.16 (-2.58, 0.28)	-0.47 (-1.70, 0.75)	-1.48 (-2.49, -0.46)	Conventional acupuncture	-0.67 (-1.53, 0.23)	-0.27 (-2.23, 1.74)	0.96 (-1.22, 3.05)	-1.50 (-2.38, -0.63)
-0.49 (-1.86, 0.86)	0.19 (-0.94, 1.35)	-0.82 (-1.72, 0.11)	0.67 (-0.23, 1.53)	Warm needle	0.39 (-1.75, 2.51)	1.62 (-0.70, 3.91)	-0.84 (-1.69, 0.03)
-0.87 (-3.36, 1.52)	-0.20 (-2.52, 2.03)	-1.20 (-3.42, 0.95)	0.27 (-1.74, 2.23)	-0.39 (-2.51, 1.75)	Acupoint catgut embedding	1.23 (-1.66, 4.11)	-1.23 (-3.41, 0.92)
-2.10 (-4.62, 0.49)	-1.42 (-3.82, 1.08)	-2.43 (-4.74, -0.04)	-0.96 (-3.05, 1.22)	-1.62 (-3.91, 0.70)	-1.23 (-4.11, 1.66)	Sham acupuncture	-2.46 (-4.74, -0.15)
0.34 (-1.03, 1.73)	1.03 (-0.12, 2.18)	0.02 (-0.95, 1.01)	1.50 (0.63, 2.38)	0.84 (-0.03, 1.69)	1.23 (-0.92, 3.41)	2.46 (0.15, 4.74)	Fire needle

Notes: The above data represent the confidence interval. The bold font indicates that there was a statistically significant difference between the two treatments.

improving WOMAC pain scores. Probability ranking results in improving WOMAC pain scores showed that fire needle > warm needle > electro-acupuncture > western medicine > moxibustion > conventional acupuncture > sham acupuncture > sham moxibustion. Moreover, electro-acupuncture was superior to western medicine and sham moxibustion while fire needle and warm needle were superior to western medicine and sham moxibustion in improving WOMAC stiffness scores. Probability ranking results in improving WOMAC stiffness scores showed that fire needle > warm needle > electro-acupuncture > conventional acupuncture > western medicine > moxibustion > sham moxibustion. Further analysis revealed that fire needle, warm needle and electro-acupuncture were all superior to conventional acupuncture and western medicine in improving WOMAC joint function scores. Probability ranking results in improving WOMAC joint function scores showed that fire needle > electro-acupuncture > warm needle > moxibustion > western medicine > conventional acupuncture > sham acupuncture > sham moxibustion. Electro-acupuncture was more effectively improved VAS scores compared to western medicine, conventional acupuncture and sham acupuncture, while fire needle was superior to conventional acupuncture and sham acupuncture in improving VAS scores. Probability ranking results in improving VAS scores showed that fire needle > electro-acupuncture > moxibustion > warm needle > western medicine > conventional acupuncture > acupoint catgut embedding > sham acupuncture. These results indicate that the fire needle had the best performance among the tested treatments in KOA treatment. Quality analysis results showed that the included studies had a medium quality. Thus, the application of the aforementioned interventions should be customized to the characteristics and condition of patients, and the probability ranking results only serve as a reference to clinicians.

Meta-analysis of previous online studies on the subject found that⁵⁷ warm needle and electro-acupuncture were probably the best acupuncture modalities for treating KOA. In this study, different conclusions were drawn. Our results indicate that fire needle, electro-acupuncture, and warm needle ranked top of all tested therapies. The fire needle regulates IL-1 signal transduction pathways to balance articular cartilage synthesis and decomposition. In this way, it reduces inflammation and joint injury, promotes local blood circulation and alleviates clinical symptoms in patients.^{58,59} Electro-acupuncture therapy is also one of the most effective KOA treatments.⁶⁰ Studies^{61,62} have reported that electro-acupuncture can reduce the

Table 11 Ranking Probability Table of VAS Scores

Intervention	Rank 1	Rank 2	Rank 3	Rank 4	Rank 5	Rank 6	Rank 7	Rank 8
Moxibustion	0.01	0.02	0.04	0.09	0.19	0.3	0.19	0.17
Western medicine	0.04	0.14	0.22	0.32	0.23	0.05	0.01	0
Electro-acupuncture	0	0	0	0.01	0.05	0.2	0.42	0.33
Conventional acupuncture	0.09	0.46	0.34	0.09	0.03	0.01	0	0
Warm needle	0.01	0.04	0.15	0.33	0.31	0.14	0.03	0
Acupoint Catgut embedding	0.15	0.23	0.18	0.12	0.11	0.07	0.06	0.08
Sham acupuncture	0.71	0.13	0.06	0.04	0.03	0.02	0.01	0.01
Fire needle	0	0	0	0.01	0.07	0.22	0.29	0.41

Notes: The bold font represents the probability of ordering the therapy.

expression of inflammatory cytokines in knee joints and inhibit inflammatory responses to achieve therapeutic effects. Warm needle suppresses inflammatory responses and alleviates clinical symptoms in KOA patients by inhibiting the expression of MMP-3 and TNF- α in joints.⁶² Warm needle up-regulates the expression of osteoprotegerin (OPG), down-regulates the expression of receptor activator of NF- κ B Ligand (RANKL), and increases the ratio of OPG/RANKL, thereby reducing bone resorption in subchondral bone and inhibiting the destruction of subchondral bone in KOA.⁶³

This study has some limitations: First, most of the included studies were not described in detail in the aspects of allocation concealment and blinding methods, and

experimental designs were not rigorously evaluated which decreases the quality of results presented here. Second, sample sizes, type, dosage and treatment course of western medicine in the included literatures were not consistent, leading to potential heterogeneity. Third, the included studies had some publication bias and small sample effect, which decreases the reliability of our results.

In conclusion, this network meta-analysis show that the fire needle is superior to warm needle and electro-acupuncture, while warm needle and electro-acupuncture is better than conventional acupuncture, western medicine, sham moxibustion, sham acupuncture in overall curative effect. In clinical practice, appropriate treatments should be selected while considering the patient's situation. Due to the


**Figure 7** WOMAC pain score comparison corrected funnel plots of different acupuncture treatments for KOA.

Table 12 Incidences of Adverse Events in Included Studies

Included Studies	Adverse Events
Zhang 2011 ¹¹	Moxibustion group: none Western medicine group: 3 cases (sour regurgitation 1, ausea 1, epigastric pain 1)
Qiu 2006 ²⁰	Electro-acupuncture group: 2 cases (local hematoma) Western medicine group: 9 cases (epigastric discomfort, sour regurgitation)
Yin 2017 ²⁴	Electro-acupuncture group: 1 case (subcutaneous hemorrhage) Western medicine group: 2 cases (gastrointestinal discomfort, mild constipation)
Chen 2012 ³¹	Warm needle group: none Western medicine group: 1 case(stomach discomfort)
Jiang 2013 ³²	Warm needle group: none Western medicine group: none
Shen 2017 ³⁷	Conventional acupuncture group: 2 cases (fainting during acupuncture treatment) Western medicine group: 6 cases (gastrointestinal discomfort, headache and dizziness, facial edema)
Zhang 2016 ³⁹	Warm needle group: 2 cases (fainting during acupuncture treatment, sticking of needle in acupuncture) Conventional acupuncture group: 8 cases (constipation, diarrhea, nausea, dizziness)
Lin 2018 ⁴⁰	Conventional acupuncture group: 2 cases (pain and hematoma at the acupuncture site) Sham acupuncture group: 1 case (pain at the acupuncture site)
Zhao 2014 ⁴¹	Moxibustion group: 10 cases (reddening of skin) sham moxibustion group: none
Lu 2011 ⁴⁴	Fire needle group: none Warm needle group: none
Lin 2012 ⁴⁸	Fire needle group: none Conventional acupuncture group: none
He 2018 ⁴⁹	Electro-acupuncture group: none Fire needle group: none

limitations associated with this study, future large scale, multi-center, high-quality randomized controlled trials are needed to validate results of this study.

Abbreviations

KOA, knee osteoarthritis; NSAIDs, Non-steroidal anti-inflammatory drugs; RCTs, Randomized controlled trials; WOMAC, Western Ontario and McMaster Universities Osteoarthritis Index; VAS, Visual Analogue Scale; MCMC, Markov Chain Monte Carlo; PSRF, potential scale reduction factor; ACR, American College of Rheumatology; TCM, traditional Chinese medicine; OARSI, Osteoarthritis Research Society International; OPG, osteoprotegerin; RANKL, receptor activator of NF- κ B Ligand.

Ethical Publication Statement

We confirm that we have read the Journal's position on issues involving ethical publication and affirm that this

report is consistent with those guidelines. Since this study is a literature analysis and does not involve patient privacy, informed consent of patients and approval of the ethics committee are not required.

Acknowledgments

The authors would like to thank Freescience (www.home-for-researchers.com) for English editing of this manuscript. Wei Liu, Yihua Fan and Yuanhao Wu have contributed equally to this work and are co-first authors.

Funding

This study was financially supported by the National Natural Science Foundation of China (No.82074377), the National Key Research and Development Project (No.2018YFC1705201), Tianjin Key Specialty Program (20210602-1), the Special Scientific Research for Traditional Chinese Medicine (No.201507001-07), and

High-Level Talent Selection and Training Project of Tianjin Health and Family Planning Industry (02005wuyuanhao).

Disclosure

The authors report no conflicts of interest in this work.

References

- Dawson J, Linsell L, Zondervan K, et al. Impact of persistent hip or knee pain on overall health status in elderly people: a longitudinal population study. *Arthritis Rheum*. 2005;53(3):368–374. doi:10.1002/art.21180
- Palmer KT, Reading I, Calnan M, Linaker C, Coggon D, Palmer KT, Reading I, Calnan M, et al. Does knee pain in the community behave like a regional pain syndrome? Prospective cohort study of incidence and persistence. *Ann Rheum Dis*. 2007;66(9):1190–1194. doi:10.1136/ard.2006.061481
- Avendaño-Coy J, Comino-Suárez N, Grande-Muñoz J, et al. Extracorporeal shockwave therapy improves pain and function in subjects with knee osteoarthritis: a systematic review and meta-analysis of randomized clinical trials. *Int J Surg*. 2020;82:64–75. doi:10.1016/j.ijssu.2020.07.055
- Lu YH, Shi XB. Current status and progress of epidemiological studies on knee osteoarthritis. *Chine J Traditional Med Traumatology Orthopedics*. 2012;20(6):81–84.
- Blagojevic M, Jinks C, Jeffery A, et al. Risk factors for onset of osteoarthritis of the knee in older adults: a systematic review and meta-analysis. *Osteoarthritis Cartilage*. 2010;18(1):24–33. doi:10.1016/j.joca.2009.08.010
- McAlindon TE, Bannuru RR, Sullivan MC, et al. OARSI guidelines for the non-surgical management of knee osteoarthritis. *Osteoarthritis Cartilage*. 2014;22(3):363–388. doi:10.1016/j.joca.2014.01.003
- Comino-Suárez N. Guidelines for Diagnosis and Treatment of Osteoarthritis. *Chine J Rheumatol*. 2010;14(6):416–419.
- Berman BM, Lao L, Langenberg P, Lee WL, Gilpin AMK, Hochberg MC, Berman BM, Lao L, Langenberg P, et al. Effectiveness of acupuncture as adjunctive therapy in osteoarthritis of the knee: a randomized, controlled trial. *Ann Intern Med*. 2004;141(12):901–910. doi:10.7326/0003-4819-141-12-200412210-00006
- Tukmachi E, Jubb R, Dempsey E, et al. The effect of acupuncture on the symptoms of knee osteoarthritis—an open randomised controlled study. *Acupunct Med*. 2004;22(1):14–22. doi:10.1136/aim.22.1.14
- McGrory B, Weber K, Lynott JA, et al. American Academy of Orthopaedic Surgeons. The American Academy of Orthopaedic Surgeons Evidence-Based Clinical Practice Guideline on Surgical Management of Osteoarthritis of the Knee. *J Bone Joint Surg Am*. 2016;98(8):688–692. doi:10.2106/JBJS.15.01311
- Zhang QJ, Cao LH, Li ZD, et al. Clinical effect and safety observation of moxibustion and celecoxib in the treatment of knee osteoarthritis. *Chine J Traditional Med Traumatology Orthopedics*. 2011;19(1):13–15.
- Zhou YL, Li J, Hou WG, et al. Clinical observation on moxibustion treatment of knee osteoarthritis. *Shanghai J Acupuncture Moxibustion*. 2014;33(12):1086–1088.
- Zhou JY. Double blind and double simulated clinical randomized controlled study of moxibustion treatment of knee osteoarthritis. *J Chengdu Univ Traditional Chin Med*. 2017.
- Ren XM, Cao JJ, Shen XY, et al. Moxibustion for knee osteoarthritis: a randomized controlled study. *Chin Acupuncture Moxibustion*. 2011;31(12):1057–1061.
- Zhou T, Chen CF, Qian YL, et al. Clinical Effect of Low frequency electro-acupuncture on knee osteoarthritis. *Shanghai J Traditional Chine Med*. 2015;49(8):56–57.
- Zhang H, Lv ZT, Lei C, et al. Clinical Efficacy of electro-acupuncture on Knee Osteoarthritis and Effect of Serum GRO- α and VEGF. *J Clin Acupuncture Moxibustion*. 2018;34(10):22–24.
- Liu N, Cai CM, Fu YT, et al. Clinical Observation on 41 Cases of Knee Osteoarthritis Treated by electro-acupuncture and Excision. *World J Integrated Traditional Western Med*. 2020;15(2):344–347.
- Guo YM, Liang YY, Gu JQ, et al. Electro-acupuncture around acupuncture treatment of early and middle knee osteoarthritis. *Jilin J Chine Med*. 2016;36(1):91.
- Wu WH, Tang J, Wu YP, et al. Evaluation of modern rehabilitation of the curative effect of electro-acupuncture on Xiyuan(EX-LE5) point in the treatment of knee osteoarthritis. *Shanghai J Traditional Chine Med*. 2015;49(06):63–65.
- Qiu ML, Dai QP, Che T, et al. Clinical study of electro-acupuncture at knee eye point in the treatment of knee osteoarthritis. *J Traditional Chine Orthopedics Traumatol*. 2006;18(3):15–16.
- Gang JH, Mi YQ, Wang HM. Clinical efficacy of electro-acupuncture versus meloxicam in the treatment of early -and mid-stage knee osteoarthritis: a randomized controlled study. *Chin Acupuncture Moxibustion*. 2016;36(05):467–470.
- Gao J, Ouyang BS, Zhang Y, et al. Comparison of curative effect between electro-acupuncture and warming acupuncture moxibustion on knee osteoarthritis of kidney deficiency and myeloid deficiency. *Chin Acupuncture Moxibustion*. 2012;32(5):395–398.
- Wang J. electro-acupuncture and warm needle in the treatment of blood stasis type knee joint osteoarthritis. *Lab Med Clin*. 2017;14(4):508–509.
- Yin XZ. Effect of electro-acupuncture on functional improvement of knee osteoarthritis. *Chine General Pract*. 2017;20(S3):388–389.
- Ju ZY, Wang Y, Chen CL, et al. Clinical study of electro-acupuncture on knee osteoarthritis. *Shanghai J Acupuncture Moxibustion*. 2017;36(09):1111–1115.
- Wu JL. Clinical observation of electro-acupuncture on knee osteoarthritis. *J Heilongjiang Univ Traditional Chin Med*. 2012.
- Chen ZX. Effect of warm needle on pain and quality of life of senile knee osteoarthritis. *Clin J Chine Med*. 2018;10(5):86–87+89.
- Liu RB, Huang Z, Wu Q. Clinical Observation of Warming Acupuncture and moxibustion on Knee Osteoarthritis of qi-stagnation and blood stasis type. *J Guangzhou Univ Traditional Chine Med*. 2014;9(1). doi:10.1186/1749-8546-9-2
- Tan HY, Feng YL. Clinical Effect of Warming Acupuncture and Acupuncture on the Dubei(ST35) and Neixiyan(EX-LE4) in the Treatment of Knee Osteoarthritis. *Clin J Chine Med*. 2016;8(14):23–24.
- Ma SJ, Yang K, Li FQ. Clinical observation on treating 54 cases of knee osteoarthritis with warm needle. *Zhejiang J Traditional Chine Med*. 2015;50(10):751.
- Chen ZG, Wu LH, Chen MJ, et al. Long-term effect of warm needle on knee osteoarthritis. *Shanghai J Acupuncture Moxibustion*. 2012;31(5):339–341.
- Jiang B, Hou XQ, Tang ZY. Clinical study on warm needle treatment of knee osteoarthritis. *Chine J Traditional Med Traumatology Orthopedics*. 2013;21(11):18–21.
- Ren ZJ. Clinical study on warm needle treatment of knee osteoarthritis. *China Health Standard Management*. 2018;9(8):94–96.
- Zhang J. Therapeutic effect of warm needle on knee osteoarthritis of deficiency and cold type. *China Health Standard Management*. 2016;7(12):128–129.
- Pan YY, Tang XJ, Cui WQ, et al. Clinical observation of acupoint catgut embedding in the treatment of knee osteoarthritis. *Chine Med Modern Distance Educ China*. 2020;18(09):90–92.
- Ji LL. Clinical study on the effect of acupuncture and moxibustion on serum cytokines in patients with knee osteoarthritis. *J Nanjing Univ Traditional Chine Med*. 2012.

37. Shen J. A comparative study of acupuncture and moxibustion therapy and diclofenac sodium emulsion in the treatment of senile knee osteoarthritis. *J Clin Acupuncture Moxibustion*. 2017;33(7):22–25.
38. Zhang SL, He YY, He SM. Clinical Observation of 28 Cases of Knee Osteoarthritis Treated by Acupuncture. *J Med Theory Pract*. 2016;29(22):3084–3085.
39. Miao GD, Jiang ZF, Tian ZJ. Observation on the curative effect of acupuncture and moxibustion on knee osteoarthritis. *Hebei J Traditional Chine Med*. 2014;36(5):725–726.
40. Lin LL, Li YT, Tu JF, et al. Effectiveness and feasibility of acupuncture for knee osteoarthritis: a pilot randomized controlled trial. *Clin Rehabil*. 2018;32(12):1666–1675. doi:10.1177/0269215518790632
41. Zhao L, Cheng K, Wang L, et al. Effectiveness of moxibustion treatment as adjunctive therapy in osteoarthritis of the knee: a randomized, double-blinded, placebo-controlled clinical trial. *Arthritis Res Ther*. 2014;16(3):R133. doi:10.1186/ar4590
42. Zhang GB, Wang WL, Zhang J, et al. A comparative study of fire needle and warm needle in the treatment of knee osteoarthritis. *Shanghai J Acupuncture Moxibustion*. 2013;32(4):299–301.
43. Zheng M, Hong KD, Chen SJ, et al. Observation on the curative effect of fire needle on knee osteoarthritis. *Chine Med Modern Distance Educ China*. 2016;14(5):103–106.
44. Lu DJ, Wang Q, Mei SW. Clinical comparative study of fire needle and warm needle in the treatment of knee osteoarthritis. *J New Chine Med*. 2011;43(10):94–96.
45. Li T, Ren J, Li LY. Study on the curative effect of fire needle on early and middle stage of knee osteoarthritis. *J Xinjiang Med Univ*. 2020;43(5):629–633.
46. Tang LT. Comparison and observation on the clinical effect of fire needle and warm needle therapy in treating knee osteoarthritis of yang deficiency and cold coagulation type. *J Guangxi Univ Chine Med*. 2017.
47. Song WP. Clinical Observation on the Treatment of Knee Osteoarthritis of Yang-deficiency and Cold Coagulation type with fire needle. *Yunnan J Traditional Chine Med Materia Med*. 2020;41(6):64–66.
48. Lin JC. *Clinical Observation on the Curative Effect of Fire Needle on Knee Osteoarthritis*. Jiangsu: Nanjing University of Traditional Chinese Medicine; 2011.
49. He TF, Bing XH, Zhou LY, et al. Comparison of curative effect of fire needle and electro-acupuncture on knee osteoarthritis. *Shanghai J Acupuncture Moxibustion*. 2018;37(6):675–679.
50. Fan CL, Zhang Q, Xu JH, et al. Clinical Observation of 54 Cases of Knee Osteoarthritis of Yang Deficiency and Cold Coagulation type treated by fire needle. *World J Integrated Traditional Western Med*. 2016;11(4):524–526+530.
51. Bannuru RR, Osani MC, Vaysbrot EE, et al. OARS guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthritis Cartilage*. 2019;27(11):1578–1589. doi:10.1016/j.joca.2019.06.011
52. Hepper CT, Halvorson JJ, Duncan ST, et al. The efficacy and duration of intra-articular corticosteroid injection for knee osteoarthritis: a systematic review of level I studies. *J Am Acad Orthop Surg*. 2009;17(10):638–646. doi:10.5435/00124635-200910000-00006
53. Han Y, Huang H, Pan J, et al. Meta-analysis Comparing Platelet-Rich Plasma vs Hyaluronic Acid Injection in Patients with Knee Osteoarthritis. *Pain Med*. 2019;20(7):1418–1429. doi:10.1093/pm/pnz011
54. Li J, Li YX, Luo LJ, et al. The effectiveness and safety of acupuncture for knee osteoarthritis: an overview of systematic reviews. *Medicine*. 2019;98(28):e16301.
55. Li-Jiang J, Jing-Yu X, Yang X-J, et al. A study of acupoint selection rules of Moxibustion and Auricular point therapy in the treatment of urinary retention after anorectal surgery based on data mining. *Asian Toxicol Res*. 2021;3(1):4.
56. Corbett MS, Rice SJ, Madurasinghe V, et al. Acupuncture and other physical treatments for the relief of pain due to osteoarthritis of the knee: network meta-analysis. *Osteoarthritis Cartilage*. 2013;21(9):1290–1298. doi:10.1016/j.joca.2013.05.007
57. Li S, Xie P, Liang Z, et al. Efficacy comparison of five different acupuncture methods on pain, stiffness, and function in osteoarthritis of the knee: a network meta-analysis. *J Ningxia Med Univ*. 2018;2018:1638904.
58. Wang WJ, Geng P, Chen DS, et al. Effects of fire needle on IL-1 Signal Transduction Pathway of Knee Osteoarthritis. *Pract Clin J Integrated Traditional Chin Western Med*. 2015;15(5):29–30+41.
59. Wang Y, Xie X, Zhu X, et al. Fire-Needle Moxibustion for the Treatment of Knee Osteoarthritis: a Meta-Analysis. *Evid Based Complementary Alternative Med*. 2016;2016:1392627.
60. Kang HR, Lee YS, Kim SH, et al. Effectiveness and safety of electrical moxibustion for knee osteoarthritis: a multicenter, randomized, assessor-blinded, parallel-group clinical trial. *Complement Ther Med*. 2020;53:102523. doi:10.1016/j.ctim.2020.102523
61. Kang HR, Jung CY, Lee SD, et al. Efficacy and safety of electrical moxibustion for knee osteoarthritis: study protocol for a randomized controlled trial. *Trials*. 2018;19(1):159. doi:10.1186/s13063-018-2514-x
62. Yue P, Gao L, Chen M, et al. Effect of Warm-needle-moxibustion on Behavior Reactions and TNF- α and MMP-3 Contents in Knee Cartilage of Rabbits with Knee Osteoarthritis. *Acupuncture Res*. 2016;41(3):235–239.
63. Ma YY, Liu D, Liu JW, et al. Effect of warm needle on expression of OPG and RANKL protein in subchondral bone of knee osteoarthritis patients. *J Ningxia Med Univ*. 2020;42(7):683–687.

Journal of Pain Research

Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication. The manuscript

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>

management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress