


# Case Report: Fatigue and Bleeding in a Polymedicated Patient Using Several Herbal Supplementations, Detected with g-Nomic<sup>®</sup> Software

Abel Saldarreaga Marin <sup>1</sup>Marc Cendros<sup>2</sup>Carlos J Ciudad <sup>3</sup>Ana Sabater <sup>2</sup>

<sup>1</sup>Bahía de Cádiz-La Janda Health District, Mobile Emergency Unit of the Andalusian Health Care, Service of Cádiz, Cádiz, Spain; <sup>2</sup>Eugenomic, Barcelona, Spain; <sup>3</sup>Biochemistry and Molecular Biology Unit, School of Pharmacy, University of Barcelona, Barcelona, Spain

**Abstract:** This was a case report of severe fatigue and bleeding in a 65-year-old man with ischemic heart disease who was wearing a stent and taking multiple medications for hypertension and diabetes. The use of a drug interaction and personalized prescription software (g-Nomic<sup>®</sup>) revealed potential interactions, involving acetylsalicylic acid and several non-pharmaceutical products including ginger, blueberry extracts, pineapple juice, docosahexaenoic acid and liquorice. Correction of these interactions resulted in complete remission of the reported side effects. This supports the idea that non-pharmaceuticals potentiated the effects of acetylsalicylic acid on haemostasis, producing the bleeding that would have caused fatigue. It is important to use appropriate tools to detect drug interactions that also take into account commonly used non-pharmaceutical products. Drug interactions can be considered illnesses by themselves.

**Keywords:** pharmacogenetics, SNP, drug-drug interaction, drug-lifestyle interactions, drug-herb, software, bleeding, fatigue, polymedication, medication risk, pharmacogenetics software, personalized prescription

## Introduction

Drug treatment can be affected by significant variability in the degree of response and the occurrence of side effects. Adverse drug reactions are responsible for considerable morbidity and mortality. Sometimes these undesirable effects are intrinsic to the drug, but in many cases they are due to drug-drug interactions.<sup>1,2</sup> It is known that drug-drug interactions can trigger adverse effects of medicines that would have been safe if they had not been used in combination. In addition, other substances that are not usually considered pharmaceuticals, like herbal products, may also cause drug-lifestyle interactions. Recognising and managing these interactions can be an invaluable process in achieving safer therapy.

We present a case of a polymedicated patient who was taking apparently harmless substances and dietary supplements and who presented with bleeding and fatigue. This situation was resolved by identifying and managing drug interactions using g-Nomic<sup>®</sup> personalised prescription software.

## Case Presentation

A 65-year-old Caucasian man with hypertension, type 2 diabetes, elevated LDL-cholesterol and ischemic cardiomyopathy, wearing a stent, presented unexplained fatigue with physical activity and coagulation defects consisting of gingival bleeding and frequent bruising.

Correspondence: Ana Sabater  
Eugenomic, c/ Londres 6, Barcelona,  
08029, Spain  
Tel +34-93-292-2963  
Email asabater@eugenomic.com

## Analytical Explorations

All biochemical parameters were within the normality except for elevated LDL-c levels. Hepatic and renal functions were normal (Alanine transaminase -ALT 17 U/L, aspartate transaminase AST 21 u/L, gamma-glutamyltransferase 24 U/L, creatinine 0.94 mg/dL, blood urea 35 mg/dL, glucose 95 mg/dL, glycosylated haemoglobin 6.1%). Prothrombin time (seconds): 10.2" (8.9–13.3). Normalized prothrombin time (INR): 0.90 (0.80–1.20).

At the moment of the consultation, the patient had the following treatment:

Telmisartan 40mg/d, metformin 850 mg/d, empagliflozin 40mg/d, bisoprolol 2.5/d, atorvastatin 40mg/d, clopidogrel 75mg/d and acetylsalicylic acid (ASA) 100mg/d.

## Examination

Except for fatigue and bleeding, the patient showed good general condition; he was conscious, oriented and collaborative, eupnoic at rest and afebrile. Cardiopulmonary

auscultation and abdominal examination did not result in pathological findings. Basal oxygen saturation: 98%. Blood pressure: 138 mm Hg (systolic)/76 mm Hg (diastolic). Heart rate; 85 bpm. Respiratory rate at rest: 14 bpm. Temperature: 35.8 °C.

The patient's medication regime was evaluated using g-Nomic® software<sup>3</sup> to check for interactions and possible genetic effects. In Figure 1 it is shown all the initial drugs entered in the personalized prescription software.

In addition, the same personalized prescription software was used to check for possible substances that could contribute to interactions even if they are not considered drugs per se, referred to in the software as lifestyle habits. After interviewing the patient about them, it was identified that the patient consumed 3 grams of docosahexaenoic acid (DHA) /day, blueberry and ginger extract supplements, and daily consumption of pineapple juice. In addition, the patient quit smoking nine years ago, and since then has been consuming liquorice ever since to support abstinence. Figure 2 shows the list of lifestyle

My Patients

Patient data

Drugs

LifeStyle (46)

Interactions (3)

Genes

PGX Report (1)

Risks (4)

Active ingredients / Brand name drug list

1 - 7 of 7

Add Active Ingredient (AI)

Name

Surname

Med.history Num/Ref.

Patient

Active ingredient	Brand name drug	Medication start	Medication end	Stop tacking	Delete/ Stop taking
Acetylsalicylic acid		31.05.2021	N/A	<input type="checkbox"/>	Delete Stop taking
Atorvastatin		31.05.2021	N/A	<input type="checkbox"/>	Delete Stop taking
Bisoprolol		31.05.2021	N/A	<input type="checkbox"/>	Delete Stop taking
Clopidogrel		31.05.2021	N/A	<input type="checkbox"/>	Delete Stop taking
Empagliflozin		31.05.2021	N/A	<input type="checkbox"/>	Delete Stop taking
Metformin		31.05.2021	N/A	<input type="checkbox"/>	Delete Stop taking
Telmisartan		31.05.2021	N/A	<input type="checkbox"/>	Delete Stop taking

Add Active Ingredient (AI)

Active ingredients / Brand name drug list

1 - 7 of 7

**Figure 1** All initial drugs entered in the personalized prescription software.

My Patients	Patient data	Drugs	LifeStyle (46)	Interactions (3)	Genes	PGX Report (1)	Risks (4)
-------------	--------------	-------	----------------	------------------	-------	----------------	-----------

LifeStyle				1 - 20 of 46 >
Name	<input type="text"/>	Surname	<input type="text"/>	Med.history Num/Ref. <input type="text"/> Abel Case <input type="text"/>
Active ingredient ^		Verify interaction ^		
Alcohol (Etanol)		Verify interaction		
Aloe vera (Aloe barbadensis)		Verify interaction		
Alpha-Lipoic Acid (ALA)		Verify interaction		
Aluminum		Verify interaction		
Arginine (Arginine)		Verify interaction		
Astragalus		Verify interaction		
Berberine (Berberis sp.)		See PGX Report tab to see interaction		
Black pepper (Piper nigrum)		See PGX Report tab to see interaction		
Blond plantain (Plantago ovata)		Verify interaction		
Blueberry (Vaccinium corymbosum, V. myrtillus)		Verify interaction		
Cat's claw (Uncaria tomentosa)		Verify interaction		
Chamomile (Matricaria chamomilla)		See PGX Report tab to see interaction		
Cinnamon (Cinnamomum verum)		Verify interaction		
Danshen (Salvia miltiorrhiza)		Verify interaction		
Embli (Phyllanthus emblica)		Verify interaction		
Feverfew (Tanacetum parthenium)		Verify interaction		
Filipendula ulmaria		Verify interaction		
Ginger (Zingiber officinale)		Verify interaction		
Ginkgo biloba-leaf extracts		Verify interaction		
Grapefruit (Citrus paradisi)		Verify interaction		
Horse-chestnut (Aesculus hippocastanum)		Verify interaction		
Karela (Momordica charantia)		Verify interaction		
Licorice (Glycyrrhiza glabra)		Verify interaction		
Linum usitatissimum		Verify interaction		
Methylthioninium chloride		See PGX Report tab to see interaction		
Milk thistle (Silybum marianum)		See PGX Report tab to see interaction		
Noni (Morinda citrifolia)		Verify interaction		
Oats bran (Avena sativa)		Verify interaction		
Omega 3 fatty acids		Verify interaction		
Pectin		Verify interaction		
Pine bark		Verify interaction		
Pineapple -juice		See PGX Report tab to see interaction		
Potassium		Verify interaction		
Primrose oil (Oenothera spp.)		Verify interaction		
Quinine		See PGX Report tab to see interaction		
Red Clover (Trifolium pratense)		See PGX Report tab to see interaction		
Red raspberry (Rubus idaeu)		See PGX Report tab to see interaction		
Resveratrol		Verify interaction		
Saint John's Wort (Hypericum perforatum)		Verify interaction		
Schisandra (Schisandra chinensis)		See PGX Report tab to see interaction		
Sulfisoxazole		See PGX Report tab to see interaction		
Tamarind (Tamarindus indica)		Verify interaction		
Turmeric (Curcuma longa)		See PGX Report tab to see interaction		
Vitamin B3 (niacin)		Verify interaction		
Willow Bark		Verify interaction		
Xiaoke Pill		Verify interaction		

**Figure 2** Shows the list of all lifestyle habits that could interact with patient's medication, reported by the personalized prescription software.

habits that may interact negatively with the patient's prescription entered into the personalized prescription software. The physician should ask the patient if he/she takes any of these substances on a regular basis in his/her daily life. If so, the doctor should click on the hyperlink to make this lifestyle habits part of the prescription as a whole.

## Actions and Outcome

After evaluating the possible interactions between medication and lifestyle, some measures were taken to avoid those interactions. Figure 3 shows drug-drug interactions as well as drug-lifestyle interactions. Figure 4 shows the possible interactions due to inhibitions and inductions caused by patient's poly medication. It is known that substances with inhibitory capacity can also impair enzyme activity even in the absence of loss-of-function variants. Therefore, a person who would be classified as a normal metaboliser based on a genetic test alone would experience a phenoconversion to a poor metaboliser when exposed to a strong inhibitor. Similarly, enzyme inducers can increase enzyme activity beyond what a genetic test might indicate. This reasoning can be extended to drug-herb and drug-lifestyle interactions as potential modulators of drug metabolism and thus drug response.

The patient was advised to discontinue the consumption of ginger, blueberry extracts and pineapple juice. The dose of omega-3 fatty acids was reduced from 3g/day DHA to 1g/day. Liquorice was maintained, but the patient was told to minimize its use.

As the patient was already being treated with a beta-blocker to control blood pressure, telmisartan was temporarily discontinued to minimise the side effects of the combination of two antihypertensive drugs.

Figure 5 shows the primary genes to be considered when checking the specific medication for to this clinical case. After assessing all drug interactions, no genetic testing of the patient was considered necessary.

The patient was scheduled for a follow-up evaluation two months later.

At the next visit, there was a clear improvement in both his clinical condition and general wellbeing. The coagulation defects had disappeared with no further gingival bleeding or skin bruising.

Blood pressure was normal (BP: 134/68 mmHg) despite the discontinuation of telmisartan, and it was considered that reinstating the antihypertensive was not justified. Furthermore, glycaemia was also correct (HbA1c:

5.4%) and it was decided to discontinue metformin leaving only Empagliflozin to manage the patient's diabetes.

The patient's general condition and well-being had also increased enormously: he did not complain of fatigue and had started exercising outdoors and working out at the gym three times a week.

## Discussion

Drug interactions seem to have been the cause of the patient's worsening condition, as the improvement was remarkable after addressing the detected potential risks of interactions.

The main concern in this patient was the excessive bleeding. Several factors could have contributed to altered haemostasis, including impaired renal function, which could also explain the increased blood pressure. ASA has an anticoagulant action and may also reduce renal function, contributing to both said problems, and its effects could have been potentiated by other substances. Ginger has hypoglycaemic action, but has also been reported to inhibit thromboxane synthesis, so it may potentially interact with acetylsalicylic acid to potentiate the effects of this drug.<sup>4</sup> Berries from the *Vaccinium* genus contain small amounts of salicylates, and berry juice in large quantities (more than 1L/day) or berry extracts may increase the hypoprothrombinaemia effects of acetylsalicylic acid.<sup>5</sup> While 1L a day is too large to be a likely cause of interaction, the patient was taking an extract supplement, which usually contains higher amounts of active compounds than those present on juice or the berry itself. Pineapple juice, which the patient consumed daily, contains bromelain, a substance that can inhibit cytochrome P450 2C9,<sup>6</sup> which is the main metabolic pathway for acetylsalicylic acid.<sup>7</sup> Therefore, regular intake of pineapple juice could have been an additional factor leading to higher exposure to salicylates.

All of this would contribute to increase the effects of salicylates on both platelet aggregation and blood pressure.

Hypoprothrombinaemia may have also been increased by omega-3 fatty acids, as the patient was consuming large doses of DHA.<sup>8</sup>

Furthermore, liquorice has been reported to reduce the effects of the antihypertensive medication by increasing liquid retention.<sup>9</sup>

In this case, none of these products would have been considered dangerous on their own, as they are consumed by a large amount of people without incident. They would have been overlooked unless the personalised prescribing software had reported them as potential risks of interaction



My Patients	Patient data	Drugs	LifeStyle (41)	Interactions (8)	Genes	PGX Report (5)	Risks (4)
Interactions Drug/Drug and Drug/Lifestyle							1 - 8 of 8
Name		Surname		Med.history Num/Ref.		Patient	
Active ingredient	Category interaction Drug/Lifestyle interaction Drug/Drug interaction						
Metformin Telmisartan	<p>Angiotensin II receptor inhibitors may increase the effect of hypoglycemic agents and facilitate hypoglycaemia. <b>MANAGEMENT OPTIONS:</b> Blood glucose concentrations should be monitored more carefully at the beginning or the end of a concomitant treatment. <b>Ref:</b> Toujeo (insulin glargine) [FDA product information] 03/2018</p>						
Empagliflozin Telmisartan	<p>Angiotensin II receptor inhibitors may increase the effect of hypoglycemic agents and facilitate hypoglycaemia. <b>MANAGEMENT OPTIONS:</b> Blood glucose concentrations should be monitored more carefully at the beginning or the end of a concomitant treatment. <b>Ref:</b> Toujeo (insulin glargine) [FDA product information] 03/2018</p>						
Acetylsalicylic acid Telmisartan	<p>NSAIDs reduce the efficacy of antihypertensive drugs because they reduce the renal function. The effect is gradual and therefore if the use of NSAIDs in the short term it is unlikely to have clinically relevant consequences. <b>MANAGEMENT OPTIONS:</b> The risk of interaction can be minimized by using acetaminophen instead of an NSAID, using nonacetylated salicylates instead of an NSAID or using the NSAID at low doses and for the shortest possible time. During a combined therapy, it is warranted to monitor blood pressure.</p>						
Ginger (Zingiber officinale) Metformin	<p>Both ginger and metformin can increase insulin levels and/or lower blood glucose levels, which by additive effect can lead to hypoglycemia. <b>MANAGEMENT OPTIONS:</b> In the concomitant use, it is recommended to monitor blood glucose levels more closely. <b>Ref:</b> Zadayan G, Fuhr U. Phenotyping studies to assess the effects of phytopharmaceuticals on in vivo activity of main human cytochrome p450 enzymes. Planta Med. 2012 Sep;78(13):1428-57.</p>						
Bisoprolol Licorice (Glycyrrhiza glabra)	<p>Licorice can cause fluid retention and reduce the effect of antihypertensives. <b>MANAGEMENT OPTIONS:</b> It is recommended to avoid the concomitant use of antihypertensive with licorice, even in infusion doses. <b>Ref:</b> Stockley, Ivan H. Stockley's Herbal Medicines Interactions. A guide to the interactions of herbal medicines, dietary supplements and nutraceuticals with conventional medicines. Great Britain: Pharmaceutical Press; 2009. p273</p>						
Acetylsalicylic acid Clopidogrel Ginger (Zingiber officinale)	<p>It has been reported that ginger (Zingiber officinale) inhibits the thromboxane synthetase activity with which it can interact with anticoagulants in a significant way. <b>MANAGEMENT OPTIONS:</b> It is recommended to avoid the association and, if necessary, monitor the INR more closely. <b>Ref:</b> Tsai HH et al. A review of potential harmful interactions between anticoagulant/antiplatelet agents and Chinese herbal medicines. PLoS One. 2013 May 9;8(5):e64255.</p>						
Acetylsalicylic acid Clopidogrel Omega 3 fatty acids	<p>The administration of omega-3 ethyl esters with antiplatelet drugs result, by a sinergic effect, in a potentation in the action of antiplatelet agents. <b>MANAGEMENT OPTIONS:</b> Watch for signs of excessive platelet antiaggregation.  Cohen MG, Rossi JS, Garbarino J et al. Insights into the inhibition of platelet activation by omega-3 polyunsaturated fatty acids: beyond aspirin and clopidogrel. Thromb Res. 2011 Oct;128(4):335-40.</p>						
Acetylsalicylic acid Blueberry (Vaccinium corymbosum, V. myrtillus)	<p>Long-term consumption of cranberry juice in large quantities (approximately 1L/day or more), as well as blueberry concentrate supplements, can potentiate the action of aspirin. There is no evidence that coagulation is affected to a clinically relevant extent with daily consumption of up to half a liter. <b>MANAGEMENT OPTIONS:</b> Due to the narrow therapeutic index of aspirin, it is prudent that the consumption of cranberry juice is rather low when using this drug. <b>Ref:</b> Srinivas NR. Cranberry juice ingestion and clinical drug-drug interaction potentials; review of case studies and perspectives. J Pharm Pharm Sci. 2013;16(2):289-303.</p>						

**Figure 3** Drug interactions and drug – lifestyle reported by the personalized prescription software.

My Patients
Patient data
Drugs
LifeStyle (41)
Interactions (8)
Genes
PGX Report (5)
Risks (4)

Pharmacogenetic Report
1 - 3 of 3

Show Secondary
Genes/Drugs interactions
Inhibitions and inductions
Specific remarks

Name
Surname
Med.history Num/Ref.
Patient

Active ingredient	Brand name drug	Evidence icon	Gene	Relevant information
Pineapple - juice			CYP2C9	Pineapple - juice is a potent inhibitor of the CYP2C9 enzyme, so the bioavailability of Acetylsalicylic acid may be increased by a factor greater than 5. It is recommended to replace Pineapple - juice, with another of similar effect that is not a potent enzyme inhibitor CYP2C9 to avoid toxicity due to overdose, except when there are guidelines that define the dose reduction for this drug.
Licorice (Glycyrrhiza glabra)			ABCB1 CYP3A4	Being Licorice (Glycyrrhiza glabra), weak inducer of the efflux transport protein Pgp -MDR1, encoded by the gene ABCB1, a slightly lower plasma levels of expected will be achieved of the drugs Atorvastatin, so it would be recommended to increase the dose by 20-30%. It is difficult to generalize and must see if there is specific information for each drug on drug monographs. Because Licorice (Glycyrrhiza glabra) is a weak inducer of the CYP3A4 gene/s, it can accelerate the metabolism of drugs that are substrates of it, i.e. Clopidogrel*. Being a prodrug/s, its transformation to the active form can be accelerated. Usually this has no negative effects. However, if the pathway that is induced is purely for elimination and does not contribute to the formation of active metabolites, the result would instead be a reduced efficacy. In some cases accelerated bioactivation can lead to overdose peaks. It is recommended to check the specific messages for more complete information. Since Licorice (Glycyrrhiza glabra) is/are weak inducer/s of the gene CYP3A4 it may accelerate the metabolism of drugs that are substrates of it: Atorvastatin, Bisoprolol. Therefore it will require higher doses to achieve the same therapeutic effect. It is recommended to increase the dose by 20-30%.
Acetylsalicylic acid			CYP2C19	Because Acetylsalicylic acid is a weak inducer of the CYP2C19 gene/s, it can accelerate the metabolism of drugs that are substrates of it, i.e. Clopidogrel*. Being a prodrug/s, its transformation to the active form can be accelerated. Usually this has no negative effects. However, if the pathway that is induced is purely for elimination and does not contribute to the formation of active metabolites, the result would instead be a reduced efficacy. In some cases accelerated bioactivation can lead to overdose peaks. It is recommended to check the specific messages for more complete information.

Show Secondary
Genes/Drugs interactions
Inhibitions and inductions
Specific remarks

Figure 4 Pharmacogenetics interactions described by the personalized prescription software.

with the patients' medication. The proactive warning prompted the physician to ask the patient about the consumption of these products. Herbal products or extracts used as supplements, such as blueberry extracts, would perhaps have been considered at some point, while

consumption habits, such as pineapple juice, would have remained unnoticed.

No specific cause of the fatigue was identified with certainty, but it was likely to be a side effect of the blood loss.

My Patients

Patient data

Drugs

LifeStyle (43)

Interactions (5)

Genes

PGX Report (5)

Risks (4)

Enter results of the genetic study. Request study if you did not do it.

\*\*\*\*\*Included 95 genes for simulation\*\*\*\*\*

### Add Other Genes Results

Name  Surname  Med.history Num/Ref.

Active ingredient	Brand name drug	Evidence icon	Recommended genes according to active ingredients	Add genotype/phenotype results
Acetylsalicylic acid		3	<i>CYP2C9</i>	Add Results
Atorvastatin		3	<i>BCRP</i>	Add Results
		2	<i>CYP3A4</i>	Add Results
		2	<i>SLCO1B1</i>	Add Results
Bisoprolol		4	<i>CYP3A4</i>	Add Results
Clopidogrel		1	<i>CYP2C19</i>	Add Results
Empagliflozin		4	<i>SLCO1B1</i>	Add Results
			<i>UGT1A9</i>	Add Results
		4	<i>UGT2B7</i>	Add Results

### Add Other Genes Results

**Figure 5** Pharmacogenetics markers described by the personalized prescription software.

In this case, the extracts the patient used, containing blueberry and ginger, could have contributed to bleeding, in addition to DHA and pineapple juice. These substances in combination were considered a likely trigger for the gingival bleeding.<sup>9</sup> It is not possible to know the relative contribution of these substances to the interaction, nor if a clinically meaningful interaction would still have happened without the presence of the whole polymedication. Caution is recommended whenever natural products are used in combination with drugs affecting haemostasis.

The patient's condition improved as the drug regimen was reduced and simplified, meaning that some drugs may

have been unnecessary in the first place. For example, the hypertension could have been secondary to the use of liquorice and salicylate potentiation, and in this case, discontinuation of these substances would have been more helpful in controlling blood pressure than starting telmisartan.

## Conclusion

This case report serves to illustrate two concepts: I) the importance of recognizing the interactions of existing compounds in commonly consumed products, in this particular case with anticoagulants. It is therefore advisable to use drug interaction software that take into account lifestyle habits. II)

Be aware of the increased risk of drug-drug interactions in patients consuming multiple natural products not considered drugs, which may be the cause of many drug-drug interactions. While being aware of the dangers of polymedication with multiple therapeutic drugs, other substances such as herbal products and dietary supplements, should not be disregarded. Reducing the drug regimen may occasionally lower the amount and severity of drug interactions, provided that the remaining drugs are enough to control the pathology. This work emphasizes the need for personalized approaches for patient empowerment in both clinical settings and home health care.

## Consent Form

Consent was obtained from the patient to allow publication of his medical record. The patient was informed that his case of polymedication would be presented as a case report to the scientific community in order to let other healthcare professionals know what his treatment was like and how it has evolved.

The patient signed a consent form to authorise his doctor to record his medical history for the purpose of a scientific publication on his health conditions. He agreed to have his medical records recorded for the named purpose of a scientific publication. No ethics statement from the institution was required, the sole statement from doctor was needed.

## Disclosure

Marc Cendrós and Ana Sabater work for EUGENOMIC S. L. and are directly involved in the development of the g-Nomic® the personalized prescription software, but neither they nor anyone at Eugenomic has had any role

in the management and clinical evaluation of this case beyond assisting the medical professionals in using the software. In addition, Mrs Ana Sabater has a patent g-Nomic issued to Eugenomic. The authors declare no other conflicts of interest in this work.

## References

1. Nikolic B, Jankovic S, Stojanov O, Popovic J. Prevalence and predictors of potential drug-drug interactions. *Cent Eur J Med.* 2014. doi:10.2478/s11536-013-0272-4
2. McQuaid KR, Laine L. Systematic review and meta-analysis of adverse events of low-dose aspirin and clopidogrel in randomized controlled trials. *Am J Med.* 2006;119(8):624–638. doi:10.1016/j.amjmed.2005.10.039
3. Sabater A, Ciudad CJ, Cendros M, Dobrokhoto D, Sabater-Tobella J. G-nomic: a new pharmacogenetics interpretation software. *Pharmgenomics Pers Med.* 2019. doi:10.2147/PGPM.S203585
4. Tsai HH, Lin HW, Lu YH, Chen YL, Mahady GB. A review of potential harmful interactions between anticoagulant/antiplatelet agents and chinese herbal medicines. *PLoS One.* 2013. doi:10.1371/journal.pone.0064255
5. Srinivas NR. Cranberry juice ingestion and clinical drug-drug interaction potentials; review of case studies and perspectives. *J Pharm Pharm Sci.* 2013;16(2):289. doi:10.18433/j3ng6z
6. Hidaka M, Nagata M, Kawano Y, et al. Inhibitory effects of fruit juices on cytochrome P450 2C9 activity in vitro. *Biosci Biotechnol Biochem.* 2008;72(2):406–411. doi:10.1271/bbb.70511
7. Agundez J, Martinez C, Perez-Sala D, et al. Pharmacogenomics in Aspirin Intolerance. *Curr Drug Metab.* 2009;10(9):998–1008. doi:10.2174/138920009790711814
8. Cohen MG, Rossi JS, Garbarino J, et al. Insights into the inhibition of platelet activation by omega-3 polyunsaturated fatty acids: beyond aspirin and clopidogrel. *Thromb Res.* 2011;128(4):335–340. doi:10.1016/j.thromres.2011.04.023
9. Stockley's herbal medicines interactions: a guide to the interactions of herbal medicines, dietary supplements and nutraceuticals with conventional medicines. *Choice Rev.* 2010. Available from: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2016/204353s020lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/204353s020lbl.pdf). Accessed August 2, 2021. doi:10.5860/choice.47-2357

### Pharmacogenomics and Personalized Medicine

### Publish your work in this journal

Pharmacogenomics and Personalized Medicine is an international, peer-reviewed, open access journal characterizing the influence of genotype on pharmacology leading to the development of personalized treatment programs and individualized drug selection for improved safety, efficacy and sustainability. This journal is indexed

on the American Chemical Society's Chemical Abstracts Service (CAS). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/pharmacogenomics-and-personalized-medicine-journal>

Dovepress