

Novel 1,3,4-Oxadiazole Derivatives of Pyrrolo [3,4-*d*]Pyridazinone Exert Anti-Inflammatory Activity without Acute Gastrototoxicity in the Carrageenan-Induced Rat Paw Edema Test

Marta Szandruk-Bender ¹
Anna Merwid-Ląd¹
Benita Wiatrak¹
Maciej Danielewski¹
Stanisław Dzimira ²
Danuta Szkudlarek³
Łukasz Szczukowski⁴
Piotr Świątek⁴
Adam Szeląg ¹

¹Department of Pharmacology, Wrocław Medical University, Wrocław, Poland;

²Department of Pathology, Wrocław University of Environmental and Life Sciences, Wrocław, Poland; ³Foundation of the Wrocław Medical University, Wrocław Medical University, Wrocław, Poland; ⁴Department of Chemistry of Drugs, Wrocław Medical University, Wrocław, Poland

Correspondence: Marta Szandruk-Bender
Department of Pharmacology, Wrocław Medical University, Mikulicza-Radeckiego 2, Wrocław, 50-345, Poland
Tel +48717841438
Fax +48717840094
Email marta.szandruk@umed.wroc.pl

Background and Purpose: Due to the risk of gastrointestinal damage and various tissue toxicity associated with non-steroidal anti-inflammatory drugs (NSAIDs) use, investigating new anti-inflammatory agents with efficacy comparable to that of NSAIDs but reduced toxicity is still a major challenge and a clinical need. Based on our previous study, new 1,3,4-oxadiazole derivatives of pyrrolo[3,4-*d*]pyridazinone, especially 6-butyl-3,5,7-trimethyl-1-[[4-[[4-(4-nitrophenyl)piperazin-1-yl]methyl]-5-thioxo-1,3,4-oxadiazol-2-yl]methoxy]pyrrolo[3,4-*d*]pyridazin-4-one and 6-butyl-1-[[4-[[4-(4-chlorophenyl)-4-hydroxy-1-piperidyl]methyl]-2-thioxo-1,3,4-oxadiazol-5-yl]methoxy]-3,5,7-trimethyl-pyrrolo[3,4-*d*]pyridazin-4-one (hereafter referred to as the compounds **10b** and **13b**, respectively) seem to be promising anti-inflammatory agents. This study aimed to elucidate the effects of these two new derivatives on the course of experimental rat inflammation, liver and kidney function, and gastric mucosa.

Methods: The anti-inflammatory effect of compounds **10b** and **13b** was evaluated using the carrageenan-induced paw edema test in rats. The increase in paw volume (paw edema), prostaglandin E₂ (PGE₂), tumor necrosis factor- α (TNF- α) and myeloperoxidase (MPO) levels, histological alterations, and inflammatory cell infiltration in paw tissue were determined. Serum alanine transaminase (ALT) and aspartate transaminase (AST) activities, serum urea and creatinine levels, as well as changes in gastric mucosa, were measured as indicators of hepatic, renal, and gastric toxicity.

Results: Pretreatment with both novel derivatives at 10 mg/kg and 20 mg/kg doses reduced paw edema, counteracted the increased PGE₂ and TNF- α levels, reduced the influx of inflammatory cells, and decreased histopathological alterations in paw tissue. Compound **13b** at a dose of 20 mg/kg was more effective than indomethacin in reversing the increased TNF- α levels and reducing the influx of inflammatory cells. Only compound **13b** at all studied doses (5, 10, or 20 mg/kg) counteracted the increased MPO level in paw tissue. Both compounds neither caused alterations in ALT, AST, urea, creatinine parameters nor gastric mucosal lesions.

Conclusion: New compounds exert an anti-inflammatory effect, presumably via inhibiting inflammatory mediators release and inflammatory cell infiltration. Moreover, both possess a more favorable benefit–risk profile than indomethacin, especially compound **13b**.

Keywords: carrageenan, anti-inflammatory agents, pyrrolo[3,4-*d*]pyridazinone, 1,3,4-oxadiazole, inflammatory mediators, toxicity

Introduction

Every living organism strives to achieve homeostasis.¹ Many highly specialized systems and defensive mechanisms allow the organism to adapt to the surrounding

environment and react in the event of exposure to endogenous or exogenous, infectious or non-infectious harmful stimuli. Inflammation, or inflammation response, is the organism's defensive reaction to remove or sequester the cause of the disturbance, repair the tissue, and finally restore homeostasis.²

Inflammation is characterized by such macroscopic symptoms as redness, edema, heat, pain, and loss of tissue function, which reflect elevated cellular metabolism, vasodilatation, and increased vascular permeability allowing leakage of plasma components and extravasation of immune cells.³ Plasma proteins and leukocytes, mainly neutrophils that are normally confined to the blood vessels gain access through the postcapillary venules to the extravascular tissues at the site of injury, thereby inducing edema. These vascular and cellular reactions are mediated by endogenous substances, known as inflammatory mediators, released at the site of the injury by tissue-resident immune cells, mostly macrophages, dendritic cells, and mast cells.⁴ Among many inflammatory mediators, proinflammatory cytokines and bioactive lipids – particularly tumor necrosis factor- α (TNF- α) and prostaglandin E₂ (PGE₂) – are considered to be the main ingredients that initiate and govern the inflammatory process.² In addition to typical inflammatory markers, the enzyme myeloperoxidase (MPO), released upon neutrophil activation, deserves special attention as a protein with proinflammatory properties independent of its enzymatic activity. These cytokine-like properties can modulate the activation state of leukocytes during inflammatory diseases.^{5,6}

Although the inflammatory response is one of the organism's defensive mechanisms against harmful factors, it is unpleasant for the patient, often causing suffering and sometimes not leading to recovery either. In principle, a controlled inflammatory response in appropriate amounts is beneficial to the organism, though it can become detrimental if dysregulated due to its tissue-damaging potential.² Usually, the inflammatory response is terminated once the triggering insult is eliminated and damaged tissue is repaired. If inflammatory response is prolonged, inefficient, or excessive, a chronic inflammatory state may ensue. This process, characterized by persistent production of proinflammatory cytokines and proinflammatory lipids, may lead to aberrant tissue remodeling, irreversible damage, and chronic disorders such as inflammatory bowel disease, atherosclerosis, asthma, or neurodegenerative disorders, which inevitably lead to impaired quality of life, loss of time from work or education, disability, and

untimely death.^{7,8} While many anti-inflammatory agents are available today, including non-steroidal anti-inflammatory drugs (NSAIDs), glucocorticoids, and biological agents, all of them have some limitations in terms of safety and efficacy.^{3,9} Therefore, there is still an unmet clinical need and a challenge for medicinal chemists to develop more effective and safer agents to treat the signs and symptoms of acute inflammation, thereby preventing its evolving into a chronic condition that leads to irreversible changes.

In this context, one of the promising strategies used in medicinal chemistry to identify new compounds is to combine a structure with known anti-inflammatory properties with a moiety capable of enhancing this activity. The pyrrolo[3,4-*d*]pyridazinone core is a structure that exerts anti-inflammatory activity.^{10,11} In turn, diversely substituted five-membered rings of 1,3,4-oxadiazole-2-thione, which is a bioisosteric analog of the carboxyl group, exhibit various biological activities – including anti-inflammatory activity.^{12–15} Moreover, anti-inflammatory compounds possessing this five-membered ring show decreased gastrotoxicity.^{13,14} Drugs used worldwide, eg, ibuprofen, diclofenac, or naproxen, were modified similarly. The obtained derivatives of the mentioned drugs, including in their structure a 1,3,4-oxadiazole-2-thione ring, show significant anti-inflammatory activity and diminished gastrointestinal adverse effects.^{13,14,16}

Inspired by this research, we designed, synthesized, and investigated the series of novel derivatives of pyrrolo[3,4-*d*]pyridazinone linked with 1,3,4 oxadiazole-2-thione pharmacophore.^{12,17,18} According to in silico and in vitro assays we performed, new molecules strongly inhibit cyclooxygenase (COX) activity, show superior affinity towards isoform COX-2, and some of them act as selective COX-2 inhibitors.^{12,17} In our previous in vivo study, the most promising novel pyrrolo[3,4-*d*]pyridazinone derivatives – 6-butyl-3,5,7-trimethyl-1-[[4-[[4-(4-nitrophenyl)piperazin-1-yl]methyl]-5-thioxo-1,3,4-oxadiazol-2-yl]methoxy]pyrrolo[3,4-*d*]pyridazin-4-one and 6-butyl-1-[[4-[[4-(4-chlorophenyl)-4-hydroxy-1-piperidyl]methyl]-2-thioxo-1,3,4-oxadiazol-5-yl]methoxy]-3,5,7-trimethylpyrrolo[3,4-*d*]pyridazin-4-one (hereafter referred to as the compounds **10b** and **13b**, respectively) – exerted dose-dependent antinociceptive activity with reduced gastrotoxicity in noxious stimuli induced models of pain, ie, the tail-flick and formalin test.¹⁸ It is well known that nociception and inflammation are functionally linked at multiple levels. Tissue damage is detected by both

nociceptors that enable pain sensation in the affected area and by tissue-resident cells, which induce an inflammatory response. Exudate formation, tissue edema, and inflammatory mediators are responsible for inflammatory pain, and nociception replenishes inflammatory sensors in tissue homeostasis monitoring.^{7,19} Since compounds **10b** and **13b** counteract inflammatory nociception in the late phase of the formalin test, compound **13b** even in a more efficient way than indomethacin,¹⁸ we decide to assess whether these novel compounds also exert anti-inflammatory action.

The current study was undertaken to elucidate the effect of pretreatment with newly synthesized pyrrolo[3,4-*d*]pyridazinone derivatives, compounds **10b** or **13b**, on the course of acute carrageenan-induced paw inflammation. Additionally, the levels of inflammatory mediators, such as PGE₂, TNF- α , and MPO, as well as the magnitude of infiltration with inflammatory cells are going to be checked to explain the possible mechanisms of action of the newly synthesized compounds. Moreover, this study was aimed at assessing the influence of new derivatives on liver and kidney function and gastric mucosa.

Materials and Methods

Drugs and Chemicals

The studied compounds – two novel derivatives of pyrrolo[3,4-*d*]pyridazinone named **10b** and **13b** – were prepared and characterized as reported earlier.¹² Analysis of the ¹H NMR, ¹³C NMR, MS, FT-IR, and elemental analysis as well as physicochemical features showed spectroscopic and analytical properties of the newly obtained derivatives to be in agreement with their assigned structure. Indomethacin, in subst. and λ -carrageenan, in subst. were purchased from Sigma Aldrich (Steinheim, Germany); carboxymethylcellulose (CMC), in subst. and formalin 37%, sol. were obtained from PolAura (Olsztyn, Poland); pentobarbital sodium + pentobarbital 133.3 mg/mL + 26.7 mg/mL, sol. was purchased from Biowet (Puławy, Poland); medetomidine hydrochloride 1 mg/mL, sol. was supplied from OrionPharma (Warszawa, Poland); normal saline was obtained from Polpharma (Starogard Gdański, Poland). Other chemicals used were included in the commercially available kits.

Animals

The Wistar rats (210–260 g) were obtained from the Animal Research Center at Wrocław Medical University (Wrocław, Poland). All animals were accustomed to the laboratory

condition for 7 days before commencing the experiments. The rats were housed, two per cage, in polypropylene cages with enrichments in a standard laboratory environment with a 12/12 h light/dark cycle, a humidity of 55–60% and a temperature of 21–24°C, with water ad libitum and free access to standard animal feed (Agropol, Motycz, Poland), except for the single procedure of deprivation.

Ethical Considerations

The current study was carried out with the permission (Resolution No. 101/2018 of 12.12.2018) of the Local Ethics Committee for Animal Experiments in Wrocław at Hirszfeld Institute of Immunology and Experimental Therapy of Polish Academy of Sciences (Wrocław, Poland). The animal care and all experimental procedures were by the applicable international, national, and institutional guidelines, including the Act of 15 January 2015 on the protection of animals used for scientific and educational purposes (Journal of Laws of 2015, item 266) and the EU directive 2010/63/EU.

Study Protocol

After seven days of adaptation, the animals were randomly allotted to nine groups (twelve animals per group) organized as follows:

- One group pretreated with 0.5% CMC solution (vehicle) intragastrically (i.g.) and injected subplantarily (s.pl.) with normal saline (control group, C);
- One group pretreated with 0.5% CMC i.g. and injected s.pl. with 1% carrageenan solution (carrageenan group, Car);
- One group pretreated with indomethacin at a dose of 10 mg/kg i.g. and injected s.pl. with 1% carrageenan solution (indomethacin group, Ind);
- And 6 groups pretreated i.g. with compound **10b** or **13b** at the doses of 5, 10, or 20 mg/kg and injected s.pl. with 1% carrageenan solution (**10b**-5, **10b**-10, **10b**-20, **13b**-5, **13b**-10, **13b**-20 groups, respectively).

The 0.5% CMC solution and studied substances suspended in 0.5% CMC solution were given in a single dose by a gastric tube (FST, Foster City, CA, USA) in a volume of 4 mL/kg. Prior to the administration of the appropriate substance, rats were food-deprived for 12 h. Doses of tested compounds and indomethacin were selected based on the earlier works.^{11,18,20,21} One hour after appropriate substance administration, the carrageenan-induced paw edema test (carrageenan-induced inflammation) was performed. After the test,

blood for biochemical assays was taken from the tail vein, centrifuged (15 min at 4000 rpm), and the obtained serum samples were kept at -80°C until further analysis. The rats were then sacrificed by intramuscular injection of medetomidine (0.5 mg/kg) followed by intraperitoneal injection of pentobarbital (200 mg/kg), and the right hind paw and the stomach were immediately dissected. Afterward, inflammatory exudate of the carrageenan-injected paw was collected for histopathological assessment. Then, one part of each right hind paw was fixed in 4% buffered formalin for histopathological analysis, and the soft tissue from the remaining part was homogenized (Homogenizer PRO250, PRO Scientific Inc., Oxford, CT, USA), with the obtained supernatants stored at -80°C for inflammatory markers evaluation. The excised stomachs were opened along the greater curvature, gently cleared of the contents by dipping in normal saline, macroscopically examined, and fixed in 4% buffered formalin for histopathological assessment.

Carrageenan-Induced Paw Edema Test

Inflammation in rats was induced by carrageenan injection according to the procedure described by Winter et al.²² After 1 h of appropriate substance administration, rats were injected once with 1% carrageenan solution in normal saline under the subplantar aponeurosis area of the right hind paw in a volume of 0.1 mL, except the control group in which rats were injected with normal saline given by the same route and in equivalent volume. The right hind paw of each rat was marked with ink at the level of the lateral malleolus and paw volume was measured up to this mark. The volume of the injected paw of each animal was measured plethysmometrically (Plethysmometer 37140, UgoBasile, Gemonio, Italy) before (time 0) and at 1, 2, 3, and 6 h after (time 1, 2, 3, 6) carrageenan or saline subplantar injection. For each time point, measurement was repeated three times and the average was then calculated. All measurements were performed by the same investigator to reduce any potential inter-operator variability. The paw edema was expressed as the relative increase in paw volume quantified by measuring the difference between the paw volume before and at 1, 2, 3, and 6 h after carrageenan or normal saline injection according to the equation:

$$\text{paw edema}(ml) = V_t - V_0$$

where V_t is the paw volume at t h after (time 1, 2, 3, 6) carrageenan or normal saline injection (mL); V_0 is the paw volume before (time 0) carrageenan or normal saline

injection (mL). The paw edema measured in the time intervals was used to calculate the percentage of paw edema inhibition using the following equation:

$$\text{inhibition of paw edema}(\%) = \left(1 - E_t/E_{car}\right) \times 100$$

where E_t is the edema volume of treated animals (mL); E_{car} is the edema volume of carrageenan-injected animals (mL).

Assessment of PGE₂, TNF- α , and MPO Levels in Paw Tissue

The concentrations of PGE₂, TNF- α , and MPO were measured in obtained supernatants with enzyme-linked immunosorbent assay (ELISA) kits: Rat PGE₂ ELISA Kit, Rat TNF- α ELISA Kit, Rat MPO ELISA Kit, (Cloud-Clone Corp., Katy, TX, USA) following the manufacturer's instructions. All concentrations were expressed as pg/mL.

Microscopic Assessment of Paw Tissue

The formalin-fixed paw specimens were embedded in paraffin, sectioned to 4 μm slices, mounted on the glass slides, stained by the routine hematoxylin-eosin (H&E) method, and examined using an Olympus BX53 light microscope combined with an Olympus UC90 camera (Olympus, Germany) at 200x magnification. Histopathological evaluation of inflammation indicatives was performed in a blinded fashion by an experienced pathologist using the 0-5-point scale presented previously by Mert et al²³ and described in detail in the legend for Table 1.

Microscopic Assessment of Paw Exudate

Cell blocks from inflammatory exudates of the paw were prepared by a tissue clot method (by allowing a clot to form in the lumen of the fine needle aspiration tip).²⁴ The clot was then transferred directly to 4% buffered formalin for fixation, embedded in paraffin, and cut into 4 μm -thick slices, which were mounted on the glass slides and stained by the routine H&E method. All slides were assessed in a blinded way by an independent pathologist for the presence of inflammatory cells (neutrophils, lymphocytes, monocytes) using an Axiolab 5 light microscope combined with an Axiocam 208 color camera (Zeiss, Jena, Germany) at 400x magnification in 40 high power fields (HPF).

Assessment of Serum Biochemical Parameters

The activities of alanine transaminase (ALT), aspartate transaminase (AST), and urea and creatinine levels in rat

Table 1 The Impact of Compounds **10b** and **13b** on Microscopic Changes of Paw Tissue and the Number of Inflammatory Cells in Exudate in H&E Staining. Indomethacin Was Used as a Reference Drug

Group	Microscopic Changes in H&E Staining (0–5 Points)	Number of Inflammatory Cells in 1 HPF of Exudate Sample
C	0	0
Car	4.20 ± 0.20***	40.10 ± 1.50***
Ind	1.80 ± 0.20***###	18.30 ± 1.25***###
10b-5	3.35 ± 0.2***^^	39.80 ± 0.68***^^
10b-10	3.10 ± 0.23***^^	36.30 ± 1.51***^^
10b-20	2.80 ± 0.25***^^	30.40 ± 1.12***###^^
13b-5	3.30 ± 0.21***^^	34.20 ± 1.65***^^
13b-10	2.90 ± 0.28***^^	21.60 ± 1.19***###
13b-20	2.00 ± 0.26***###	9.20 ± 1.38***###^^

Notes: Experimental groups: C – control group; Car – carrageenan group; Ind – group receiving 10 mg/kg indomethacin; **10b-5**, **10b-10**, and **10b-20** – groups receiving, respectively, 5, 10, or 20 mg/kg of compound **10b**; **13b-5**, **13b-10**, and **13b-20** – groups receiving, respectively, 5, 10, or 20 mg/kg of compound **13b**. Scoring scale of microscopic assessment of paw tissue damage: (0) = no inflammation; (1) = mild inflammation; (2) = mild or moderate inflammation; (3) = moderate inflammation; (4) = moderate or severe inflammation; (5) = severe inflammation. ***p<0.001 vs control group; #p<0.05, ###p<0.01, ####p<0.001 vs carrageenan group; ^p<0.05, ^^p<0.01, ^^p<0.001 vs indomethacin group.

Abbreviation: HPF, high power field.

serum were measured in a certified laboratory using Architect c4000 equipment and commercial ALT (Ref 8L92-20 8L92-40), AST (Ref 8L91-20 8L91-40), Urea (Ref 7D75-20 7D75-30) and Creatinine (Ref 7D64-20) Architect Abbott Kits (Abbott, Wiesbaden, Germany) following the manufacturer's instructions.

Macro- and Microscopic Assessment of Gastric Mucosa

The damage to gastric mucosa was evaluated macro- and microscopically. The severity of macroscopic mucosal changes was assessed using the 0-3-point scale described in the legend for Table 2, according to the criteria previously presented by Szabo et al.²⁵ Afterwards, formalin-fixed stomach specimens were

embedded in paraffin and cut into 4 µm thick sections, which were mounted on the glass slides and stained by the routine H&E method. Then, the microscopic analysis was performed using an Olympus BX53 light microscope combined with an Olympus UC90 camera (Olympus, Germany). Histopathological changes of stomach tissue sections were examined at 100x magnification and assessed using the 0-3-point scale described in the legend for Table 2.

Statistical Analysis

All data are presented as mean values ± standard error of the mean (SEM). The one-way analysis of variance (ANOVA) and multiple comparisons with Scheffe's post hoc test were used to analyze the statistical significance of differences among studied groups. The multi-criteria

Table 2 The Impact of Compounds **10b** and **13b** on Gastric Mucosa. Indomethacin Was Used as a Reference Drug

Group	Macroscopic Lesions (0–3 Points)	Microscopic Lesions in H&E Staining (0–3 Points)
C	0	0
Car	0	0
Ind	2.25 ± 0.19***	2.35±0.17***
10b-5	0 ^{^^}	0 ^{^^}
10b-10	0.15 ± 0.08 ^{^^}	0.20±0.08 ^{^^}
10b-20	0.53 ± 0.11 ^{^^}	0.36±0.11 ^{^^}
13b-5	0 ^{^^}	0 ^{^^}
13b-10	0 ^{^^}	0.05 0.05 ^{^^}
13b-20	0.34 ± 0.07 ^{^^}	0.32±0.07 ^{^^}

Notes: Experimental groups: C – control group; Car – carrageenan group; Ind – group receiving 10 mg/kg indomethacin; **10b-5**, **10b-10**, and **10b-20** – groups receiving, respectively, 5, 10, or 20 mg/kg of compound **10b**; **13b-5**, **13b-10**, and **13b-20** – groups receiving, respectively, 5, 10, or 20 mg/kg of compound **13b**. Scoring scale of macroscopic assessment of gastric mucosa: (0) = no damage; (1) = 1–4 small petechiae; (2) = 5 or more petechiae or hemorrhagic streaks up to 4 mm; (3) = erosions longer than 5 mm or confluent hemorrhages. Scoring scale of microscopic assessment of gastric mucosa: (0) = no damage; (1) = mild changes; (2) = moderate changes; (3) = severe changes. Data are presented as mean values ± SEM (n=12). ***p<0.001 vs control group; ^^p<0.001 vs indomethacin group.

decision analysis (MCDA) using the weighted sum model (WSM) was executed to compare the pharmacological and toxicological properties of the studied compounds. Equal weights were set for each bioassay performed. All statistical analyses were performed using GraphPad Prism version 8.0 (GraphPad Software, San Diego, CA, USA) and Statistica version 13.3 (StatSoft, Kraków, Poland) with a p -value < 0.05 considered as the significance level.

Results

The Effects of Pyrrolo[3,4-*d*]Pyridazinone Derivatives on the Carrageenan-Induced Paw Edema in Rats

To assess the anti-inflammatory activity of the novel pyrrolo[3,4-*d*]pyridazinone derivatives – compounds **10b** and **13b** – carrageenan-induced paw edema was performed. The absolute rat paw volume and relative increase in paw volume (paw edema) before and at 1, 2, 3, and 6 h after carrageenan or normal saline injection in all experimental groups are shown in Figures 1A and B, respectively. Detailed data of the absolute rat paw volume (mean values \pm SEM including the level of significance between groups) are presented in [Supplementary Material, Table S1](#). Detailed data of the paw edema (mean values \pm SEM including the level of significance between groups and the percentage of paw edema inhibition) are presented in [Table 3](#). There were no significant differences in rat paw volume before the carrageenan or normal saline injection between the experimental groups ($p = \text{NS}$). Subplantar injection of

0.1 mL of 1% carrageenan solution into the rat hind paw caused severe discernible inflammation with a significant increase in paw volume (paw edema) at 1, 2, 3, and 6 h after injection ($p < 0.001$ vs control group in all comparisons; [Figure 2A and B](#)). The maximum paw edema was observed in each studied group at 6 h after carrageenan administration. Pretreatment of rats with the studied compounds, **10b** or **13b**, resulted in an inhibition of paw edema starting from 2 h after carrageenan injection ([Figure 2D and E](#)). Pretreatment with compound **10b** at a dose of 10 or 20 mg/kg partly reversed the carrageenan-induced increase in paw volume at 2, 3, and 6 h ($p < 0.001$ vs carrageenan group and $p < 0.001$ vs control group in all comparisons). At the low dose (5 mg/kg), compound **10b** partly inhibited the carrageenan-induced edema only at 2 h ($p < 0.05$ vs carrageenan group and $p < 0.001$ vs control group). At all examined doses, compound **13b** partly reversed the increase in paw volume at 2, 3, and 6 h after carrageenan injection ($p < 0.001$ vs carrageenan group for 10 and 20 mg/kg dose; $p < 0.01$, $p < 0.001$, and $p < 0.05$ vs carrageenan group for 5 mg/kg dose at 2, 3, and 6 h, respectively). For all studied doses of compound **13b** at 2, 3, and 6 h the differences versus the control group were still significant ($p < 0.001$ in all cases). Administration of reference drug indomethacin before carrageenan injection also caused a significant inhibition of paw edema ([Figure 2C](#)) at 2, 3, and 6 h of carrageenan test, wherein at 3 h indomethacin completely reversed the increase in paw volume ($p < 0.001$ vs carrageenan group; $p = \text{NS}$ vs control group) and at 2 and 6 h indomethacin partly




Figure 1 The impact of compounds **10b** and **13b** on the paw volume (**A**) and the increase in paw volume (paw edema) (**B**) after carrageenan injection. Indomethacin was used as a reference drug. Experimental groups: C – control group receiving 0.5% CMC intragastrically (i.g.) and normal saline subplantarly (s.pl.); Car – carrageenan group receiving 0.5% CMC i.g. and 1% carrageenan solution s.pl.; Ind – indomethacin group receiving 10 mg/kg of indomethacin i.g. and 1% carrageenan solution s.pl.; **10b**-5, **10b**-10, and **10b**-20 – groups receiving, respectively, 5, 10, or 20 mg/kg of investigated compound **10b** i.g. and 1% carrageenan solution s.pl.; **13b**-5, **13b**-10, and **13b**-20 – groups receiving, respectively, 5, 10, or 20 mg/kg of investigated compound **13b** i.g. and 1% carrageenan solution s.pl. Data are presented as mean values \pm SEM ($n=12$).

Table 3 The Impact of Compounds **10b** and **13b** on the Increase in Paw Volume (Paw Edema) and the Percentage of Paw Edema Inhibition After Carrageenan Injection. Indomethacin Was Used as a Reference Drug

Group	1 h	2 h	3 h	6 h
	Increase in Paw Volume (Paw Edema; mL) Paw Edema Inhibition (%)			
C	0.07 ± 0.05	0.09 ± 0.04	0.15 ± 0.04	0.11 ± 0.04
Car	0.56 ± 0.02***	1.16 ± 0.03***	1.46 ± 0.03***	2.01 ± 0.03***
Ind	0.37 ± 0.05 (33.56%)*	0.41 ± 0.06 (64.36%)*###	0.42 ± 0.06 (71.22%)*###	0.71 ± 0.06 (64.41%)*###
10b -5	0.53 ± 0.04 (4.36%)*	0.88 ± 0.05 (24.18%)*###	1.31 ± 0.05 (10.02%)*###	1.64 ± 0.06 (18.01%)*###
10b -10	0.49 ± 0.04 (12.62%)*	0.74 ± 0.03 (36.27%)*###	0.86 ± 0.03 (40.78%)*###	1.24 ± 0.05 (37.74%)*###
10b -20	0.43 ± 0.06 (22.54%)*	0.57 ± 0.05 (50.88%)*###	0.62 ± 0.04 (57.53%)*###	1.04 ± 0.05 (48.09%)*###
13b -5	0.51 ± 0.04 (8.77%)*	0.83 ± 0.04 (28.45%)*###	1.02 ± 0.05 (29.98%)*###	1.59 ± 0.07 (20.50%)*###
13b -10	0.47 ± 0.04 (15.86%)*	0.66 ± 0.05 (42.90%)*###	0.68 ± 0.05 (53.43%)*###	1.00 ± 0.08 (50.25%)*###
13b -20	0.37 ± 0.03 (33.15%)*	0.49 ± 0.04 (57.56%)*###	0.55 ± 0.04 (62.33%)*###	0.76 ± 0.04 (62.19%)*###


Notes: Experimental groups: C – control group; Car – carrageenan group; Ind – group receiving 10 mg/kg indomethacin; **10b**-5, **10b**-10, and **10b**-20 – groups receiving, respectively, 5, 10, or 20 mg/kg of compound **10b**; **13b**-5, **13b**-10, and **13b**-20 – groups receiving, respectively, 5, 10, or 20 mg/kg of compound **13b**. Data are presented as mean values ± SEM (n=12). *p<0.05, **p<0.01, ***p<0.001 vs control group; #p<0.05, ##p<0.01, ###p<0.001 vs carrageenan group; ^p<0.01, ^^p<0.001 vs indomethacin group.

reversed the carrageenan-induced paw edema (p<0.001 vs carrageenan group; p<0.01 and p<0.001 vs control group, respectively). Neither the studied compounds nor indomethacin exerted significant activity at 1 h after carrageenan injection (p = NS vs carrageenan group and p<0.001 vs control group in all cases). The maximum inhibitory effect in the treated groups compared to the carrageenan group was observed at 3 h of carrageenan test after administration of indomethacin at a dose of 10 mg/kg (71.2% inhibition of paw edema) as well as after the administration of compound **10b** or **13b** at the high dose (57.5% and 62.3%, respectively). Moreover, pretreatment with compound **10b** or **13b** at the high dose inhibited paw edema at all studied time points in a manner comparable to that of indomethacin. There were no significant differences in paw edema between the group receiving 20 mg/kg of compound **10b** or **13b** and the indomethacin-receiving group (p = NS). The action of compound **13b** given at a dose of 10 mg/kg

at 2 h was also comparable to that provided by indomethacin (p = NS).

The Effects of Pyrrolo[3,4-d]Pyridazinone Derivatives on PGE₂, TNF-α, and MPO Levels in the Carrageenan-Injected Rat Paw

The enzyme-linked immunosorbent assay tests were performed to evaluate the effects of compounds **10b** and **13b** on the levels of PGE₂, TNF-α, and MPO, the increase of which takes place in the inflammatory response. The concentrations of these proinflammatory parameters (mean ± SEM) in paw tissue are presented in Figure 3A–C. Injection of carrageenan into the rat paw caused a significant increase in the PGE₂, TNF-α, and MPO levels in paw tissue in comparison to the control group (p<0.001, p<0.001, and p<0.01, respectively). Pretreatment with compound **10b** at a dose of 10 mg/kg or 20 mg/kg prevented the increase in the PGE₂ level in paw tissue compared to the

**Figure 2** The effects of compounds **10b** and **13b** on the carrageenan-induced paw edema in rats. Indomethacin was used as a reference drug. Experimental groups: control group (A); carrageenan group (B); group receiving 10 mg/kg indomethacin (C); group receiving 20 mg/kg compound **10b** (D); group receiving 20 mg/kg compound **13b** (E).




Figure 3 The impact of compounds **10b** and **13b** on PGE₂ (**A**), TNF-α (**B**), and MPO (**C**) concentrations in paw tissue. Indomethacin was used as a reference drug. Experimental groups: C – control group receiving 0.5% CMC intragastrically (i.g.) and normal saline subplantarily (s.pl.); Car – carrageenan group receiving 0.5% CMC i.g. and 1% carrageenan solution s.pl.; Ind – indomethacin group receiving 10 mg/kg of indomethacin i.g. and 1% carrageenan solution s.pl.; **10b**-5, **10b**-10, and **10b**-20 – groups receiving, respectively, 5, 10, or 20 mg/kg of investigated compound **10b** i.g. and 1% carrageenan solution s.pl.; **13b**-5, **13b**-10, and **13b**-20 – groups receiving, respectively, 5, 10, or 20 mg/kg of investigated compound **13b** i.g. and 1% carrageenan solution s.pl. Data are presented as mean values ± SEM (n=12). Differences *p<0.05 vs control group; **p<0.01 vs control group; ***p<0.001 vs control group; #p<0.05 vs carrageenan group; ##p<0.01 vs carrageenan group; ###p<0.001 vs carrageenan group; ^p<0.05 vs indomethacin group were deemed statistically significant.

group of carrageenan-injected animals ($p<0.01$ and $p<0.001$, respectively). PGE₂ concentrations in these two treated groups were not different from the control group ($p = \text{NS}$). Administration of compound **10b** at a dose of 10 mg/kg or 20 mg/kg partly reversed the alteration in TNF-α tissue concentration ($p<0.05$, $p<0.001$ vs carrageenan group; $p<0.001$, $p<0.05$ vs control group, respectively). The low dose of compound **10b** (5 mg/kg) did not exert significant activity against the carrageenan-induced increase in PGE₂ and TNF-α tissue level ($p = \text{NS}$ vs carrageenan group; $p<0.01$ and $p<0.001$ vs control group, respectively). The concentrations of MPO in groups receiving compound **10b** in all studied doses were not significantly different from that in the carrageenan group, but they did not differ significantly from the control group either ($p = \text{NS}$). Pretreatment with compound **13b** at a dose of 10 mg/kg or 20 mg/kg

counteracted the increased PGE₂, TNF-α, and MPO levels in paw tissue compared to the group of carrageenan-injected animals ($p<0.01$ for PGE₂ for the 10 mg/kg dose and $p<0.001$ for others). The difference to the control group was insignificant in all these cases ($p = \text{NS}$). Administered at the low dose (5 mg/kg), compound **13b** protected from the increase of MPO concentration ($p<0.01$ vs carrageenan group and $p = \text{NS}$ vs control group), partly reversed the altered TNF-α level ($p<0.01$ vs carrageenan and control group) and did not affect PGE₂ concentration ($p = \text{NS}$ vs carrageenan group and $p<0.001$ vs control group). Indomethacin (10 mg/kg) given before carrageenan injection prevented the increase of PGE₂ and MPO tissue levels ($p<0.001$, $p<0.01$ vs carrageenan group, $p = \text{NS}$ vs control group) and partly reversed the increased TNF-α level ($p<0.01$ vs carrageenan group and $p<0.001$ vs control group). Compound **10b** or **13b**

administered at a medium or high dose normalized the PGE₂ concentration in a manner comparable to that of indomethacin ($p = \text{NS}$). Following pretreatment with compound **10b** or **13b** at each dose tested, the TNF- α and MPO levels were normalized in a manner comparable to that of indomethacin ($p = \text{NS}$). Additionally, compound **13b** given at a dose of 20 mg/kg counteracted the increased TNF- α level more effectively than indomethacin ($p < 0.05$), causing a nearly 2-fold greater decrease in the TNF- α level in comparison to indomethacin (59.42% vs 31.20%).

The Effects of Pyrrolo[3,4-*d*]Pyridazinone Derivatives on Carrageenan-Induced Histopathological Alterations in Paw

To investigate whether the studied compounds counteract carrageenan-induced changes in the paw tissue, histopathological analysis of paw tissue was performed. Table 1 shows the scoring of paw tissue samples. Microscopic examination of paw tissue of carrageenan-injected rats showed massive inflammation with pronounced interstitial and intermuscular edema, inflammatory cell infiltration, and loss of normal muscle paw architecture (Figure 4B) compared to the control group with no histological damage ($p < 0.001$;

Figure 4A). Infiltration of inflammatory cells, including enormous numbers of neutrophils, lymphocytes, and sparse mast cells, was localized in connective tissue and between muscle bundles (Figure 4B). Following pretreatment with the studied compounds at a dose of 10 or 20 mg/kg, there was an explicit recovery of carrageenan-injected paw tissue with decreased edema and dispersion of muscle bundles and reduced inflammatory cell infiltration (Figure 4E, F, H and I). Scoring of paw tissue samples indicated that in comparison to the carrageenan group, compound **10b** or **13b** administered at the medium or high dose remarkably diminished the carrageenan-induced alterations in paw tissue ($p < 0.05$ and $p < 0.01$ for compound **10b**; $p < 0.01$ and $p < 0.001$ for compound **13b**, respectively). The tissue sections from the hind paws of rats receiving compound **13b** at a dose of 20 mg/kg or indomethacin showed a weak inflammatory reaction with nearly normal paw tissue histological architecture (Figure 4C and I) while the effect of compound **13b** at the high dose was comparable to that of indomethacin ($p = \text{NS}$). Pretreatment with compound **10b** or **13b** at the low dose (5 mg/kg) showed only a slight improvement in edema formation and inflammatory cell infiltration (Figure 4D and G), and this effect was insignificant when compared to the carrageenan group ($p = \text{NS}$).




Figure 4 The photomicrographs of paw tissue after hematoxylin-eosin staining demonstrated that compounds **10b** and **13b** diminished histological alterations induced by carrageenan injection. Indomethacin was used as a reference drug. Experimental groups: control group (A); carrageenan group (B); group receiving 10 mg/kg indomethacin (C); group receiving 5 mg/kg compound **10b** (D); group receiving 10 mg/kg compound **10b** (E); group receiving 20 mg/kg compound **10b** (F); group receiving 5 mg/kg compound **13b** (G); group receiving 10 mg/kg compound **13b** (H); group receiving 20 mg/kg compound **13b** (I); magnification 200 \times .

The Effects of Pyrrolo[3,4-*d*]Pyridazinone Derivatives on Carrageenan-Induced Paw Exudate

To assess the composition of the inflammatory infiltration, a histopathological examination of the exudate of carrageenan-injected paws was performed. Table 1 shows the scoring of paw exudate samples. Carrageenan injection into the rats' right hind paws caused massive infiltration of inflammatory cells, significantly greater than in the control group ($p < 0.001$; Figure 5A and B). The cellular composition of the exudate comprised enormous number of polymorphonuclear leukocytes (neutrophils), lymphocytes, and few monocytes (Figure 5B). Pretreatment with compound **10b** at the low or medium dose did not affect the composition of the exudate ($p = \text{NS}$, Figure 5D and E) and only at the high dose (20 mg/kg) reduced the number of inflammatory cells ($p < 0.001$, Figure 5F), while the administration of compound **13b** at each studied dose reduced the number of inflammatory cells in comparison to the carrageenan group ($p < 0.05$, $p < 0.001$, and $p < 0.001$, respectively; Figure 5G–I). The effect of compound **13b** at the medium dose was comparable to that provided by a reference drug, indomethacin ($p = \text{NS}$), and at the high

dose was even greater than that of indomethacin ($p < 0.001$, Figure 5C, H and I).

The Effects of Pyrrolo[3,4-*d*]Pyridazinone Derivatives on the Liver and Kidney Function Parameters

The serum activities of ALT and AST, as well as serum urea and creatinine concentrations, were measured to investigate the function of liver and kidney in rats pretreated with a single dose of pyrrolo[3,4-*d*]pyridazinone derivatives. These results as mean \pm SEM are presented in Table 4. Local administration of carrageenan did not significantly increase the serum ALT or AST activities and did not change the serum urea or creatinine levels in comparison to the control group ($p = \text{NS}$). Intragastrical administration of either the **10b** or **13b** compound at all studied doses (5, 10, or 20 mg/kg) did not significantly alter the ALT or AST activities as well as the urea or creatinine levels in comparison to the control group ($p = \text{NS}$), except for compound **10b** given at the low dose (5 mg/kg) which significantly decreased the ALT activity ($p < 0.05$ vs control). After a single 10 mg/kg intragastrical dose, indomethacin significantly increased the serum AST activity and serum urea concentration compared to the control group ($p < 0.05$).




Figure 5 The photomicrographs of exudate of carrageenan-injected paw after hematoxylin-eosin staining demonstrated that the studied compounds reduced inflammatory cell infiltration. Indomethacin was used as a reference drug. Experimental groups: control group (A); carrageenan group (B); group receiving 10 mg/kg indomethacin (C); group receiving 5 mg/kg compound **10b** (D); group receiving 10 mg/kg compound **10b** (E); group receiving 20 mg/kg compound **10b** (F); group receiving 5 mg/kg compound **13b** (G); group receiving 10 mg/kg compound **13b** (H); group receiving 20 mg/kg compound **13b** (I); magnification 400 \times .

Table 4 The Effects of Compounds **10b** and **13b** on the Serum Biochemical Parameters in Rats. Indomethacin Was Used as a Reference Drug

Group	ALT [U/l]	AST [U/l]	Urea [mg/dl]	Creatinine [mg/dl]
C	36.00 ± 1.92	118.75 ± 3.97	36.38 ± 0.82	0.21 ± 0.02
Car	37.00 ± 2.24	132.00 ± 7.34	39.63 ± 0.94	0.23 ± 0.02
Ind	37.63 ± 2.54	151.25 ± 10.24*	44.63 ± 3.83*	0.21 ± 0.03
10b -5	27.00 ± 0.96*	115.13 ± 2.95	32.13 ± 1.55	0.25 ± 0.02
10b -10	28.50 ± 0.63	116.50 ± 3.95	35.13 ± 1.37	0.21 ± 0.01
10b -20	32.38 ± 2.66	129.75 ± 4.05	37.00 ± 1.51	0.21 ± 0.02
13b -5	28.88 ± 1.09	120.50 ± 4.67	30.13 ± 0.90	0.24 ± 0.01
13b -10	31.38 ± 1.80	124.38 ± 3.50	30.50 ± 0.78	0.24 ± 0.01
13b -20	31.75 ± 1.18	132.50 ± 2.90	34.25 ± 1.11	0.24 ± 0.01

Notes: Experimental groups: C – control group; Car – carrageenan group; Ind – group receiving 10 mg/kg indomethacin; **10b**-5, **10b**-10, and **10b**-20 – groups receiving, respectively, 5, 10, or 20 mg/kg of compound **10b**; **13b**-5, **13b**-10, and **13b**-20 – groups receiving, respectively, 5, 10, or 20 mg/kg of compound **13b**. Data are presented as mean values ± SEM (n=12). *p<0.05 vs the control group.

The Effects of Pyrrolo[3,4-*d*]Pyridazinone Derivatives on Gastric Mucosa in Macro- and Microscopic Assessment

Macro- and microscopic evaluation was carried out to characterize the gastric safety profile of the compounds studied. The presence and severity of macroscopically visible mucosal lesions (petechiae, hemorrhagic erosions) were scored as indicators of ulcerogenic activity (Table 2). There was no injury to gastric mucosa in the control or carrageenan group (Figure 6A and B). The findings of these ulcerogenic liability studies demonstrated that novel derivatives in all studied

doses caused negligible mucosal lesions as compared to the control group (p = NS, Figure 6D–G), whereas indomethacin at a dose of 10 mg/kg caused significant mucosal injuries ranging from hyperemia to hemorrhagic lesions covered with coagulated blood (p<0.001; Figure 6C). The macroscopic changes were reflected in the microscopic appraisal (Table 2). The stomach tissue of rats pretreated with the studied compounds as well as control rats showed no histopathological changes with intact mucosa, submucosa, and musculosa (p = NS; Figure 7A, F, G, H and I). Subplantar administration of carrageenan caused no significant injury to




Figure 6 The macroscopic appearance of the gastric mucosa revealed that the studied compounds caused negligible mucosal lesions. Indomethacin was used as a reference drug. Experimental groups: control group (A); carrageenan group (B); group receiving 10 mg/kg indomethacin (C); group receiving 10 mg/kg compound **10b** (D); group receiving 20 mg/kg compound **10b** (E); group receiving 10 mg/kg compound **13b** (F); group receiving 20 mg/kg compound **13b** (G).




Figure 7 The microphotographs of stomach tissue after hematoxylin-eosin staining revealed that the studied compounds neither altered the normal stomach tissue architecture nor induced ulcerations. Indomethacin was used as a reference drug. Experimental groups: control group (A); carrageenan group (B); group receiving 10 mg/kg indomethacin (C–E); group receiving 10 mg/kg compound **10b** (F); group receiving 20 mg/kg compound **10b** (G); group receiving 10 mg/kg compound **13b** (H); group receiving 20 mg/kg compound **13b** (I); magnification 100 \times .

the stomach tissue in microscopic analysis ($p = \text{NS}$; Figure 7B). Meanwhile, following indomethacin administration, the stomach tissue was characterized by appreciable damage to the protective mucosal layer with local thinning of the mucosa and damage to the superficial layer with some visible crater-like cavities, focal necrosis of gastric mucosa, submucosal edema, and congestion of mucosal and submucosal blood vessels ($p < 0.001$; Figure 7C–E).

Multi-Criteria Decision Analysis

The results obtained from each bioassay (carrageenan-induced paw edema test, ELISA, microscopic assessment of paw tissue and paw exudate, assessment of serum biochemical parameters, and macro- and microscopic assessment of gastric mucosa) were analyzed by MCDA to compare the studied pharmacological and toxicological properties, as well as the risks and benefits of pretreatment with the new compounds. The MCDA results (Figure 8) indicated that the most favorable profile of action, ie, the strongest anti-inflammatory effect with the lowest risk of hepatic, renal, and gastric toxicity, was found for compound **13b** at a dose of 20 mg/kg. Moreover, compounds **10b** and **13b** exerted a more favorable effect than indomethacin when administered at both the same dose as indomethacin (10 mg/kg) and at a higher dose than indomethacin (20 mg/kg). In the studied dose range, compound **13b** exerted a more favorable effect than compound **10b**.




Figure 8 Multi-criteria decision analysis (MCDA) of the anti-inflammatory effect and hepatic, renal, and gastric toxicity of the studied compounds **10b** and **13b**.

Discussion

The word carrageenan, derived from the Irish word “carragin” meaning Irish moss (*Chondrus crispus* L.), refers not only to this species of algae but also to its mucopolysaccharide extract, discovered by the British pharmacist Stanford in 1862.²⁶ This mucopolysaccharide extract consists of several types of carrageenans, including lambda carrageenan, which is used in an experimental model of acute inflammation as a potent inflammatory-triggering agent.^{26,27} Carrageenan-induced inflammation (also called carrageenan-induced paw edema), originally described by Winter,²² is a valuable tool widely used to assess the potential anti-

inflammatory activity of any novel substance.^{27–29} It is noteworthy that inhibition of carrageenan-induced inflammation is highly predictive of anti-inflammatory drug activity in human inflammatory disorders. Moreover, the doses of anti-inflammatory agents in this model correlate well with effective doses in patients.^{30,31} Thus, this model has a vital role in novel drug development.

Carrageenan injection into the rat hind paw elicits an inflammatory response in a biphasic manner. The first phase (0–1 h after injection) mediated by histamine, serotonin, and bradykinin is followed by a second phase (2–6 h after injection), which is attributed to infiltration of polymorphonuclear leukocytes, mainly neutrophils, and production of prostaglandins (PGs), especially PGE₂ and various proinflammatory cytokines such as IL-1, IL-6, and TNF- α .^{32,33} The first phase causes local vasodilatation and rapid increase in the vascular permeability with consequent edema formation, and the second phase enables edema maintenance due to PGs' potent vasoactive action and their ability to recruit inflammatory cells. Therefore, this animal model, as it happened in this study, displays hallmark signs of acute inflammation (redness, edema, heat, pain, loss of function), which develop instantaneously following subplantar injection of carrageenan solution into the rat hind paw. In carrageenan-induced inflammation, the inflammatory response is quantified by an increase in paw volume (paw edema), which can be modulated by inhibitors of specific molecules within the inflammatory cascade.²⁷ In the present study, the inflammatory response indicated by the paw edema lasted up to 6 h following carrageenan injection. Both studied compounds given at a dose of 10 or 20 mg/kg partly inhibited paw edema starting at 2 h after carrageenan injection, and this effect peaked at 3 h and continued for up to 6 h. At the low dose (5 mg/kg), compound **10b** partly suppressed paw edema only at 2 h, while compound **13b** at 2, 3, and 6 h after carrageenan injection. Since novel pyrrolo [3,4-*d*]pyridazinone derivatives suppressed the second phase of carrageenan-induced paw edema, it can be assumed that they exert an anti-inflammatory effect by inhibiting the release of prostaglandins and proinflammatory cytokines and reducing inflammatory cell infiltration. The action of compounds **10b** and **13b** given at the high dose was comparable to the effect provided by indomethacin, which also decreased edema at 2, 3, and 6 h with maximal inhibitory effect at 3 h. At 3 h, indomethacin completely reversed the increase in paw volume, whereas at 6 h it reversed the increase only partly. This diminution

of inhibitory effect at 6 h can be explained by the fact that COX-2 expression increases after 3 h following carrageenan injection and reaches maximal expression at 6 h.³⁴ In our previous work, we demonstrated that new pyrrolo [3,4-*d*]pyridazinone derivatives strongly inhibit cyclooxygenase with a better affinity towards COX-2 isoform, and some of them, including compound **13b**, act as selective COX-2 inhibitors.¹² Thus, compound **10b** as a preferential COX-2 inhibitor and indomethacin as COX-1/COX-2 inhibitor showed a greater diminution in the inhibitory effect than compound **13b**, a selective COX-2 inhibitor, whose diminution of inhibitory effect at 6 h was the lowest. Neither the new compounds nor indomethacin inhibited the early phase of edema formation. This is consistent with findings reported by other authors who have shown that the second phase of edema formation is sensitive to non-steroidal anti-inflammatory agents, including indomethacin.^{28,35}

Although many components have been implicated in the course of the inflammatory response, a lot of studies placed prostanoids synthesis, especially PGE₂, as a crucial element and important link in the chain of events leading to the carrageenan-invoked inflammatory response.^{28,36,37} PGE₂ is well established as a mediator of acute inflammation, which regulates multiple aspects of inflammation and multiple functions of different immune cells. PGE₂ can promote local vasodilatation, fluid and protein extravasation, and the local attraction of neutrophils and macrophages from the bloodstream to the site of tissue injury and their activation, thereby leading to general symptoms of inflammation.⁷ Nevertheless, this molecule is also involved in the detrimental transition from acute to chronic inflammation and its maintenance, which may lead to chronic inflammatory diseases. PGE₂ can convert a short-term inflammatory response into a long-term process, primarily by enhancing proinflammatory cytokine release cascade, contributing to the differentiation and activation of Th1 and Th17 cells, and contributing to aberrant tissue remodeling.^{38,39} Thus, normalization of the increased PGE₂ levels may be crucial not only in the removal of symptoms of acute inflammation but also in preventing acute inflammation from becoming a chronic inflammatory state. As aforementioned, a decrease in the PGE₂ level may contribute to the attenuation of the second phase of carrageenan-induced inflammation. In the current study, we have indicated that pretreatment with both novel 1,3,4-oxadiazole derivatives of pyrrolo[3,4-*d*]pyridazinone at the medium or high dose counteracted the increased

PGE₂ level in paw tissue compared to the group of carrageenan-injected rats, and this action was comparable to that of indomethacin. This is in agreement with the results described by Ozyazici et al, showing that some other 1,3,4-oxadiazole derivatives based on compounds with known anti-inflammatory action reduced PGE₂ production in LPS-stimulated RAW 264.7 cells similarly to indomethacin.⁴⁰ It is known that PGE₂ constitutes a potent sensitizing agent able to modulate the nociceptive pathway via peripheral and central mechanisms and is regarded as an essential mediator of hyperalgesia.⁷ In our previous study, we found that at the medium and high doses, compound **13b**, but not **10b**, has an antinociceptive effect, which may be due to the inhibition of nociceptor sensitization by decreasing the PGE₂ level.¹⁸ This discrepancy between the action of compound **10b** on the PGE₂ level reported in our earlier paper¹⁸ and the current study may result from the fact that while both cyclooxygenase isoforms are involved in the inflammatory response, COX-2 is the prevalent isoform involved in the nociceptor sensitization and hyperalgesia.⁷

In addition to prostaglandins, proinflammatory cytokines are the other molecules contributing to the development of edema during the second and subsequent hours after injection of carrageenan.³³ One of the most important cytokines that govern the development and maintenance of inflammation is TNF- α .⁴¹ TNF- α can activate macrophages and upregulate other proinflammatory cytokines and endothelial adhesion molecules, thus promoting the adhesion of neutrophils and lymphocytes to endothelial cells and their extravasation.^{2,42,43} Moreover, TNF- α contributes to tissue damage and multiorgan failure and is considered an important mediator of the development of various chronic inflammatory diseases.⁴² In the current study, inhibition of the inflammatory response by the new compounds was accompanied by a decrease in the TNF- α level. Compound **10b** at the medium or high dose partly reversed alteration in TNF- α tissue concentration as compared with carrageenan-injected rats, while compound **13b** at the same doses completely counteracted the increased TNF- α level in paw tissue. It is noteworthy that the action of compound **13b** given at the high dose was even greater than the effect of indomethacin. Moreover, we found that, contrary to compound **10b** and indomethacin, the suppressive effect of compound **13b** on TNF- α release was greater than its effect on PGE₂, which supports the possibility that compound **13b** has a greater inhibitory effect on proinflammatory cytokines release

than either compound **10b** or indomethacin. Our results are in line with previous findings showing that 1,3,4-oxadiazole derivatives can decrease the tissue TNF- α level in the carrageenan-induced paw edema test.⁴⁴ Likewise, Mogilski et al¹¹ reported that pyrrolo[3,4-*d*]pyridazinone derivatives decreased the TNF- α level in LPS-activated RAW264.7 macrophage, a cell line frequently used for the screening of anti-inflammatory activity of new compounds.⁴²

Local neutrophil infiltration and activation also contribute to the inflammation caused by tissue damage,⁴⁵ including carrageenan-induced inflammation.⁴² After carrageenan injection, neutrophils are both the first and the primary cells recruited to the site of inflammation; their infiltration can be identified by measuring the levels of myeloperoxidase, one of the major enzymes released from activated neutrophils.^{5,42} Interestingly, accumulating evidence indicates that MPO serves not only as an index of neutrophil infiltration into inflamed tissues but also displays cytokine-like properties.⁵ Proinflammatory properties of MPO are independent of its enzymatic and bactericidal activity and occur as a result of its ability to interact with an integrin Mac-1 (CD11b/CD18). The linking of MPO to Mac-1 leads to the modulation of intracellular neutrophil signaling pathways and thereby to neutrophil activation and extravasation. Indeed, MPO evokes a neutrophil response similar to those triggered by proinflammatory cytokines, especially TNF- α .⁵ Moreover, MPO may increase Mac-1 expression, cause further MPO release from neutrophils, and delay their intrinsic apoptosis.^{46,47} This MPO-dependent feed-forward loop amplifies the response of neutrophils, thereby prolonging inflammation, causing local tissue damage, and leading to chronic inflammatory conditions.⁴⁶ As such, the reduction of the MPO level is regarded as another crucial condition to alleviate the second phase of carrageenan-induced inflammatory response.⁴⁸ In the present study, we indicated that compared to the carrageenan-injected rats, pretreatment with compound **13b** at all doses tested counteracted the increased MPO level in paw tissue, and this effect was similar to that provided by indomethacin. The achieved results concur well with our earlier findings that pretreatment with compound **13b**, but not **10b**, at the doses of 10 and 20 mg/kg decreased the MPO level in the inflammatory phase of the formalin test.¹⁸ Moreover, Cidade et al⁴⁴ demonstrated that some other 1,3,4-oxadiazole derivatives decreased the MPO level in the carrageenan-induced paw edema test.

The action of the above-mentioned inflammatory mediators, PGE₂, TNF- α , and MPO, results in plasma exudation and migration of peripheral blood leukocytes into the injured area.^{2,5,7,42} It has been pointed out that after the injection of carrageenan, the total number of exudate leukocytes increases with time, and over 96% of exudate leukocytes are neutrophils.⁴⁹ Palaska et al⁵⁰ found that various 1,3,4-oxadiazole-2-thione derivatives reduced the total number of exudate leukocytes. In our study, histological evaluation of paw exudates showed that subplantar carrageenan injection caused massive infiltration of inflammatory cells, including neutrophils, lymphocytes, and monocytes. Compared to the carrageenan group, pretreatment with compound **10b** at the high dose or with compound **13b** at each studied dose reduced the influx of all inflammatory cells into the inflamed tissue. The effect of compound **13b** at the medium dose was comparable to that elicited by the reference drug, indomethacin, and at the high dose was even greater than that of indomethacin. The more pronounced effect exhibited by compound **13b** could be explained by the fact that, contrary to compound **10b**, compound **13b** decreased the MPO level, leading to the interruption of the MPO-dependent loop with a consequent reduction in neutrophil activation and migration into the inflamed tissue.

The histological analysis of the paw tissue indicated that carrageenan-induced inflammation is linked to intense edema characterized by increased infiltration of inflammatory cells and loss of normal muscle architecture in the inflamed paw tissue, consistent with the findings reported by other authors.⁵¹ The results obtained in our study show that both compounds at a dose of 10 or 20 mg/kg decreased the edema formation and the elevated level of cellular infiltration in rat paw tissue induced by carrageenan injection. Histopathology of the paw tissue also revealed that pretreatment with compound **13b** at the high dose resulted in a nearly normal paw tissue architecture with minimal inflammation and an effect comparable to that of indomethacin. These data support the results of paw edema measurement, changes in biochemical parameters, and histology of exudates and verify and confirm the anti-inflammatory effect of the studied compounds against acute inflammation. To our best knowledge, our study is the first to evaluate the effect of pyrrolo[3,4-*d*]pyridazinone derivatives on inflammatory cell infiltration and tissue alterations induced by carrageenan injection in experimental inflammation in rats, indicating that pretreatment with these new compounds prevented not only the

increase in inflammatory cell migration into the inflamed tissue but also morphological alterations.

The liver and kidneys are important essential organs for drug metabolism, storage, and excretion, making them particularly vulnerable to drug-related damage. Moreover, inhibition of COX-1 and COX-2 derived prostaglandins by most commonly used anti-inflammatory drugs, ie, NSAIDs, may cause serious adverse effects, eg, gastrointestinal irritation, erosions, ulceration, and bleeding as well as glomerular filtration impairment, and sodium and water retention.^{9,52,53} It is noteworthy that it is not only the NSAIDs' mechanism of action but also their chemical structure that may contribute to their toxicity.⁹ Typical NSAIDs consist of an acidic moiety linked to an aromatic functional group. The majority of NSAIDs contain a free carboxyl group as an acidic moiety, which may elicit gastric mucosal damage by local irritation in the ion trapping mechanism and through a reduction of mucosal surface hydrophobicity.⁵⁴ The chemical structure of NSAIDs may also contribute to their hepatotoxicity. Although the exact mechanism of NSAIDs' hepatotoxicity is not completely understood, it is assumed that an acidic group of NSAIDs or reactive adducts of NSAID metabolites may interact with host proteins and lead to cellular injury in susceptible patients.⁵⁵ Considering that the tested compounds act by inhibiting the COX pathway, it seems justified to make a preliminary assessment of the effect of pretreatment with a single dose of these new compounds on the markers of toxicity to the liver, kidneys, and gastric mucosa. An increase in serum ALT and AST activities may be interpreted as a result of hepatocyte injury or changes in their membrane permeability, indicating severe hepatocellular damage. Additionally, ALT and AST are useful indicators for identifying inflammation or necrosis of the liver.⁵⁶ Thereby, serum ALT and AST activity are considered predictors of possible hepatic toxicity.⁵⁷ In some studies, elevated levels of ALT have been observed already several hours after administration of a single dose of NSAID.^{58,59} Physiologically, urea and creatinine are filtered out of the blood by the kidneys. In renal disorders, when the kidneys are unable to excrete urea and creatinine properly, they are retained in the blood, and their levels become elevated.⁵⁶ Urea and creatinine levels are thus regarded as biomarkers predicting possible renal toxicity.⁶⁰ The results of the present study revealed that, unlike indomethacin, pretreatment with compound **10b** or **13b** – even at a high dose – did not alter ALT and AST activities or urea and creatinine levels compared

to the control rats. This suggests that these new compounds are not hepato- or nephrotoxic at the doses tested. As far as we know, this work is the first to evaluate the effect of 1,3,4-oxadiazole derivatives on hepatic and renal toxicity markers. It cannot be excluded that the replacement of the free carboxyl group with 1,3,4-oxadiazole-2-thione may contribute to reducing the hepatotoxicity of the designed compounds. Similarly, in contrast to indomethacin, which caused mucosal injury ranging from hyperemia to hemorrhagic lesions, both studied compounds given in all studied doses caused negligible macroscopic mucosal lesions and no histopathological changes, with the mucosa, submucosa, and musciosa remaining intact. In the previously published paper,¹⁸ the same compounds – **10b** and **13b** – were superior in gastric safety profile to indomethacin in macro- and microscopic evaluation in mice. This implies greater safety of our newly synthesized compounds on the gastric mucosa. Our results corroborated with other authors' findings showing that a 1,3,4-oxadiazole-2-thione ring incorporated into a given structure results in lower gastrotoxicity.^{13–15} Presumably, the conversion of the free carboxyl group to the five-membered 1,3,4-oxadiazole-2-thione, bioisosteric ring results in no local irritation and increases selectivity to COX-2, thereby reducing gastrotoxicity. More detailed research is needed to evaluate the safety profile of these two new compounds after repeated administration since many anti-inflammatory agents are used for long-term treatment.

Conclusion

The results provided herein point to the conclusion that the novel pyrrolo[3,4-*d*]pyridazinone derivatives, **10b** and **13b**, exert anti-inflammatory activity, and their mechanism of action might be related to the decrease of the PGE₂, TNF- α , and MPO levels and the reduction of inflammatory cell infiltration in inflamed tissues. Even though the new pyrrolo[3,4-*d*]pyridazinone derivatives are slightly less effective than the reference drug indomethacin, they do not cause gastric mucosal injuries and hepato-, or nephrotoxicity like it does, which is their indisputable advantage. Based on the findings of multi-criteria decision analysis, it may be inferred that the new compounds, **10b** and **13b**, have a more favorable benefit–risk profile than indomethacin. Compound **13b** at the high dose (20 mg/kg) reveals the most favorable benefit–risk profile among all doses of the studied compounds. Accordingly, the application of compound **13b** might be considered a promising

therapeutic strategy that could be useful in the management of various inflammatory diseases. Additionally, the results achieved in this study prove that the new 1,3,4-oxadiazole derivatives of pyrrolo[3,4-*d*]pyridazinone represent a promising template for further development towards potent and safe anti-inflammatory agents. Nevertheless, further investigations are needed to confirm and clarify the mechanism involved in the anti-inflammatory effect of the compounds examined.

Acknowledgments

The authors are grateful to Paulina Jawień and Joanna Kwiatkowska for support with experiments conduction. We are also thankful to Kamila Czyszczon, and Maja Lipińska for their assistance with animal maintenance and care.

Author Contributions

All authors contributed to data analysis, drafting or revising the article, have agreed on the journal to which the article will be submitted, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors declare no conflicts of interest in this work.

References

1. Goldstein DS. How does homeostasis happen? Integrative physiological, systems biological, and evolutionary perspectives. *Am J Physiol Regul Integr Comp Physiol*. 2019;316(4):R301–R317. doi:10.1152/ajpregu.00396.2018
2. Medzhitov R. Origin and physiological roles of inflammation. *Nature*. 2008;454(7203):428–435. doi:10.1038/nature07201
3. Chen Z, Ni W, Yang C, et al. Therapeutic effect of *Anomum villosum* on inflammatory bowel disease in rats. *Front Pharmacol*. 2018;9:639. doi:10.3389/fphar.2018.00639
4. Gilroy D, De Maeyer R. New insights into the resolution of inflammation. *Semin Immunol*. 2015;27(3):161–168. doi:10.1016/j.smim.2015.05.003
5. Lau D, Mollnau H, Eiserich JP, et al. Myeloperoxidase mediates neutrophil activation by association with CD11b/CD18 integrins. *Proc Natl Acad Sci U S A*. 2005;102(2):431–436. doi:10.1073/pnas.0405193102
6. Haegens A, Heeringa P, van Suylen RJ, et al. Myeloperoxidase deficiency attenuates lipopolysaccharide-induced acute lung inflammation and subsequent cytokine and chemokine production. *J Immunol*. 2009;182(12):7990–7996. doi:10.4049/jimmunol.0800377
7. Leuti A, Fazio D, Fava M, Piccoli A, Oddi S, Maccarrone M. Bioactive lipids, inflammation and chronic diseases. *Adv Drug Deliv Rev*. 2020;159:133–169.
8. Ferrero-Miliani L, Nielsen OH, Andersen PS, Girardin SE. Chronic inflammation: importance of NOD2 and NALP3 in interleukin-1 β generation. *Clin Exp Immunol*. 2007;147(2):227–235.

9. Bindu S, Mazumder S, Bandyopadhyay U. Non-steroidal anti-inflammatory drugs (NSAIDs) and organ damage: a current perspective. *Biochem Pharmacol.* 2020;180:114147. doi:10.1016/j.bcp.2020.114147
10. Malinka W, Redzicka A, Lozach O. New derivatives of pyrrolo [3,4-*d*]pyridazinone and their anticancer effects. *Farmaco.* 2004;59(6):457–462. doi:10.1016/j.farmac.2004.03.002
11. Mogilski S, Kubacka M, Redzicka A, et al. Antinociceptive, anti-inflammatory and smooth muscle relaxant activities of the pyrrolo[3,4-*d*]pyridazinone derivatives: possible mechanisms of action. *Pharmacol Biochem Behav.* 2015;133:99–110. doi:10.1016/j.pbb.2015.03.019
12. Szczukowski Ł, Redzicka A, Wiatrak B, et al. Design, synthesis, biological evaluation and in silico studies of novel pyrrolo[3,4-*d*]pyridazinone derivatives with promising anti-inflammatory and anti-oxidant activity. *Bioorg Chem.* 2020;102:104035. doi:10.1016/j.bioorg.2020.104035
13. Manjunatha K, Poojary B, Lobo PL, Fernandes J, Kumari NS. Synthesis and biological evaluation of some 1,3,4-oxadiazole derivatives. *Eur J Med Chem.* 2010;45(11):5225–5233. doi:10.1016/j.ejmech.2010.08.039
14. Palkar MB, Singhai AS, Ronad PM, et al. Synthesis, pharmacological screening and in silico studies of new class of diclofenac analogues as a promising anti-inflammatory agents. *Bioorg Med Chem.* 2014;22(10):2855–2866. doi:10.1016/j.bmc.2014.03.043
15. Koksai M, Ozkan-Dagliyan I, Ozyazici T, et al. Some novel mannich bases of 5-(3,4-dichlorophenyl)-1,3,4-oxadiazole-2(3H)-one and their anti-inflammatory activity. *Arch Pharm.* 2017;350(9):1700153. doi:10.1002/ardp.201700153
16. Avci A, Taşci H, Kandemir Ü, Can ÖD, Gökhan-Kelekçi N, Tozkoparan B. Synthesis, characterization, and in vivo pharmacological evaluation of novel mannich bases derived from 1,2,4-triazole containing a naproxen moiety. *Bioorg Chem.* 2020;100:103892. doi:10.1016/j.bioorg.2020.103892
17. Wakulik K, Wiatrak B, Szczukowski Ł, et al. Effect of novel pyrrolo [3,4-*d*]pyridazinone derivatives on lipopolysaccharide-induced neuroinflammation. *Int J Mol Sci.* 2020;21(7):2575. doi:10.3390/ijms21072575
18. Szandruk-Bender M, Wiatrak B, Szczukowski Ł, et al. Novel 1,3,4-oxadiazole derivatives of pyrrolo[3,4-*d*]pyridazinone exert antinociceptive activity in the tail-flick and formalin test in rodents and reveal reduced gastrotoxicity. *Int J Mol Sci.* 2020;21(24):9685. doi:10.3390/ijms21249685
19. Basbaum AI, Bautista DM, Scherrer G, Julius D. Cellular and molecular mechanisms of pain. *Cell.* 2009;139(2):267–284. doi:10.1016/j.cell.2009.09.028
20. Florentino IF, Silva DPB, Galdino PM, et al. Antinociceptive and anti-inflammatory effects of *Memora nodosa* and allantoin in mice. *J Ethnopharmacol.* 2016;186:298–304. doi:10.1016/j.jep.2016.04.010
21. Malinka W, Redzicka A, Jastrzębska-Więsek M, et al. Derivatives of pyrrolo[3,4-*d*]pyridazinone, a new class of analgesic agents. *Eur J Med Chem.* 2011;46(10):4992–4999. doi:10.1016/j.ejmech.2011.08.006
22. Winter CA, Risley EA, Nuss GW. Carrageenin-induced edema in hind paw of the rat as an assay for antiinflammatory drugs. *Proc Soc Exp Biol Med.* 1962;111(3):544–547. doi:10.3181/00379727-111-27849
23. Mert T, Yaman S. Pro-inflammatory or anti-inflammatory effects of pulsed magnetic field treatments in rats with experimental acute inflammation. *Environ Sci Pollut Res Int.* 2020;27(25):3154–31554. doi:10.1007/s11356-020-09401-z
24. Jain D, Mathur SR, Iyer VK. Cell blocks in cytopathology: a review of preparative methods, utility in diagnosis and role in ancillary studies. *Cytopathology.* 2014;25(6):356–371.
25. Szabo S, Trier JS, Brown A, Schnoor J, Homan HD, Bradford JC. A quantitative method for assessing the extent of experimental gastric erosions and ulcers. *J Pharmacol Methods.* 1985;13(1):59–66. doi:10.1016/0160-5402(85)90068-3
26. Necas J, Bartosikova L. Carrageenan: a review. *Vet Med.* 2013;58(4):187–205. doi:10.17221/6758-VETMED
27. Fehrenbacher JC, Vasko MR, Duarte DB. Models of inflammation: Carrageenan- or complete Freund's Adjuvant (CFA)-induced edema and hypersensitivity in the rat. *Curr Protoc Pharmacol.* 2012; Chapter 5:Unit5.4.
28. Abu-Taweel GM, Mohsen GAM, Antonisamy P, et al. *Spirulina* consumption effectively reduces anti-inflammatory and pain related infectious diseases. *J Infect Public Health.* 2019;12(6):777–782. doi:10.1016/j.jiph.2019.04.014
29. Zhang H, Shang C, Tian Z, et al. Diallyl disulfide suppresses inflammatory and oxidative machineries following carrageenan injection-induced paw edema in mice. *Mediators Inflamm.* 2020;2020:8508906. doi:10.1155/2020/8508906
30. Otterness IG, Wiseman EH, Gans DJ. A comparison of the carrageenan edema test and ultraviolet light-induced erythema test as predictors of the clinical dose in rheumatoid arthritis. *Agents Actions.* 1979;9(2):177–183. doi:10.1007/BF02024731
31. Ben Khedir S, Mzid M, Bardaa S, Moalla D, Sahnoun Z, Rebai T. In vivo evaluation of the anti-inflammatory effect of *Pistacia lentiscus* fruit oil and its effects on oxidative stress. *Evid Based Complement Alternat Med.* 2016;2016:6108203. doi:10.1155/2016/6108203
32. Karim N, Khan I, Khan W, et al. Anti-nociceptive and anti-inflammatory activities of asparacosin involve selective cyclooxygenase 2 and inflammatory cytokines inhibition: an in-vitro, in-vivo, and in-silico approach. *Front Immunol.* 2019;10:581. doi:10.3389/fimmu.2019.00581
33. Loram LC, Fuller A, Fick LG, Cartmell T, Poole S, Mitchell D. Cytokine profiles during carrageenan-induced inflammatory hyperalgesia in rat muscle and hind paw. *J Pain.* 2007;8(2):127–136. doi:10.1016/j.jpain.2006.06.010
34. Posadas I, Bucci M, Roviezzo F, et al. Carrageenan-induced mouse paw oedema is biphasic, age-weight dependent and displays differential nitric oxide cyclooxygenase-2 expression. *Br J Pharmacol.* 2004;142(2):331–338. doi:10.1038/sj.bjp.0705650
35. Vysakh A, Jayesh K, Helen LR, Jyothis M, Latha MS. Acute oral toxicity and anti-inflammatory evaluation of methanolic extract of *Rotula aquatica* roots in Wistar rats. *J Ayurveda Integr Med.* 2020;11(1):45–52. doi:10.1016/j.jaim.2017.09.007
36. Nantel F, Denis D, Gordon R, et al. Distribution and regulation of cyclooxygenase-2 in carrageenan-induced inflammation. *Br J Pharmacol.* 1999;128(4):853–859. doi:10.1038/sj.bjp.0702866
37. Zhang J, Wang H, Wang T, et al. Anti-inflammatory activity of *Yanshu* spraying agent in animal models. *Exp Ther Med.* 2013;5(1):73–76. doi:10.3892/etm.2012.761
38. Nakanishi M, Rosenberg DW. Multifaceted roles of PGE2 in inflammation and cancer. *Semin Immunopathol.* 2013;35(2):123–137. doi:10.1007/s00281-012-0342-8
39. Aoki T, Narumiya S. Prostaglandins and chronic inflammation. *Trends Pharmacol Sci.* 2012;33(6):304–311. doi:10.1016/j.tips.2012.02.004
40. Ozyazici T, Gurdal EE, Orak D, et al. Synthesis, anti-inflammatory activity, and molecular docking studies of some novel mannich bases of the 1,3,4-oxadiazole-2(3H)-thione scaffold. *Arch Pharm.* 2020;353(7):e2000061. doi:10.1002/ardp.202000061
41. Li CL, Tan LH, Wang YF, et al. Comparison of anti-inflammatory effects of berberine, and its natural oxidative and reduced derivatives from *Rhizoma coptidis* in vitro and in vivo. *Phytomedicine.* 2019;52:272–283. doi:10.1016/j.phymed.2018.09.228
42. Cai C, Chen Y, Zhong S, et al. Anti-inflammatory activity of N-butanol extract from *Ipomoea stolonifera* in vivo and in vitro. *PLoS One.* 2014;9(4):e95931. doi:10.1371/journal.pone.0095931
43. Heeba GH, Mahmoud ME, El Hanafy AA. Anti-inflammatory potential of curcumin and quercetin in rats: role of oxidative stress, heme oxygenase-1 and TNF- α . *Toxicol Ind Health.* 2014;30(6):551–560. doi:10.1177/0748233712462444

44. Cidade AF, Vasconcelos PA, Silva DPB, et al. Design, synthesis and pharmacological evaluation of new anti-inflammatory compounds. *Eur J Pharmacol.* 2016;791:195–204. doi:10.1016/j.ejphar.2016.08.033
45. Kumar KP, Nicholls AJ, Wong CHY. Partners in crime: neutrophils and monocytes/macrophages in inflammation and disease. *Cell Tissue Res.* 2018;371(3):551–565. doi:10.1007/s00441-017-2753-2
46. Othman A, Sekheri M, Filep JG. Roles of neutrophil granule proteins in orchestrating inflammation and immunity. *FEBS J.* 2021. doi:10.1111/febs.15803
47. El Kebir D, József L, Pan W, et al. 15-epi-lipoxin A4 inhibits myeloperoxidase signaling and enhances resolution of acute lung injury. *Am J Respir Crit Care Med.* 2009;180(4):311–319. doi:10.1164/rccm.200810-1601OC
48. Bezerra Rodrigues Dantas L, Silva ALM, Da Silva Júnior CP, et al. Nootkatone inhibits acute and chronic inflammatory responses in mice. *Molecules.* 2020;25(9):2181. doi:10.3390/molecules25092181
49. Sin YM, Pook SH, Tan TM, Pettersson A, Kara AU, The WF. Changes in glutathione and its associated enzymes during carrageenan-induced acute inflammation in mice. *Comp Biochem Physiol.* 1997;116(3):191–195.
50. Palaska E, Sahin G, Kelicen P, Durlu NT, Altinok G. Synthesis and anti-inflammatory activity of 1-acylthiosemicarbazides, 1,3,4-oxadiazoles, 1,3,4-thiadiazoles and 1,2,4-triazole-3-thiones. *Farmaco.* 2002;57(2):101–107. doi:10.1016/S0014-827X(01)01176-4
51. Begum MM, Islam A, Begum R, et al. Ethnopharmacological inspections of organic extract of *Oroxylum indicum* in rat models: a promising natural gift. *Evid Based Complement Alternat Med.* 2019;2019:1562038. doi:10.1155/2019/1562038
52. Naughton CA. Drug-induced nephrotoxicity. *Am Fam Physician.* 2008;78(6):743–750.
53. Sadek B, Hamruoni AM, Adem A. Anti-inflammatory agents of the carbamoylmethyl ester class: synthesis, characterization, and pharmacological evaluation. *J Inflamm Res.* 2013;6:35–43. doi:10.2147/JIR.S39743
54. Bjarnason I, Scarpignato C, Holmgren E, Olszewski M, Rainsford KD, Lanás A. Mechanisms of damage to the gastrointestinal tract from nonsteroidal anti-inflammatory drugs. *Gastroenterology.* 2018;154(3):500–514. doi:10.1053/j.gastro.2017.10.049
55. Sriuttha P, Sirichanchuen B, Permsuwan U. Hepatotoxicity of non-steroidal anti-inflammatory drugs: a systematic review of randomized controlled trials. *Int J Hepatol.* 2018;2018:5253623. doi:10.1155/2018/5253623
56. Saidu Y, Bilbis LS, Lawal M, Isezuo SA, Hassan SW, Abbas AY. Acute and sub-chronic toxicity studies of crude aqueous extract of *Albizia chevalieri* Harms (Leguminosae). *Asian J Biochem.* 2007;2(4):224–236. doi:10.3923/ajb.2007.224.236
57. Ozer J, Ratner M, Shaw M, Bailey W, Schomaker S. The current state of serum biomarkers of hepatotoxicity. *Toxicology.* 2008;245(3):194–205. doi:10.1016/j.tox.2007.11.021
58. Junior SDA, Pereira PM, de Souza Tótolí V, et al. Incorporation of indomethacin into a mesoporous silica nanoparticle enhances the anti-inflammatory effect indomethacin into a mesoporous silica. *Eur J Pharm Sci.* 2021;157:105601. doi:10.1016/j.ejps.2020.105601
59. Deng X, Stachlewitz RF, Liguori MJ, et al. Modest inflammation enhances diclofenac hepatotoxicity in rats: role of neutrophils and bacterial translocation. *J Pharmacol Exp Ther.* 2006;319(3):1191–1199. doi:10.1124/jpet.106.110247
60. Kim SY, Moon A. Drug-induced nephrotoxicity and its biomarkers. *Biomol Ther.* 2012;20(3):268–272. doi:10.4062/biomolther.2012.20.3.268

Journal of Inflammation Research

Dovepress

Publish your work in this journal

The Journal of Inflammation Research is an international, peer-reviewed open-access journal that welcomes laboratory and clinical findings on the molecular basis, cell biology and pharmacology of inflammation including original research, reviews, symposium reports, hypothesis formation and commentaries on: acute/chronic inflammation; mediators of inflammation; cellular processes; molecular

mechanisms; pharmacology and novel anti-inflammatory drugs; clinical conditions involving inflammation. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-inflammation-research-journal>