Open Access Full Text Article

Usability of Norm Values on Results from a Simulator Device and Cognitive Tests in Traffic Medicine

Kersti Samuelsson ()¹, Ewa Wressle ()²

¹Department of Rehabilitation Medicine and Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden; ²Department of Acute Internal Medicine and Geriatrics in Linköping, and Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden

Correspondence: Kersti Samuelsson, Department of Rehabilitation Medicine and Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden, Email Kersti.samuelsson@regionostergotland.se

Introduction: Driving is an essential everyday task for most adults, and fitness to continue car-driving after a brain injury/disease is a common issue in rehabilitation settings. There is no consensus on how this assessment should be performed and thus further research and development are of great value. The aim was to study the usability of cut-off values, based on recently developed norm values for a driving simulator device (CyberSiM), as well as cognitive tests, for patients already considered fit-to-drive after a standardized traffic medical investigation.

Methods: The study had a retrospective case-control design. Norm results (n = 129) were compared with patient results (n = 126) divided into two age groups (\leq 59 years and \geq 60 years). Results from Useful Field of View, Trail Making Test, Nordic Stroke Driver Screening Assessment as well as a simulator device (CyberSiM) were compared.

Results: The group of patients considered fit-to-drive after a traffic medicine assessment had worse results on all cognitive tests compared with norms. Results on CyberSiM subtests II and III did not differ from norms. The proportion of patients within suggested cut-off limits (mean±2SD norm) and considered fit to drive (mean±2SD norm) were highest (75–95%) for all three subtests of CyberSiM and for Useful Field of View in both age groups.

Conclusion: Availability of norm values in decision on continued driving is of value when interpreting the results of cognitive assessments sensitive to age, but it must be handled with care because many factors are important for individuals' ability to drive. **Keywords:** acquired brain injury, age, attention, behavior

Introduction

It is well established that safe driving, an important activity for daily living, successful occupational function, social participation, and well-being, is heavily reliant on functioning cognitive processes.¹ However, there is a lack of evidence in how to predict driving capacity after brain injury/disease^{2,3} and most studies conclude that no single assessment tool is sufficient to decide on driving fitness.^{3–7} Most studies recommend to collect information from different areas of cognitive ability by analyzing test results from several cognitive assessment tools and preferably in addition observe behavior in simulated or real traffic activities.⁸ To improve the ability to interpret results from cognitive functions.⁹ Another aspect of age that might affect the decision from a traffic medicine evaluation, is that the elderly often have more driving experience than younger persons, meaning that actual driving tasks can often happen automatically, which might compensate for some cognitive impairment.¹⁰

Focused, sustained and divided attention, processing speed, executive function, and memory are described as important functions for safe driving.^{11–13} To assess the relevant cognitive functions, several tests, such as Useful Field of View (UFOV),^{14,15} Trail Making Test (TMT),¹⁶ and the Nordic version of Stroke Driver Screening Assessment

© 2022 Samuelson and Wressle. This work is published and licensed by Dove Medical Press Limited. The full terms of this license are available at https://www.dovepress. accessing the work you hereby accept the Terms. Non-commercial uses of the work are permitted without any further permission from Dove Medical Press Limited, provided the work is properly attributed. For permission for commercial uses of this work, please see paragraphs 42, and 5 of our Terms (https://www.dovepress.com/terms.php). (NorSDSA)¹⁷ are often used in the Nordic countries in decisions on continued driving. These assessment tools are available to occupational therapists, who are often engaged in traffic medical assessments.

A simulator device might contribute with information on cognitive areas other than cognitive tests and could therefore be of value in decision based on assessments made in clinical practice.^{7,18,19} A new type of a simulator tool has recently been developed and tested at five different traffic medicine units in Sweden. The device (CyberSiM) has shown good internal consistency and promising results in adding important information to the decision discriminating patients considered fit or unfit to drive.⁶ The simulator device includes cognitive assessment of attention, processing speed, visuospatial function, simultaneous capacity and executive function including inhibition.⁷ Because age is known to have an impact on cognition,^{20,21} norm values have been produced for the CyberSiM device as well as for some other cognitive tests.^{6,22}

The aim was to study the usability of cut-off values, based on recently developed norm values for a driving simulator device (CyberSiM), as well as cognitive tests, for patients already considered fit-to-drive after a standardized traffic medical investigation.

Methods

Design

A retrospective case-control study was done based on existing data from two other studies.^{6,7} A cut-off for comparing results for patients already considered fit-to drive with a norm population, was set at ± 2 standard deviations (SD) from the mean.^{16,22}

Participants

Healthy individuals (n=129) were used as controls. Cases included patients referred to four different traffic medicine units in Sweden (n=126) for an assessment of continued driving ability after a brain injury/disease (B license).

Data from five different cognitive assessment tools, as well as results from a simulator device, were used to compare the results for the control and patient groups. Based on expected age differences²⁰ and the limited number of participants, the two groups were divided into two according to age: \leq 59 years versus \geq 60 years.

Data Collection

The study-specific protocol has been used in two former studies.^{6,7} The protocol included descriptive information on age, gender, educational level (≤ 9 years, 10–12 years, >12 years) and estimated driving/year in Swedish miles (<1000, 1000–2000, >2000). Occupational therapists at each unit collected all patient data, and a psychology student collected all norm data. All data related to one participant were collected by the same therapist on one occasion.

Assessment Tools

CyberSiM: the Simulator Test

CyberSiM consists of a software which is installed on a laptop and attached to a steering wheel.⁶ Paddles located on the left- and right-hand side of the steering wheel are used to respond to visual stimuli on the screen (Figure 1), no pedals are included. The participants are asked to steer the car on the simulated road at a constant speed (40 km/h) and at the same time react to stimuli shown on the left or right or both sides of the screen. Before each subtest, a short exercise is used to check that the participants understand each subtest. Participants are asked to steer the car on the right of the road and avoid obstacles at the side of the road. The examiner follows the session via a separate computer (Figure 1).

Each time a visual stimulus (an arrow pointing to the left or right) pops up on the screen, participants are asked to respond, by pressing the correct paddle as fast as possible (Figure 1). Three subtests (I, II and III) are included, representing increasing cognitive challenges. The time-frame for the entire session is 30–40 minutes.

Subtest I: A simple stimuli-response task is presented and the participant is expected to press the right paddle whenever an arrow appears on the screen.



Figure I CyberSiM. The steering wheel with two paddles (www.Cybersim.com).

Subtest II: An arrow pointing to the left or right on the left or right side of the screen is presented and the participant responds by pressing the corresponding paddle depending on the arrow direction.

Subtest III: Two arrows are presented at the same time, one on each side of the screen. When the arrows are pointing in opposite directions, the participant should not respond (inhibited response); when both arrows point in the same direction, the participant should press the paddle representing the direction of the two arrows.

Mean reaction time (in milliseconds), missed and wrong responses are reported. A wobbling algorithm is calculated to illustrate how well the participant manage to keep the car on the right side of the road and within road markings. The algorithm is based on the number of the intersection of the road markings and how long.

Useful Field of View

UFOV 6.1.4 is a computer-based screening assessment for fitness to drive,^{14,15} consisting of three subtests assessing the precision of visual processing under increasingly complex tasks. Briefly presented targets has to be detected, identified and localized by the participant by responding on a touch screen or by using a mouse. UFOV adjusts the duration of stimuli presentation depending on the participant's responses.¹⁴ The process of tracking the perceptual threshold goes on until a stable estimate of 75% correct is calculated.

Subtest 1: Processing speed is required for this subtest. The task is to detect, identify, and localize briefly presented vehicles centrally on the screen.

Subtest 2: Divided attention is required for this subtest. The task is to simultaneously identify a centrally located vehicle as well as a peripherally located vehicle.

Subtest 3: Selective attention is required for this subtest. The task is identical to subtest 2 except that the vehicle in the periphery is embedded in 47 triangles.

Results are reported in milliseconds. A total score as well as scores for every subtests are provided. A high validity (Cohen's d = 0.945) when UFOV was compared with driving measures was shown in a meta-analysis study.²³ In addition, UFOV has been shown to have a high level of prediction of crash risk.^{24–26}

Trail Making Test

The TMT version used in this study is a paper-based test consisting of two subtests.¹⁶ Results are presented as the time taken to complete each subtest.

Subtest 1: TMT-A The participant is asked to draw lines sequentially connecting 25 numbers as quickly as possible without lifting the pencil.

Subtest 2: TMT-B The participant is asked to draw lines alternating between 13 numbers and 12 letters in two sequences.²⁷

TMT is considered to assess visual searching, scanning, processing speed, mental flexibility, and executive functions.¹⁶ TMT-A is considered to require visuoperceptual function and TMT-B primarily working memory and secondarily task-switching ability.²⁸

Nordic Stroke Driver Screening Assessment

The NorSDSA¹⁷ is based on the British Stroke Driver Screening Assessment (SDSA).²⁹ The Nordic version is adapted for right-hand traffic, and some road signs were replaced.

The NorSDSA is a paper-based test including four subtests:

Subtest 1: This dot-cancellation test includes dots lined up in groups of three, four, or five and the participant is asked to mark all groups with four dots. Results are expressed as time taken in seconds, missed groups of four, and crossed out groups of three or five, so-called false results.

Subtest 2: This direction test consists of a 4×4 square matrix and 16 stimulus cards showing a lorry (represented by a large arrow) and a car (represented by a small arrow) traveling in different directions. Directional arrows are placed along the side and top of the matrix. The participant then places the stimulus cards in the square that corresponds correctly to one small and one large arrow. Maximum time is 5 minutes and results are expressed in points, one point given for each correctly placed vehicle.

Subtest 3: This compass test includes arrows representing eight compass directions and two cars leaving or driving onto a roundabout on two of eight different roads. The participant is asked to correctly place 16 out of 28 cards. Maximum time is 5 minutes and results are expressed in points, one point given for each correctly placed card.

Subtest 4: This road sign recognition test includes 12 cards depicting different traffic situations. The participant is asked to correctly place 12 out of 20 cards with traffic signs, on top of the corresponding traffic situation. One point is given for each correctly placed card.

According to the NorSDSA manual, dot cancellation requires visual scanning and visual perception, sustained and selective attention as well as speed. The direction test requires visuospatial functions, and the compass test scanning and processing of visual material. Road sign recognition requires visual scanning and processing as well as verbal and visuospatial memory. NorSDSA provides a weighted total score based on results for dot cancellation, compass and road sign recognition subtests. Higher score represent better result.¹⁷

Some studies have shown that SDSA as well as NorSDSA correctly classifies a sufficient number of unsuitable drivers,³⁰ but have been reported inaccurate for neurological conditions other than stroke.^{31,32}

On-Road Observation: Patient Group

An on-road observation with specially educated occupational therapists was performed after the cognitive tests, when additional information was needed and not in cases when decision considering fitness for driving was obvious without this observation. The patients undergoing on-road assessment (67%) drove on a standardized road trip, lasting for about 60 minutes. Different traffic situations and demands were included in the on-road observation. Risk situations and misbehavior in relation to actions taken were documented.

Final Decision

The final decision on whether a patient should be considered fit or unfit to continue car-driving was always made in a team consensus discussion, among a specialist physician, an occupational therapist and a psychologist. Cognitive test results as well as results from an on-road assessment (when performed) were the basis for this discussion. Norm data adjusted for a Nordic population were not available for any of the included tests at the time for data collection or the final decision. The on-road observation was performed and included in the final decision whenever additional or confirmatory information was needed. The simulator test results were not included in the final decision.

Statistics

The Statistical Package for the Social Sciences (SPSS) version 23.0 was used for statistical analyses. Descriptive statistics were used for the sociodemographic data. Parametric and non-parametric statistical methods were used to analyze data based on the results from normal distribution analyses and type of variables analyzed. The accepted level of significance was p < 0.05. Based on a number of studies describing a wide range of variability in test results, a cut-off for using norm values for comparison was set at ±2 standard deviations (SD) from the mean.^{16,22,33}

Results

A comparison of demographic data for the norm versus the patient group showed significant differences for gender, age, level of education, and estimated driving per year (Table 1). A within-group analysis taking gender and educational level into account showed no differences in the test results (norm group, p > 0.05; patient group, p > 0.05), thus this was not taken into further account.

Test results for the patient group based on age and results for the final team assessment (fit/unfit to drive) were compared showing significant differences between groups for all tests except TMT-A for the oldest group and for some of the results for CyberSiM wobbling, misses, and faults (Table 2).

A comparison between age-related norm values versus patients who were considered unfit to drive showed significant differences for all test results and for both age groups (p < 0.05).

	Patients (n=129)	Norms (n=126)	Chi-Squared Test	Þ
Gender (%)				
Men	74	43	24.9	<0.001
Women	26	57		
Age (years), mean ± SD (range)	61 ±14 (23-88)	48 ±19 (20-85)	F=28.9	<0.001*
Diagnosis (%)				
Stroke	50			
MCI	14			
ТВІ	12			
MS/PD	8			
Subarachnoid hemorrhage	4			
Brain tumor	3			
Other	9			
Education (%)				
≤9 years	28	6	70.3	<0.001
10–12 years	48	17		
≥12 years	24	77		
Estimated driving per year (%)				
<1000 Swedish miles/year	33	47	5.6	ns
1000–2000 Swedish miles/year	45	36		
>2000 Swedish miles/year	22	17		

Table	Demographic	Data for	the	Participants
-------	-------------	----------	-----	--------------

Note: * Student's t-test.

Abbreviations: MCI, mild cognitive impairment; MS, multiple sclerosis; PD, Parkinson disease;; SD, standard deviation; TBI, traumatic brain injury.

	≤59 Years, Mean (SD)			≥60 Years, Mean (SD)		
	Fit to Drive (n=40)	Unfit to Drive (n=12)	p value	Fit to Drive (n=45)	Unfit to Drive (n=32)	p value
UFOV total, msec	168 (105)	338 (249)	0.001	363 (255)	611 (334)	<0.001
TMT-A, sec	36 (10)	48 (14)	0.002	57 (56)	68 (39)	ns
TMT-B, sec	89 (36)	146 (55)	0.001	125 (72)	193 (84)	0.002
NorSDSA, score	2.3 (1.2)	-0.06 (1.5)	<0.001	1.10 (1.6)	-0.55 (1.9)	<0.001
CyberSiM I, msec	615 (169)	784 (267)	0.011	776 (196)	938 (249)	0.003
Wobbling, score	0.56 (0.3)	0.76 (0.3)	ns	0.64 (0.2)	0.99 (0.7)	0.002
Misses, number	0.23 (0.5)	0.36 (0.7)	ns	0.51 (0.7)	1.87 (3.4)	0.011
Faults, number	0.33 (0.6)	0.27 (0.6)	ns	0.36 (1.2)	0.71 (1.3)	ns
CyberSiM II, msec	1042 (181)	1230 (394)	0.026	1027 (172)	1280 (234)	<0.001
Wobbling, score	0.62 (0.4)	0.81 (0.4)	ns	0.73 (0.3)	1.11 (0.8)	0.007
Misses, number	0.10 (0.4)	0.25 (0.6)	ns	0.18 (0.4)	1.69 (2.6)	<0.001
Faults, number	1.38 (1.4)	2.67 (2.7)	0.029	2.73 (2.3)	3.45 (3.0)	ns
CyberSiM III, msec	1375 (321)	1811 (507)	0.001	1386 (271)	1656 (310)	<0.001
Wobbling, score	0.60 (0.3)	0.80 (0.3)	ns	0.81 (0.3)	1.08 (0.6)	0.009
Misses, number	0.10 (0.4)	0.50 (1.2)	ns	0.24 (1.1)	1.10 (1.9)	0.014
Faults, number	0.85 (1.0)	1.92 (2.4)	0.031	1.24 (1.6)	2.79 (2.7)	0.003

Table 2 Comparison Between Scores for Patients Who are Fit to Drive versus Unfit to Drive

Note: Student's t-test.

Abbreviations: NorSDSA, Nordic Stroke Driver Screening Assessment; SD, standard deviation; UFOV, Useful Field of View.

Results for the norm groups versus patients fit to drive showed no differences in the results for CyberSiM II and III for the oldest age group; other test results differed significantly. Test results on the age groups for norms, fit to drive, and unfit to drive are illustrated with medians, 1st and 3rd quartiles, minimums, and maximums in Figures 2–5.



Figure 2 UFOV total score: norm data, fit to drive, unfit to drive.





Cut-off values based on norm values (mean \pm 2SD) were applied to study the extent, to which patients who were evaluated as fit to drive were within set limits, expressed as proportion of patients. Results varied between the tests (Table 3). For the oldest age group, UFOV total and CyberSiM I–III had the highest conformity with the norms. For patients <59 years of age, CyberSiM I–III had the highest conformity. The highest proportion of results within the set cut-off limits for both age groups was identified for the CyberSiM reaction time (75–95%). For CyberSiM wobbling and misses, conformity with norms was 83–100% (Table 3). TMT-A had the lowest proportion of patients fit to drive within the defined interval (\pm 2SD): 46% and 55% respectively for the different age groups.



Figure 4 NorSDSA total score: norm data, fit to drive, unfit to drive.

Discussion

Several studies report the value of using standardized cognitive tests in clinical practice on traffic medicine issues.³⁴ The aim for this study was to analyze the usability of cut-off values, based on recently developed norm values for a driving simulator device (CyberSiM), as well as cognitive tests, for patients already considered fit-to-drive after a standardized traffic medical investigation. The comparison between norms (mean \pm 2SD) and patients considered fit to drive showed that 75–95% of patients performed within the stipulated limits for all three subtests of CyberSiM (reaction time) and for both age groups.

The proportion of patients who were within the cut-off limit for UFOV total was 82% for the oldest patient group and 71% for the younger group. For TMT-B, the proportion was about 70%. TMT-A and NorSDSA were less compliant compared with norm values (between 46% and 65%). NorSDSA has been shown to be inaccurate for neurologic conditions other than stroke, which might explain the low compliance for the study group, which included a diverse range of diagnoses.^{31,32}

UFOV total score represents a summary of the results from the three different subtests, which together aim to address three different types of attention (processing speed, divided attention and selective attention). Another form of attention described as being of utmost importance in driving is sustained attention.¹³ Sustained attention is not addressed in any of the cognitive tests included here. The relatively short period of time it takes to perform, for example, the UFOV assessment, and the fact that the test person is well aware of the different stimuli and the response needed, and is thus well prepared and focused, must be taken into consideration. CyberSiM takes longer to perform, and thus demands sustained attention. In addition, it includes different "driving scenarios" and unexpected events as well as inhibited responses, all of which might be more in concordance with demands on attention in a real traffic context. An interesting finding was the lack of differences in the results based on age group for CyberSiM reaction time (Figures 2–5). This

66



Figure 5 (A-C) CyberSiM I-III: norm data, fit to drive, unfit to drive. * =outlier.

indicates that CyberSiM is not as sensitive to age as the other tests. The reason for this lack of difference could be related to the fact that sustained attention is less sensitive to age than other forms of attention³⁵ and/or that the activity is equally familiar to everyone regardless of age.

Much of what we do when driving is repetitive and hence predictable, and thus might be expected to be an automated skill when it comes to the actual driving task.¹³ Thus the actual driving skill, is seldom the problem after a brain injury/ disease unless a muscular weakness or paralysis is involved.

Referrals on cognitive conditions for continued driving after a brain injury/or disease often come from physicians to occupational therapists or psychologists. Because psychologists are significantly fewer and often have other responsibilities within health care, most driving assessments are performed by occupational therapists. Immediate response to a stimuli repeated for a short period of time can be captured using traditional tests such as UFOV. However, driving puts demands on sustained attention; driving in real life means that unpredictable events might arise at any time and thus the driver must make decisions on action or no action. To capture this, we need tests that last over time and with unforeseen elements and elements that require both action and no action.

UFOV could be used as a basic attention test, but the results should be handled with care and not as a stand-alone test for making conclusions on driving. A simulator assessment such as the CyberSiM could contribute with information about attention, and especially sustained attention, in a more complex context. In addition, a simulator device might contribute with observed aspects of behavior, multitasking, and active multiple responses in terms of reaction and inhibition for different stimuli.^{6,7}

	Cut-Off Limit	≤59 Years (%)	Cut-Off Limit	≥60 Years (%)
UFOV total, msec	<172	68	<508	82
TMT-A, sec	<34	46	<45	55
TMT-B, sec	<93	68	<121	71
NorSDSA, score	>1.9	65	>0.57	63
CyberSiM I, msec	<772	75	<877	76
Wobbling, score	<0.68	85	<0.85	89
Misses, number	0	83	<2	98
Faults, number	0	73	0	82
CyberSiM II, msec	<1242	95	<1304	89
Wobbling, score	<0.71	83	<1.12	93
Misses, number	0	93	0	84
Faults, number	<3	80	<3	53
CyberSiM III, msec	<1547	85	<1630	78
Wobbling, score	<3.22	100	<1.29	93
Misses, number	<3	100	0	89
Faults, number	<2	83	<3	84

Table 3 The Proportion of Patients Considered Fit to Drive and Within the Cut-Off Limits Based on the Mean ± 2SD (Norm)

Abbreviations: NorSDSA, Nordic Stroke Driver Screening Assessment; TMT, Trail making test; SD, standard deviation; UFOV, Useful Field of View.

Although age-related norms might contribute to the evaluation of continued driving after a brain injury/disease, one should pay attention to the fact that age in itself is not the best indicator of driving. Irrespective of age, there are differences in, for example functional ability and driving experience that affect driving competence.³⁴

Strengths and Limitations

All data were collected by well-informed and experienced personnel, which should be seen as a strength together with the standardized procedures used. The simulator device was not included in the final decisions for the patient group, and no norm values were available at the time of the study.

A limitation was the small number of participants and a wide spread in the results, especially in the oldest groups. Using ± 2 SD as the cut-off limit could be seen as an arbitrary decision, however a lot of calculated alternatives were performed and rejected, finally resulting in this permissive approach, which seems to capture the team decisions in a reasonable manner. The generalizability is limited due to the relatively small patient group and the large spread in results. More studies are needed to verify the results of the study.

Conclusions

The use of different standardized assessment tools in evaluating continued car-driving in patients after brain injury/ disease is important and recommended. The availability of norm values when interpreting the results of different assessments could guide decision-making, but should be handled with care. For CyberSiM and UFOV, a comparison with norm values (cut-off limit of ± 2 SD) showed that a large proportion of patients considered fit to drive were within those limits. This could indicate the value of including norms in the discussion and evaluation.

Ethics Approval and Informed Consent

No medical or other risks related to participation in the study were identified. All participants received information about the study and an invitation letter together with an informed consent form. The informed consent was signed by the participants. All data protocols were coded, and no participant could be identified by the researchers. All procedures were in accordance with the ethical standards of the Declaration of Helsinki. The study is based on data from two former studies, which have ethical approvals from the Swedish Ethical Review Authority in Linköping: 2016-181-31, 2016-271-32 (patients), and 2016-353-31 (healthy individuals).

Acknowledgments

We would like to thank all participants at the units who contributed to the data collection for the present study; Department of Rehabilitation Medicine, Jönköping, Sweden; Department of Rehabilitation Medicine, Linköping, Sweden; Mobility Centre Gothenburg, Sweden, SamRehab Västervik, Sweden, and Umeå University and University Hospital of Umeå, Sweden.

Author Contributions

Both authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

The development of CyberSiM was funded by VINNOVA (2013-04010). The data collection in the present study was funded by the Council of Östergötland, Sweden (LIO-582071 and SC-2017-00091-26) and the Medical Research Council of Southeast Sweden (FORSS-654091; 755541; 605951; 850271).

Disclosure

The authors declare no conflicts of interest in this work.

References

- 1. Chihuri S, Mielenz TJ, DiMaggio CJ, et al. Driving cessation and health outcomes in older adults. J Am Ger Soc. 2016;64(2):332-341. doi:10.1111/jgs.13931
- 2. Ortoleva C, Brugger C, Van der Linden M, Walder B. Prediction of driving capacity after traumatic brain injury: a systematic review. J Head Trauma Rehabil. 2012;27(4):302–313. doi:10.1097/HTR.0b013e3182236299
- Classen S, Levy C, McCarthy D, Mann WC, Lanford D, Waid-Ebbs JK. Traumatic brain injury and driving assessment: an evidence-based literature review. Am J Occup Ther. 2009;64(5):580–591. doi:10.5014/ajot.63.5.580
- Larsson H, Lundberg C, Falkmer T, Johansson K. A Swedish survey of occupational therapists' involvement and performance in driving assessments. Scand J Occup Ther. 2007;14(4):215–220. doi:10.1080/11038120601110983
- 5. Schultheis M, Whipple E. Driving after traumatic brain injury: evaluation and rehabilitation interventions. *Curr Phys Med Rehabil Rep.* 2014;2 (3):176–183. doi:10.1007/s40141-014-0055-0
- Samuelsson K, Tropp M, Lundqvist A, Wressle E. Development, concurrent validity and internal consistency of a simulator tool for assessing continued car driving after a brain injury/disease. Br J Occup Ther. 2019;82(9):544–552. doi:10.1177/0308022619836935
- 7. Samuelsson K, Wressle E. Fitness to drive after acquired brain injury: results from patient cognitive screening and on-road assessment compared with age-adjusted norm values. *Scand J Psychol.* 2021;63(1):55–63. doi:10.1111/sjop.12774
- Patomella A-H, Tham K, Johansson K, Kottorp A. P-drive on-road: internal scale validity and reliability of an assessment of on-road driving performance in people with neurological disorders. *Scand J Occup Ther.* 2010;17(1):86–93. doi:10.3109/11038120903071776
- 9. Murman DL. The impact of age on cognition. Semin Hear. 2015;36(3):111-121. doi:10.1055/s-0035-1555115
- 10. Jeter K. Examining the role of cognition in driving: comparisons between driver groups and the development of the Maynooth on-road driving assessment [Master's thesis]. Ireland: National University of Ireland, Maynooth; 2016.
- Roca J, Castro C, López-Ramón M, et al. Measuring vigilance while assessing the functioning of the three attentional networks: the ANTI-vigilance task. J Neurosci Methods. 2011;198(2):312–324. doi:10.1016/j.jneumeth.2011.04.014
- 12. Asimakopulos J, Boychuck Z, Sondergaard D, et al. Assessing executive function in relation to fitness to drive: a review of tools and their ability to predict safe driving. *Aust Occup Ther J*. 2012;59(6):402–427. doi:10.1111/j.1440-1630.2011.00963.x
- 13. Groeger JA. Understanding Driving: Applying Cognitive Psychology to a Complex Everyday Task. Hove, East Sussex, UK: Psychology Press; 2000.
- 14. Visual Awareness Research Group. UFOV User's Guide Version 6.1.4. Birmingham, AL: Visual Awareness; 2009.
- 15. Marshall SC, Molnar F, Man-Son-Hing M, et al. Predictors of driving ability following stroke: a systematic review. *Top Stroke Rehabil*. 2007;14 (1):98–114. doi:10.1310/tsr1401-98
- 16. Tombaugh TN. Trail making test A and B: normative data stratified by age and education. Arch Clin Neuropsychol. 2004;19(2):203-214. doi:10.1016/S0887-6177(03)00039-8
- 17. Lundberg C. Manual for the Nordic Stroke Driver Assessment (Norsdsa). Blidö, Sweden: NorSDSA Produktion Görel Caneman and Catarina Lundberg; 2003. Swedish.
- Hird M, Egeto P, Fischer C, et al. A systematic review and meta-analysis of on-road simulator and cognitive driving assessment in Alzheimer's disease and mild cognitive impairment. J Alzheimers Dis. 2016;53(2):713–729. doi:10.3233/JAD-160276
- 19. Matas N, Nettelbeck T, Burns N. Assessment of driving simulator validity and acceptability for older adult drivers. Adv Transp Stud. 2016;39:69–88.
- 20. Salthouse TA. Selective review of cognitive aging. J Int Neuropsychol Soc. 2010;16(5):754-760. doi:10.1017/S1355617710000706

- Bergman I, Johansson K, Almkvist O, Lundberg C. Health-adjusted neuropsychological test norms based on 463 older Swedish car drivers. Scand J Psychol. 2016;57(2):93–107. doi:10.1111/sjop.12273
- 22. Selander H, Wressle E, Samuelsson K. Cognitive prerequisites for fitness to drive: norm values for the TMT, UFOV and NorSDSA tests. *Scand J* Occup Ther. 2020;27(3):231–239. doi:10.1080/11038128.2019.1614214
- 23. Ball K, Clay O, Wadley V, Roth D Predicting driving performance in older adults with the useful field of view test: a meta-analysis. In Rizzo M, Lee JD, McGehee DV, editors. Proceedings of the Third International Driving Symposium on Human Factors in Driver Assessment, Training and Vehicle Design; June 27–30, 2005.; Rockport, Maine, Iowa City: University of Iowa; 2005: 51–57.
- McManus B, Cox MK, Vance DE, Stavrinos D. Predicting motor vehicle collisions in a driving simulator in young adults using the useful field of view assessment. *Traffic Inj Prev.* 2015;16(8):818–823. doi:10.1080/15389588.2015.1027339
- 25. Clay O, Wadley V, Edwards J, Roth D, Roenker D, Ball K. Cumulative meta-analysis of the relationship between useful field of view and driving performance in older adults: current and future implications. *Optom Vis Sci.* 2005;82(8):724–731. doi:10.1097/01.opx.0000175009.08626.65
- 26. Ball K, Roenker D, Wadley V, et al. Can high-risk older drivers be identified through performance-based measures in a Department of motor vehicles setting? J Am Ger Soc. 2006;54(1):77–84. doi:10.1111/j.1532-5415.2005.00568.x
- 27. Vaucher P, Herzig D, Cardoso I, Herzog MH, Mangin P, Favrat B. The trail making test as a screening instrument for driving performance in older drivers: a translational research. *BMC Geriatr.* 2014;14(1):123. doi:10.1186/1471-2318-14-123
- 28. Sanchez-Cubillo I, Perianez J, Adrover-Roig D, et al. Construct validity of the trail making test: role of task-switching, working memory, inhibition/interference control, and visuomotor abilities. J Int Neuropsychol Soc. 2009;15(3):438–450. doi:10.1017/S1355617709090626
- 29. Nouri F, Lincoln N. Validation of a cognitive assessment: predicting driving performance after stroke. *Clin Rehabil.* 1992;6(4):275-281. doi:10.1177/026921559200600402
- Lundberg C, Caneman G, Samuelsson S, Hakamies-Blomqvist L, Almkvist O. The assessment of fitness to drive after a stroke: the Nordic stroke driver screening assessment. Scand J Psychol. 2003;44(1):23–30. doi:10.1111/1467-9450.00317
- 31. Lincoln NB, Radford KA, Nouri FM. Stroke Drivers Screening Assessment: Revised Manual. Nottingham, UK: University of Nottingham; 2012.
- 32. Selander H, Johansson K, Lundberg C, Falkmer T. The Nordic stroke driver screening assessment as predictor for the outcome of an on-road test. Scand J Occup Ther. 2010;17(1):10–17.
- 33. Edwards JD, Ross LA, Wadley VG, et al. The useful field of view test: normative data for older adults. Arch Clin Neuropsychol. 2006;21(4):275–286. doi:10.1016/j.acn.2006.03.001
- 34. Mathias J, Lucas L. Cognitive predictors of unsafe driving in older drivers: a meta-analysis. *Int Psychogeriatr.* 2009;21(4):637–653. doi:10.1017/S1041610209009119
- 35. Better Health While Aging. Ways that memory & thinking change with normal aging [homepage on the internet]. Available from https:// betterhealthwhileaging.net/how-brain-function-changes-with-normal-cognitive-aging. Accessed September 29, 2021.

Medical Devices: Evidence and Research

Dovepress

Publish your work in this journal

Medical Devices: Evidence and Research is an international, peer-reviewed, open access journal that focuses on the evidence, technology, research, and expert opinion supporting the use and application of medical devices in the diagnosis, monitoring, treatment and management of clinical conditions and physiological processes. The identification of novel devices and optimal use of existing devices which will lead to improved clinical outcomes and more effective patient management and safety is a key feature of the journal. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit http://www.dovepress.com/testimonials.php to read real quotes from published authors.

Submit your manuscript here: https://www.dovepress.com/medical-devices-evidence-and-research-journal