

# Increased Demand for Ketamine Infusions and Associated Complexities

Evan Peskin<sup>1</sup>, Jeffrey Gudin<sup>2</sup>, Michael E Schatman <sup>3,4</sup>

<sup>1</sup>Department of Pain Management - Insight Institute of Neurosurgery & Neuroscience, Flint, MI, USA; <sup>2</sup>Professor of Anesthesiology, Perioperative Medicine and Pain Management - University of Miami, Miller School of Medicine, Miami, FL, USA; <sup>3</sup>Department of Anesthesiology, Perioperative Care and Pain Medicine, NYU Grossman School of Medicine, New York, NY, USA; <sup>4</sup>Department of Population Health – Division of Medical Ethics, NYU Grossman School of Medicine, New York, NY, USA

Correspondence: Michael E Schatman, Department of Anesthesiology, Perioperative Care, and Pain Medicine, NYU Grossman School of Medicine, 550 1st Ave, New York, NY, 10016, USA, Tel +1 425-647-4880, Email Michael.Schatman@NYULangone.org

There has been a significant recent trend toward offering ketamine therapies for a multitude of medical and mental health diagnoses.<sup>1</sup> The treated conditions have included major depressive disorder,<sup>2–5</sup> post-traumatic stress disorder,<sup>6–10</sup> postpartum and peripartum depression,<sup>11–14</sup> anxiety,<sup>15–18</sup> substance abuse disorders,<sup>19–22</sup> eating disorders,<sup>23–25</sup> chronic pain,<sup>26–31</sup> and mood disorders<sup>32</sup> among others, in a continually expanding list.<sup>33</sup> This interest has been associated with widespread openings of clinics in the United States and world-wide that offer ketamine therapies. In addition to variability of guidelines and organizational oversight on a state-by-state basis, patients may experience a vastly different standard of care between individual treatment sites. Although a growing body of evidence does support potential benefits of ketamine for some of the disorders above, a prospective evaluation and retrospective analysis should be conducted in order to support the safe, widespread, and clinically effective use of this controlled substance.

Ketamine infusions are associated with several complexities that make them challenging to offer broadly to patients, particularly if they are to be considered safe and effective. Ketamine treatment, by definition, includes an active pharmacologic infusion with recognized hemodynamic and physiologic implications that can vary widely based on total delivered dosage, rate of infusion, consideration of medical comorbidities (particularly cardiovascular), patient volume and catecholaminergic status, and individual pharmacodynamic response.<sup>34</sup> Studies have suggested that discrete pharmacogenomic factors are involved in response to ketamine including a brain derived neurotrophic factor (BDNF) gene Val66Met polymorphism, cytochrome P450 family 2 subfamily B member 6 (CYP2B6) levels, and particular allelic variants.<sup>35–37</sup> The clinical impact of the treatment may potentially be affected by the route of administration, utilization of a racemic versus enantiomeric compound, and augmentation of secondary pharmacologic agents such as benzodiazepines or antiemetics.<sup>38–40</sup> In addition, there are staffing, resource, safety, and cost issues that require consideration. Each of the factors above should be weighed when planning and administering therapeutic interventions involving ketamine.

An initial requirement in considering ketamine therapy is the setting of the treatment. From a safety standpoint, one might consider inpatient administration if higher or bolus dosing is needed. Academic or inpatient facilities may have more regulatory oversight, have undergone more extensive evaluation by medical governing bodies, and have restrictive policies and procedures regarding appropriate candidates and safe administration of ketamine treatments. Unfortunately, while there are currently no data, our suspicion is that healthcare insurers are and will likely remain reluctant to cover inpatient hospitalization for ketamine treatments. Outpatient settings offering ketamine infusions can vary immensely, ranging from experienced centers that have previously specialized in infusions such as chemotherapy, to established mental healthcare facilities that may or may not have offered infusions previously, to completely de novo ketamine specialized centers employing staff with a wide range of experience and competencies. Even within the realm of inpatient care, site of service can vary from intensive care and post-anesthesia care to step-down units and general medical floors. To date, no comparative studies examining various clinical settings for these infusions have been published, and it should be recognized that inpatient

management alone likely does not have the same impact as the training and experience of the providers administering the infusion, the resources and preparation of the individual site, and the parameters of the infusion itself. Therefore, inpatient versus outpatient status of a patient for ketamine infusions, or more specifically the defined clinical setting, warrants further examination.

Another facet that warrants discussion is the associated cost and financial structuring of the infusions. An infusion billed as an inpatient service for a patient actively managed in an Intensive Care Unit will likely fall within the monetary jurisdictions of reimbursed insurance coverage. An infusion for a chronic pain patient in the community who has severe neuropathic peripheral pain secondary to prior chemotherapy nonresponsive to traditional conservative and pharmacologic care may fall within a personal out-of-pocket payment structure. Therefore, infusions may be more financially feasible and should be considered earlier for patients managed in an inpatient setting who require multimodal pain management or those with preexisting histories of complex pain presentations. This example addresses only neuropathic pain and not the newly developing mental health indications, and these generalities do not apply to all individuals and insurance coverage plans. Review of the current outpatient landscape for medical and mental health indications for ketamine infusions has demonstrated that patients are often required to pay a higher proportion of the associated cost matched to comparative therapies, creating a substantial financial barrier to access. The FDA-recognized indications for intranasal Spravato<sup>®</sup> (S-ketamine)<sup>41,42</sup> offer valuable support to the efficacy of the enantiomeric formulation, but the drastically more cost-effective generic racemic ketamine lacks FDA on-label recognition for the same indication. As research has demonstrated, both mental health diagnoses and chronic pain are not limited to a particular socioeconomic class or patient population. Ensuring equal and equitable access is imperative for maximizing the theoretical benefit of this treatment<sup>43,44</sup>; such a lack of equitable access may be interpreted as a violation of the bioethical principle of “justice”.

A third factor associated with ketamine therapy that should be acknowledged is that the drug has historically been used primarily for intraoperative anesthetic management of patients undergoing surgical intervention and secondarily for pain management, neurologic conditions, and mental health indications.<sup>45</sup> Understandably, due to this imbalance, there is disparate volume and quality of scientific medical literature comparing infusion protocols for these varied medical and mental health diagnoses suggested to be responsive to ketamine.<sup>46</sup> Research concerning mental health conditions including several depressive disorders, anxiety, suicidality, and post-traumatic stress disorder have provided a growing body of concordant evidence, while secondary indications such as substance abuse, addiction, and eating disorders are still in very early stages of scientific exploration. Routes of administration, including oral tablets, intranasal sprays, intramuscular injections, and intravenous infusions or boluses have been individually presented in various case reports without larger powered, controlled scientific trials comparing outcomes. Arguably, certain routes of administration, such as intranasal delivery, are fraught with more potential for variable drug delivery, with issues such as nasal scar tissue or patient competence in operating inhaler devices resulting in decreased exposure with already limited bioavailability.<sup>47,48</sup>

Unique concerns for ketamine therapy for mental health indications bring additional complexity for study, including the efficacy and durability of outcomes when offering the infusion alone versus pairing it with traditional mental health therapies such as cognitive behavioral therapy and motivational interviewing. It is paramount that we develop a standard of care to ensure patient safety and optimal outcomes. Cardiovascular complications, psychosis, or death attributed to ketamine administration outside of an inpatient setting are rare, although have been reported.<sup>49–52</sup> A more recent trend is home mail-order prescribing of ketamine capsules for oral administration in non-healthcare settings, which creates another host of ethical and legal issues.<sup>53</sup>

The United States has witnessed significant growth in ketamine clinics by a wide and diverse group of clinicians, treating disorders ranging from depression to PTSD to pain. To advance the science and protect patient safety, the scientific community must collect and publish meaningful data to better elucidate optimal dosages, infusion rates, clinical settings, adjuvant therapeutics, expected side effects, relative contraindications, and further prospective responsive conditions amenable to treatment. It is likely that there will be a need for treatment in both inpatient and outpatient settings, and whether there are any advantages to either will certainly depend on the training and experience of the providers. As the virgin fields of Psychedelic Medicine and Psychedelic-Assisted Therapy expand, maintaining a high degree of scientific ethics and standards represents a viable pathway to the most sustainable and beneficial practices.

Ketamine therapy is not and should not be touted as first-line treatment without impartial investigation and study. It would be helpful if the health insurance industry supported this potential emerging therapy while the evidence basis expands. Although scarce, research related to the administration of ketamine is underway and should help frame safe treatment guidelines and promote payor policies to support patient access. Arthur Schopenhauer is quoted as opining, “all truth passes through three stages. First, it is ridiculed. Second, it is violently opposed. Third, it is accepted as being self-evident”<sup>54</sup> (p. 451). Only time – and scientific effort – will determine whether ketamine treatments and ketamine assisted therapy will be recognized as viable medical and psychological treatments.

## Disclosure

Dr Evan Peskin is a co-founder of a mental healthcare organization that treats depression, PTSD, anxiety and uses ketamine-assisted therapy. He does not actively participate in any direct clinical care through this company. Dr Michael E Schatman is a research consultant for Modoscript and a member of the scientific Steering Committee of Collegium Pharmaceutical, outside the submitted work. The authors report no other conflicts of interest in this work.

## References

1. FinancialNewsMedia.com. Use of ketamine for mental disorders rising while new therapies emerge; 2021. Available from: <https://www.pnews.wire.com/news-releases/use-of-ketamine-for-mental-disorders-rising-while-new-therapies-emerge-301426549.html>. Accessed January 26, 2023.
2. Dean RL, Hurducas C, Hawton K, et al. Ketamine and other glutamate receptor modulators for depression in adults with unipolar major depressive disorder. *Cochrane Database Syst Rev*. 2021;9(9):CD011612. PMID: 34510411; PMCID: PMC8434915. doi:10.1002/14651858.CD011612.pub3
3. Murrough JW, Iosifescu DV, Chang LC, et al. Antidepressant efficacy of ketamine in treatment-resistant major depression: a two-site randomized controlled trial. *Am J Psychiatry*. 2013;170(10):1134–1142. PMID: 23982301; PMCID: PMC3992936. doi:10.1176/appi.ajp.2013.13030392
4. Zarate CA, Singh JB, Carlson PJ, et al. A randomized trial of an N-methyl-D-aspartate antagonist in treatment-resistant major depression. *Arch Gen Psychiatry*. 2006;63(8):856–864. PMID: 16894061. doi:10.1001/archpsyc.63.8.856
5. Jelen LA, Stone JM. Ketamine for depression. *Int Rev Psychiatry*. 2021;33(3):207–228. PMID: 33569971. doi:10.1080/09540261.2020.1854194
6. Jumaili WA, Trivedi C, Chao T, Kubosumi A, Jain S. The safety and efficacy of ketamine NMDA receptor blocker as a therapeutic intervention for PTSD review of a randomized clinical trial. *Behav Brain Res*. 2022;424:113804. PMID: 35181391. doi:10.1016/j.bbr.2022.113804
7. Norbury A, Rutter SB, Collins AB, et al. Neuroimaging correlates and predictors of response to repeated-dose intravenous ketamine in PTSD: preliminary evidence. *Neuropsychopharmacol*. 2021;46:2266–2277. doi:10.1038/s41386-021-01104-4
8. Liriano F, Hatten C, Schwartz TL. Ketamine as treatment for post-traumatic stress disorder: a review. *Drugs Context*. 2019;8(8):212305.
9. Abdallah CG, Roache JD, Gueorguieva R, et al. Dose-related effects of ketamine for antidepressant-resistant symptoms of posttraumatic stress disorder in veterans and active duty military: a double-blind, randomized, placebo-controlled multi-center clinical trial. *Neuropsychopharmacology*. 2022;47(8):1574–1581.
10. Ross C, Jain R, Bonnett CJ, Wolfson P. High-dose ketamine infusion for the treatment of posttraumatic stress disorder in combat veterans. *Ann Clin Psychiatry*. 2019;31(4):271–279.
11. Alipoor M, Loripoor M, Kazemi M, Farahbakhsh F, Sarkoobi A. The effect of ketamine on preventing postpartum depression. *J Med Life*. 2021;14(1):87–92. PMID: 33767791; PMCID: PMC7982256. doi:10.25122/jml-2020-0116
12. Ma JH, Wang SY, Yu HY, et al. Prophylactic use of ketamine reduces postpartum depression in Chinese women undergoing cesarean section. *Psychiatry Res*. 2019;279:252–258. PMID: 31147085. doi:10.1016/j.psychres.2019.03.026
13. Chen-Li D, Lui LMW, Rosenblat JD, et al. Ketamine as potential treatment for postpartum depression: a narrative review. *Ann Clin Psychiatry*. 2022;34(4):264–274. PMID: 36282614. doi:10.12788/acp.0082
14. Li Q, Wang S, Mei X. A single intravenous administration of a sub-anesthetic ketamine dose during the perioperative period of cesarean section for preventing postpartum depression: a meta-analysis. *Psychiatry Res*. 2022;310:114396. PMID: 35278826. doi:10.1016/j.psychres.2022.114396
15. Glue P, Neehoff S, Sabadel A, et al. Effects of ketamine in patients with treatment-refractory generalized anxiety and social anxiety disorders: exploratory double-blind psychoactive-controlled replication study. *J Psychopharmacol*. 2020;34(3):267–272. PMID: 31526207. doi:10.1177/0269881119874457
16. Tully JL, Dahlén AD, Haggarty CJ, Schiöth HB, Brooks S. Ketamine treatment for refractory anxiety: a systematic review. *Br J Clin Pharmacol*. 2022;88(10):4412–4426. PMID: 35510346; PMCID: PMC9540337. doi:10.1111/bcp.15374
17. Taylor JH, Landeros-Weisenberger A, Coughlin C, et al. Ketamine for social anxiety disorder: a randomized, placebo-controlled crossover trial. *Neuropsychopharmacology*. 2018;43(2):325–333. PMID: 28849779; PMCID: PMC5729569. doi:10.1038/npp.2017.194
18. Banov MD, Young JR, Dunn T, Szabo ST. Efficacy and safety of ketamine in the management of anxiety and anxiety spectrum disorders: a review of the literature. *CNS Spectr*. 2020;25(3):331–342. PMID: 31339086. doi:10.1017/S1092852919001238
19. Ivan Ezquerro-Romano I, Lawn W, Krupitsky E, Morgan CJA. Ketamine for the treatment of addiction: evidence and potential mechanisms. *Neuropharmacology*. 2018;142:72–82. PMID: 29339294. doi:10.1016/j.neuropharm.2018.01.017
20. Walsh Z, Mollaahmetoglu OM, Rootman J, et al. Ketamine for the treatment of mental health and substance use disorders: comprehensive systematic review. *B J Psych Open*. 2021;8(1):e19. PMID: 35048815; PMCID: PMC8715255. doi:10.1192/bjo.2021.1061
21. Worrell SD, Gould TJ. Therapeutic potential of ketamine for alcohol use disorder. *Neurosci Biobehav Rev*. 2021;126:573–589. PMID: 33989669. doi:10.1016/j.neubiorev.2021.05.006
22. Grabski M, McAndrew A, Lawn W, et al. Adjunctive ketamine with relapse prevention-based psychological therapy in the treatment of alcohol use disorder. *Am J Psychiatry*. 2022;179(2):152–162. PMID: 35012326. doi:10.1176/appi.ajp.2021.21030277

23. Mills IH, Park GR, Manara AR, Merriman RJ. Treatment of compulsive behaviour in eating disorders with intermittent ketamine infusions. *QJM*. 1998;91(7):493–503. PMID: 9797933. doi:10.1093/qjmed/91.7.493
24. Schwartz T, Trunko ME, Feifel D, et al. A longitudinal case series of IM ketamine for patients with severe and enduring eating disorders and comorbid treatment-resistant depression. *Clin Case Rep*. 2021;9(5):e03869. PMID: 34026123; PMCID: PMC8117825. doi:10.1002/ccr3.3869
25. Ragnhildstveit A, Slayton M, Jackson LK, et al. Ketamine as a novel psychopharmacotherapy for eating disorders: evidence and future directions. *Brain Sci*. 2022;12(3):382. PMID: 35326338; PMCID: PMC8963252. doi:10.3390/brainsci12030382
26. Cohen SP, Bhatia A, Buvanendran A, et al. Consensus guidelines on the use of intravenous ketamine infusions for chronic pain from the American Society of Regional Anesthesia and Pain Medicine, the American Academy of Pain Medicine, and the American Society of Anesthesiologists. *Reg Anesth Pain Med*. 2018;43(5):521–546. PMID: 29870458; PMCID: PMC6023575. doi:10.1097/AAP.0000000000000808
27. Orhurhu VJ, Roberts JS, Ly N, Cohen SP. Ketamine in acute and chronic pain management. In: *StatPearls [Internet]*. Treasure Island (FL): StatPearls Publishing; 2022. PMID: 30969646.
28. Guclu CY. Ketamine for chronic pain. In: Saiz-Sapena N, Granell-Gil M editors. *Ketamine Revisited - New Insights into NMDA Inhibitors*. IntechOpen; 2022. doi:10.5772/intechopen.104874
29. Hocking G, Cousins M. Ketamine in chronic pain management: an evidence-based review. *Anesth Analg*. 2003;97(6):1730–1739. doi:10.1213/01.ANE.0000086618.28845.9B
30. Yang Y, Maher DP, Cohen SP. Emerging concepts on the use of ketamine for chronic pain. *Expert Rev Clin Pharmacol*. 2020;13(2):135–146. PMID: 31990596. doi:10.1080/17512433.2020.1717947
31. Dahan A, Olofsen E, Sigtermans M, et al. Population pharmacokinetic-pharmacodynamic modeling of ketamine-induced pain relief of chronic pain. *Eur J Pain*. 2011;15(3):258–267. PMID: 20638877. doi:10.1016/j.ejpain.2010.06.016
32. Sanacora G, Frye MA, McDonald W, et al. A consensus statement on the use of ketamine in the treatment of mood disorders. *JAMA Psychiatry*. 2017;74(4):399–405. doi:10.1001/jamapsychiatry.2017.0080
33. Gautam CS, Mahajan SS, Sharma J, Singh H, Singh J. Repurposing potential of ketamine: opportunities and challenges. *Indian J Psychol Med*. 2020;42(1):22–29. PMID: 31997862; PMCID: PMC6970308. doi:10.4103/IJPSYM.IJPSYM\_228\_19
34. Mion G, Villeveille T. Ketamine pharmacology: an update (pharmacodynamics and molecular aspects, recent findings). *CNS Neurosci Ther*. 2013;19(6):370–380. PMID: 23575437; PMCID: PMC6493357. doi:10.1111/cns.12099
35. Hashimoto K. A BDNF Val66Met polymorphism and ketamine-induced rapid antidepressant action. *Clin Psychopharmacol Neurosci*. 2012;10(1):59–60. PMID: 23431323; PMCID: PMC3569155. doi:10.9758/cpn.2012.10.1.59
36. Cook-Sather SD, Adamson PC, Li J, Hakonarson H. CYP2B6\*6 or not CYP2B6\*6—that remains a question for precision medicine and ketamine! *Anesthesiology*. 2016;125(6):1085–1087. PMID: 27763886; PMCID: PMC5123751. doi:10.1097/ALN.0000000000001399
37. Meshkat S, Rodrigues NB, Di Vincenzo JD, et al. Pharmacogenomics of ketamine: a systematic review. *J Psychiatr Res*. 2021;145:27–34. PMID: 34844049. doi:10.1016/j.jpsychires.2021.11.036
38. Feeney A, Hoepfner BB, Freeman MP, et al. Effect of concomitant benzodiazepines on the antidepressant effects of ketamine: findings from the RAPID intravenous ketamine study. *J Clin Psychiatry*. 2022;84(1):22m14491. PMID: 36383742. doi:10.4088/JCP.22m14491
39. Maher DP, Chen L, Mao J. Intravenous ketamine infusions for neuropathic pain management: a promising therapy in need of optimization. *Anesth Analg*. 2017;124(2):661–674. PMID: 28067704. doi:10.1213/ANE.0000000000001787
40. Albott CS, Shiroma PR, Cullen KR, et al. The antidepressant effect of repeat dose intravenous ketamine is delayed by concurrent benzodiazepine use. *J Clin Psychiatry*. 2017;78(3):e308–e309. PMID: 28394513. doi:10.4088/JCP.16111277
41. United States Food and Drug Administration. Press Release: FDA approves new nasal spray medication for treatment-resistant depression; Available only at a certified doctor’s office or clinic; 2019. Available from: <https://www.fda.gov/news-events/press-announcements/fda-approves-new-nasal-spray-medication-treatment-resistant-depression-available-only-certified>. Accessed December 31, 2022.
42. SPRAVATO® (esketamine) nasal spray [package insert]. Titusville, NJ: Janssen Pharmaceuticals, Inc; 2020. Available from: [https://www.access.data.fda.gov/drugsatfda\\_docs/label/2020/211243s004lbl.pdf](https://www.access.data.fda.gov/drugsatfda_docs/label/2020/211243s004lbl.pdf). Accessed December 31, 2022.
43. Macintyre A, Ferris D, Gonçalves B, Quinn N. What has economics got to do with it? The impact of socioeconomic factors on mental health and the case for collective action. *Human Soc Sci Commun*. 2018;4(1):1–5.
44. Prego-Domínguez J, Khazaeipour Z, Mallah N, Takkouche B. Socioeconomic status and occurrence of chronic pain: a meta-analysis. *Rheumatology*. 2021;60(3):1091–1105. PMID: 33276382. doi:10.1093/rheumatology/keaa758
45. Kurdi MS, Theerth KA, Deva RS. Ketamine: current applications in anesthesia, pain, and critical care. *Anesth Essays Res*. 2014;8(3):283–290. PMID: 25886322; PMCID: PMC4258981. doi:10.4103/0259-1162.143110
46. Grinspoon P. Ketamine for treatment-resistant depression: when and where is it safe? *Harvard Health*. 2022. Available from: [www.health.harvard.edu/blog/ketamine-for-treatment-resistant-depression-when-and-where-is-it-safe-202208092797](http://www.health.harvard.edu/blog/ketamine-for-treatment-resistant-depression-when-and-where-is-it-safe-202208092797).
47. Fortuna A, Schindowski K, Sonvico F. Intranasal drug delivery: challenges and opportunities. *Front Pharmacol*. 2022;13:868986. PMID: 35308245; PMCID: PMC8924485. doi:10.3389/fphar.2022.868986
48. Ulusoy S, Bayar Muluk N, Karpischenko S, et al. Mechanisms and solutions for nasal drug delivery - a narrative review. *Eur Rev Med Pharmacol Sci*. 2022;26(2Suppl):72–81. PMID: 36524914. doi:10.26355/eurrev\_202212\_30487
49. Orhurhu VJ, Vashisht R, Claus LE, Cohen SP. Ketamine Toxicity. In: *StatPearls [Internet]*. Treasure Island (FL): StatPearls Publishing; 2022. PMID: 31082131.
50. Chaves TV, Wilffert B, Sanchez ZM. Overdoses and deaths related to the use of ketamine and its analogues: a systematic review. *Am J Drug Alcohol Abuse*. 2022;1–10. PMID: 36410032. doi:10.1080/00952990.2022.2132506
51. Schak KM, Vande Voort JL, Johnson EK, et al. Potential risks of poorly monitored ketamine use in depression treatment. *Am J Psychiatry*. 2016;173(3):215–218. PMID: 26926127. doi:10.1176/appi.ajp.2015.15081082
52. Palamar JJ, Rutherford C, Keyes KM. Trends in ketamine use, exposures, and seizures in the United States up to 2019. *Am J Public Health*. 2021;111(11):2046–2049. PMID: 34618543; PMCID: PMC8630483. doi:10.2105/AJPH.2021.306486
53. Landrum RE, Banov MD. The online, at-home ketamine experience: a clinician’s dilemma. *Psychiatric Times*. 2022;39(10). Available from: <https://www.psychiatristimes.com/view/the-online-at-home-ketamine-experience-a-clinicians-dilemma>.
54. Fitzhenry RI, editor. *The Harper Book of Quotations*. 3rd ed. New York, NY: Harper; 1993.

Journal of Pain Research

Dovepress

### Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>