

Clinical Outcomes Among Patients With Sickle Cell Disease and Transfusion-Dependent Beta-Thalassemia Treated With Allogeneic Hematopoietic Stem Cell Transplantation: A Systematic Literature Review

Rajeshwari Nair¹ , Nanxin Li¹, Suzan Imren¹, Puja Kohli¹, Krzysztof Lach², Lucía Zhu², Mirella Dudzic²

¹Vertex Pharmaceuticals Incorporated, Boston, MA, USA; ²Maple Health Group LLC, New York, NY, USA

Correspondence: Rajeshwari Nair, Vertex Pharmaceuticals Incorporated, 50 Northern Ave, Boston, MA, 02210, USA, Email Rajeshwari_Nair@vrtx.com

Background: The purpose of this study is to synthesize evidence on disease-specific outcomes in patients with sickle cell disease (SCD) or transfusion-dependent beta-thalassemia (TDT) following allogeneic hematopoietic stem cell transplant (allo-HSCT).

Methods: A systematic literature review (SLR) was conducted in MEDLINE and Embase to identify publications up to May 2023, including patients with SCD or TDT treated with allo-HSCT. Occurrence of vaso-occlusive crises (VOCs) including acute pain, acute chest syndrome, priapism, and splenic sequestration in SCD, and red blood cell transfusion (RBCT) requirements in TDT were the main outcomes of interest. Transplant-related outcomes such as graft-versus-host disease (GVHD) and graft failure/rejection were summarized in the studies that reported main outcomes. Proportion of patients experiencing VOCs or RBCTs, GVHD, and graft failure/rejection after allo-HSCT were aggregated and descriptively reported with range across studies.

Results: Thirty-one SCD studies met inclusion criteria. Twenty-nine studies assessed for VOC and pain crisis events after allo-HSCT; 11 studies reported ≥ 1 VOCs after allo-HSCT in 6.9% of the 2,760 patients. Graft failure was reported in 14.4% (0.9%–18.8%, 14 studies) of patients, graft rejection in 5.5% (1.6%–100.0%, 12 studies) of patients, acute GVHD in 22.4% (1.6%–50.0%, 19 studies) of patients, and chronic GVHD in 20.4% (3.3%–57.1%, 14 studies) of patients. Seventy-eight TDT studies met inclusion criteria. Fifty-six studies reported that 8.8% of the 3,107 patients required RBCTs after allo-HSCT. Graft failure was reported in 5.4% (1.1%–80.0%, 21 studies) of patients, graft rejection in 7.5% (0.5%–42.9%, 50 studies) of patients, acute GVHD in 28.4% (5.2%–100.0%, 57 studies) and chronic GVHD in 15.2% (1.3%–50.0%, 51 studies) of TDT patients.

Conclusion: Based on this SLR, after allo-HSCT, a portion of patients with SCD continue to experience VOCs and a portion of patients with TDT continue to require RBCTs, in addition to experiencing GVHD and graft failure or rejection.

Keywords: allogeneic HSCT, sickle cell disease, vaso-occlusive crisis, beta-thalassemia, transfusion dependent

Background

Sickle cell disease (SCD) and beta-thalassemia are among the most common inherited monogenic disorders globally.¹ It is estimated that 5.0%–7.0% of the world's population is a carrier of a significant hemoglobin variant.² Despite great progress in the understanding of the pathophysiology of these conditions, the burden of disease remains high as evidenced by the significant morbidity and early mortality in affected patients.^{2–4}

SCD is an autosomal recessive disorder caused by a single point variant in the *HBB* gene that encodes the beta-globin, resulting in production of sickle hemoglobin.^{5,6} SCD is characterized by recurrent vaso-occlusive crises (VOCs), progressive vasculopathy, and chronic hemolytic anemia which subsequently leads to end-organ damage and early

mortality.^{6–8} VOCs are a hallmark clinical feature of SCD that manifest as debilitating pain and leads to additional organ complications and increased mortality.^{9,10} Current treatments for SCD primarily consist of pain management and hydroxyurea which may reduce some acute and chronic complications in some patients.^{11,12}

Beta-thalassemia is an autosomal recessive disorder caused by pathogenic variants in the *HBB* gene, resulting in severely reduced or absent production of beta-chains in the adult hemoglobin.¹³ Beta-thalassemia is characterized by ineffective erythropoiesis, chronic anemia, and dysregulated iron metabolism.^{14,15} Transfusion-dependent beta-thalassemia (TDT) is a severe form of beta-thalassemia which causes severe anemia and organ damage.¹⁶ Patients with TDT require lifelong, regular red blood cell transfusions (RBCTs) to survive, and iron chelation therapy to control iron overload and minimize risk of end-organ toxic effects and damage due to hemosiderosis.^{16,17}

Allogeneic hematopoietic stem cell transplantation (allo-HSCT) is a potentially curative option for patients with SCD or TDT.^{14,18} Prior to receipt of allo-HSCT, nearly every patient with SCD experiences ≥ 1 VOC annually and up to 2/3rd patients experience ≥ 3 VOCs per year.^{19,20} Patients with TDT were observed to receive a median of 17.4 RBCTs per year and required regular iron chelation therapy prior to allo-HSCT.²¹ Average life expectancy for patients with SCD and beta-thalassemia on chronic therapies is observed to be about 40 years and 50 years, respectively, in recent times.^{22,23} Eligibility to receive allo-HSCT is limited by the availability of HLA-matched donors and pre-existing comorbidities such as sickling-related or iron-overload-related organ damage among SCD and TDT patients, respectively.^{24,25} Recent advances in transplant techniques such as haploidentical donors or use of reduced intensity conditioning expands eligibility for allo-HSCT. Graft-versus-host disease (GVHD), graft rejection, complications associated with immunosuppression such as infections, and transplant-related mortality are some of the known complications of allo-HSCT.^{25–29} There is a lack of synthesis of the data on disease-specific outcomes after allo-HSCT in SCD and TDT, whether patients become VOC free or transfusion independent (TI) after allo-HSCT to enable evaluation of the effectiveness of allo-HSCT. Impact on quality of life and functional recovery after allo-HSCT are also important measures in assessing the effectiveness of the transplant, however these factors are beyond the scope of this review.

The primary objective of this systematic literature review (SLR) was to identify published literature on patients with SCD or TDT treated with allo-HSCT and synthesize the occurrence of disease-specific outcomes such as VOCs and RBCT requirements after allo-HSCT, respectively. Additionally, transplant-related outcomes such as GVHD and graft failure/rejection were also summarized to characterize the benefits and limitations of allo-HSCT.

Methods

Literature Search Strategy

This SLR was conducted on clinical outcomes after allogeneic HSCT among patients with SCD and TDT, according to the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) guidelines.^{30,31} This review was registered with the international prospective systematic review register (PROSPERO # CRD42023445828).³² Please refer to [Supplementary Table 1](#) for the PRISMA 2020 checklist.

A health sciences librarian performed extensive literature searches in MEDLINE (via PubMed) and Embase (via Elsevier) from inception of the databases to May 2023. Only English language human studies were included in the review. No filters were used for any other parameters. Bibliographies of other literature reviews were hand-searched to identify additional eligible studies. Abstracts for major conferences such as American Society of Hematology (ASH), Tandem Meetings (the American Society for Transplantation and Cellular Therapy [ASTCT] and the Center for International Blood and Marrow Transplant Research [CIBMTR]), European Hematology Association (EHA), European Society for Blood and Marrow Transplantation (EBMT) that were indexed in MEDLINE were also included in the review. Search strategy was developed by researching appropriate subject headings (ie, MeSH, Emtree terms) and free search terms and are available in [Supplementary Table 2](#). The strategy was validated by comparing search results to a sample set of published studies and review papers that were expected to be identified for this literature review.

Study Selection Criteria

Citations identified in the database searches were exported to an EndNote library, where duplicate citations were identified and removed. Citations were then exported to a literature review screening platform called DistillerSR®,

which was used for all phases of the study selection process.³³ Each citation was evaluated against the predetermined inclusion criteria in two screening phases.

Inclusion and exclusion criteria were established a priori. English language studies with patients with SCD or beta-thalassemia of all ages treated with allo-HSCT were eligible to be screened for inclusion. Studies that reported data on at least one of the pre-determined clinical outcomes after allo-HSCT were eligible for full text screening. There was no restriction applied by donor type, stem cell source, conditioning regimen, study design, or geographical location, but studies reporting data on <5 patients were excluded. Commentaries, letters, notes, editorials, pre-clinical studies, and expert opinion were excluded. Previously published SLRs were included only for bibliography check purposes.

Each title and abstract were independently reviewed against the eligibility criteria by two reviewers. Citations proceeded to full-text review if at least one disease-specific SCD or TDT outcome was reported; citations reporting only allo-HSCT-related outcomes were excluded. In cases where the applicability of the inclusion criteria to the title/abstract was unclear and/or the abstract was not available for review, the citations were included to ensure that all potentially relevant studies are captured. Each full-text publication was independently reviewed against the eligibility criteria by two reviewers. Citations were included if at least one disease-related clinical outcome was reported and were excluded if only transplant-related outcomes were reported. Studies providing insufficient information for the evaluation of the eligibility criteria were excluded at this stage to ensure that only relevant publications are captured. The number of publications excluded during the full-text review phase along with their respective primary reason for exclusion was recorded and presented in a PRISMA diagram in Figure 1. For both screening phases, any disagreements between the two reviewers regarding the eligibility of a study were discussed until a consensus was met. No unresolved conflicts required a third reviewer.




Figure 1 PRISMA flow diagram for selection of sickle cell disease (SCD) and transfusion-dependent beta-thalassemia (TDT) studies for SLR.

Note: Date of search execution was May 12, 2023.

Abbreviations: PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; SCD, sickle cell disease; SLR, systematic literature review; TDT, transfusion-dependent beta-thalassemia.

Data Extraction

Data extraction was performed by a single individual for each included study and 100.0% of the outcomes were independently verified by a second reviewer once completed. Any discrepancies or missing information identified by the second reviewer were discussed by both reviewers until a consensus was reached. Unresolved conflicts were decided by a third reviewer. If multiple publications reported the same outcomes from the same study cohort, then full-text publications were prioritized over conference abstracts; if multiple abstracts were published then the latest abstract was prioritized; and larger sample size publication using the same cohort was prioritized over smaller sample size publication. If multiple publications reported different outcomes from the same study cohort, then only the publication that met a greater number of inclusion criteria was used for data extraction.

Data were extracted into an Excel workbook developed and validated for this review. Pre-determined disease-related outcomes, such as occurrence of VOC including pain crisis, acute chest syndrome (ACS), priapism, and splenic sequestration in SCD, transfusion independence in TDT, and transplant-related outcomes such as graft failure/rejection and GVHD for both SCD and TDT, were extracted as available in each included study. For this synthesis, primary graft failure or graft failure events were reported as graft failure, and secondary graft failure or graft rejection events were harmonized to report as graft rejection. Only disease- and transplant-related outcomes after first allo-HSCT were synthesized from SCD and TDT studies included in this review. Additional key elements such as study, population, and transplant characteristics were also extracted.

Data Syntheses

Key results of this SLR were synthesized following the reporting standards outlined in the expanded 2020 PRISMA checklist and presented by disease type (SCD or TDT). Outcomes reported as transfusion independence (TI) and transfusion dependence (TD) were harmonized to report proportion of patients achieving TI after allo-HSCT for all included TDT studies. Proportion was calculated for each outcome by pooling results across studies and dividing the number of patients with the outcome by the total number of patients. Additionally, for each outcome, the proportion of patients experiencing the outcome in each study was also calculated. Results for each outcome was reported in this SLR as the proportion from all studies, and the median proportion with range across proportions calculated for each study. The median and range for proportions considers the variability in sample size across studies. Studies reporting zero frequency for a given outcome were summarized in the review but not included in calculation of proportions. A meta-analysis was not planned for this review due to potential for small number of studies and heterogeneous study characteristics and definitions, such as study population and definition of outcomes. Synthesized results for proportion of patients with SCD who experienced VOC and proportion of patients with TDT who became transfusion independent after allo-HSCT were presented as bubble plots to illustrate the temporal distribution of outcomes.

Assessment of Study Quality

The Newcastle Ottawa Scale (NOS) for cohort and case-control studies,³⁴ and the National Institutes of Health (NIH) quality assessment tools for pre-post studies without control group, case series, and cross-sectional studies³⁵ were used to assess the risk of bias in included studies. Only studies published as full-text articles were subject to quality appraisal. Quality assessment was performed by one reviewer and about 10.0% of these assessments were verified by a second reviewer. Discrepancy between reviewers was resolved by achieving consensus or by a third reviewer.

Results

A comprehensive literature search for the review identified 5,418 non-duplicate full-text articles and abstracts on SCD and TDT treated with allo-HSCT, in MEDLINE and Embase. After title and abstract screening, 1,331 SCD and TDT publications were eligible for full-text screening. Among the full-text screened publications, 120 SCD and 162 TDT publications reported data on clinical outcomes and were included in the review. Results presented here are a focused synthesis of evidence on disease-related outcomes: SCD (VOCs including acute pain crisis, ACS, priapism, splenic sequestration) and TDT (transfusion independence). Thirty-one unique SCD publications (1997–2023) and 78 unique

TDT publications (1986–2023) were synthesized to report results on occurrence of disease-specific and transplant-related outcomes after allo-HSCT (Figure 1 and [Supplementary Tables 3 and 4](#)).

Sickle Cell Disease (SCD)

Study Characteristics

The majority of 31 SCD publications included pediatric patients ($n = 31$, 100.0%),^{36–65} patients with HbSS genotype ($n=18$ of 19 studies that reported genotype, 94.7%),^{37–39,42,44,45,47,49–51,54,56–58,60,62,63,66} and patients underwent HLA-matched sibling donor (MSD) HSCT ($n = 22$ of 29 studies that reported donor type, 75.9%).^{37,40–47,49,51,52,54–61,63–65} Thirteen of the 29 (44.8%) studies included patients treated with myeloablative conditioning.^{37,40,41,46,47,49,55,56,60–64} Twenty-five of the 31 studies reported engraftment data on a total of 2,534 patients.^{36–40,42–49,51,54–60,62,63,65,66} Most were non-interventional studies by design ($n = 24$ of 31, 77.4%),^{36,37,40–52,54–57,61–65} sample sizes varied from 6 to 1,641 patients, more than half included patients from the United States ($n = 16$ of 31, 51.6%)^{38,39,41,44,45,48–50,53,56–61,64,66} and about one-third studies included patients from Europe ($n = 11$ of 31, 35.5%).^{37,40,43,46,47,51,62–66}

VOCs After Allo-HSCT

Acute Pain

Twenty-nine SCD studies assessed occurrence of events referred to as VOC after allo-HSCT among a total of 3,181 patients.^{37–55,57,58,60–67} Terms used to indicate VOC varied considerably among included studies: 11 studies reported as recurrent/refractory/frequent/multiple VOC,^{38,42,44,46,51,52,58,60,62,64,65} 8 as pain crisis or vaso-occlusive pain crisis,^{39,41,43,45,47,61,63,66} 1 as sickling recurrent events,⁴² 3 as vaso-occlusive disease or crisis,^{40,48,55} 1 as painful VOC or ACS,⁴⁶ and 1 as vaso-occlusive events (VOEs) that included acute pain episode, ACS, hepatic or splenic sequestration, and priapism.⁴⁹ Of the 29 studies, 2 assessed VOC that required hospitalization or emergency room (ER) visit,^{49,50} 2 assessed VOC that occurred despite hydroxyurea treatment or were not prevented by medical treatment,^{51,54} and 1 assessed VOE that required intervention in medical setting indicated as ≥ 24 hour hospital stay, or at least 2 visits to a day unit or ER over 72 hours with both requiring intravenous treatment.⁴⁹ Six studies included frequency in their terms to indicate VOCs prior to allo-HSCT: ≥ 3 per year ($n = 3$),^{53,54,59} ≥ 2 per year ($n = 1$),³⁶ >2 per year for 2 years ($n = 1$),⁶⁶ and multiple VOCs per year ($n = 1$).⁵⁸ VOC occurrence after allo-HSCT was more frequently assessed as clinical outcomes in studies published after 2012 and was captured in studies with relatively larger sample sizes (Figure 2). Twenty-two of the 29 studies included patients with SCD,^{38–43,45–52,55,57,59–63,65} and 7 studies included patients with severe or high-risk SCD.^{37,44,53,54,58,64,66}

One hundred ninety-one (6.9%) of 2,760 patients had ≥ 1 VOCs during follow-up after allo-HSCT as reported in 11 studies (Across studies: median: 4.0%, range: 0.4%–32.4%).^{37,40,41,47–50,58,61,64,66} (Figure 2) Of the 25 studies that reported engraftment data in 2,534 patients in the overall SLR, 2,169 patients (85.6%) achieved neutrophil and/or platelet engraftment after allo-HSCT (range - 81.3%–100.0%; one study reported engraftment in 64.7% patients³⁹). Three of these studies reported occurrence of VOC among patients with stable engraftment after allo-HSCT.^{48,49,58} Median duration of follow-up for studies that reported ≥ 1 VOCs after allo-HSCT ranged from 17 months to 95 months.

Eighteen of the 29 studies assessed for VOCs reported that none of the 421 SCD patients experienced any VOC after allo-HSCT with a median duration of follow-up ranging from 19 months to 60 months.^{38,39,42–46,51–55,57,59,60,62,63,65}

Acute Chest Syndrome, Priapism, Splenic Sequestration

SCD studies reported occurrence of ACS or priapism after allo-HSCT; ACS ($n = 3$ studies; 640 patients),^{36,49,61} or priapism ($n = 2$ studies; 385 patients) (Table 1).^{44,61} None of the studies reported occurrence of splenic sequestration after allo-HSCT. Occurrence of ACS or priapism was relatively lower compared to acute pain VOCs after allo-HSCT. ACS was reported in 0.82% after graft failure,³⁶ 1.1% with unknown graft status,⁶¹ and 1.2% after graft rejection⁴⁹ of patients after allo-HSCT at a median follow-up of 48 months, 50 months and 24 months, respectively. Priapism was reported in 0.3%⁶¹ and 3.3%⁴⁴ patients after allo-HSCT at a median follow-up of 50 months and 41 months, respectively.




Figure 2 Roportion of vaso-occlusive crisis (VOC) reported in studies among patients with SCD after allogeneic HSCT. **Abbreviations:** HSCT, hematopoietic stem cell transplant; SCD, sickle cell disease; VOC, vaso-occlusive crisis.

Transfusion-Dependent Beta-Thalassemia (TDT)

Study Characteristics

The majority of TDT studies included pediatric patients (n = 75 of 76 studies that reported age, 98.6%),^{21,25,38,55,67–137} assessed patients treated with MSD HSCT (n=44 of 74 studies that reported donor type, 59.5%),^{21,55,69,70,72,74–81,83–86,89,90,93,94,98,101–103,105,107–111,113,115–121,124,131,133,134,138–140} and myeloablative conditioning (n = 46 of 72 studies that reported conditioning

Table 1 Proportion of Patients With SCD Experiencing Acute Chest Syndrome and Priapism, After Allogeneic HSCT

Author, Year	Sample Size	Transplant Type and Conditioning Regimen	Population	Frequency of Patients with Complication After Allo-HSCT	Median Follow-Up
Acute Chest Syndrome					
Alzahrani et al, 2021 ³⁶	122	MRD, non-MAC	Pediatric, adult	1 (0.82%)	48 months
Stenger et al, 2019 ^{61*}	355	MSD, MAC	Pediatric, adult	4 (1.1%)	50 months
Leonard et al, 2023 ⁴⁹	163	MRD, MUD, Cord blood MAC, non-MAC	Pediatric, adult	2 (1.2%)	24 months
Priapism					
Stenger et al, 2019 ^{61*}	355	MSD, MAC	Pediatric, adult	1 (0.3%)	50 months
Hsieh et al, 2014 ⁴⁴	30	MSD, non-MAC	Adolescents, adult	1 (3.3%)	41 months

Note: *Stenger et al, study was conducted using data from Center for International Blood and Marrow Transplant Research (CIBMTR). 66% patients were administered MAC, 59% patients had MSD allo-HSCT.
Abbreviations: HSCT, hematopoietic stem cell transplant; MAC, myeloablative conditioning; MSD, matched sibling donor; MRD, matched related donor; MUD, matched unrelated donor.

regimen, 63.9%).^{70–72,75,77–84,88–93,96,98–102,104,105,107,108,110,111,116,117,120,122,124,126–128,131,132,134,135,138,139} Forty-seven of the 78 included studies reported on patients with thalassemia-major,^{25,55,69–71,73,74,78,81–84,86,87,91–93,95–98,100,102,104–106,110–117,119,120,128–130,132,133,135,137,141} 28 reported on patients with TDT,^{21,67,72,76,77,79,80,85,88,89,94,99,101,103,107,109,118,121–126,131,134,139,140,142} and 3 reported on patients with beta-thalassemia.^{75,108,127} Most were non-interventional studies by design (n = 73 of 78, 93.6%)^{21,55,68–72,74–80,82–116,118–135,137–141} and had sample sizes varying from 5 to 328 patients. The top three countries from 78 studies were China (n = 15, 19.2%),^{25,76,77,79,80,95,106,122,124,128,129,133,135,137,139} Italy (n = 13, 16.7%),^{21,68,73,75,86,98,104,108,111,113,120,126,131} and India (n = 7, 8.9%).^{67,78,82,87,109,112,118}

Transfusion Outcomes After Allo-HSCT

Seventy-eight studies assessed and reported RBCT requirements at the end of the study follow-up period after allo-HSCT. Fifty-five studies assessed transfusion independence (TI),^{25,38,55,67,69,71,73,74,76–81,86–88,90–93,95–104,106,109,112–119,121,123,124,126–129,132–134,137,139,141,142} 20 studies assessed transfusion dependence (TD),^{21,68,70,72,75,82–85,89,94,105,107,108,120,122,125,131,135,140} and 3 studies assessed both.^{110,111,130} TI was most frequently reported as absence of regular RBCTs and TD was reported as dependence on RBCTs. Only 4 studies reported the duration of RBCT-free period to be classified as TI. Two studies defined this duration as at least 1 year after allo-HSCT,^{91,92} one study defined this duration as no RBCTs starting 2 months after allo-HSCT,¹²⁹ and one other study defined the duration as absence of RBCTs after engraftment.¹¹⁶

Studies included in this review assessed transfusion outcomes in a total of 3,848 patients. A total of 3,784 patients (98.3%) achieved neutrophil and/or platelet engraftment after allo-HSCT. Twenty-one patients (0.6%) died before engraftment. Among engrafted patients, 3,445 (91.1%) patients were alive and TI after allo-HSCT (Across studies: median proportion - 86.6%; range - 20.0%–99.5%) (Figure 3). Fifty-six studies reported that 8.8% patients with graft failure resumed regular RBCTs after allo-HSCT. Median duration of follow-up after allo-HSCT was 6 months to 288 months. A sensitivity analysis excluding studies with <6 months of minimum follow-up after allo-HSCT showed similar TI rates as for overall analysis.^{25,75,76,80,81,88,95,102,106,107,109,112,119,125,128,131,134,140} This approach was implemented to account for the possibility of biased classification of patients with RBCT support immediately after allo-HSCT, as transfusion dependent.




Figure 3 Proportion of transfusion independence reported at end of the study follow-up period among patients with TDT after allogeneic HSCT.

Abbreviations: HSCT, hematopoietic stem cell transplant; TDT, transfusion-dependent beta-thalassemia; TI, transfusion independence.

Transplant-Related Outcomes

SCD

Graft failure or graft rejection were assessed in 27 of the 31 studies (Table 2)^{36–40,42–49,51,53–60,62–66} Fourteen studies reported graft failure in 14.4% patients (Across studies: median: 9.5%; range: 0.9%–18.8%).^{36,38,39,44,45,47–49,56,59,62,64,66,143} Twelve studies reported graft rejection in 5.5% patients (Across studies: median: 6.9%; range: 1.6%–100.0%).^{36,39,40,45,47,53,57–60,63,143}

Acute or chronic GVHD was assessed in 27 of the 31 included studies (Table 2).^{36,37,39,40,42,43,45–49,51–54,56–66} Acute GVHD events occurred in 22.4% patients as reported in 19 studies (Across studies: median: 27.3%; range: 1.6%–50.0%).^{36,37,39,40,43,45,46,49,51–54,56,59,60,63–66} Fourteen studies reported chronic GVHD events in 20.4% patients (Across studies: median: 10.7%; range: 3.3%–57.1%).^{46,48,49,52–54,56,61–66,143}

TDT

Graft failure or graft rejection were assessed in 58 of the 78 studies (Table 2).^{21,38,55,67,69–75,77–79,81,83,84,86,87,89–91,94,96,97,99,102–110,112–120,122–124,126,129–131,133–135,137,140–142} Twenty-one studies reported graft failure events in 5.4% patients (Across studies: median – 8.4%; range - 1.1%–80.0%).^{21,67,74,77,79,81–86,99,102,104,106–108,116,119,124,142} Fifty studies reported graft rejection in 7.5% patients (Across studies: median – 10.0%; range - 0.5%–42.9%).^{21,69–73,75,76,78–81,83–87,89,90,94,96,97,99,102,104–111,115,117–120,122–126,130,131,133,135,138–141}

Acute or chronic GVHDs were assessed in 66 of the 78 studies (Table 2).^{21,55,67,70–82,84,86,88,89,91–93,95–113,115–124,126–129,131–135,137,139,140,142} Fifty-seven studies reported acute GVHD in 28.4% patients (Across studies: median – 26.3%; range - 5.2%–100.0%).^{21,55,67,70–76,78–81,84,88,89,91–93,95–106,108–113,115–119,121–124,128,129,132–135,137,139,140,142} Fifty-one studies reported chronic GVHD in 15.2% patients (Across studies: median – 16.4%; range - 1.3%–50.0%).^{21,67,70,72–80,82,84,86,89,92,93,95–99,101–105,107–109,111,113,116–120,122,123,127–129,131–135,137,140,142}

Table 2 Proportion of Patients With SCD or TDT Experiencing Graft Failure and Graft-Versus-Host Disease After Allogeneic HSCT

Outcome	Number of Studies that Assessed the Outcome	Number of Studies Reporting Patients Experiencing the Outcome After Allo-HSCT	Proportion of Patients Experiencing the Outcome [#]	Study-Level Proportion of Patients Experiencing the Outcome, Median (Range) ^{**}
Sickle Cell Disease				
Graft failure	24	14 (58.3%)	14.4%	9.5% (0.9% - 18.8%)
Graft rejection	16	12 (75.0%)	5.5%	6.9% (1.6% - 100.0%)
Acute GVHD	26	19 (73.1%)	22.4%	27.3% (1.6% - 50.0%)
Chronic GVHD	24	14 (58.3%)	20.4%	10.7% (3.3% - 57.1%)
Transfusion-Dependent Beta-Thalassemia				
Graft failure	78	21 (26.9%)	5.4%	8.4% (1.1% - 80.0%)
Graft rejection	78	50 (64.1%)	7.5%	10.0% (0.5% - 42.9%)
Acute GVHD	61	57 (93.4%)	28.4%	26.3% (5.2% - 100.0%)
Chronic GVHD	60	51 (85.0%)	15.2%	16.4% (1.3% - 50.0%)

Notes: [#] Among studies reporting patients experiencing the outcome after allo-HSCT. ^{**} Data is presented as median proportion and range of proportion across studies.

Abbreviations: GVHD, graft-versus-host disease; HSCT, hematopoietic stem cell transplant; SCD, sickle cell disease; TDT, transfusion-dependent beta-thalassemia.

Risk of Bias Assessment

Twenty SCD studies and 55 TDT studies were assessed for risk of bias using validated and reliable tools for study quality assessment.^{25,36,38,39,42,45,47–51,53,55–60,62,63,65–71,73–81,83,84,86,89–95,97,99–101,105,107–112,115–120,123–132,134,140} Key sources of bias in assessed studies were likely to stem from incomparability between cohorts/cases and controls, as well as issues with reporting of patient follow-up. Several studies did not control for important confounding factors, such as age resulting in study groups that were often incomparable. Another source of uncertainty was related to patient follow-up as there was lack of description of patients lost to follow-up and rationale for choice of follow-up period to assess for disease outcomes.

Discussion

We performed a systematic literature review to synthesize evidence on occurrence of VOCs and RBCT requirements following allo-HSCT in patients with SCD or TDT, respectively. Overall, 6.9% of patients with SCD continue to experience VOCs after allo-HSCT and 8.8% of patients with TDT continue to require regular RBCTs beyond the anticipated transfusion support after allo-HSCT, based on an unadjusted analysis of the included studies. SCD and TDT studies included in this SLR also reported considerable burden due to graft failure, graft rejection, and acute and chronic GVHD after allo-HSCT. Taken together these data provide important context for clinicians when considering allo-HSCT as a treatment option. These results could serve as a rationale for future research to identify patients with SCD who experienced chronic pain prior to their allo-HSCT and are at risk to continue to have chronic pain or acute exacerbations of their chronic pain after allo-HSCT, or patients with TDT who will be at risk for graft failure/rejection and will resume regular transfusions after allo-HSCT. Such findings are likely to inform treatment and management options for patients after allo-HSCT.

To our knowledge, this represents the first SLR on disease-related outcomes in patients with SCD or TDT after allo-HSCT that includes results from both clinical trials and real-world studies; previously published Cochrane reviews on allo-HSCT outcomes in SCD and TDT focused only on clinical trials.^{144,145} A study by Krishnamurti et al using data from the Center for International Blood and Marrow Transplant Research (CIBMTR) registry observed recurrence of pain crisis requiring hospitalization or treatment in 8.7% of SCD patients after allo-HSCT.¹⁴⁶ Prior history of VOC, older age, graft failure, and donor types were noted as important predictors for recurrence of pain crisis following successful allo-HSCT. Studies reporting acute pain events after allo-HSCT indicate the possibility of other etiologies of pain in these patients such as chronic pain due to avascular necrosis of a joint or leg ulcers, neuropathic pain or chronic pain without evident pathology.^{146,147} Another study by Leonard et al, that was included in this SLR, reported that the mean rate of VOE post allo-HSCT was lower in engrafted patients (0–12 months after allo-HSCT: 0.8 per 100-patient years; 12–24 months after allo-HSCT: 0.1 per 100-patient years) compared to those patients with graft rejection (0–12 months after allo-HSCT: 1.1 per 100-patient years; 12–24 months after allo-HSCT: 0.8 per 100-patient years).⁴⁹ Results from the above mentioned studies indicate that patients with SCD who have risk factors such as older age at the time of allo-HSCT, higher pain burden before allo-HSCT, or do not have stable donor engraftment after allo-HSCT could be at higher risk for continued experience of VOCs or pain crisis. Additional studies will be needed to understand the complex etiology of pain events in patients with SCD following allo-HSCT.

Similarly, TDT studies included in this SLR noted that graft failure or graft rejection resulted in thalassemia recurrence and patients resuming regular RBCTs after allo-HSCT. This SLR noted that patients with TDT who had sustained engraftment remained transfusion free, including those with stable mixed chimerism. Two studies reported RBCTs among engrafted patients however these patients subsequently became transfusion independent.^{21,107} Lin et al, reported that 1 patient received one RBCT after allo-HSCT for transient anemia. Merli et al, reported that 7 patients needed RBCT for 3–12 months after allo-HSCT - 2 patients for thalassemia and 5 patients for other reasons including gastrointestinal bleeding. Studies that reported RBCT requirements due to graft failure or graft rejection in patients following allo-HSCT speculated that older age, higher RBCT burden pre-allo-HSCT, higher Pesaro risk score (particularly group III who are patients at higher risk for graft rejection), alloimmunization due to prior RBCTs, and disease-related immunocompromised status as possible reasons for graft failure or graft rejection after allo-HSCT in patients with TDT.^{67,69–71,79,87,90,104,107,126}

In this SLR, variability in VOC occurrence and RBCT requirements following allo-HSCT could not be assessed by age groups, donor type, conditioning regimen, or variable follow-up period as the studies included mixed populations and did not consistently report results by these subgroups. However, it is important to reiterate that post- allo-HSCT VOCs in patients with SCD and RBCTs in patients with TDT maybe related to patient- and/or allo-HSCT- related factors.

Despite advances in prophylaxis and management, GVHD remains a critical complication after allo-HSCT with potential for significant morbidity and mortality. Additionally, GVHD treatment such as corticosteroids could result in adverse events such as infections due to prolonged use of immunosuppression.¹⁴⁸ Acute GVHD that affects up to 70% patients also results in poor prognosis after allo-HSCT.¹⁴⁹ Severe acute GVHD (Grade III–IV) is reported in up to 14% patients despite standard GVHD prophylaxis. These patients require aggressive immunosuppressive treatment, which increases the risk of infections and end organ damage and puts these patients at higher risk for developing chronic GVHD.¹⁴⁹ Chronic GVHD, a late complication of allo-HSCT, affects an estimated 30.0%–50.0% of allo-HSCT recipients and has the potential to substantially impact quality of life in affected patients.^{150–154} Studies have observed chronic GVHD to be a leading cause of mortality among long-term (≥ 2 years) survivors of allo-HSCT.^{155,156}

Results from this review also highlight the challenge associated with assessing the exact incidence of VOCs following allo-HSCT due to the lack of standardized outcome definitions and consistent outcome reporting. A prior SLR by Zaidi et al, conducted to assess frequency of VOCs in patients with SCD treated with standard of care in real-world settings, reported similar variability in VOC definitions.²⁰ Specifically, that review reported differences in VOC definitions across studies in terms of events that were classified as VOC and follow-up time for reporting VOCs. This current review also observed differences in VOC reporting among studies of patients with SCD after allo-HSCT. This review also observed potential for variable effectiveness findings of allo-HSCT in TDT due to differences in definition of transfusion independence/dependence, lack of consistent reporting of time from allo-HSCT to beginning of RBCT-free period, as well as duration of RBCT-free period that would deem patients to be transfusion independent. It is critical to develop and implement standardized definitions for outcomes to avoid reporting inconsistent or heterogeneous treatment effects of allo-HSCT.

There are some limitations to the current review. There was considerable heterogeneity in inclusion criteria for studies, age groups, conditioning regimens, donor sources, and follow-up period. Variability in outcome definitions for VOCs in SCD studies and RBCT requirements in TDT studies could impact comparisons of outcome results between studies; a meta-analysis was not conducted due to the likely lack of feasibility due to this clinical and methodological heterogeneity, and due to limitation from pooling of unadjusted results. Graft failure and graft rejection were summarized as reported in included studies. There is potential for misclassification of events between these categories if studies reported without differentiation of failure and rejection. Transplant outcomes were summarized only from publications that reported VOCs and RBCTs for SCD and TDT, respectively. There is potential for underestimation of transplant outcomes based on the study inclusion criteria for this review. Lack of standardized diagnostic practices and representation of geographies with highest burden of SCD and TDT such as sub-Saharan Africa or South Asia could potentially result in evidence gaps on the effectiveness of allo-HSCT and impact generalizability of results to these areas. Finally, only English language studies were included, so there is potential for publication bias even though studies were included from all countries.

In conclusion, synthesis of available evidence on disease-related outcomes shows that on average 1 in 14 patients with SCD continue to experience VOCs or acute pain episodes despite having stable donor-derived hematopoiesis, and on average 1 in 11 patients with TDT, mostly those with graft failure or rejection, continue to require RBCTs after allo-HSCT. This highlights that allo-HSCT may not be consistently successful in all transplanted patients, with the potential for residual disease burden in addition to significant transplant-related burden from graft failure/rejection, acute/chronic GVHD, infections, life-threatening organ damage, and mortality.

Acknowledgments

We would like to thank Nathan Blow for editorial support and Meaghan Muir for literature search strategy under the guidance of the authors. Both NB and MM are employees of Vertex Pharmaceuticals Incorporated and may own stock or stock options in the company.

Disclosure

Rajeshwari Nair, Nanxin Li, Suzan Imren, and Puja Kohli are employees of Vertex Pharmaceuticals Incorporated and may own stock or stock options in the company. Lucia Zhu was an employee of Maple Health Group at the time of conduct of the study, Krzysztof Lach and Mirella Dudzic are consultants of Maple Health Group, LLC, which received funding from Vertex Pharmaceuticals Incorporated in connection with this research. The authors report no other conflicts of interest in this work.

References

1. Harteveld CL, Achour A, Arkesteijn SJG, et al. The hemoglobinopathies, molecular disease mechanisms and diagnostics. *Int J Lab Hematol*. 2022;44(Suppl 1):28–36. doi:10.1111/ijlh.13885
2. McGann PT, Nero AC, Ware RE. Clinical Features of β -Thalassemia and Sickle Cell Disease. *Adv Exp Med Biol*. 2017;1013:1–26. doi:10.1007/978-1-4939-7299-9_1
3. Udeze C, Evans KA, Yang Y, et al. Economic and clinical burden of managing transfusion-dependent β -thalassemia in the United States. *J Med Econ*. 2023;26(1):924–932. doi:10.1080/13696998.2023.2235928
4. Udeze C, Evans KA, Yang Y, et al. Economic and Clinical Burden of Managing Sickle Cell Disease with Recurrent Vaso-Occlusive Crises in the United States. *Adv Ther*. 2023;40(8):3543–3558. doi:10.1007/s12325-023-02545-7
5. Piel FB, Steinberg MH, Rees DC. Sickle Cell Disease. *N Engl J Med*. 2017;376(16):1561–1573. doi:10.1056/NEJMra1510865
6. Kato GJ, Piel FB, Reid CD, et al. Sickle cell disease. *Nat Rev Dis Primers*. 2018;4:18010. doi:10.1038/nrdp.2018.10
7. Quinn CT. Minireview: clinical severity in sickle cell disease: the challenges of definition and prognostication. *Exp Biol Med*. 2016;241(7):679–688. doi:10.1177/1535370216640385
8. Sundd P, Gladwin MT, Novelli EM. Pathophysiology of Sickle Cell Disease. *Annu Rev Pathol*. 2019;14:263–292. doi:10.1146/annurev-pathmechdis-012418-012838
9. Ware RE, de Montalembert M, Tshilolo L, Abboud MR. Sickle cell disease. *Lancet*. 2017;390(10091):311–323. doi:10.1016/s0140-6736(17)30193-9
10. Gardner K, Douiri A, Drasar E, et al. Survival in adults with sickle cell disease in a high-income setting. *Blood*. 2016;128(10):1436–1438. doi:10.1182/blood-2016-05-716910
11. Elendu C, Amaechi DC, Alakwe-Ojimba CE, et al. Understanding Sickle cell disease: causes, symptoms, and treatment options. *Medicine*. 2023;102(38):e35237. doi:10.1097/md.00000000000035237
12. McGann PT, Ware RE. Hydroxyurea therapy for sickle cell anemia. *Expert Opin Drug Saf*. 2015;14(11):1749–1758. doi:10.1517/14740338.2015.1088827
13. Cao A, Galanello R. Beta-thalassemia. *Genet Med*. 2010;12(2):61–76. doi:10.1097/GIM.0b013e3181cd68ed
14. Taher AT, Musallam KM, Cappellini MD. β -Thalassemias. *N Engl J Med*. 2021;384(8):727–743. doi:10.1056/NEJMra2021838
15. Origa R. β -Thalassemia. *Genet Med*. 2017;19(6):609–619. doi:10.1038/gim.2016.173
16. Jobanputra M, Paramore C, Laird SG, McGahan M, Telfer P. Co-morbidities and mortality associated with transfusion-dependent beta-thalassaemia in patients in England: a 10-year retrospective cohort analysis. *Br J Haematol*. 2020;191(5):897–905. doi:10.1111/bjh.17091
17. Lal A, Wong T, Keel S, et al. The transfusion management of beta thalassemia in the United States. *Transfusion*. 2021;61(10):3027–3039. doi:10.1111/trf.16640
18. Gluckman E, Cappelli B, Bernaudin F, et al. Sickle cell disease: an international survey of results of HLA-identical sibling hematopoietic stem cell transplantation. *Blood*. 2017;129(11):1548–1556. doi:10.1182/blood-2016-10-745711
19. Evensen CT, Treadwell MJ, Keller S, et al. Quality of care in sickle cell disease: cross-sectional study and development of a measure for adults reporting on ambulatory and emergency department care. *Medicine*. 2016;95(35):e4528. doi:10.1097/MD.00000000000004528
20. Zaidi AU, Glaros AK, Lee S, et al. A systematic literature review of frequency of vaso-occlusive crises in sickle cell disease. *Orphanet J Rare Dis*. 2021;16(1):460. doi:10.1186/s13023-021-02096-6
21. Merli P, Ruggeri A, Algeri M, et al. Clinical Outcomes after Allogeneic Hematopoietic Stem Cell Transplantation in Patients with Transfusion-Dependent β -Thalassemia Treated at the Bambino Gesù Children's Hospital, Rome, Italy. *Blood*. 2019;134:969. doi:10.1182/blood-2019-123440
22. Farmakis D, Giakoumis A, Angastiniotis M, Eleftheriou A. The changing epidemiology of the ageing thalassaemia populations: a position statement of the Thalassaemia International Federation. *Eur J Haematol*. 2020;105(1):16–23. doi:10.1111/ejh.13410
23. Jiao B, Johnson KM, Ramsey SD, Bender MA, Devine B, Basu A. Long-term survival with sickle cell disease: a nationwide cohort study of Medicare and Medicaid beneficiaries. *Blood Adv*. 2023;7(13):3276–3283. doi:10.1182/bloodadvances.2022009202
24. Walters MC, De Castro LM, Sullivan KM, et al. Indications and Results of HLA-Identical Sibling Hematopoietic Cell Transplantation for Sickle Cell Disease. *Biol Blood Marrow Transplant*. 2016;22(2):207–211. doi:10.1016/j.bbmt.2015.10.017
25. Li C, Mathews V, Kim S, et al. Related and unrelated donor transplantation for β -thalassemia major: results of an international survey. *Blood Adv*. 2019;3(17):2562–2570. doi:10.1182/bloodadvances.2019000291
26. Hsieh MM, Fitzhugh CD, Tisdale JF. Allogeneic hematopoietic stem cell transplantation for sickle cell disease: the time is now. *Blood*. 2011;118(5):1197–1207. doi:10.1182/blood-2011-01-332510
27. Shenoy S. Hematopoietic stem cell transplantation for sickle cell disease: current practice and emerging trends. *Hematology Am Soc Hematol Educ Program*. 2011;2011:273–279. doi:10.1182/asheducation-2011.1.273
28. Cappellini MD, Motta I. New therapeutic targets in transfusion-dependent and -independent thalassemia. *Hematology Am Soc Hematol Educ Program*. 2017;2017(1):278–283. doi:10.1182/asheducation-2017.1.278
29. Baronciani D, Angelucci E, Potschger U, et al. Hemopoietic stem cell transplantation in thalassemia: a report from the European Society for Blood and Bone Marrow Transplantation Hemoglobinopathy Registry, 2000–2010. *Bone Marrow Transplant*. 2016;51(4):536–541. doi:10.1038/bmt.2015.293

30. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021;372:n71. doi:10.1136/bmj.n71
31. Page MJ, Moher D, Bossuyt PM, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ*. 2021;372:n160. doi:10.1136/bmj.n160
32. Nair R, Li N, Imren S, et al. Systematic Literature Review of Allogeneic Hematopoietic Stem Cell Transplantation Clinical, Humanistic, and Economic Outcomes in Sickle Cell Disease and Transfusion-Dependent Beta-Thalassemia. National Institute for Health and Care Research (NIHR) PROSPERO: International prospective register of systematic reviews. Updated 18 September 2023. Available from: https://www.crd.york.ac.uk/prospero/display_record.php?ID=CRD42023445828. Accessed June, 2023.
33. DistillerSR Inc. DistillerSR. Version 2.35. 2024.
34. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*. 2010;25(9):603–605. doi:10.1007/s10654-010-9491-z
35. Ma LL, Wang YY, Yang ZH, Huang D, Weng H, Zeng XT. Methodological quality (risk of bias) assessment tools for primary and secondary medical studies: what are they and which is better? *Mil Med Res*. 2020;7(1):7. doi:10.1186/s40779-020-00238-8
36. Alzahrani M, Damlaj M, Jeffries N, et al. Non-myeloablative human leukocyte antigen-matched related donor transplantation in sickle cell disease: outcomes from three independent centres. *Br J Haematol*. 2021;192(4):761–768. doi:10.1111/bjh.17311
37. Bernaudin F, Dalle JH, Bories D, et al. Long-term event-free survival, chimerism and fertility outcomes in 234 patients with sickle-cell anemia younger than 30 years after myeloablative conditioning and matched-sibling transplantation in France. *Haematologica*. 2020;105(1):91–101. doi:10.3324/haematol.2018.213207
38. Bolaños-Meade J, Cooke KR, Gamper CJ, et al. Effect of increased dose of total body irradiation on graft failure associated with HLA-haploidentical transplantation in patients with severe haemoglobinopathies: a prospective clinical trial. *Lancet Haematol*. 2019;6(4):e183–e193. doi:10.1016/s2352-3026(19)30031-6
39. Bolaños-Meade J, Fuchs EJ, Luznik L, et al. HLA-haploidentical bone marrow transplantation with posttransplant cyclophosphamide expands the donor pool for patients with sickle cell disease. *Blood*. 2012;120(22):4285–4291. doi:10.1182/blood-2012-07-438408
40. García Morin M, Cela E, Marco Sánchez JM, et al. 48 HLA-identical related hematopoietic stem cell transplants in sickle cell disease: improving results during the last decade. *Bone Marrow Transplant*. 2022;57:245–246.
41. Guilcher GMT, Dalal S, Suresh T, et al. Clinical Outcomes and Long-Term Follow-up of Children with Mixed Myeloid Chimerism Following Hematopoietic Cell Transplantation (HCT) for Sickle Cell Disease (SCD). *Transplant Cell Ther*. 2023;29(2):S296–S297. doi:10.1016/S2666-6367(23)00460-8
42. Guilcher GMT, Monagel DA, Nettel-Aguirre A, et al. Nonmyeloablative Matched Sibling Donor Hematopoietic Cell Transplantation in Children and Adolescents with Sickle Cell Disease. *Biol Blood Marrow Transplant*. 2019;25(6):1179–1186. doi:10.1016/j.bbmt.2019.02.011
43. Hartz B, Schrum J, Grosse R, Kordes U, Bleeke M, Müller I. Conditioning with treosulfan, fludarabine, thiopeta and ATG for matched family donor transplantation in children with sickle cell anemia. *Bone Marrow Transplant*. 2014;49:S318–S9.
44. Hsieh MM, Fitzhugh CD, Weitzel RP, et al. Nonmyeloablative HLA-matched sibling allogeneic hematopoietic stem cell transplantation for severe sickle cell phenotype. *JAMA*. 2014;312(1):48–56. doi:10.1001/jama.2014.7192
45. Iannone R, Casella JF, Fuchs EJ, et al. Results of minimally toxic nonmyeloablative transplantation in patients with sickle cell anemia and beta-thalassemia. *Biol Blood Marrow Transplant*. 2003;9(8):519–528. doi:10.1016/s1083-8791(03)00192-7
46. Isgro A, Sodani P, Marziani M, et al. Reduction of intramedullary apoptosis after stem cell transplantation in black African variant of pediatric sickle cell anemia. *Mediterr J Hematol Infect Dis*. 2014;6(1):e2014054. doi:10.4084/mjhid.2014.054
47. Kogel F, Hakimeh D, Sodani P, et al. Allogeneic hematopoietic stem cell transplantation from sibling and unrelated donors in pediatric patients with sickle cell disease-A single center experience. *Pediatr Transplant*. 2021;25(2):e13892. doi:10.1111/petr.13892
48. Krishnamurti L, He Z, Deng Y, Hamidi R, Flagg A, Shah NC. Vaso-Occlusive Pain Requiring Hospitalization or Treatment Is Rare Following Successful Hematopoietic Cell Transplantation for Sickle Cell Disease. *Transplant Cell Ther*. 2021;29(2):S291. doi:10.1016/S2666-6367(23)00453-0
49. Leonard A, Furstenu D, Abraham A, et al. Reduction in vaso-occlusive events following stem cell transplantation in patients with sickle cell disease. *Blood Adv*. 2023;7(2):227–234. doi:10.1182/bloodadvances.2022008137
50. Mahesri M, Schneeweiss S, Globe D, et al. Clinical outcomes following bone marrow transplantation in patients with sickle cell disease: a cohort study of US Medicaid enrollees. *Eur J Haematol*. 2021;106(2):273–280. doi:10.1111/ejh.13546
51. Marzollo A, Calore E, Tumino M, et al. Treosulfan-Based Conditioning Regimen in Sibling and Alternative Donor Hematopoietic Stem Cell Transplantation for Children with Sickle Cell Disease. *Mediterr J Hematol Infect Dis*. 2017;9(1):e2017014. doi:10.4084/mjhid.2017.014
52. Mohamed SY, Alzahrani H, Ibrahim KT, et al. Pre-Conditioning Steroids and Hydroxyurea Followed By Reduced Toxicity Conditioning (RTC) with ATG-Bu-FLU Is Safe and Effective in Allo-Sib-HSCT in Adolescents and Adults with Sickle Cell Anemia (SCA). *Biol Blood Marrow Transplant*. 2015;21(2):S38–S40. doi:10.1016/j.bbmt.2014.11.032
53. Ngwube A, Shah N, Godder K, Jacobsohn D, Hulbert ML, Shenoy S. Abatacept is effective as GVHD prophylaxis in unrelated donor stem cell transplantation for children with severe sickle cell disease. *Blood Adv*. 2020;4(16):3894–3899. doi:10.1182/bloodadvances.2020002236
54. Ozdogu H, Boga C, Yeral M, et al. Excellent outcomes of allogeneic transplantation from peripheral blood of HLA-matched related donors for adult sickle cell disease with ATLG and posttransplant cyclophosphamide-containing regimen: an update work. *Bone Marrow Transplant*. 2020;55(8):1647–1651. doi:10.1038/s41409-020-0805-9
55. Park BK, Kim HS, Kim S, et al. Allogeneic hematopoietic stem cell transplantation in congenital hemoglobinopathies with myeloablative conditioning and rabbit anti-thymocyte globulin. *Blood Res*. 2018;53(2):145–151. doi:10.5045/br.2018.53.2.145
56. Patel A, Wilkerson K, Chen H, et al. Reduction in the Prevalence of Thrombotic Events in Sickle Cell Disease after Allogeneic Hematopoietic Transplantation. *Transplant Cell Ther*. 2022;28(5):277.e1–277.e6. doi:10.1016/j.jctc.2022.02.010
57. Saraf SL, Ghimire K, Patel P, et al. Improved health care utilization and costs in transplanted versus non-transplanted adults with sickle cell disease. *PLoS One*. 2020;15(2):e0229710. doi:10.1371/journal.pone.0229710
58. Saraf SL, Oh AL, Patel PR, et al. Nonmyeloablative Stem Cell Transplantation with Alemtuzumab/Low-Dose Irradiation to Cure and Improve the Quality of Life of Adults with Sickle Cell Disease. *Biol Blood Marrow Transplant*. 2016;22(3):441–448. doi:10.1016/j.bbmt.2015.08.036

59. Shah NC, Bhoopatiraju S, Abraham A, et al. Granulocyte Colony-Stimulating Factor Is Safe and Well Tolerated following Allogeneic Transplantation in Patients with Sickle Cell Disease. *Transplant Cell Ther.* 2022;28(3):174.e1–174.e5. doi:10.1016/j.jtct.2021.12.016
60. Soni S, Gross TG, Rangarajan H, Baker KS, Sturm M, Rhodes M. Outcomes of matched sibling donor hematopoietic stem cell transplantation for severe sickle cell disease with myeloablative conditioning and intermediate-dose of rabbit anti-thymocyte globulin. *Pediatr Blood Cancer.* 2014;61(9):1685–1689. doi:10.1002/pbc.25059
61. Stenger E, Phelan R, Shaw BE, et al. Excellent Overall Survival and Low Incidence of Late Effects in Patients Undergoing Allogeneic Hematopoietic Cell Transplant for Sickle Cell Disease: a Report from the Center for International Blood and Marrow Transplant Research (CIBMTR). *Blood.* 2019;134:697. doi:10.1182/blood-2019-128797
62. Strocchio L, Zecca M, Comoli P, et al. Treosulfan-based conditioning regimen for allogeneic haematopoietic stem cell transplantation in children with sickle cell disease. *Br J Haematol.* 2015;169(5):726–736. doi:10.1111/bjh.13352
63. Vermynen C, Cornu G, Ferster A, et al. Haematopoietic stem cell transplantation for sickle cell anaemia: the first 50 patients transplanted in Belgium. *Bone Marrow Transplant.* 1998;22(1):1–6. doi:10.1038/sj.bmt.1701291
64. Walters MC, Patience M, Leisenring W, et al. Collaborative multicenter investigation of marrow transplantation for sickle cell disease: current results and future directions. *Biol Blood Marrow Transplant.* 1997;3(6):310–315.
65. Lucarelli G, Isgrò A, Sodani P, et al. Hematopoietic SCT for the Black African and non-Black African variants of sickle cell anemia. *Bone Marrow Transplant.* 2014;49(11):1376–1381. doi:10.1038/bmt.2014.167
66. de la Fuente J, Dhedin N, Koyama T, et al. Haploidentical Bone Marrow Transplantation with Post-Transplantation Cyclophosphamide Plus Thiotepa Improves Donor Engraftment in Patients with Sickle Cell Anemia: results of an International Learning Collaborative. *Biol Blood Marrow Transplant.* 2019;25(6):1197–1209. doi:10.1016/j.bbmt.2018.11.027
67. Shah SA, Shah KM, Patel KA, et al. Unrelated Umbilical Cord Blood Transplant for Children with β -Thalassemia Major. *Indian J Hematol Blood Transfus.* 2015;31(1):9–13. doi:10.1007/s12288-014-0391-3
68. Angelucci E, Muretto P, Nicolucci A, et al. Effects of iron overload and hepatitis C virus positivity in determining progression of liver fibrosis in thalassemia following bone marrow transplantation. *Blood.* 2002;100(1):17–21. doi:10.1182/blood.v100.1.17
69. Aydogdu S, Toret E, Aksoy BA, et al. Comparison of Hematopoietic Stem Cell Transplantation Results in Patients with β -Thalassemia Major from Three Different Graft Types. *Hemoglobin.* 2021;45(1):25–29. doi:10.1080/03630269.2021.1872611
70. Ball LM, Lankester AC, Giordano PC, et al. Paediatric allogeneic bone marrow transplantation for homozygous beta-thalassaemia, the Dutch experience. *Bone Marrow Transplant.* 2003;31(12):1081–1087. doi:10.1038/sj.bmt.1704066
71. Bazarbachi A, Hatoum H, Mugharbel A, et al. Hematopoietic stem cell transplantation in Lebanon: first comprehensive report. *Bone Marrow Transplant.* 2008;42:S96–s102. doi:10.1038/bmt.2008.128
72. Benakli M, Redhouane AN, Belhadj R, et al. A retrospective long-term study from allogeneic haematopoietic stem cells transplantation in major β -thalassaemia: the Algerian experience. *Haematologica.* 2017;102:84–85.
73. Bernardo ME, Zecca M, Piras E, et al. Treosulfan-based conditioning regimen for allogeneic haematopoietic stem cell transplantation in patients with thalassaemia major. *Br J Haematol.* 2008;143(4):548–551. doi:10.1111/j.1365-2141.2008.07385.x
74. Boulad F, Giardina P, Gillio A, et al. Bone marrow transplantation for homozygous beta-thalassemia. The Memorial Sloan-Kettering Cancer Center experience. *Ann N Y Acad Sci.* 1998;850:498–502. doi:10.1111/j.1749-6632.1998.tb10532.x
75. Caocci G, Efficace F, Ciotti F, et al. Prospective assessment of health-related quality of life in pediatric patients with beta-thalassemia following hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant.* 2011;17(6):861–866. doi:10.1016/j.bbmt.2010.09.011
76. Chen H, Li XY, Zhan LP, et al. Prediction, management, and prognosis of mixed chimerism after hematopoietic stem cell transplantation in transfusion-dependent pediatric thalassemia patients. *Pediatr Transplant.* 2020;24(8):e13876. doi:10.1111/petr.13876
77. Cheuk DK, Mok AS, Lee AC, et al. Quality of life in patients with transfusion-dependent thalassemia after hematopoietic SCT. *Bone Marrow Transplant.* 2008;42(5):319–327. doi:10.1038/bmt.2008.165
78. Choudhary D, Sharma SK, Gupta N, et al. Treosulfan-thiotepa-fludarabine-based conditioning regimen for allogeneic transplantation in patients with thalassemia major: a single-center experience from north India. *Biol Blood Marrow Transplant.* 2013;19(3):492–495. doi:10.1016/j.bbmt.2012.11.007
79. Fang J, Huang S, Chen C, et al. Umbilical cord blood transplantation in Chinese children with beta-thalassemia. *J Pediatr Hematol Oncol.* 2004;26(3):185–189. doi:10.1097/00043426-200403000-00008
80. Fang J, Huang S, Chen C, Zhou D, Wu Y, Bao R. Allogeneic peripheral blood stem cell transplantation in beta-thalassemia. *Pediatr Hematol Oncol.* 2002;19(7):453–458. doi:10.1080/08880010290097260
81. Farzana T, Shamshi TS, Irfan M, Ansari SH, Baig MI, Shakoar N. Peripheral blood stem cell transplantation in children with beta-thalassemia major. *J Coll Physicians Surg Pak.* 2003;13(4):204–206.
82. Fouzia N, Sindhuvi E, Kavitha M, et al. Long Term Outcome of Allogeneic Hematopoietic Stem Cell Transplantation for Beta Thalassemia Major. *Blood.* 2016;128(22):4606. doi:10.1182/blood.V128.22.4606.4606
83. Frappaz D, Gluckman E, Souillet G, et al. Bone marrow transplantation (BMT) for thalassemia major (TM). The French experience. *Prog Clin Biol Res.* 1989;309:207–216.
84. Galambrun C, Pondarré C, Bertrand Y, et al. French multicenter 22-year experience in stem cell transplantation for beta-thalassemia major: lessons and future directions. *Biol Blood Marrow Transplant.* 2013;19(1):62–68. doi:10.1016/j.bbmt.2012.08.005
85. Gassas A, O'Boyle F, Chaudhury S, et al. Complex Clonal Evolution Can Occur Following Transplantation for Transfusion Dependent Thalassaemia in the Context of Mixed Myeloid Chimerism and Reduced Conditioning Regimens. *Blood.* 2021;138:2906. doi:10.1182/blood-2021-153519
86. Gaziev J, Isgrò A, Sodani P, et al. Optimal Outcomes in Young Class 3 Patients With Thalassemia Undergoing HLA-Identical Sibling Bone Marrow Transplantation. *Transplantation.* 2016;100(4):925–932. doi:10.1097/TP.0000000000000928
87. George B, Mathews V, Sindhuvi E, et al. Outcome of Mixed Chimerism in Children Undergoing Allogeneic Stem Cell Transplant (HSCT) for Thalassemia Major. *Biol Blood Marrow Transplant.* 2012;18(2):S231. doi:10.1016/j.bbmt.2011.12.085
88. Ghanem KM, Hariss N, Koussa S, et al. Setting-up a transplant service for thalassemia in a middle-income country. *Bone Marrow Transplant.* 2019;53:484–485.

89. Goussetis E, Peristeri I, Kitra V, et al. HLA-matched sibling stem cell transplantation in children with β -thalassemia with anti-thymocyte globulin as part of the preparative regimen: the Greek experience. *Bone Marrow Transplant.* 2012;47(8):1061–1066. doi:10.1038/bmt.2011.219
90. Graphacos S, Kitra V, Peristeri J, Goussetis E, Papadakis V. Haemopoietic transplantation for thalassaemic children: the Greek experience. *Bone Marrow Transplant.* 1997;19(SUPPL 2):68–69.
91. Hamidieh AA, Hamidi Z, Behfar M, et al. Relationship between endocrine changes and bone markers in pediatric thalassemic patients after hematopoietic stem cell transplantation. *Minerva Pediatr.* 2021;73(5):414–419. doi:10.23736/s2724-5276.16.04553-9
92. Hamidieh AA, Hamidi Z, Nedaeifard L, et al. Changes of bone density in pediatric patients with β -thalassemia major after allogeneic hematopoietic stem cell transplantation. *Arch Iran Med.* 2013;16(2):88–92.
93. Hamidieh AA, Tayebi S, Moeninia F, et al. T2(*) MRI changes in the heart and liver of ex-thalassemic patients after hematopoietic stem cell transplantation. *Hematol Oncol Stem Cell Ther.* 2014;7(3):103–108. doi:10.1016/j.hemonc.2014.06.002
94. Ho WL, Lin KH, Wang JD, et al. Financial burden of national health insurance for treating patients with transfusion-dependent thalassemia in Taiwan. *Bone Marrow Transplant.* 2006;37(6):569–574. doi:10.1038/sj.bmt.1705287
95. Hong X, Chen Y, Lu J, Lu Q. Addition of ruxolitinib in Graft-versus-Host disease prophylaxis for pediatric β -Thalassemia major patients after allogeneic stem cell transplantation: a retrospective cohort study. *Pediatr Transplant.* 2023;27(2):e14466. doi:10.1111/ptr.14466
96. Hussein AA, Al-Zaben A, Khattab E, Haroun A, Frangoul H. Hematopoietic stem cell transplantation from non-sibling matched family donors for patients with thalassemia major in Jordan. *Pediatr Transplant.* 2016;20(1):120–123. doi:10.1111/ptr.12622
97. Hussein AA, Al-Zaben AI, Khattab E, Haroun A, Frangoul HA. Allogeneic Hematopoietic Stem Cell Transplantation Using a Reduced Intensity Conditioning Regimen for Adolescents and Young Adults with Class 3 Beta-Thalassemia. *Biol Blood Marrow Transplant.* 2016;22(3):S248. doi:10.1016/j.bbmt.2015.11.670
98. Inati A, Gaziev J, Abbas HA, et al. Outcome Of Bone Marrow Transplantation In Lebanese Children With β -Thalassemia Major. *Blood.* 2013;122(21):5505. doi:10.1182/blood.V122.21.5505.5505
99. Jaing TH, Hung IJ, Yang CP, et al. Unrelated cord blood transplantation for thalassaemia: a single-institution experience of 35 patients. *Bone Marrow Transplant.* 2012;47(1):33–39. doi:10.1038/bmt.2011.39
100. Jaing TH, Hung IJ, Yang CP, Chen SH, Sun CF, Chow R. Rapid and complete donor chimerism after unrelated mismatched cord blood transplantation in 5 children with beta-thalassemia major. *Biol Blood Marrow Transplant.* 2005;11(5):349–353. doi:10.1016/j.bbmt.2005.02.003
101. Jitpirasakun S, Pooliam J, Sriwichakorn C, Sanpakit K, Nakavachara P. Differences in longitudinal growth patterns of children and adolescents with transfusion-dependent hemoglobin E/ β -thalassemia and those achieving successful hematopoietic stem-cell transplantation. *Int J Hematol.* 2022;115(4):575–584. doi:10.1007/s12185-021-03279-4
102. Karasu G, Uygur V, Hazar V, et al. Hematopoietic stem cell transplantation in patients with thalassemia major: better in younger. *Bone Marrow Transplant.* 2019;54:128–129.
103. Kleinschmidt K, Foell J, Pfistering B, et al. Haploidentical T-Cell Depleted HSCT Represents a Safe Therapeutic Alternative to MSD Transplantation in Patients with Transfusion-dependent β -Thalassemia Major. *Bone Marrow Transplant.* 2020;55(Suppl):161–162.
104. La Nasa G, Giardini C, Argioli F, et al. Unrelated donor bone marrow transplantation for thalassemia: the effect of extended haplotypes. *Blood.* 2002;99(12):4350–4356. doi:10.1182/blood.v99.12.4350
105. Lawson SE, Roberts IA, Amrolia P, Dokal I, Szydlo R, Darbyshire PJ. Bone marrow transplantation for beta-thalassaemia major: the UK experience in two paediatric centres. *Br J Haematol.* 2003;120(2):289–295. doi:10.1046/j.1365-2141.2003.04065.x
106. Li C, He Y, Liao J, et al. Unrelated Cord Blood Following Haploidentical HSCT with Post-Transplant Cyclophosphamide Improves Results of Alternative Donor Transplant in Thalassemia Major. *Blood.* 2015;126(23):5531. doi:10.1182/blood.V126.23.5531.5531
107. Lin KH, Lin KS, Feig SA. Marrow transplantation for thalassemia. *Bone Marrow Transplant.* 1986;1(2):115–120.
108. Lucarelli G, Galimberti M, Polchi P, et al. Marrow transplantation in patients with thalassemia responsive to iron chelation therapy. *N Engl J Med.* 1993;329(12):840–844. doi:10.1056/nejm199309163291204
109. Mehta DMP, Khushoo V. Is Old (Fludarabine/Busulfan/Cyclophosphamide/rAntiThymocyteGlobulin) Conditioning Still Gold for Allogeneic Transplants in Transfusion Dependent Beta-Thalassemia of All Risk Categories in 21st Century? *Indian J Hematol Blood Transfus.* 2023;39(4):1–9. doi:10.1007/s12288-023-01646-1
110. Mentzer WC, Cowan MJ. Bone marrow transplantation for beta-thalassemia: the University of California San Francisco experience. *J Pediatr Hematol Oncol.* 2000;22(6):598–601. doi:10.1097/00043426-200011000-00030
111. Mulas O, Caocci G, Efficace F, et al. Long-term health-related quality of life in patients with β -thalassemia after unrelated hematopoietic stem cell transplantation. *Bone Marrow Transplant.* 2022;57(12):1833–1836. doi:10.1038/s41409-022-01823-2
112. Mullanfiroze K, Uppuluri R, Subburaj D, et al. Matched unrelated donor HSCT for thalassemia major using treosulfan based conditioning protocol for children: a single center experience from India. *Pediatr Hematol Oncol J.* 2017;2(1):7–11. doi:10.1016/j.phoj.2017.05.001
113. Muretto P, Angelucci E, Lucarelli G. Reversibility of cirrhosis in patients cured of thalassemia by bone marrow transplantation. *Ann Intern Med.* 2002;136(9):667–672. doi:10.7326/0003-4819-136-9-200205070-00009
114. O'Boyle F, Karnik L, Bradshaw A, Szydlo RM, de la Fuente J. Fludarabine/Treosulfan/Thiotepa/ATG Conditioning for Matched and Single Antigen Mismatched Unrelated Bone Marrow Transplantation in Haemoglobinopathies Is Feasible and Leads to Early and Sustained Engraftment with No Long-Term Toxicity or GvHD. *Blood.* 2016;128(22):2468. doi:10.1182/blood.V128.22.2468.2468
115. Oostenbrink LVE, Pool ES, Jol-van der Zijde CM, et al. Successful mismatched hematopoietic stem cell transplantation for pediatric hemoglobinopathy by using ATG and post-transplant cyclophosphamide. *Bone Marrow Transplant.* 2021;56(9):2203–2211. doi:10.1038/s41409-021-01302-0
116. Pakakasama S, Hongeng S, Chaisiripoomkere W, Chuansumrit A, Sirachainun N, Jootar S. Allogeneic peripheral blood stem cell transplantation in children with homozygous beta-thalassemia and severe beta-thalassemia/hemoglobin E disease. *J Pediatr Hematol Oncol.* 2004;26(4):248–252. doi:10.1097/00043426-200404000-00007
117. Rahal I, Galambrun C, Bertrand Y, et al. Late effects after hematopoietic stem cell transplantation for β -thalassemia major: the French national experience. *Haematologica.* 2018;103(7):1143–1149. doi:10.3324/haematol.2017.183467

118. Raj R, Swaminathan VV, Meena S, et al. Donor Characteristics Predict the Success of Allogeneic Hematopoietic Stem Cell Transplantation in Thalassemia Major: a Single-Center Analysis of 250 Patients. *Indian J Hematol Blood Transfus.* **2022**;38(2):411–415. doi:10.1007/s12288-021-01487-w
119. Resnick IB, Aker M, Tsigotis P, et al. Allogeneic stem cell transplantation from matched related and unrelated donors in thalassemia major patients using a reduced toxicity fludarabine-based regimen. *Bone Marrow Transplant.* **2007**;40(10):957–964. doi:10.1038/sj.bmt.1705826
120. Santarone S, Pepe A, Meloni A, et al. Secondary solid cancer following hematopoietic cell transplantation in patients with thalassemia major. *Bone Marrow Transplant.* **2018**;53(1):39–43. doi:10.1038/bmt.2017.214
121. Schuster F, Kyrilopoulou O, Ghosh S, et al. Treosulfan-based allogeneic haematopoietic stem cell transplantation in children, adolescents and young adults with transfusion-dependent thalassemia. *Bone Marrow Transplant.* **2019**;54:610–611.
122. See WQ, Tung JY, Cheuk DK, et al. Endocrine complications in patients with transfusion-dependent thalassaemia after haemopoietic stem cell transplantation. *Bone Marrow Transplant.* **2018**;53(3):356–360. doi:10.1038/s41409-017-0052-x
123. Shenoy S, Walters MC, Ngwube A, et al. Unrelated Donor Transplantation in Children with Thalassemia using Reduced-Intensity Conditioning: the URTH Trial. *Biol Blood Marrow Transplant.* **2018**;24(6):1216–1222. doi:10.1016/j.bbmt.2018.01.023
124. Shing MM, Yuen PM, Li CK, et al. A more immunosuppressive pre-transplant conditioning may be required for Chinese patients with thalassaemia. *Bone Marrow Transplant.* **1996**;17(6):907–910.
125. Sial N, Ahmed P, Wattoo FH, Ali N. Association of early donor chimerism status with survival outcomes in post allogeneic haematopoietic stem cell transplant patients of nonmalignant diseases. *J Pak Med Assoc.* **2022**;72(3):464–470. doi:10.47391/jpma.1519
126. Sodani P, Isgrò A, Gaziev J, et al. Purified T-depleted, CD34+ peripheral blood and bone marrow cell transplantation from haploidentical mother to child with thalassemia. *Blood.* **2010**;115(6):1296–1302. doi:10.1182/blood-2009-05-218982
127. Sumboonnanonanda A, Sanpakit K, Piyaphanee N. Renal tubule function in beta-thalassemia after hematopoietic stem cell transplantation. *Pediatr Nephrol.* **2009**;24(1):183–187. doi:10.1007/s00467-008-0949-0
128. Sun L, Wang N, Chen Y, et al. Unrelated Donor Peripheral Blood Stem Cell Transplantation for Patients with β -Thalassemia Major Based on a Novel Conditioning Regimen. *Biol Blood Marrow Transplant.* **2019**;25(8):1592–1596. doi:10.1016/j.bbmt.2019.03.028
129. Sun Q, Wu B, Lan H, et al. Haploidentical haematopoietic stem cell transplantation for thalassaemia major based on an FBCA conditioning regimen. *Br J Haematol.* **2018**;182(4):554–558. doi:10.1111/bjh.15438
130. Thuret I, Pondarré C, Loundou A, et al. Complications and treatment of patients with β -thalassemia in France: results of the National Registry. *Haematologica.* **2010**;95(5):724–729. doi:10.3324/haematol.2009.018051
131. Torlontano G, Di Girolamo G, Di Bartolomeo F, et al. Bone marrow transplantation in children and adults for thalassemia major and other high risk congenital diseases (Fanconi's disease and chronic granulomatous disease). *Haematologica.* **1990**;75 Suppl 5:107–110.
132. Uygun V, Tayfun F, Akcan M, et al. Quality of life assessment in hematopoietic stem cell transplantation performed on thalassemia major patients. *Pediatr Hematol Oncol.* **2012**;29(5):461–471. doi:10.3109/08880018.2012.693577
133. Wen J, Haque Q, Pei F, et al. Transplant Outcomes in Beta-Thalassemia Major Patients Receiving Combined Granulocyte Colony-Stimulating Factor-Primed Bone Marrow and Cord Blood Graft Compared to Granulocyte Colony-Stimulating Factor-Primed Bone Marrow Alone. *Acta Haematol.* **2018**;140(1):20–29. doi:10.1159/000490407
134. Yesilipek MA, Hazar V, Küpesiz A, Kizilörs A, Uguz A, Yegin O. Peripheral blood stem cell transplantation in children with beta-thalassemia. *Bone Marrow Transplant.* **2001**;28(11):1037–1040. doi:10.1038/sj.bmt.1703284
135. Yu U, Wang X, Zhang X, et al. Outcomes of haplo-cord transplantation in children with beta thalassemia major. *HemaSphere.* **2020**;4:718.
136. Li CK, Shing MM, Chik KW, et al. Haematopoietic stem cell transplantation for thalassaemia major in Hong Kong: prognostic factors and outcome. *Bone Marrow Transplant.* **2002**;29(2):101–105. doi:10.1038/sj.bmt.1703340
137. The outcomes of haploidentical hematopoietic stem cell transplantation for thalassemia major based on a FBCA conditioning regimen. **2018**.
138. Li Q, Luo J, Zhang Z, et al. G-CSF-Mobilized Blood and Bone Marrow Grafts as the Source of Stem Cells for HLA-Identical Sibling Transplantation in Patients with Thalassemia Major. *Biol Blood Marrow Transplant.* **2019**;25(10):2040–2044. doi:10.1016/j.bbmt.2019.06.009
139. Li CK, Chik KW, Wong GW, Cheng PS, Lee V, Shing MM. Growth and endocrine function following bone marrow transplantation for thalassemia major. *Pediatr Hematol Oncol.* **2004**;21(5):411–419. doi:10.1080/08880010490457132
140. Shin SH, Park SS, Park S, et al. Non-myeloablative matched sibling stem cell transplantation with the optional reinforced stem cell infusion for patients with hemoglobinopathies. *Eur J Haematol.* **2020**;105(4):387–398. doi:10.1111/ejh.13455
141. Pryce A, O'Boyle F, Lund K, et al. Degree of matching determines disease-free survival and alloreactive complications in unrelated blood and marrow transplantation for haemoglobinopathies. *Bone Marrow Transplant.* **2019**;54:352.
142. Patel DA, Hanna R, Karnik L, et al. Non-Myeloablative Bone Marrow Transplant with Post-Transplant Cyclophosphamide Plus Thiotepa Improves Donor Engraftment in Patients with Transfusion Dependent Thalassemia: results of an International Consortium (VGC2). *Biol Blood Marrow Transplant.* **2020**;26(3):S221–S222. doi:10.1016/j.bbmt.2019.12.540
143. Bernaudin F, Socie G, Kuentz M, et al. Long-term results of related myeloablative stem-cell transplantation to cure sickle cell disease. *Blood.* **2007**;110(7):2749–2756. doi:10.1182/blood-2007-03-079665
144. Oringanje C, Nemecek E, Oniyangi O. Hematopoietic stem cell transplantation for people with sickle cell disease. *Cochrane Database Syst Rev.* **2020**;7(7):CD007001. doi:10.1002/14651858.CD007001.pub5
145. Sharma A, Jagannath VA, Puri L. Hematopoietic stem cell transplantation for people with beta-thalassaemia. *Cochrane Database Syst Rev.* **2021**;4(4):CD008708. doi:10.1002/14651858.CD008708.pub5
146. Krishnamurti L, Liang J, He Z, et al. Incidence and risk factors of pain crisis after hematopoietic cell transplantation for sickle cell disease. *Blood Adv.* **2024**;8(8):1908–1919. doi:10.1182/bloodadvances.2023010749
147. Darbari DS, Liljencrantz J, Ikechi A, et al. Pain and opioid use after reversal of sickle cell disease following HLA-matched sibling haematopoietic stem cell transplant. *Br J Haematol.* **2019**;184(4):690–693. doi:10.1111/bjh.15169
148. Flinn AM, Gennery AR. Recent advances in graft-versus-host disease. *Fac Rev.* **2023**;12:4. doi:10.12703/r/12-4
149. Rashid N, Krakow EF, Yeh AC, et al. Late Effects of Severe Acute Graft-versus-Host Disease on Quality of Life, Medical Comorbidities, and Survival. *Transplant Cell Ther.* **2022**;28(12):844e1–844e8. doi:10.1016/j.jtct.2022.08.027
150. Arai S, Arora M, Wang T, et al. Increasing incidence of chronic graft-versus-host disease in allogeneic transplantation: a report from the Center for International Blood and Marrow Transplant Research. *Biol Blood Marrow Transplant.* **2015**;21(2):266–274. doi:10.1016/j.bbmt.2014.10.021

151. Arora M, Cutler CS, Jagasia MH, et al. Late Acute and Chronic Graft-versus-Host Disease after Allogeneic Hematopoietic Cell Transplantation. *Biol Blood Marrow Transplant.* **2016**;22(3):449–455. doi:10.1016/j.bbmt.2015.10.018
152. Sarantopoulos S, Cardones AR, Sullivan KM. How I treat refractory chronic graft-versus-host disease. *Blood.* **2019**;133(11):1191–1200. doi:10.1182/blood-2018-04-785899
153. Yu J, Hamilton BK, Turnbull J, et al. Patient-reported symptom burden and impact on daily activities in chronic graft-versus-host disease. *Cancer Med.* **2023**;12(3):3623–3633. doi:10.1002/cam4.5209
154. Yu J, Khera N, Turnbull J, et al. Impact of Chronic Graft-versus-Host Disease on Patient Employment, Income, and Informal Caregiver Burden: findings From the Living With Chronic GVHD Patient Survey. *Transplant Cell Ther.* **2023**;29(7):470e1–470e9. doi:10.1016/j.jtct.2023.04.004
155. Atsuta Y, Hirakawa A, Nakasone H, et al. Late Mortality and Causes of Death among Long-Term Survivors after Allogeneic Stem Cell Transplantation. *Biol Blood Marrow Transplant.* **2016**;22(9):1702–1709. doi:10.1016/j.bbmt.2016.05.019
156. Wingard JR, Majhail NS, Brazauskas R, et al. Long-term survival and late deaths after allogeneic hematopoietic cell transplantation. *J Clin Oncol.* **2011**;29(16):2230–2239. doi:10.1200/JCO.2010.33.7212

Journal of Blood Medicine

Publish your work in this journal

The Journal of Blood Medicine is an international, peer-reviewed, open access, online journal publishing laboratory, experimental and clinical aspects of all aspect pertaining to blood based medicine including but not limited to: Transfusion Medicine; Blood collection, Donor issues, Transmittable diseases, and Blood banking logistics; Immunohematology; Artificial and alternative blood based therapeutics; Hematology; Biotechnology/nanotechnology of blood related medicine; Legal aspects of blood medicine; Historical perspectives. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/journal-of-blood-medicine-journal>

Dovepress
Taylor & Francis Group