

Tumor Compression of the Hepatic or Portal Vein Predicts the Presence of Microvascular Invasion and Satellite Nodules in Hepatocellular Carcinoma: A Retrospective Study

Qing-Bo Wang^{1,*}, Wan-Ling Luo^{1,*}, Yu-Kai Li^{1,*}, Jin Li^{1,*}, Zi-Sheng Yang^{1,*}, Kun Zhao^{2,*}, Yawhan Lakang^{1,*}, Yu-Bo Liang^{1,*}, Xing-Ming Chen^{1,*}, Jin-Xiang Zuo^{1,*}, Yang Duan³, Xi Xu³, Li-Ming Shang⁴, Yang Ke^{1,5,6}

¹Department of Hepatobiliary Surgery, The Second Affiliated Hospital of Kunming Medical University, Kunming, Yunnan Province, People's Republic of China; ²Department of Hepatobiliary Surgery, The First Affiliated Hospital of Guangxi Medical University, Nanning, Guangxi Province, People's Republic of China; ³The Second Clinical Medical College, Kunming Medical University, Kunming, Yunnan Province, People's Republic of China; ⁴Department of Hepatobiliary Surgery, Beihai People's Hospital, Beihai, Guangxi Province, People's Republic of China; ⁵Department of Surgical Education and Research, The Second Affiliated Hospital of Kunming Medical University, Kunming, Yunnan Province, People's Republic of China; ⁶Yunnan Yunke Bio-Technology Institution, Kunming, Yunnan Province, People's Republic of China

*These authors contributed equally to this work

Correspondence: Li-Ming Shang, Department of Hepatobiliary Surgery, Beihai People's Hospital, No. 83 Heping Road, Haicheng District, Beihai, 536000, Guangxi Province, People's Republic of China, Tel +86-13768512036, Email 231614649@qq.com; Yang Ke, Department of Hepatobiliary Surgery, The Second Affiliated Hospital of Kunming Medical University, No. 374 Kunrui Road, Wuhua District, Kunming, 650101, Yunnan Province, People's Republic of China, Tel +86-15808875159, Email keyang1218@126.com

Purpose: This study aimed to evaluate the association of tumor compression in the hepatic or portal vein with the presence of microvascular invasion (MVI) and satellite nodules in patients with hepatocellular carcinoma (HCC).

Patients and Methods: HCC patients at the Barcelona Clinic Liver Cancer (BCLC) stages 0-A who underwent a radical liver resection in our hospitals from January 2016 to December 2022 were collected. The tumor compression of the portal or hepatic vein in individual patients was analyzed by preoperative imaging and postoperative pathology. Their MVI, satellite nodules, overall survival (OS), and recurrence-free survival (RFS) were analyzed, and the potential risk factors for the MVI and satellite nodules of patients were analyzed by univariable and multivariable logistic analyses.

Results: A total of 390 patients were included with 333 male and 263 patients <60 years old. Of them, 51 (13.1%) HCC patients had tumor venous compression, which was not significantly associated with OS and RFS, but significantly related to higher positive rates of MVI and satellite nodules than those without tumor-venous compression (MVI, 51.0% vs 36.6%, $P = 0.025$; satellite nodules, 19.6% vs 9.1%, $P = 0.023$). Tumor venous compression was an independent risk factor for the development of MVI (OR = 1.902, 95% CI: 1.049–3.447; $P = 0.034$) and satellite nodules (OR = 2.871, 95% CI: 1.277–6.458; $P = 0.011$).

Conclusion: Preoperative tumor venous compression is an independent predictor of MVI and satellite nodules in HCC patients at BCLC stages 0-A and may serve as an imaging biomarker for determining the resection margin and treatment planning.

Keywords: hepatocellular carcinoma, microvascular invasion, perivascular, prognosis, satellite nodules, venous compression

Introduction

Hepatocellular carcinoma (HCC) is a globally prevalent malignancy with high incidence and mortality rates.^{1–3} Currently, HCC patients at very early or early stages are treated with radical resection, ablation, or liver transplantation, according to the Barcelona Clinic Liver Cancer (BCLC) staging and treatment strategy.^{4,5} Although the patients with HCC at very early or early stages have the highest post-operative 5-year overall survival (OS) rate of 75.2%, nearly half (43.9%) of these patients suffer from recurrence within 5 years post radical resection.⁶ Microvascular invasion (MVI) is

the presence of HCC cells in small blood or lymphatic vessels of the liver.⁷ Approximately, MVI can be detected in 15–57% of surgical specimens from HCC patients with liver resection and transplantation,⁸ and nearly half (49%) of HCC patients with MVI suffer from HCC recurrence within 6 months post liver resection.⁹ Furthermore, satellite nodules are small HCC lesions located in the liver parenchyma, typical surrounding within 2 cm of the primary tumor.¹⁰ Approximately, satellite nodules are detected in 19–41% of surgical specimens from HCC patients with liver tumor resection and transplantation,^{11,12} and the presence of satellite nodules is also an important predictor of HCC recurrence after liver tumor resection and liver transplantation.^{12,13} Mechanistically, satellite nodules are considered to be derived from MVI.¹⁴ MVI and satellite nodules are difficult to be pre-operatively determined by CT or MRI, and the unresected MVI and satellite nodules in the residual liver will lead to inevitable HCC recurrence of those with liver tumor resection. Therefore, accurate and preoperative evaluation of the presence of MVI and satellite nodules is of great significance for guiding personalized and precise therapies for HCC patients.¹⁵

Previous studies have explored pre-operatively non-invasive evaluation of MVI and satellite nodules of HCC in the clinic. High levels of serum alpha-fetoprotein (AFP), low levels of serum albumin, liver cirrhosis, high tumor burden score, non-smooth tumor margin, absent radiological capsule on MRI, and intratumoral artery are independent risk factors of MVI development in HCC.^{16–18} Multimodal AI models by integrating clinical, imaging, and radiomics features have shown excellent performance in predicting MVI in HCC.^{19–23} Our recent studies have also shown that low levels of blood supply in the portal veins are associated with the development of MVI in HCC.^{24,25} Furthermore, the prevalence of satellite nodules is significantly associated with the single nodular type of HCC with extranodular growth, the confluent multinodular type of HCC, and the poor differentiated HCC, compared to the early HCC with a vaguely nodular type, the single nodular type of HCC, and the well and moderately differentiated HCC.¹¹ Moreover, a maximal standardized uptake value (SUVmax) >8.8 of glucose metabolism on 18F-FDG PET/CT is a significantly independent factor for the development of distant satellite nodules >1 cm from the main tumor.²⁶

A recent study has elucidated the biomechanical process of MVI of tumors, including tumor vascular compression, intravasation, and thrombus formation.²⁷ Theoretically, tumor vascular compression is a risk factor of MVI and satellite nodules in HCC. Although tumor venous compression in patients with HCC at very early or early stages is common, yet its association with the development of MVI and satellite nodules in patients with HCC at very early and early stages remains unexplored.

This study retrospectively analyzed the association between preoperative tumor venous compression and postoperative MVI as well as satellite nodules in patients with HCC at very early and early stages. This study aimed at determining whether the presence of tumor vascular compression was a useful preoperative biomarker to predict the development of MVI and satellite nodules in the liver of HCC patients.

Methods

Patients

The study was approved by the Ethics Committee of the Second Affiliated Hospital of Kunming Medical University (approval No. PJ-SCIENCE-2024-4). Informed consent was waived due to its retrospective design. Data in the study were anonymized, and particular attention was paid to protecting patient privacy and ensuring data confidentiality and security. The study complied with the *Declaration of Helsinki* adopted by the World Medical Association.^{28–30}

Inclusion criteria were: patients with newly diagnosed HCC at very-early or early stages (BCLC stages 0 or A) underwent radical liver tumor resection at the Second Affiliated Hospital of Kunming Medical University between January 2016 and December 2022; HCC and its R0 margin were confirmed by postoperative pathology; and triphasic liver CT or MRI scans were performed within 2 weeks before liver tumor resection. The exclusive criteria included HCC patients with preoperative transcatheter arterial chemoembolization (TACE), transcatheter arterial embolization (TAE), transcatheter arterial infusion (TAI), hepatic artery infusion chemotherapy (HAIC), or radiotherapy, as well as systemic treatments with targeted therapy or immunotherapy before liver tumor resection; they had concurrent diagnoses of other malignancies; or their clinical or survival data missed.

Grouping

Tumor venous compression is defined as the presence of a tumor contacting with the hepatic or portal vein, compressing the vein to the point of luminal narrowing or even complete occlusion without a visible tumor thrombus within the venous lumen by imaging (Figure 1A). The presence of tumor venous compression in the portal veins or hepatic veins of individual HCC patients was examined using preoperative triphasic liver CT or MRI scans (Figure 1B–E), reviewed by two experienced radiologists, and the absence of tumor thrombus in the portal veins or hepatic veins in individual HCC patients was confirmed by postoperative pathology. The included HCC patients were divided into the tumor venous compression and non-venous compression groups.

Pathological Evaluation of MVI and Satellite Nodules

The pathological examination was conducted by two experienced pathologists. The surgical liver specimens were fixed in 10% formalin, and paraffin-embedded. The liver tissue sections (4 μ m) were serially sectioned with an interval of 10 mm and stained with hematoxylin and eosin. The presence of MVI in individual sections was examined and defined as a cluster of cancer cells in the microvascular lumen.³¹ The presence of satellite nodules was analyzed and defined as small cancerous foci located in the liver parenchyma within 2 cm from the main tumor, which was separated by non-tumor liver tissues.³² When it was difficult to distinguish MVI from satellite nodule by pathology, a diagnosis of satellite nodule was made.¹⁴




Figure 1 Venous compression of HCC. (A) A schematic diagram illustrates the process of venous compression by HCC. (B and C) A case of HCC with tumor venous compression of the right portal vein branch on CT (B) and MRI (C). (D and E) A case of HCC with tumor venous compression of the middle hepatic vein on CT (D) and MRI (E). Yellow arrowheads indicate the typical sites of tumor venous compression.

Abbreviation: HCC, hepatocellular carcinoma.

Data Collection

The data at admission were obtained from the medical record system of the Second Affiliated Hospital of Kunming Medical University, and they included age, which was stratified into ≤ 60 or > 60 years old;³³ sex;³⁴ preoperative laboratory parameters containing hepatitis B or C virus infection,²⁴ the albumin-bilirubin (ALBI) grade, which is a simpler, more objective, and discriminatory method to assess liver function in HCC patients than Child-Pugh score and stratified into 1, 2, 3 grades,³⁵ and serum AFP levels, which were stratified into ≤ 400 ng/mL or > 400 ng/mL;^{36,37} preoperative imaging parameters including liver cirrhosis, the maximum tumor diameter, and the number of macroscopic tumor nodules.^{38,39}

Post-operative data were obtained from all patients when they were subjected to a standard follow-up until December 2024, including every 1–2 months for the first 6 months after discharge, every 3–4 months thereafter until 2 years post-surgery, and every 5–6 months after 2 years.^{24,40} During the each follow-up visiting, all patients were subjected to physical examination, liver function tests, tumor marker tests, chest X-ray or CT, and at least one abdominal imaging of liver ultrasound, or triphasic liver CT or MRI. The tumor recurrence was defined as imaging evidence of HCC.⁴¹ HCC patients with recurrence were treated with liver transplantation, repeat liver resection, radio-frequency ablation, TACE, targeted therapy, immunotherapy, or best supportive care as appropriate.^{42–45} Recurrence-free survival (RFS) was calculated from the date of hepatectomy to the date of HCC recurrence or death. OS was determined from the date of hepatectomy to the date of death from any cause.

Statistical Analysis

Data were analyzed using SPSS 26.0 (SPSS, Chicago, IL, USA). Continuous variables with normal distribution were expressed as mean \pm standard deviation; continuous variables with skewed distribution were expressed as median (interquartile range, IQR); discrete variables were expressed as number and percentage. Inter-group difference was analyzed by Student's *t*-tests, Mann–Whitney *U*-tests, or χ^2 -tests, as appropriate. The difference in survival between two groups was estimated by Kaplan-Meier method and tested by Log-rank test. The potential pre-operative risk factors for MVI and satellite nodules in HCC were analyzed by univariable logistic regression and individual factors with $P < 0.1$ from univariable logistic regression were further analyzed by the multivariable logistic regression using backward Likelihood Ratio (LR) method.⁴⁶ A two-tailed *P*-value of < 0.05 indicated statistical significance.

Results

Baseline Characteristics

During the studying period, there were 861 HCC patients with liver tumor resection at the Second Affiliated Hospital of Kunming Medical University. Of them, 371 patients were excluded because of HCC at BCLC stages B-C (Figure 2), 39 due to pre-operatively locoregional therapies or systemic therapies, 29 without survival data, 14 with comorbid life-threatening cardiac, cerebral, renal, or other hepatic diseases, 13 with R1/R2 resection, and 5 with other concurrent malignancies. The remaining 390 patients were included into the data analysis.

Analysis of triphasic liver CT or MRI images unveiled that there were 51 (13.1%) HCC patients with the evidence of tumor portal vein or hepatic vein compression (Table 1). Of them, there were 14 (27.4%) patients with the portal vein compression, 21 (41.2%) patients with the hepatic vein compression, and 16 (31.4%) patients with both the portal vein and hepatic vein compression. The remaining 339 (86.9%) patients had neither the portal vein nor the hepatic vein compression in this population. There was no statistically significant difference in their demographic and clinical data between the tumor venous compression and non-venous compression groups (all $P > 0.05$).

Effect of Tumor Venous Compression on OS and RFS

The median follow-up period of those patients was 60.0 months (95% CI, 55.3–64.8 months). During the follow-up period, 156 (40.0%) patients suffered from tumor recurrence and 99 (25.4%) patients died. The median OS and RFS in the study population were > 60.0 months and 27.2 months (95% CI, 19.0–35.3 months), respectively. The 1-, 3-, and 5-year OS rates were 95.0%, 82.1%, and 74.9% in the non-venous compression group, and 94.1%, 76.0%, and 71.5% in




Figure 2 Flowchart of patient selection of the study.

the tumor venous compression group, respectively ($P = 0.514$, [Figure 3A](#)). The median OS was >60.0 months in the two groups. The 1-, 3-, and 5-year RFS rates were 72.2%, 46.4%, and 37.0% in the non-venous compression group, and 72.2%, 42.2%, and 42.2% in the tumor venous compression group, respectively ($P = 0.533$, [Figure 3B](#)). The median RFS

Table I The Demographic and Clinical Characteristics of Patients at Admission

Variables	Total n, (%)	Compression Group (n = 51)	Non-Compression Group (n = 339)	P
Age (year)				0.140
<60	263 (67.4%)	39 (76.5%)	224 (64.3%)	
≥60	127 (32.6%)	12 (23.5%)	115 (35.7%)	
Sex				0.816
Male	333 (85.4%)	43 (84.3%)	290 (85.5%)	
Female	57 (14.6%)	8 (15.7%)	49 (14.5%)	
ALBI grade				0.409
1	244 (62.6%)	29 (56.9%)	215 (63.4%)	
2	123 (31.5%)	20 (39.2%)	103 (30.4%)	
3	23 (5.9%)	2 (3.9%)	21 (6.2%)	
HBV and/or HCV infection	303 (77.7%)	36 (70.6%)	267 (78.8%)	0.191
Cirrhosis				0.075
Yes	250 (64.1%)	27 (52.9%)	223 (65.8%)	
No	140 (35.9%)	24 (47.1%)	116 (34.2%)	
AFP (ng/mL)				0.619
≤400	295 (75.6%)	40 (78.4%)	255 (75.2%)	
>400	95 (24.4%)	11 (21.6%)	84 (24.8%)	
Single nodule	381 (97.7%)	50 (98.0%)	331 (97.6%)	0.860
Maximum tumor diameter (cm)				0.332
<2	18 (4.6%)	1 (2.0%)	17 (5.0%)	
≥2	372 (95.4%)	50 (98.0%)	322 (95.0%)	

Abbreviations: AFP, alpha fetoprotein; ALBI, albumin and total bilirubin; HBV, hepatitis B virus; HCV, hepatitis C virus.




Figure 3 Effect of tumor venous compression on (A) overall survival and (B) recurrence-free survival of patients with HCC at BCLC stages 0-A after liver tumor resection. **Abbreviations:** BCLC, Barcelona Clinic Liver Cancer staging system; HCC, hepatocellular carcinoma.

was 27.2 months (95% CI, 18.6–35.7 months) in the non-venous compression group and 22.4 months (95% CI, 4.1–40.8 months) in the tumor venous compression group. Further stratification analyses indicated that there was no statistically significant difference in OS ($P = 0.593$) or RFS ($P = 0.625$) among the groups of non-venous compression, hepatic venous compression alone, portal venous compression alone, and hepatic venous plus portal venous compression (Figure 4). Apparently, the presence of tumor portal vein and/or hepatic vein compression did not significantly affect the OS and RFS of patients with HCC at very early and early stages following liver tumor resection.




Figure 4 Effect of sub-types of tumor venous compression on (A) overall survival and (B) recurrence-free survival of patients with HCC at BCLC stages 0-A after liver tumor resection. The sub-types of tumor venous compression included tumor hepatic vein compression alone, portal vein compression alone, and combined hepatic and portal vein compression.

Abbreviations: BCLC, Barcelona Clinic Liver Cancer staging system; HCC, hepatocellular carcinoma.

Association of Tumor Venous Compression with the Development of MVI

MVI is a risk factor of HCC recurrence and metastasis. Analyses of surgical liver sections revealed that there were 150 (38.5%) HCC patients with MVI and the percentage of HCC cases with MVI in the tumor venous compression group (26 cases, 51.0%) was significantly higher than 36.6% (124 cases) in the non-venous compression group ($P = 0.025$, Figure 5A), highlighting that tumor venous compression of the portal veins or hepatic veins was associated significantly with the development of MVI in HCC. The univariable logistic regression analyses of multiple demographic and clinical factors revealed that the presence of tumor venous compression, but not other factors tested, was significantly associated with the development of MVI (OR = 1.951, 95% CI: 1.078–3.528, $P = 0.027$, Table 2). Further multivariable analysis indicated that the presence of tumor venous compression was an independent risk factor of MVI development in HCC (OR = 1.902, 95% CI: 1.049–3.447, $P = 0.034$). Stratification analyses unveiled that the percentage of HCC cases with MVI in the portal vein compression group was significantly higher than those in the non-venous compression group ($P = 0.009$, Figure 5B), implying that the presence of tumor portal vein compression was significantly related to the development of MVI in HCC.

Association of Tumor Venous Compression with the Development of Satellite Nodules

Next, we analyzed the value of tumor venous compression in the predicting satellite nodules in HCC. Pathological examinations indicated satellite nodules in 41 (10.5%) patients, including 10 (19.6%) cases in the tumor venous compression group and 31 (9.1%) cases in the non-venous compression group, implying that the percentage of satellite nodules in HCC with tumor venous compression was significantly higher than that in the non-venous compression group ($P = 0.023$, Figure 6A). The univariable logistic regression analysis of the demographic and clinical factors unveiled that the presence of liver cirrhosis (OR = 2.137, 95% CI: 0.989–4.617, $P = 0.053$), higher serum AFP levels (OR = 1.94, 95% CI: 0.980–3.840, $P = 0.057$), and positive tumor venous compression (OR = 2.423, 95% CI: 1.107–5.306, $P = 0.027$, Table 3) trended to be




Figure 5 Association of tumor venous compression with microvascular invasion in patients with HCC at BCLC stages 0-A after liver resection. (A) Overall analysis. (B) Sub-group analysis according to non-tumor compression or tumor compression of the hepatic vein alone, the portal vein alone, and combined the hepatic and portal veins. **Abbreviations:** BCLC, Barcelona Clinic Liver Cancer staging system; HCC, hepatocellular carcinoma; MVI, microvascular invasion.

Table 2 Univariate and Multivariate Analysis for Microvascular Invasion

Variables	Univariate Analysis			Multivariate Analysis		
	P	OR	95% CI	P	OR	95% CI
Sex (male vs female)	0.138	1.584	0.862–2.911			
Age (≥60 year vs <60 year)	0.584	0.886	0.576–1.365			
Cirrhosis (yes vs no)	0.260	1.28	0.833–1.966			
ALBI (1 vs 2)	0.335	0.802	0.512–1.256			
ALBI (1 vs 3)	0.327	0.63	0.250–1.587			
HBV and/or HCV infection (yes vs no)	0.243	1.348	0.816–2.226			
AFP (>400 ng/ml vs ≤400 ng/ml)	0.308	1.277	0.798–2.042			
Single nodule (yes vs no)	0.317	2.248	0.461–10.967			
Maximum tumor diameter (≥2 cm vs <2 cm)	0.062	3.304	0.940–11.610	0.074	3.154	0.894–11.121
Compression (yes vs no)	0.027	1.951	1.078–3.528	0.034	1.902	1.049–3.447

Abbreviations: AFP, alpha fetoprotein; ALBI, albumin and total bilirubin; HBV, hepatitis B virus; HCV, hepatitis C virus.

significantly associated with the development of satellite nodules in HCC. Further multivariate analysis indicated that liver cirrhosis (OR = 2.527, 95% CI: 1.144–5.584, $P = 0.022$), higher serum AFP levels (OR = 2.184, 95% CI: 1.084–4.401, $P = 0.029$), and positive tumor venous compression (OR = 2.871, 95% CI: 1.277–6.458, $P = 0.011$) were independent risk factors of the development of satellite nodules in HCC. Stratification analyses revealed that the percentage of satellite nodules in HCC




Figure 6 Association of tumor venous compression with satellite nodules in patients with HCC at BCLC stages 0-A after liver resection. **(A)** Overall analysis. **(B)** Sub-group analysis according to non-tumor compression or tumor compression of the hepatic vein alone, the portal vein alone, and combined the hepatic and portal veins. **Abbreviations:** BCLC, Barcelona Clinic Liver Cancer staging system; HCC, hepatocellular carcinoma; SN, satellite nodule.

Table 3 Univariate and Multivariate Analysis for Satellite Nodule

Variables	Univariate Analysis			Multivariate Analysis		
	P	OR	95% CI	P	OR	95% CI
Sex (male vs female)	0.643	1.261	0.473–3.361			
Age (≥ 60 year vs < 60 year)	0.723	1.129	0.576–2.214			
Cirrhosis (yes vs no)	0.053	2.137	0.989–4.617	0.022	2.527	1.144–5.584
ALBI (1 vs 2)	0.444	0.742	0.346–1.593			
ALBI (1 vs 3)	0.122	2.329	0.798–6.797			
HBV and/or HCV infection (yes vs no)	0.954	1.023	0.469–2.235			
AFP (> 400 ng/ml vs ≤ 400 ng/ml)	0.057	1.94	0.980–3.840	0.029	2.184	1.084–4.401
Single nodule (yes vs no)	0.953	0.938	0.114–7.698			
Maximum tumor diameter (≥ 2 cm vs < 2 cm)	0.492	2.048	0.265–15.804			
Compression (yes vs no)	0.027	2.423	1.107–5.306	0.011	2.871	1.277–6.458

Abbreviations: AFP, alpha fetoprotein; ALBI, albumin and total bilirubin; HBV, hepatitis B virus; HCV, hepatitis C virus.

from the tumor portal vein compression group was significantly higher than that in the non-venous compression group ($P = 0.017$, [Figure 6B](#)). Thus, the presence of tumor venous compression, particularly with the tumor portal vein compression, was significantly associated with the development of satellite nodules in HCC. These results are consistent with the hypothesis that “tumor venous compression promotes vascular invasiveness of HCC”.

Discussion

MVI and satellite nodules are well-known predictors of tumor recurrence and poor survival in HCC patients.^{47–49} However, MVI and satellite nodules are only detected by post-operative microscopy of a surgical liver specimen.⁵⁰ Accordingly, there is little chance to take useful and timely action in response to positive MVI and satellite nodules before and during operation.^{51–53} Tumor venous compression is an early step of vascular invasion,²⁷ and it is common to see in the very early and early stages of HCC. Currently, little is known on whether tumor venous compression can act as a predictor for the development of MVI and satellite nodules in HCC.

Our pre-operative study of CT and MRI images found that 13.1% of HCC patients at the very early and early stages had either positive tumor portal vein and/or hepatic vein compression. Interestingly, the percentage of MVI in the tumor venous compression group, particularly in the tumor portal vein compression group was significantly higher than that in the non-venous compression group of HCC patients. Multivariable analysis revealed that tumor venous compression was the only independent risk factor of MVI development in patients with HCC at BCLC stages 0-A. Compared to the traditional predictors, like higher levels of serum AFP, tumor number, and maximum tumor diameter, the pre-operative non-invasive detection of tumor venous compression may be better to predict MVI development for HCC patients at the very early and early stages.¹⁶

Similarly, the percentage of HCC cases with satellite nodules in the tumor venous compression group, particularly in the tumor portal vein compression group, was significantly higher than that in the non-venous compression group of patients with HCC at very early and early stages. Multi-variable analysis unveiled that tumor venous compression, like liver cirrhosis and higher serum AFP levels, was an independent risk factor of satellite nodule development in patients with HCC at very early and early stages. Consistently, our preliminary studies found that liver cirrhosis was accompanied by lymphangiogenesis,⁵⁴ especially by the ones surrounding HCC, which can increase the prevalence of satellite nodules in HCC.^{55,56} To the best of our knowledge, our findings provided scientific evidence that the presence of tumor venous compression was associated significantly with the development of MVI and satellite nodules in patients with HCC at very early and early stages. Conceivably, pre-operative measurements of tumor venous compression may be valuable for

predicting the development of MVI and satellite nodules in HCC at early stages, and these results are consistent with the hypothesis that “tumor venous compression promotes vascular invasiveness of HCC”.

Why does tumor venous compression promote the development of MVI and satellite nodules in early stages of HCC? First, tumor compression surrounding blood and lymphatic vessels can impair blood flow, oxygen delivery, and therapy efficacy, which enhance tumor invasiveness.^{57,58} Second, tumor compression surrounding blood and lymphatic vessels can elevate interstitial fluid pressure in the tumor microenvironment, which results in edema and enables tumor cells and growth factors to invade nearby tissues.^{59,60} Third, tumor cells can also respond to mechanical forces through the mechanosensitive pathways, such as the YAP/TAZ, which are activated under a physical stress and promote tumor growth and invasiveness.^{59,61,62} Fourth, tumor compression surrounding blood and lymphatic vessels can disrupt their normal architecture and increase vascular permeability, which creates a more favorable environment for tumor cells to spread into surrounding tissues.²⁷

Interestingly, tumor venous compression did not significantly influence the survival of patients with HCC at very early or early stages following radical liver tumor resection, though the percentages of HCC patients with MVI and satellite nodules were significantly higher in the tumor venous compression group than in the non-venous compression group. One possible explanation is that HCC patients with tumor venous compression generally received a larger extent of liver tissue resection, which may counteract the higher percentages of HCC patients with MVI and satellite nodules. The results were also supported by the findings that radiofrequency ablation as the first line treatment of small HCC with a maximum diameter of <3 cm achieves similar long-term OS and disease-free survival between patients with perivascular HCC and non-perivascular HCC.⁶³ However, radiofrequency ablation of patients with periportal HCC causes higher local tumor progression rates and worse OS than those with non-periportal HCC.⁶⁴ The different effects of radiofrequency ablation for HCC may be attributed to the different “heat sink effect” in these studies, which takes thermal energy away from the tumor by veins and limits therapeutic outcomes.^{65,66} Given that tumor venous compression is common to see in HCC at BCLC stages 0-A and a preliminary step of vascular invasion, the different steps of vascular invasion have different influences on OS of HCC patients. It is critical to re-consider for defining the BCLC stage C by “vascular invasion”⁶⁷ because the preliminary step of vascular invasion did not affect the survival of patients with HCC at early stages following liver tumor resection. The China Liver Cancer (CNLC) staging system has utilized “vascular tumor thrombus” instead of “vascular invasion” for defining the similarly advanced BCLC stage C.⁶⁸ Obviously, “vascular tumor thrombus” is more precise than “vascular invasion” to describe the disease progression of HCC at BCLC stage C.

Clearly, the findings of this study should be interpreted with caution due to several limitations. First, the single-center design may limit the generalizability of the results, as the sample may not be representative of the broader patient population. Secondly, the relatively small sample size may change the statistical power, increasing the potential Type II errors. Thirdly, the retrospective nature of the study introduces inherent biases in patient selection and data collection, which could affect the reliability and validity of the conclusions. Fourthly, there is a potential bias in the imaging evaluation due to the difference of multiple radiologists. Therefore, further prospective, multi-center studies with a larger sample size are warranted.

Conclusion

Preoperative measurement of tumor venous compression is valuable for predicting the development of MVI and satellite nodules in patients with HCC at BCLC stages 0-A. Tumor venous compression may be a useful imaging biomarker for determining the resection margin and the treatment planning for HCC.

Abbreviations

AFP, Alpha-fetoprotein; ALBI, Albumin and total bilirubin; BCLC, Barcelona Clinic Liver Cancer; CI, confidence interval; CT, Computed Tomography; HAIC, Hepatic Artery Infusion Chemotherapy; HCC, Hepatocellular carcinoma; HBV, hepatitis B virus; HCV, hepatitis C virus; IQR, interquartile range; LR, Likelihood Ratio; MRI, Magnetic Resonance Imaging; MVI, Microvascular invasion; OS, Overall survival; RFS, Recurrence-free survival; SN, satellite nodules; SUVmax, Maximal standardized uptake values; TACE, Transcatheter Arterial Chemoembolization; TAI, Transcatheter Arterial Infusion; TAE, Transcatheter Arterial Embolization; 18F-FDG PET/CT, Fluorine-18 fluorodeoxyglucose positron emission tomography/computed tomography.

Data Sharing Statement

The datasets used and/or analyzed during the current study are available from the corresponding author Yang Ke upon a reasonable request.

Ethics Approval and Informed Consent

This study was approved by the Ethics Committee of the Second Affiliated Hospital of Kunming Medical University (approval No. PJ-SCIENCE-2024-4). Written informed consent was waived by the Ethics Committee due to the retrospective nature of the study. Data in the study were anonymized, and particular attention was paid to protecting patient privacy and ensuring data confidentiality and security. The study complied with World Medical Association *Declaration of Helsinki*.

Funding

This study was funded by grants to Y.K. from the National Natural Science Foundation of China (#82460461), the Medical Subject Leader Project of Yunnan Province (General Surgery) (#D-2024029), the Yunnan Fundamental Research Project for Excellent Young Scholars (#202401AW070003), the Young and Mid-aged Academic and Technical Leader Reserve Talent Program of Yunnan Province (#202205AC160063), the Top Talent Project of Kunming Medical University (# not available), the National College Students Innovation and Entrepreneurship Training Program (#202410678018), and the College Students Innovation and Entrepreneurship Training Program of Yunnan Province (#2024CYD022).

Disclosure

All authors declare no competing interests in this work.

References

- Filho AM, Laversanne M, Ferlay J, et al. The GLOBOCAN 2022 cancer estimates: data sources, methods, and a snapshot of the cancer burden worldwide. *Int J Cancer*. 2025;156(7):1336–1346. doi:10.1002/ijc.35278
- Han B, Zheng R, Zeng H, et al. Cancer incidence and mortality in China, 2022. *J Natl Cancer Cent*. 2024;4(1):47–53. doi:10.1016/j.jncc.2024.01.006
- Wang Y, Chen S, Xiao X, et al. Impact of apolipoprotein A1 on tumor immune microenvironment, clinical prognosis and genomic landscape in hepatocellular carcinoma. *Precis Clin Med*. 2023;6(3):pbad021. doi:10.1093/pcmedi/pbad021
- Wang QB, Li J, Zhang ZJ, et al. The effectiveness and safety of therapies for hepatocellular carcinoma with tumor thrombus in the hepatic vein, inferior vena cava and/or right atrium: a systematic review and single-arm meta-analysis. *Expert Rev Anticancer Ther*. 2025;25(5):561–570. doi:10.1080/14737140.2025.2489651
- Chatzipanagiotou OP, Catalano G, Kawashima J, et al. Minimally invasive surgery approaches in major liver resection for Hepatocellular carcinoma. *Hepatoma Res*. 2024;10:34. doi:10.20517/2394-5079.2024.77
- Liu YW, Yong CC, Lin CC, et al. Liver resection of hepatocellular carcinoma within and beyond the Barcelona Clinic Liver Cancer guideline recommendations: results from a high-volume liver surgery center in East Asia. *J Surg Oncol*. 2020;122(8):1587–1594. doi:10.1002/jso.26183
- Bosi C, Rimini M, Casadei-Gardini A, et al. Understanding the causes of recurrent HCC after liver resection and radiofrequency ablation. *Expert Rev Anticancer Ther*. 2023;23(5):503–515. doi:10.1080/14737140.2023.2203387
- Yu Y, Wang XH, Hu WJ, et al. Patterns, risk factors, and outcomes of recurrence after hepatectomy for hepatocellular carcinoma with and without microvascular invasion. *J Hepatocell Carcinoma*. 2024;11:801–812. doi:10.2147/JHC.S438850
- Chen ZH, Zhang XP, Feng JK, et al. Patterns, treatments, and prognosis of tumor recurrence after resection for hepatocellular carcinoma with microvascular invasion: a multicenter study from China. *HPB*. 2022;24(7):1063–1073. doi:10.1016/j.hpb.2021.11.016
- Wang X, Yu Y, Tao Y, et al. Clinical-radiological characteristic for predicting ultra-early recurrence after liver resection in solitary hepatocellular carcinoma patients. *J Hepatocell Carcinoma*. 2023;10:2323–2335. doi:10.2147/JHC.S434955
- Okusaka T, Okada S, Ueno H, et al. Satellite lesions in patients with small hepatocellular carcinoma with reference to clinicopathologic features. *Cancer*. 2002;95(9):1931–1937. doi:10.1002/cncr.10892
- Plessier A, Codes L, Consigny Y, et al. Underestimation of the influence of satellite nodules as a risk factor for post-transplantation recurrence in patients with small hepatocellular carcinoma. *Liver Transpl*. 2004;10(2 Suppl 1):S86–S90. doi:10.1002/lt.20039
- Li HM, Huang W, Hu C, et al. Prognostic factors for patients with proliferative hepatocellular carcinoma after liver resection. *J Hepatocell Carcinoma*. 2023;10:2059–2071. doi:10.2147/JHC.S440636
- Li C, Ouyang W, Yang T. The association of microvascular invasion with satellite nodule, tumor multiplicity, tumor encapsulation and resection margin of hepatocellular carcinoma. *J Hepatol*. 2022;77(3):890–891. doi:10.1016/j.jhep.2022.03.036
- Xiang J, Li YK, Mei SM, et al. Novel diagnostic and therapeutic strategies based on PANoptosis for hepatocellular carcinoma. *Cancer Biol Med*. 2025;22(8):928–939. doi:10.20892/j.issn.2095-3941.2025.0150

16. Chang YS, Tsai MJ, Tsai CJ, et al. A new model based on preoperative AFP, albumin, and tumor burden score for predicting microvascular invasion in early-stage HCC. *Am J Cancer Res.* 2024;14(10):4979–4988. doi:10.62347/zgrj7827
17. Zhang J, Zeng F, Jiang S, et al. Preoperative prediction model of microvascular invasion in patients with hepatocellular carcinoma. *HPB.* 2023;25(1):45–53. doi:10.1016/j.hpb.2022.08.007
18. Tang Y, Lu X, Liu L, et al. A reliable and repeatable model for predicting microvascular invasion in patients with hepatocellular carcinoma. *Acad Radiol.* 2023;30(8):1521–1527. doi:10.1016/j.acra.2023.02.035
19. Wang L, Wu M, Li R, et al. Multi-task deep learning strategy using computed tomography (CT)-based radiomics for end-to-end high efficiency prediction of microvascular invasion in hepatocellular carcinoma. *Cancers.* 2022;14(12):2956. doi:10.3390/cancers14122956
20. Panettieri E, Campisi A, Bianchi V, et al. Artificial intelligence and its role in guiding liver-directed therapy for hepatocellular carcinoma: is it ready for prime time? *Hepatoma Res.* 2022;8:41. doi:10.20517/2394-5079.2022.57
21. Zhang C, Gu J, Zhu Y, et al. AI in spotting high-risk characteristics of medical imaging and molecular pathology. *Precis Clin Med.* 2021;4(4):271–286. doi:10.1093/pcmedi/pbab026
22. Fu S, Lai H, Huang M, et al. Multi-task deep learning network to predict future macrovascular invasion in hepatocellular carcinoma. *EClinicalMedicine.* 2021;42:101201. doi:10.1016/j.eclinm.2021.101201
23. Zhang J, Zhou S, Jiang Y, et al. 3D assessment of skeletal muscle and adipose tissue for prognosis of hepatocellular carcinoma: a multicenter cohort study. *Clin Nutr ESPEN.* 2025;67:626–634. doi:10.1016/j.clnesp.2025.03.168
24. Li YK, Wu S, Wu YS, et al. Portal venous and hepatic arterial coefficients predict post-hepatectomy overall and recurrence-free survival in patients with hepatocellular carcinoma: a retrospective study. *J Hepatocell Carcinoma.* 2024;11:1389–1402. doi:10.2147/JHC.S462168
25. Li YK, Wang QB, Liang YB, et al. CT measurement of blood perfusion in hepatocellular carcinoma: from basic principle, measurement methods to clinical application. *Chin J Oncol.* 2024;46(10):940–947. doi:10.3760/cma.j.cn112152-20240605-00240
26. Ochi H, Hirooka M, Hiraoka A, et al. (18F)-FDG-PET/CT predicts the distribution of microsatellite lesions in hepatocellular carcinoma. *Mol Clin Oncol.* 2014;2(5):798–804. doi:10.3892/mco.2014.328
27. Nia HT, Munn LL, Jain RK. Physical traits of cancer. *Science.* 2020;370(6516):eaz0868. doi:10.1126/science.aaz0868
28. Liu J, Yao L, Sun J, et al. Cancer risk in relatives of BRCA1/2 pathogenic variant carriers in a large series of unselected patients with breast cancer. *Cancer Biol Med.* 2023;20(2):147–154. doi:10.20892/j.issn.2095-3941.2022.0593
29. Zhu Z, Tang H, Ying J, et al. Efficacy and safety of gemcitabine plus S-1 vs. gemcitabine plus nab-paclitaxel in treatment-naïve advanced pancreatic ductal adenocarcinoma. *Cancer Biol Med.* 2023;20(10):765–778. doi:10.20892/j.issn.2095-3941.2023.0189
30. De la Torre K, Shin WK, Lee HW, et al. Weight gain after 35 years of age is associated with increased breast cancer risk: findings from a large prospective cohort study. *Cancer Biol Med.* 2024;21(8):651–655. doi:10.20892/j.issn.2095-3941.2024.0172
31. Huang H, Wu F, Yu Y, et al. Multi-transcriptomics analysis of microvascular invasion-related malignant cells and development of a machine learning-based prognostic model in hepatocellular carcinoma. *Front Immunol.* 2024;15:1436131. doi:10.3389/fimmu.2024.1436131
32. Li J, Wang WQ, Zhu RH, et al. Postoperative adjuvant tyrosine kinase inhibitors combined with anti-PD-1 antibodies improves surgical outcomes for hepatocellular carcinoma with high-risk recurrent factors. *Front Immunol.* 2023;14:1202039. doi:10.3389/fimmu.2023.1202039
33. Han CL, Tian BW, Yan LJ, et al. The effect of age, sex, and eastern cooperative oncology group performance status on the efficacy and safety of immune checkpoint inhibitors in patients with hepatocellular carcinoma: a systematic review and meta-analysis. *Expert Rev Anticancer Ther.* 2024;24(5):303–312. doi:10.1080/14737140.2024.2341723
34. Chen J, Yang Z, Gao F, et al. Influence of sex on outcomes of liver transplantation for hepatocellular carcinoma: a multicenter cohort study in China. *Cancer Biol Med.* 2024;21(4):347–362. doi:10.20892/j.issn.2095-3941.2023.0453
35. Persano M, Casadei-Gardini A, Burgio V, et al. Five years of lenvatinib in hepatocellular carcinoma: are there any predictive and/or prognostic factors. *Expert Rev Anticancer Ther.* 2023;23(1):19–27. doi:10.1080/14737140.2023.2156340
36. Han J, Kuai W, Yang L, et al. Impact of metabolic dysfunction-associated steatotic liver disease on the efficacy of immunotherapy in patients with chronic hepatitis B-related hepatocellular carcinoma. *Cancer Biol Med.* 2024;21(9):813–825. doi:10.20892/j.issn.2095-3941.2024.0048
37. Yao C, Wu S, Kong J, et al. Angiogenesis in hepatocellular carcinoma: mechanisms and anti-angiogenic therapies. *Cancer Biol Med.* 2023;20(1):25–43. doi:10.20892/j.issn.2095-3941.2022.0449
38. Fu X, Zhang Y, Luo Q, et al. Targeting the mechano-microenvironment and liver cancer stem cells: a promising therapeutic strategy for liver cancer. *Cancer Biol Med.* 2023;20(11):816–829. doi:10.20892/j.issn.2095-3941.2023.0229
39. Peng F, Wang S, Feng Z, et al. Circulating cell-free mtDNA as a new biomarker for cancer detection and management. *Cancer Biol Med.* 2023;21(2):105–110. doi:10.20892/j.issn.2095-3941.2023.0280
40. Wang Y, Lai Y, Du L, et al. Impact of quality control circle on patient outcomes after hepatocellular carcinoma intervention: a meta-analysis. *Expert Rev Anticancer Ther.* 2023;23(8):875–882. doi:10.1080/14737140.2023.2219899
41. Ma X, Wang Z, Wang S, et al. The assessment of circulating tumor DNA associated with Wnt/ β -catenin signaling pathway as a diagnostic tool for liver cancer: a systematic review and meta-analysis. *Expert Rev Anticancer Ther.* 2024;24(3–4):155–167. doi:10.1080/14737140.2024.2312246
42. Chen J, Li H, Zhuo J, et al. Impact of immunosuppressants on tumor pulmonary metastasis: new insight into transplantation for hepatocellular carcinoma. *Cancer Biol Med.* 2024;21(11):1033–1049. doi:10.20892/j.issn.2095-3941.2024.0267
43. Zhu S, Yu Y, Yang M, et al. Hepatic artery infusion chemotherapy combined with the FOLFOX regimen for the treatment of hepatocellular carcinoma: recent advances and literature review. *Expert Rev Anticancer Ther.* 2024;24(6):423–434. doi:10.1080/14737140.2024.2346624
44. Werida RH, Abd El Baset OA, Askar S, et al. Efficacy of doxorubicin and lipiodol therapy by trans-arterial chemoembolization in hepatocellular carcinoma Egyptian patients and relation to genetic polymorphisms. *Expert Rev Anticancer Ther.* 2024;24(10):1009–1020. doi:10.1080/14737140.2024.2391364
45. Stefanini B, Ielasi L, Chen R, et al. TKIs in combination with immunotherapy for hepatocellular carcinoma. *Expert Rev Anticancer Ther.* 2023;23(3):279–291. doi:10.1080/14737140.2023.2181162
46. Fan H, Wang W, Zhang Y, et al. Current treatment paradigm and survival outcomes among patients with newly diagnosed multiple myeloma in China: a retrospective multicenter study. *Cancer Biol Med.* 2023;20(1):77–87. doi:10.20892/j.issn.2095-3941.2022.0612
47. Liu J, Zhang X, Lin J, et al. HBcrAg is associated with prognosis of hepatitis B virus-related hepatocellular carcinoma in patients after hepatectomy undergoing antiviral therapy. *Int J Cancer.* 2025;156(6):1293–1303. doi:10.1002/ijc.35224

48. Li J, Qiu Y, Liu Y, et al. Comparison of adjuvant treatment regimens for high-risk hepatocellular carcinoma: a Bayesian network meta analysis and systematic review. *Front Immunol.* 2024;15:1487353. doi:10.3389/fimmu.2024.1487353
49. Xu Q, Lan L, Zeng J, et al. The effect of microvascular invasion on hepatocellular carcinoma with portal vein tumor thrombus after hepatectomy: a retrospective study. *Cancer Control.* 2024;31:10732748241265257. doi:10.1177/10732748241265257
50. Ning C, Liu G, Zhang J, et al. Case Report: the application of associating liver partition and portal vein ligation for staged hepatectomy in patients with hepatitis b virus-related hepatocellular carcinoma after undergoing treatment with an immune checkpoint inhibitor: a report of two cases. *Front Immunol.* 2023;14:1159885. doi:10.3389/fimmu.2023.1159885
51. Feng Y, Guo K, Jin H, et al. Efficacy and safety of neoadjuvant combination immunotherapy in surgically resectable malignant solid tumors: a systematic review and meta-analysis. *Expert Rev Anticancer Ther.* 2024;24(3–4):169–181. doi:10.1080/14737140.2024.2325404
52. Guven DC, Yildirim HC, Chalabiyev E, et al. Emerging treatment strategies in hepatobiliary cancer. *Expert Rev Anticancer Ther.* 2023;23(3):243–256. doi:10.1080/14737140.2023.2183844
53. Chen W, Cheng S. Nomogram and calculator for predicting the prognosis of patients with giant hepatocellular carcinoma. *Expert Rev Anticancer Ther.* 2024;24(8):781–788. doi:10.1080/14737140.2024.2369129
54. Li JF, Wang QB, Li YK, et al. Activation of lymphangiogenesis by platelet as novel therapeutic approaches for liver cirrhosis and portal hypertension. *World J Gastroenterol.* 2025;31(23):107554. doi:10.3748/wjg.v31.i23.107554
55. Li J, Liang YB, Wang QB, et al. Tumor-associated lymphatic vessel density is a postoperative prognostic biomarker of hepatobiliary cancers: a systematic review and meta-analysis. *Front Immunol.* 2024;15:1519999. doi:10.3389/fimmu.2024.1519999
56. Li J, Wang QB, Liang YB, et al. Tumor-associated lymphatic vessel density is a reliable biomarker for prognosis of esophageal cancer after radical resection: a systemic review and meta-analysis. *Front Immunol.* 2024;15:1453482. doi:10.3389/fimmu.2024.1453482
57. Riaz F, Zhang J, Pan F. Forces at play: exploring factors affecting the cancer metastasis. *Front Immunol.* 2024;15:1274474. doi:10.3389/fimmu.2024.1274474
58. Liang YB, Li L, Liu B, et al. Research advances in the roles of high-altitude hypoxic stress in hepatocellular carcinoma. *J Sichuan Univ Med Sci.* 2024;55(6):1436–1445. doi:10.12182/20241160605
59. Feng X, Cao F, Wu X, et al. Targeting extracellular matrix stiffness for cancer therapy. *Front Immunol.* 2024;15:1467602. doi:10.3389/fimmu.2024.1467602
60. Chen K, Li Y, Wang B, et al. Patient-derived models facilitate precision medicine in liver cancer by remodeling cell-matrix interaction. *Front Immunol.* 2023;14:1101324. doi:10.3389/fimmu.2023.1101324
61. Wang Y, Ruf S, Wang L, et al. The dual roles of lamin A/C in macrophage mechanotransduction. *Cell Prolif.* 2025;58(5):e13794. doi:10.1111/cpr.13794
62. Gao Y, Gong Y, Lu J, et al. Targeting YAP1 to improve the efficacy of immune checkpoint inhibitors in liver cancer: mechanism and strategy. *Front Immunol.* 2024;15:1377722. doi:10.3389/fimmu.2024.1377722
63. Kang TW, Lim HK, Lee MW, et al. Perivascular versus nonperivascular small HCC treated with percutaneous RF ablation: retrospective comparison of long-term therapeutic outcomes. *Radiology.* 2014;270(3):888–899. doi:10.1148/radiol.13130753
64. Cao S, Lyu T, Fan Z, et al. Long-term outcome of percutaneous radiofrequency ablation for periportal hepatocellular carcinoma: tumor recurrence or progression, survival and clinical significance. *Cancer Imaging.* 2022;22(1):2. doi:10.1186/s40644-021-00442-2
65. Verdonschot KHM, Arts S, Van den Boezem PB, et al. Ablative margins in percutaneous thermal ablation of hepatic tumors: a systematic review. *Expert Rev Anticancer Ther.* 2023;23(9):977–993. doi:10.1080/14737140.2023.2247564
66. Pahk KJ, Heo J, Joung C, et al. Noninvasive mechanical destruction of liver tissue and tissue decellularisation by pressure-modulated shockwave histotripsy. *Front Immunol.* 2023;14:1150416. doi:10.3389/fimmu.2023.1150416
67. Reig M, Forner A, Rimola J, et al. BCLC strategy for prognosis prediction and treatment recommendation: the 2022 update. *J Hepatol.* 2022;76(3):681–693. doi:10.1016/j.jhep.2021.11.018
68. Zhou J, Sun H, Wang Z, et al. Guidelines for the diagnosis and treatment of primary liver cancer (2022 edition). *Liver Cancer.* 2023;12(5):405–444. doi:10.1159/000530495

Journal of Hepatocellular Carcinoma

Publish your work in this journal

The Journal of Hepatocellular Carcinoma is an international, peer-reviewed, open access journal that offers a platform for the dissemination and study of clinical, translational and basic research findings in this rapidly developing field. Development in areas including, but not limited to, epidemiology, vaccination, hepatitis therapy, pathology and molecular tumor classification and prognostication are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-hepatocellular-carcinoma-journal>

Dovepress
Taylor & Francis Group