

Effect of Intraoperative Intravenous Lidocaine on Postoperative Delirium in Elderly Patients Undergoing Posterior Lumbar Interbody Fusion [Corrigendum]

Feng X, Jia J, Sang Y, et al. *Drug Des Devel Ther.* 2026;20:1–12. <https://doi.org/10.2147/DDDT.S571032>

The authors have advised due to an error that occurred inadvertently at the time of figure assembly, the number at risk data in Figure 2 on page 7 of the published paper is incorrect.

The correct Figure 2 is shown below.




Figure 2 Kaplan–Meier curves showing time to postoperative delirium in the lidocaine and control groups. There was a significant difference between the two groups (Log rank test, $P = 0.0055$).

The authors apologize for the error and advise it does not affect the results and conclusion of the paper.

Drug Design, Development and Therapy

Dovepress
Taylor & Francis Group

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>

<https://doi.org/10.2147/DDDT.S604220>