

Intrauterine contraception after cesarean section and during lactation: a systematic review

Norman D Goldstuck¹
Petrus S Steyn²

¹Department of Obstetrics and Gynaecology, Tygerberg Hospital, Cape Town, Western Cape, South Africa; ²Reproductive Health and Fertility Regulation, Department of Obstetrics and Gynaecology, Groote Schuur Hospital and University of Cape Town, Cape Town, South Africa

Background: All postpartum women, including those who are breastfeeding or have had a cesarean section, appear potentially suited to intrauterine contraception, a long acting reversible contraceptive (LARC). Like any other method used after delivery, it should not interfere with lactation or be affected by cesarean section.

Study design: We searched the MEDLINE, PubMed, Popline, Google Scholar, and Clinicaltrials.gov databases from January 1968 through to December 2012. Studies were included if they reported event rates in women who had a cesarean section and event rates and clinical outcomes in lactating women or their infants in the breastfeeding group. Summary odds ratios were not calculated because of the diverse methods of reporting event rates in the cesarean section group and the heterogeneity of the results in the breastfeeding group.

Results: We found 26 articles on event rates in interval and post-placental intrauterine device (IUD) use, and 18 on event rates and clinical outcomes in breastfeeding IUD users. Four prospective studies and one retrospective study showed an increased expulsion rate in interval insertion. There were 19 studies, of which five were controlled in post-placental IUD insertion after cesarean section. Four studies had expulsion rates of 10 or more per 100 woman-years of use and 15 expulsion rates below 10 per 100 woman-years of use. Three studies showed that event rates for lactating IUD users are the same as those for non-lactating users. Fifteen controlled studies showed that the IUD had no effect on milk production and seven of these showed no effect on infant growth. Pharmacovigilance databases report an increased rate of IUD perforations in lactating women, while the event rate studies report that insertion is generally easier and less painful than expected. These were uncontrolled reports.

Conclusion: The IUD is a long-acting reversible method of contraception with expulsion rates of 5–15 per 100 woman-years of use when used as a post-placental method immediately after cesarean section. As an interval procedure (6 or more weeks after cesarean section) it appears to have a high expulsion rate (5% or higher) notably in older devices. The IUD does not affect breastfeeding and is easy to insert in these women, but appears to be associated with a higher perforation rate (>1 per 100). Providers should not be deterred from using this contraception method, especially in developing countries, but should be attentive to preventing these potential problems.

Keywords: long acting reversible contraception, IUD, cesarean section, post-placental, lactation

Correspondence: Norman D Goldstuck
Department of Obstetrics and Gynaecology, Tygerberg Hospital, Francie van Zijl Avenue, Tygerberg, Cape Town, Western Cape 7505, South Africa
Tel +27 823 418 200
Email nahunzh@yahoo.com

Introduction

Developing countries tend to have high birth rates.¹ In most cases, these countries are desirous of reducing them. The most effective contraceptive methods are the long acting reversible contraceptive (LARC) which are “forgettable” (user does not have to remember

to take it).² The importance of being able to breastfeed in all situations, and especially in underprivileged areas, is important for infant welfare. Women who are breastfeeding need to use a contraceptive method which has no effect on lactation.³ Women who have had a cesarean section (CS), whether for a current or previous pregnancy, also need a contraceptive method which is independent of this procedure. Lactation was used as a contraceptive before the advent of modern methods, and certain populations have used this method alone to space their children 3–4 years apart.⁴ During lactation, the suckling reflex inhibits ovulation. This is not mediated via prolactin, which inhibits ovulation at highly elevated levels due to pathological events, eg, pituitary tumors.⁴ During normal lactation, oxytocin-induced elevation of endorphins inhibits ovulation.⁴ In studies with primates, blocking this response with naloxone induces ovulation.⁴ Conversely, blocking prolactin activity with bromocriptine stops milk flow, but does not induce ovulation if the oxytocin endorphin pathway is maintained.⁴ It is important to ensure that the contraceptive method used in lactating women augments rather than diminishes lactation, and induces rather than inhibits whatever contraceptive activity lactation is providing.

This review examines the role of intrauterine contraception as a LARC method in these two conditions, and the role it can play in helping women to space out and adequately nourish their children, especially in underdeveloped countries and situations where continuing with lactation is paramount for infant health.

Methods

This study is a systematic review to evaluate possible complications in IUD users when inserted immediately after CS, and 6 weeks or longer after CS. It also evaluates the effect of lactation on IUD performance and vice versa, including the factors determining the lactatory response. The reference list for this study was generated from searching for references from the following databases:

MEDLINE: ‘Intrauterine device (IUD, IUCD)’/ ‘Intrauterine system (IUS) [MeSH]’ and ‘Cesarean section, caesarean section CS, C/S.’ [MeSH]. ‘Intrauterine device (IUD, IUCD)’/ ‘Intrauterine system (IUS)’ and ‘Lactation’ [MeSH].

POPLINE: ‘IUD’ and ‘cesarean section’; ‘IUD’ and ‘lactation’.

PubMed: ‘IUD’ and ‘cesarean section’; ‘IUD’ and ‘lactation’.

Google Scholar: ‘IUD’ and ‘post placental cesarean section’, and ‘IUD’ and ‘interval cesarean section’; ‘IUD’ and ‘lactation’; the terms were filtered so as not to overlap in the advanced search.

Clinical trials.gov: ‘intrauterine device’ AND ‘cesarean section’ and ‘intrauterine device’ AND ‘lactation’.

We did not search Wangfang Data or Weipu Data, which refer to Chinese publications, as most large Chinese studies are available in the main databases. This is nevertheless a shortcoming, but the lack of referral to any Chinese studies other than the ones we found suggests there are not likely to be any other landmark studies in this area. Our study includes peer reviewed papers from January 1968 to December 2012. The search produced 1,145 papers in total, 317 for CS “post placental” and 62 for CS “interval”, which was reduced to 262 and 44 after duplicates were eliminated. The search for “lactation” papers produced 828 papers, which was reduced to 592 excluding duplicates.

Only those papers which provided data which were directly referable to “cesarean section (caesarean section)” or “lactation” (assessed from the abstracts) were included, as the terms are used descriptively for subjects in very many IUD studies. The CS papers included only those which gave event rates (ie, analysis of IUD problems such as pregnancy, expulsion, pain and/or bleeding, and infection), expressed preferably by life table analysis, but also those which expressed data as a Pearl index or on a percentage basis. Controlled and uncontrolled studies were included. The same criteria applied to the lactation studies, but in addition studies were included which reported possible chemical changes due to the IUD on lactation. Studies of event rates required at least 20 subjects (one exception was made for a study of post-placental IUD insertion with 19). The studies of possible chemical changes due to IUD insertion were reported irrespective of the number of subjects. Event rates for both the CS and lactation studies were compared to controls whenever possible, and in one instance are tabulated separately. After imposition of these criteria there were 7 studies of interval IUD insertion post CS, 19 studies of post-placental IUD insertion after CS, 7 studies of the effect of IUDs on lactation chemistry, 14 studies on the effect of the IUD on lactation, and 3 studies on IUD event rates in lactating women. The effect of the IUD on lactation was also examined against controls whenever possible. The data were extracted by one author (NDG) and subsequently independently verified at a later date (PSS). Summary odds ratios were not calculated because of the variable methods of data reporting and some of the poorly quantifiable outcomes reported.

Results

Cesarean section

We found seven studies of interval insertion (ie, insertion at least 6 weeks after the last delivery, whether vaginal or by CS) of IUDs in women who had previously had one CS or more.^{5–11}

There were five prospective and two retrospective studies. Five of the studies, including one retrospective, showed elevated expulsion results. Unfortunately, the methods of expression of event rates were both cumulative and non-cumulative, making comparison difficult. Pregnancy rates and device removal rates for medical problems were reported as being in the range of 0–5 per 100 woman-years of use.

The type of IUDs and number of subjects in each study is given in Table 1. There was one reported perforation, possibly due to a “hyperinvoluted” uterus.¹¹ There was one study on Copper-T insertion in 76 women with previous cesarean section following surgical termination of pregnancy.¹² The IUD was inserted post-abortion at 8–11 weeks. This study was controlled and the expulsion rate was 2.6%. Since it was neither an interval nor a post-placental CS study, it does not appear in the table.

There were 19 studies of IUD insertion at the time of CS (post-placental cesarean IUD insertion) for which adequate data relative to the procedure is available.^{13–25} These studies have been carried out with different types of IUDs over the

past 40 years. The studies were all prospective and most of the data is from the People’s Republic of China and Latin America. A summary of the studies by device type is given in Table 2.

Five of the studies were controlled by comparing post-partum insertion after CS with insertion of the IUD in patients with normal vaginal deliveries. In all cases, the expulsion rate was lower in the CS group, and significantly lower in many cases.^{15,20,23,24} These studies are summarized separately in Table 3. Additionally, there are a few large, unpublished studies where details are sometimes quoted in the text as personal communications. The results from these observations were not used, as they did not meet the selection criteria. Some of the possible reasons for the large degree of variation in these results and the implications for post-placental IUD insertion after CS are discussed later.

The effect of IUDs on lactation

There have been numerous studies, both open and controlled, on the possible effect of IUDs on lactation itself,

Table 1 Interval IUD insertion after cesarean section

Type of IUD	Number of subjects	Time of insertion	Significant event rate	Comments	Study
Antigon I-45 Antigon III-17 Antigon IV-29 Antigon F-33	121	6–8 weeks	Expulsion 28.3% ^e Controls expulsion 11.8%	Prospective, cumulative per 100 woman years ^a	Wiese ⁵
Lippes loop Cu-T Multiload	318	6 weeks+	Nil	Retrospective pooled Controlled ^b	Chi and Balogh ⁶
Cu-T 200	107	6 weeks	Expulsion 6.5% [*] Removals 8.4% [*]	Prospective controlled %, not life table ^c	Gupta et al ⁷
Dalkon shield 2 MLCu 250 168 Copper 7 [®] 22 Lippes loop 7 Copper-T [®] 2 Not stated 14	215	6 weeks+	Expulsion 19 ^a	Retrospective cumulative rate, non-controlled ^d	Holloway et al ⁸
Szontagh IUD	288	Interval	Expulsion elevated (rate not specified) ^a	Controlled prospective cumulative rates not expressed	Chi et al ⁹
VCu 200 TCu 380A	400	Interval	Pregnancy [□] Removal ^o Expulsion* VCu 200 0% TCu 380A 3.1% ^a	Prospective controlled VCu vs TCu 380A Event rates for both devices within accepted limits Not cumulative	Tang and Feng ¹⁰
Cu-T 200	45	Interval	Expulsion 8.8% ^a	Not life-table 1 perforation not controlled	Parikh and Gandhi ¹¹

Notes: * $P < 0.05$ for study group versus control group; $\square P < 0.05$ for VCu (1.02%) versus TCu 380A (4.64%); $^o P < 0.05$ for VCu (1.02%) versus TCu 380A (4.13%); $^a P < 0.05$ for VCu (0%) versus TCu 380A (3.1%); b 12 month follow up; c 6 month follow up; d 24 month follow up; e 48 month follow up; a per 100 women years of use. Antigon (Svend Schroeder Co., Copenhagen, Denmark); Lippes Loop (Ortho Pharmaceutical, Raritan, NJ, USA); Dalkon shield (AH Robbins, Richmond, VA, USA); MLCu (Prosan S.A. Arnham, the Netherlands); Copper 7[®] (G.D.Searle and Co., High Wycombe, England, UK); Copper-T[®] (Ortho Pharmaceutical); Szontagh IUD (Simmelweiss University, Budapest, Hungary); VCu (Dr Y. Wu, Shanghai, People’s Republic of China); TCu 380A (Teva Pharmaceutical Industries Ltd., Petach Tikva, Israel); Copper T 200 originally marketed by Ortho Pharmaceuticals, now replaced by TCu 380A of Teva Pharmaceutical Industries Ltd.

Abbreviation: IUD, intrauterine device; vs, versus.

Table 2 Post-placental IUD insertions after cesarean section

Event rate per 100 woman years (range)						
Type of IUD	Studies	Subjects followed	Expulsion	Medical removal	Pregnancy	Comments and references
Mirena®	1	33	0	0	0	12 month follow up ¹³
MLCu 250	1	154	2.6%	7.8%	0	12 month percentage rate ¹⁴
TCu 220	3	714	(1.78–10.9)	Not stated	0–1.1	12 month cumulative rate ^{15–17}
TCu 380	5	314	(0–17.6)	(0–10.6)	0	One study had 6 month follow up only ^{18–22}
Delta T	2	135	(4.1–9.6)	7.2	2.4	^a One reported 6 month cumulative, the other 24 month percentage rates ^{23,24}
Delta loop						
Delta Beijing						
Metal ring ± catgut knots	7	3131	(1.2–9.6)	(0.8 – 7.2)	(0–7.5)	^a Cumulative rate from 6–24 months ^{24,25}

Notes: ^aReference 24 reported 9 Chinese studies also reported in 23 and 25. The data was only used once. Mirena® (Bayer, Wuppertal, Germany); MLCu 250 (Prosan S.A. Arnhem, the Netherlands); TCu 380 (Teva Pharmaceutical Industries Ltd., Petach Tikva, Israel); Delta T, Delta loop, Delta Beijing, Metal ring (Chinese government, Beijing, People's Republic of China).

Abbreviation: IUD, intrauterine device.

and on the hormonal factors responsible for initiating and maintaining lactation. Following a report of reversible galactorrhea in two IUD users,²⁶ a number of conflicting studies on the role of IUDs on prolactin, cortisol, and copper metabolism in lactating IUD users have been conducted.^{27–34}

The results of these studies are summarized in Table 4. The consequences of the IUD altering prolactin levels in lactating women appear to be ephemeral at most. The IUD is a LARC method and if it is to be useful during lactation it is important that it shows no effect on inhibiting lactation. Unlike other LARC methods, its mechanism of action is not systemic (including the levonorgestrel intrauterine system). It should not be expected to have an effect on lactation (either positive or negative). Studies of lactation in women using the IUD confirm this.^{33,35–45}

Additionally, the active agent (whether copper or levonorgestrel) should not adversely affect the quality of the milk as well as the quantity produced. The effect of the IUD on lactation and infant development in users has also been studied.^{45–48} A summary of the studies confirming that there is no adverse

effect on quantity and quality of milk production is given in Table 5.

The effect of lactation on IUD performance

Vaginal post-partum IUD insertion has its own complications. Whether lactation has a role at this stage cannot be demonstrated. The effect of established lactation on IUD insertion and performance from 6 weeks post-delivery is amenable to objective evaluation. There are three detailed studies using life table analysis on the effect of lactation on IUD performance,^{49–51} and these are given in Table 6. Two of these^{50,51} show a significantly increased medical removal rate for non-breastfeeders over those who were breastfeeding. The reasons for this may be that pain and bleeding is reduced in breastfeeders due to the full or partial lactational amenorrhea (LA) which reduces bleeding, and also the possible role of β -endorphin secretion in breastfeeding women.⁵¹ β -endorphin is a powerful natural analgesic and would be expected to decrease all types of pain. Pregnancy rates and expulsion rates were

Table 3 Comparison of IUD expulsion rate after post-placental insertion following cesarean or vaginal delivery

Type of IUD	Subjects	Expulsion rate		Comments and reference
		PPCS	PPV	
TCu 220	554	10.9	16.4 ^a	3 month cumulative ¹⁵
TCu 380A	19	0	50 ^b	Ultrasound detection cumulative at 12 months ²⁰
Delta T	52	4.1	20.5 ^b	6 month cumulative ^{23,24}
Delta loop	83	3.9	7.5	Catgut strings on IUDs cumulative at 12 months ²⁴
Delta T 24				
Delta loop 26				
Delta Beijing 33	906	5.5	23.5	Cumulative at 12 months ²⁴
Metal ring ± 3 catgut knots				

Notes: ^a $P < 0.05$; ^b $P < 0.001$. TCu 380 (Teva Pharmaceutical Industries Ltd., Petach Tikva, Israel); Delta T, Delta loop, Delta T 24, Delta Beijing, Metal ring (Chinese Government, Beijing, People's Republic of China).

Abbreviations: IUD, intrauterine device; PPCS, post-placental insertion after cesarean section; PPV, post-placental insertion after vaginal delivery.

Table 4 Prolactin levels in IUD users

Type of IUD	Subjects	Prolactin levels	Comments and reference
Lippes loop	6	Increase after insertion	Normalized next cycle ³⁴
Lippes loop C	2	Increase after insertion	Galactorrhea reversed after removal ²⁶
Copper 7® Copper-T® Soonawalla YCu	40	Increase compared to control ($P < 0.05$)	Lactating and non-lactating ²⁸
Copper (unspecified)	25	No increase	Controlled ²⁹
Inert (unspecified)	25		
Progestasert	26	No increase	27
Copper 7	25	No increase	Controlled ³⁰
Lippes loop	17		
Progestasert	14	No increase	Controlled-lactating ³³
Copper-T	21		

Notes: Lippes Loop (Ortho Pharmaceutical, Raritan, NJ, USA); Copper 7® (G.D. Searle and Co., High Wycombe, England, UK); Copper-T® (Ortho Pharmaceutical); Soonawalla YCu (Prof RP Soonawala, Indian Council of Medical Research, Mumbai, India); Progestasert (Alza Corporation, Palo Alto, CA, USA). Copper T 200 originally marketed by Ortho Pharmaceuticals, now replaced by TCu 380A of Teva Pharmaceutical Industries Ltd.

Abbreviation: IUD, intrauterine device.

not significantly different for those who were breastfeeding and those who were not.

Discussion

A number of large studies have noted that IUD insertion is relatively easy and pain free in women who are lactating, and that lactation reduces the need for cervical dilation.⁵¹ A study

of 6493 women also found that this was especially pronounced in women with lactational amenorrhea; ie, currently breast-feeding, which would be expected in the presence of increased β -endorphin levels, which also makes insertion less painful.^{50,52} The force required to insert an IUD in lactating women is also lower than in non-lactating women for the Copper-T® (Ortho Pharmaceuticals, Raritan, NJ, USA), TCu 380A, ParaGard®, (Teva Pharmaceutical Industries Ltd., Petach Tikva, Israel), MLCu 250®, 375® (Prosan S.A., Arnham, the Netherlands), and Nova-T 200® (Bayer, Wuppertal, Germany) IUDs.⁵³

The risk of perforation during IUD insertion in women who are lactating is increased approximately 10 fold. This was reported in a study of the Lippes loop (Ortho Pharmaceutical, Raritan, NJ, USA), Copper 7® (G.D. Searle and Co., High Wycombe, England, UK), Dalkon shield® (A.H. Robins Company, Richmond, VA, USA), Safe-T-Coil® (Julius Schmidt Laboratories, Little Falls, NJ, USA), Copper-T®, and Progestasert® (Alza Corporation, Palo Alto, CA, USA) devices.⁵² The reported rate of perforation has been quoted as being from 1 in 350, to 1 in 2,600 insertions.^{54,55} Lactation also appears to be a major factor in perforations with the Mirena® (Bayer, Wuppertal, Germany). Of 701 perforations reported by pharmacovigilance centers, 192 (42%) were in Mirena® users who were breastfeeding at the time.⁵⁶

The reasons for the high rate of expulsion in interval IUD insertions after CS are not apparent. Five of seven studies of interval IUD insertion in women who had previously undergone CS showed elevated expulsion rates. These studies were mainly on older devices; there was no information on the

Table 5 The effect of the IUD on lactation

Type of IUD	Studies	Subjects	Observations and results	Comments and references
Lippes loop D	1	185	No change in duration of lactation and amenorrhea versus controls	PPV insertion ³⁵
Progestasert	2	100	Quantity and quality of milk significantly higher versus controls	PPV insertion ^{37,38}
TCu 380A TCu 200B	1 33 29	62	No increased copper in milk	33
Mirena®	1	163	No changes in lactation or infant development versus TCu 380A	157 in TCu 380A group insertion 6–8 weeks post-partum ⁴⁶
TCu 380A	2	831	No change in lactation versus progesterone vaginal ring	42,43
LNG – IUD	2	90	Milk production and infant development same as controls LNG in breast milk 0.1% of daily serum level	10 ug and 30 ug releasing IUDs inserted 6 weeks post-partum ^{39,47}
Cu – IUD	1	38	No change in volume and composition of breast milk versus Implanon	Insertion 6 weeks post-partum. IUD type not specified ⁴⁸
Cu – IUD	1	68	No alteration of duration of lactation or LAM	Interval insertion DMPA and POP controls ³⁶
MLCu 250	3	169	No change in milk quality and quantity and infant development	Interval insertion comparison with POP and DMPA ^{38,40,41}
MLCu 375	1	41	No change in milk quality and infant development	Interval insertion comparison with POP ⁴⁵

Notes: Lippes Loop (Ortho Pharmaceutical, Raritan, NJ, USA); Progestasert (Alza Corporation, Palo Alto, CA, USA); TCu 380 (Teva Pharmaceutical Industries Ltd., Petach Tikva, Israel); Mirena® (Bayer, Wuppertal, Germany); MLCu (Prosan S.A., Arnham, the Netherlands).

Abbreviations: IUD, intrauterine device; DMPA, depo-medroxyprogesterone acetate; PPV, post-placental vaginal; LAM, lactational amenorrhea; LNG, levonorgestrel; POP, progestogen only pills.

Table 6 IUD event rates in breastfeeding women

Type of IUD	Subjects	Expulsion	Event rates per 100 woman years		
			Medical removal	Pregnancy	Comments and references
Lippes loop D	173	3.4 ^a	0.6 ^a	0.0 ^a	3 month rate interval insertion ⁴⁹
Copper-T [®]	109	2.1 ^a	0.0 ^a	0.0 ^a	3 month rate interval insertion ⁴⁹
TCu 380A	590	13 ^a	0.9 ^b	1 ^a	No distinction between interval and PPV insertions 6 month rate ⁵⁰
TCu 380A	1032	3.9 ^a	2.9 ^b	0.3 ^a	12 month rate all interval insertions ⁵¹

Notes: ^aNot significant versus non breastfeeding; ^bP<0.05 versus non breastfeeding (significantly lower than non-breastfeeders). Lippes Loop (Ortho Pharmaceutical, Raritan, NJ, USA); Copper 7[®] (G.D.Searle and Co., High Wycombe, England, UK); TCu 380 (Teva Pharmaceutical Industries Ltd., Petach Tikva, Israel).

Abbreviations: IUD, intrauterine device; PPV, post-partum vaginal insertion.

Mirena[®], Nova-T[®] and Gyne-Fix[®] (Control Limited, Ghent, Belgium) IUDs, which are in common use today, and limited information on the TCu 380A[®], the current benchmark for copper-containing IUDs. It would be interesting to determine the attitude of inserters to see if there is any variation in their practical technique. Alternatively, it may be possible that the uterine scar is a factor, as it is in vaginal birth after CS (VBAC). A wealth of information on this topic exists in many databases. It would be worthwhile to extract and re-examine this data.

The variation in expulsion rates after post-placental CS can be explained by the large number of different devices and techniques used. Some promising ideas are already being evaluated, so this should become a highly viable option.

The CS rate is increasing worldwide and very rapidly in some countries.⁵⁷ While the rate itself might be lower in most developing countries, the absolute numbers are high because of the high birth rates. In addition, the need for women to breastfeed in these countries is especially important because of the limited financial ability to use infant feeding formulas and to use them adequately when they are adopted.

For these women to be able to space their families adequately, it is important that they are able to use a LARC method which does not interfere with breastfeeding or breast milk, is cost-effective, and the use of which does not conflict with some of the consequences of childbirth, eg, CS.

There is no one reversible birth control method, short or long-acting, which is perfect in this regard. The IUD comes fairly close to this, with a few provisos. LARC methods should be introduced as soon after delivery as possible, and certainly before the mother leaves the facility because of the problems of follow up. The problems of the post-placental insertion of IUDs after vaginal births are well known, in particular, premature device expulsion.

Fortunately, the rates of expulsion after CS post-placental insertions appear to be significantly lower than when the IUD is used immediately after vaginal birth following delivery of the placenta, making insertion of IUDs immediately after CS this a much more viable option. There are many reasons

for this and the design of IUDs especially for this purpose should both solve the expulsion problem and make this a very practical LARC method.

In order not to deprive women who have had previous cesarean deliveries of the IUD, we suggest that extra care should be taken in these subjects to ensure fundal IUD placement. Puerperal insertion also requires expertise to avoid these excessive expulsions.³ The newer IUDs may be less likely to be associated with some of these problems, and that is a subject for further examination.

Earlier studies suggesting that the IUD could raise prolactin levels^{26,34} and possibly affect lactation have not been confirmed. Both the current copper-releasing and hormone-releasing IUDs appear to have no effect on the quantity, quality, and duration of breastfeeding. In lactating women, IUD event rates of expulsion, pregnancy, and medical removal are not adversely different from non-lactating women, with the exception that medical removal rates are lower.⁵¹ Numerous studies have shown that insertion of IUDs in lactating women requires less cervical dilation and lower force to actually place the device. This is probably due to higher circulating β -endorphin levels and the softer consistency of lactating and especially puerperal uterine muscle. These factors may make insertion easier, but they also make perforation more likely since the inserter will perceive less resistance to insertion and the acceptor will experience less pain during the procedure. A previous CS and the presence of lactation require careful attention to detail to avoid the problems of expulsion and perforation, respectively. Despite these problems, neither a previous CS nor lactation is a World Health Organization contraindication to IUD insertion.⁵⁸

Conclusion

Women who have had a CS and/or are breastfeeding are good candidates for intrauterine contraception such as LARCs. Post-placental CS IUD insertion is especially valuable as it appears to present fewer problems than after vaginal delivery. While IUDs do not affect lactation, and insertion of an IUD in

lactating women is generally less painful and better tolerated, there are still attendant risks, such as perforation.

In women who have previously had a CS, IUD use in the presence of uterine scar appears to be associated with a higher risk of expulsion. Both these groups of women may use the IUD as a LARC method in appropriate circumstances. Extra care should be taken when placing IUDs in these women in order to try and mitigate potential IUD related problems.

Disclosure

This study was not funded by any external source. Norman D Goldstuck has never received funds or gifts in kind from any outside agency. Petrus S Steyn has lectured for various pharmaceutical companies. The authors report no other conflicts of interest in this work.

References

- Cohen JE. Human population: the next half century. *Science*. 2003;302(5648):1172–1175.
- Winner B, Peipert JF, Zhao Q, et al. Effectiveness of long-acting reversible contraception. *N Eng J Med*. 2012;366(21):1998–2007.
- Saarikosi S. Contraception during lactation. *Ann Med*. 1993;25(2):181–184.
- Short RV. Breastfeeding. *Sci Am*. 1984;250(4):35–41.
- Wiese J. A study of four Antigon models and factors in use-effectiveness of the IUD. *Contraception*. 1976;14(6):687–701.
- Chi IC, Balogh S. Interval insertion of intrauterine device in women with previous cesarean section. *Contraception*. 1984;30(3):209–214.
- Gupta I, Mahajan U, Lata P, Gupta AN. Copper intrauterine contraceptive device insertion 6 weeks after cesarean section. *Indian J Med Res*. 1985;82:214–216.
- Holloway GA, Goldstuck ND, Bowen D. Interval insertion of an intrauterine contraceptive device following cesarean section. *Int J Gynecol Obstet*. 1989;29(3):243–248.
- Chi IC, Zhou SW, Balogh S, Ng K. IUD insertions following cesarean deliveries examined. 1983; *111th Annual Meeting of the American Public Health Association*.
- Tang YH, Feng ZJ. [Comparative study of clinical effects of VCU IUD and TCu 380A IUD were used on women who once been done caesarean section]. *Zhonghua Yi Xue Za Zhi*. 2012;92(17):1209–1211. Chinese.
- Parikh V, Gandhi AS. Safety of copper-T as contraceptive after caesarean section. *J Indian Med Assoc*. 1989;87(5):113–115.
- Gupta I, Mahajan U, Sawney H. Concurrent copper T insertion with medical termination of pregnancy in women with previous caesarean section delivery. *Indian J Med Res*. 1988;87:450–452.
- Puzey M. Mirena at caesarean section. *Eur J Contracept Reprod Health Care*. 2005;10(3):164–167.
- Ruiz – Velasco V, Garcia C, Castro H. Cesarean section IUD insertion. *Contracept Deliv Syst*. 1982;3(1):21–24.
- Lara R, Sanchez RA, Aznar R. [Application of intrauterine device through the incision of the cesarean section]. Aplicacion del dispositivo intrauterino a traves de la incision de la cesarea. *Ginecol Obstet Mex*. 1989;57:23–27. Spanish.
- Alvarez Pelago J, Borbolla Sala ME. [IUD insertion during cesarean section and its most frequent complications]. DIU transcesarea y sus complicaciones mas frecuentes. *Ginecol Obstet Mex*. 1994;62:330–335. Spanish.
- Van Kets H, Thiery M, Van der Pas H. IUD insertion during cesarean section. *Adv Contracept*. 1985;1(4):337–339.
- Celen S, Sucak A, Yildez Y, Danisman N. Immediate postplacental insertion of an intrauterine contraceptive device during cesarean section. *Contraception*. 2011;84(3):240–243.
- Nelson AL, Chen S, Eden R. Intraoperative placement of the Copper-T-380 intrauterine device in women undergoing elective cesarean delivery: a pilot study. *Contraception*. 2009;80(1):81–83.
- Letti Müller AL, Lopes Ramos JG, Martins-Costa SH, et al. Transvaginal ultrasonographic assessment of the expulsion rate of intrauterine devices inserted in the immediate postpartum period: a pilot study. *Contraception*. 2005;72(3):192–195.
- Levi E, Cantillo E, Ades V, Banks E, Murthy A. Immediate post-placental IUD insertion at caesarean delivery: a prospective cohort study. *Contraception*. 2012;86(2):102–105.
- Celen S, Moroy P, Sucak A, Aktulay A, Danisman N. Clinical outcomes of early postplacental insertion of intrauterine contraceptive devices. *Contraception*. 2004;69(9):279–282.
- Chi IC, Su-Wen Z, Balogh S, Ng K. Post-cesarean section insertion of intrauterine devices. *Am J Pub Health*. 1984;74(11):1281–1282.
- Chi IC, Gao JI, Siemens AJ, Waszak CS. IUD insertion at cesarean section – the Chinese experience. *Adv Contracept*. 1986;(2):145–145.
- Liu BH, Zhang LY, Zhang BR. Intrauterine contraceptive device insertion with suture fixation at cesarean section. *Chin Med J*. 1983;96(2):141–144.
- Horn H, Finkelstein M, Ben-David M. Possible effect of intrauterine device on galactorrhoea in women. *Contraception*. 1975;12(4):419–424.
- Spellacy WN, Buhi WC. A prospective study of plasma prolactin levels in women using the progesterone releasing intrauterine device (P-IUD). *Contraception*. 1979;19(1):91–94.
- Mehta S, Pawar V, Joshi J, et al. Serum prolactin levels in women using copper IUDs. *Contraception*. 1977;15(3):327–334.
- Ransom WTM, Chapman RS, Hull MGR. Serum prolactin concentrations related to copper or inert intrauterine devices (IUDs) in women. *Contraception*. 1979;20(5):441–446.
- Wenof M, Aubert JM, Reyniak JV. Serum prolactin levels in short-term and long-term use of inert plastic and copper intrauterine devices. *Contraception*. 1979;19(1):21–27.
- Askalani H, Badraoui MH, Mahrous I, et al. Serum cortisol level in women using the Progestasert system. *Contracept Deliv Syst*. 1981;2(1):133–138.
- Badraoui MH, Askalani H, Mahrous I, et al. Serum-prolactin in lactating women using the Progestasert system. *Contracept Deliv Syst*. 1981;2(1):121–126.
- Rodrigues da Cunha AC, Doren JG, Cantuaria AA. Intrauterine device and maternal copper metabolism during lactation. *Contraception*. 2001;63(1):37–39.
- Fortuny A, Pajol-Amat P, Calaf J, Mackinnon PCB, Robyn C. Plasma levels of LH, FSH and prolactin in women during and after insertion of an intrauterine device. *J Reprod Fert*. 1973;35:628–629.
- Hingorani V, Uma Bai GR. Lactation and lactational amenorrhoea with post-partum IUCD insertions. *J Reprod Fert*. 1970;23:513–515.
- Prema K. Duration of lactation and return of menstruation in lactating women using hormonal contraception and IUDs. *Contracept Deliv Syst*. 1982;3(1):39–46.
- Badraoui MH, Askalani H, Mahrons H, Serour G, Hefnawi F. Lactation pattern in Egyptian women using the Progestasert system. *Contracept Deliv Syst*. 1982;3(1):53–60.
- Badraoui MH, Hefnawi F, Bahgat R, et al. Contraception during lactation. *Reproduccion*. 1982;6:9–18.
- Heikkila M, Haukkamaa M, Luukainen T. Levonorgestrel in milk and plasma of breastfeeding women with a levonorgestrel-releasing IUD. *Contraception*. 1982;25(1):41–49.
- Delgado Betancourt J, Sandoval JC, Sanchez F, Vallesteros De Cano P, De La Luz Bantista M, Jimenez F. Influence of Exluton (progestogen-only OC) and the Multiload Cu 250 IUD on lactation. *Contracept Deliv Syst*. 1984;5(1):91–95.

41. Sinchai W, Sethavanich S, Asavapiriyant S, et al. Effects of a progestogen-only pill (Exluton) and an intrauterine device (Multiload Cu250) on breastfeeding. *Adv Contracept*. 1995;11(2):143–155.
42. Chen JH, Wu SC, Shao WQ, et al. The comparative trial of TCU 380A IUD and progesterone releasing vaginal ring used by lactating women. *Contraception*. 1998;57(6):371–379.
43. Sivin I, Diaz S, Croxatto HB, et al. Contraceptives for lactating women: a comparative trial of a progesterone-releasing vaginal ring and the copper T 380A IUD. *Contraception*. 1997;55(4):225–232.
44. Laukaran VH. Contraceptive choices for lactating woman: suggestions for postpartum family planning. *Stud Fam Plan*. 1981;12(4):156–163.
45. Bjarnadottir RI, Gottfredsdottir H, Sigurdardottir K, Geirsson RT, Dieben TO. Comparative study of the effects of a progestogen-only pill containing desogestrel and an intrauterine contraceptive device in lactating women. *BJOG*. 2001;108:1174–1180.
46. Shaamash AH, Sagen GH, Hussein MH, Shaaban MM. A comparative study of the levonorgestrel – releasing intrauterine system Mirena versus the Copper T 380A intrauterine device during lactation: breastfeeding performance, infant growth and infant development. *Contraception*. 2005;72(5):346–351.
47. Heikkila M, Luukkainen T. Duration of breastfeeding and development of children after insertion of a levonorgestrel-releasing intrauterine contraception device. *Contraception*. 1982;25(3):279–292.
48. Taneepanichskul S, Reinprayoon D, Thaithumyanon P, Praisuwana P, Tosukhowong P, Dieben T. Effects of the etonogestrel – releasing implant Implanon and a non-medicated intrauterine device on the growth of breast-fed infants. *Contraception*. 2006;73(4):368–371.
49. Cole LP, McCann M, Higgins JE, Waszak CS. Effects of breastfeeding on IUD performance. *Am J Pub Health*. 1983;73(4):384–388.
50. Chi IC, Potts M, Wilkens LR, Champion CB. Performance of the Copper T-380A intrauterine device in breastfeeding women. *Contraception*. 1989;39(6):603–618.
51. Farr G, Rivera R. Interactions between intrauterine contraceptive device and breast-feeding status at time of intrauterine contraceptive device insertion: analysis of TCU 380A acceptors in developing countries. *Am J Obstet Gynecol*. 1992;167(1):144–151.
52. Chi IC, Wilkens LR, Champion CB, Machemer RE, Rivera R. Insertion pain and other IUD insertion-related rare events for breastfeeding and non breastfeeding woman – a decade's experience in developing countries. *Adv Contracept*. 1989;2(5):101–119.
53. Goldstuck ND, Holloway G. IUD insertion forces: effects of recent childbirth and lactation. *Adv Contracept*. 1988;4(2):159–169.
54. Chi IC, Kelly E. Is lactation a risk factor of IUD – and sterilization – related perforation? A Hypothesis. *Int J Gynecol Obstet*. 1984;22(4):315–317.
55. Heartwell SF, Schlesselman S. Risk of uterine perforation among users of intrauterine devices. *Obstet Gynecol*. 1983;61(1):31–36.
56. Van Grootheest K, Sachs B, Harrison-Woolrych M, Caduff – Janosa P, van Puijenbroek E. Uterine perforation with the levonorgestrel – releasing intrauterine device: analysis of reports from four national pharmacovigilance centres. *Drug Saf*. 2011;34(1):83–88.
57. Roberts CL, Algert CS, Ford JB, Todd AL, Morris JM. Pathways to a rising caesarean section rate: a population-based cohort study. *BMJ Open*. 2012;2(5):e001725.
58. World Health Organization. Selected practice recommendations for contraceptive use. 2nd edition. 2008. Available from: <http://www.who.int/reproductivehealth/publications/en/>. Accessed November 4, 2013.

International Journal of Women's Health

Publish your work in this journal

The International Journal of Women's Health is an international, peer-reviewed open-access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of women's healthcare including gynecology, obstetrics, and breast cancer. The manuscript management system is completely online and includes

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-womens-health-journal>

Dovepress

a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.