

Evidence for differences between *B. bruxellensis* strains originating from an enological environment

Vincent Renouf^{1,2}
Cécile Miot-Sertier²
Marie-Claire Perello²
Gilles de Revel²
Aline Lonvaud-Funel²

¹Laffort, Bordeaux, France; ²UMR
Œnologie, INRA-Université Bordeaux,
France

Abstract: The aim of this paper is to study and compare the physiological diversity of different strains of a wine spoilage yeast species: *Brettanomyces bruxellensis*. The minimum inhibitory concentrations of several drugs on different *B. bruxellensis* strains were scored on solid nutrient media. This revealed variations in resistance among the *B. bruxellensis* strains. Their capacity to develop in different wine and must environments: pH, ethanol, and SO₂ concentrations, were evaluated by measuring the direct incubation survival rate. The results, compared with those obtained for other wine yeast species, confirmed the remarkable resistance of *B. bruxellensis* strains to various conditions which inhibit the growth of other species. Nevertheless some differences were observed among the *B. bruxellensis* strains, thus confirming their physiological diversity. A comparison of their volatile phenol production revealed intraspecific heterogeneity among *B. bruxellensis* strains. *B. bruxellensis* is one of the microbial species most resistant to environmental constraints in wine. It is the best adapted to growing in wine and spoiling it by volatile phenol production. However, different *B. bruxellensis* strains exhibit varying characteristics, particularly their capacity to produce volatile phenols. This implies that certain strains are more prejudicial than others. Further studies are required to determine the molecular causes of this intraspecific diversity.

Keywords: *Brettanomyces bruxellensis*, strain diversity, physiology, volatile phenols

Introduction

Among microbial deteriorations of wine, the production of volatile phenols, 4-ethylphenol and 4-ethylguaiacol, by *Brettanomyces bruxellensis* yeast is certainly the most dreaded by winemakers and the most criticized by consumers. A great deal of research has focused on this subject.¹ Previous work investigated the origins of *B. bruxellensis*, the conditions it required to develop in wine,^{2,3} and its interactions with other microorganisms.⁴ Several techniques for eliminating these microorganisms were also studied.⁵ Fast, effective detection methods were also developed.⁶

Microbial identification had always been restricted to the species level, until a recently-reported macrorestriction analysis of the chromosomal DNA by pulsed-field gel electrophoresis (REA-PFGE) method made it possible to differentiate among *B. bruxellensis* strains.⁷ Distinguishing among strains is particularly useful as it makes it possible to clarify how contamination occurs and study intraspecific diversity.⁸ Studies of differences between *Brettanomyces* isolates were essential^{9,10} to explain the differences in spoilage due to volatile phenol synthesis observed during production.

The goal of this work is to characterize the physiology of the *B. bruxellensis* strains isolated at different stages, from grapes on the vine to bottled wine. These *B. bruxellensis* strains were also compared with other yeast genera. Firstly, we measured the resistance of each strain to some inhibitory compounds, and then we tested their capacity to establish colonies in grape must and wine, at various levels of pH, alcohol, and SO₂. Finally, volatile phenol production in grape must was evaluated in the laboratory.

Correspondence: Vincent Renouf
UMR Œnologie 1219 INRA-Université
Bordeaux 2 Victor Ségalen, INRA, ISVV,
351, cours de la libération, 33405 Talence
cedex, France
Tel +33 556 865 304
Fax +33 556 863 050
Email vincentrenouf@yahoo.fr

Materials and methods

Microorganisms

B. bruxellensis strains used in this study are listed in Table 1. They were isolated from a Bordeaux domain at different stages in the winemaking process, from grapes through to bottled wine. Wines samples were used to inoculate solid nutritive medium specific to the non-*Saccharomyces* yeast species. This medium is composed of glucose (20 g/L), yeast extract (10 g/L), bactotryptone (10 g/L), agar (20 g/L), biphenyl (150 mg/L), cloramphenicol (100 mg/L), cycloheximide (500 mg/L), pH = 5.0 with H₃PO₄ and aerobic incubation at 25 °C for 10 days. After colonies were tested by the species-specific polymerase chain reaction (PCR) species developed by Ibeas¹¹ in order to identify *B. bruxellensis* isolates at the species level. After positive colonies of *B. bruxellensis* were compared from point of the strain typing by REA-PFGE developed by Miot-Sertier.⁷ This method is the most relevant method to type strains of *B. bruxellensis* isolated throughout the winemaking process^{1,7,12} by providing a specific PFGE pattern for each strain. For each different pattern obtained by the REA-PFGE method during our monitoring a strain were conserved at -80 °C in the IOEB (Institute of Oenology of Bordeaux) collection. The list and the nomenclature of these strains are provided in Table 1.

Microorganism cultures

Starter cultures were prepared in a grape-juice medium consisting of 200 mL/L commercial grape juice, 40 mL/L

ethanol, 1.5 g/L malt extract, 1.5 g/L yeast extract, 0.5 g/L (NH₄)₂SO₄, 0.2 g/L MgSO₄. The pH was adjusted to 5.0 using KOH 5N. Growth was monitored by epifluorescence, using the protocol developed by Millet and Lonvaud-Funel.¹³ For each experiment, microbial cells were collected at the end of the exponential growth phase.

Measurement of minimum inhibitory concentrations

Drug sensitivity was assayed by measuring the minimum inhibitory concentration (MIC) in a medium similar to that used for the starter cultures, with 20 g/L of agar, containing different drug concentrations between 10 g/L and 10 mg/L (10, 5, 2.5, 1.25, 0.6, 0.4, 0.25, 0.2, 0.15, 0.100, 0.050, 0.025, 0.010 g/L). The initial drug solutions were prepared in ethanol (70% v/v). 10 µL viable yeasts were plated on the Petri dishes using the Steer's equipment, at an initial concentration of 10⁶ cells/mL. In all cases, drug resistance was scored by the presence or absence of observation colonies on the agar surface after ten days' incubation at 25 °C. A growth positive control was prepared without any drugs for each strain. Triplicate assays were carried out at each concentration.

The drug tested were Congo red (Sigma-Aldrich, St. Louis, MO), Calcofluor white (Sigma-Aldrich), and vanillin (Sigma-Aldrich). Vanillin has a phenolic function that interacts with enzymes involved in essential redox reactions.¹⁴ Its amphiphilic properties disturb the phospholipid organization in the cytoplasm membrane.¹⁵ Calcofluor white binds microfibrils of cellulose or chitin, thus disrupting their assembly and leading to cell lysis.¹⁶ This inhibitory activity is similar to that of Congo red, which also affects the cell-wall organization.¹⁷

Measuring the yeasts' development capacity

The strains' capacity to establish themselves at varying levels of pH, ethanol, and free SO₂, was evaluated by the relationship, τ₄₈, between the population counted two hours after direct inoculation of cells pre-cultured and collected at the end of the exponential phase into must or wine previously sterilized by filtration (0.22 µm). The concentration was adjusted to 10³ cells/mL in the medium inoculated. The population was counted after forty-eight hours of incubation at 25 °C.

Two pH values were tested, 3 and 3.6, in grape must (glucose + fructose = 200 g/L, total phenolic index = 35, L-malic-acid = 4.2 g/L). Ethanol was tested at 12.5% v/v and 15% v/v in homogenized wine (pH = 3.6, total phenolic index = 76, free SO₂ = 4 mg/L, total SO₂ = 22 mg/L). The two

Table 1 List of strains used in this study

| Species | Designation | Origin |
|-----------------------------------|----------------------------|--|
| <i>Brettanomyces bruxellensis</i> | *IOEBL0411 | Fermenting must during alcoholic fermentation |
| | IOEBL0407 | Wine during aging |
| | IOEBL0447 | Fermenting must during malolactic fermentation |
| | IOEBL0462 | Wine during aging |
| | IOEBL0468 | Grapes |
| | IOEBL0469 | Grapes |
| | IOEBL0506 | Bottled wine |
| | IOEBL0522 | Bottled wine |
| | CLIB300 | Belgian beer |
| | <i>Candida cantarellii</i> | IOEBL0416 |
| <i>Metschnikowia fructicola</i> | IOEBL0416 | Must after pressing |
| <i>Piichia anomala</i> | IOEBL0534 | Fermenting must |
| <i>Rhodotorula mucilaginosa</i> | CLIB370 | Grapes |
| <i>Saccharomyces cerevisiae</i> | Levain | Industrial strain |
| | 522 Davis | |

Notes: *IOEBL, Yeast collection of the Institute of Enology of Bordeaux, Talence, France; CLIB, collection de levures d'intérêt biotechnologique, INRA, Thiverval Grignon, France.

SO₂ levels tested were 10 mg/L and 30 mg/L free SO₂ in wine collected at the end of malolactic fermentation (MLF) (pH = 3.6, ethanol = 12.5% v/v, total phenolic index = 78). All these media were sterilized by filtration (0.22 µm).

Measuring volatile phenol production

Volatile phenol production was evaluated in grape-juice cultures (pH = 3.6, glucose + fructose = 200 g/L, L-malic acid = 4.2 g/L, total phenol index = 35). Microbial cells collected at the end of exponential growth phase were inoculated at an initial concentration of 10³ CFU/mL. Alcoholic fermentation (AF) was monitored by measuring glucose and fructose, using an enzymatic kit (Boehringer Mannheim GmbH, Mannheim, Germany). In each assay, volatile phenols were measured when AF was completed. Each assay in triplicate was compared with negative control without microbial cells. Volatile phenols were measured by dichloromethane extraction from 50 mL grape juice, followed by gas chromatography.¹⁸ The temperature was 25 °C.

Results

Minimum inhibitory concentrations

Table 2 lists the MIC of the species and strains tested. Three types of resistance were distinguished. Firstly, some species were highly sensitive to the drugs used: *C. cantarelli* and *M. fructicola*. *S. cerevisiae* exhibited intermediate resistance and two species were highly resistant: *B. bruxellensis* and *P. anomala*. There was considerable intraspecific diversity within the *B. bruxellensis* species in terms of resistance to Calcofluor white and vanillin, whereas the MICs for the

other two drugs were the same for each strain. The same strain (IOEBL 0447) was the most resistant to Calcofluor white and vanillin, but it was difficult to identify a correlation between these two resistances, as the IOEBL 0468 strain was one of the most resistant to vanillin and one of the most sensitive to Calcofluor white, and inversely for IOEBL 0522.

Capacity to develop following inoculation

B. bruxellensis had a greater capacity than other species to resist direct inoculation in wine (Table 3) and was less affected by variations in pH, alcohol content, and SO₂. Unsurprisingly, *S. cerevisiae* was the species best suited to developing in grape juice. Its development rate was three times higher at pH = 3.6 than at pH = 3, whereas this value did not change significantly for *B. bruxellensis* strains. After AF, the survival rate of *S. cerevisiae* decreased by 100-fold at an ethanol content of 12.5% v/v, and 1000-fold at 15% v/v, while it remained relatively stable for *B. bruxellensis* strains. Unlike the other species tested, the latter exhibited strong resistance to SO₂. In two cases, the cell-count decreased by 10-fold (*R. mucilaginosa*) or 100-fold (*M. fructicola*) after forty-eight hours of incubation in wine containing only 10 mg/L free SO₂.

The key observation concerned the significant differences between *B. bruxellensis* strains. The CLIB300 strain was the least and IOEBL 0447 the most resistant to direct incubation under all conditions. The latter was more resistant to inoculation in must than in post-fermentation wines, irrespective of pH. This was not the case for the IOEBL 0462 strain, which developed twice as well at pH = 3.6 than pH = 3.0.

Table 2 Comparison of minimum inhibitory concentrations for different drugs on solid medium

| | | Calcofluor white (g/L) | Congo red (g/L) | Vanillin (g/L) |
|---------------------------------|-----------|------------------------|-----------------|----------------|
| <i>B. bruxellensis</i> | CLIB300 | 0.6 | 2.5 | 1.25 |
| | IOEBL0468 | 0.3 | 2.5 | 5 |
| | IOEBL0469 | 0.3 | 2.5 | 1.25 |
| | IOEBL0447 | 1.25 | 2.5 | 5 |
| | IOEBL0407 | 0.3 | 2.5 | 1.25 |
| | IOEBL0462 | 1.25 | 2.5 | 2.5 |
| | IOEBL0453 | 0.3 | 2.5 | 1.25 |
| | IOEBL0506 | 0.3 | 2.5 | 2.5 |
| | IOEBL0522 | 1.25 | 2.5 | 1.25 |
| <i>S. cerevisiae</i> | 522 Davis | 0.3 | 1.25 | 1.25 |
| <i>Metschnikowia fructicola</i> | IOEBL0530 | 0.3 | 0.3 | 0.3 |
| <i>Rhodotorula mucilaginosa</i> | CLIB370 | 0.6 | 2.5 | 2.5 |
| <i>Candida cantarelli</i> | IOEBL0416 | 0.15 | 0.3 | 0.6 |
| <i>Pichia anomala</i> | IOEBL0534 | 0.6 | 2.5 | 2.5 |

Table 3 Comparison of the capacity to develop following inoculation

| | Must pH = 3 | Must pH = 3.6 | Wine after AF and before MLF pH = 3.6 ethanol content = 12.5% v/v | Wine after AF and before MLF pH = 3.6 ethanol content = 15% v/v | Post-fermentation wine pH = 3.6 ethanol content = 13% v/v, free SO ₂ = 10 mg/L H ₂ SO ₄ | Post-fermentation wine pH = 3.6 ethanol content = 13% v/v, free SO ₂ = 30 mg/L H ₂ SO ₄ |
|---------------------------------|-------------|---------------|---|---|--|--|
| <i>B. bruxellensis</i> | | | | | | |
| CLIB300 | 1.6 ± 0.7 | 5.8 ± 2.3 | 1.5 ± 0.5 | 0.7 ± 0.1 | 1.8 ± 0.4 | 0.6 ± 0.2 |
| IOEBL0468 | 4.2 ± 0.3 | 4.4 ± 0.7 | 1.9 ± 1.0 | 1.5 ± 0.2 | 0.8 ± 0.1 | 0.7 ± 0.3 |
| IOEBL0469 | 5.3 ± 2.6 | 9.5 ± 1.6 | 6.5 ± 3.9 | 1.6 ± 0.9 | 1.7 ± 0.4 | 0.9 ± 0.1 |
| IOEBL0447 | 24.1 ± 2.9 | 27.1 ± 1.7 | 4.6 ± 2.6 | 5.4 ± 0.9 | 9.8 ± 1.4 | 3.4 ± 0.3 |
| IOEBL0407 | 12.1 ± 1.9 | 13.0 ± 1.1 | 1.2 ± 0.4 | 1.4 ± 0.4 | 1.3 ± 0.4 | 0.3 ± 0.2 |
| IOEBL0462 | 11.0 ± 1.2 | 21.1 ± 6.8 | 2.7 ± 1.0 | 0.8 ± 0.1 | 4.5 ± 0.8 | 2.4 ± 0.4 |
| IOEBL0453 | 15.4 ± 1.2 | 8.6 ± 2.6 | 1.4 ± 0.5 | 1.3 ± 0.1 | 0.9 ± 0.1 | 0.5 ± 0.3 |
| IOEBL0506 | 1.2 ± 0.3 | 1.30 ± 0.03 | 2.0 ± 0.8 | 1.6 ± 0.2 | 2.4 ± 1.1 | 0.6 ± 0.2 |
| IOEBL0522 | 1.2 ± 0.5 | 1.0 ± 0.1 | 1.7 ± 1.4 | 1.6 ± 0.2 | 4.7 ± 2.3 | 5.1 ± 0.2 |
| <i>S. cerevisiae</i> | | | | | | |
| 522 Davis | 32.7 ± 4.7 | 93.9 ± 2.5 | 0.2 ± 0.05 | 0.06 ± 0.02 | 0.15 ± 0.06 | 0.08 ± 0.02 |
| <i>Metschnikowia fructicola</i> | | | | | | |
| IOEBL0530 | 1.0 ± 0.7 | 27.0 ± 1.8 | 0.7 ± 0.2 | 0.06 ± 0.04 | 0.04 ± 0.03 | 0.02 ± 0.01 |
| <i>Rhodotorula mucilaginosa</i> | | | | | | |
| CLIB370 | 0.1 ± 0.06 | 2.0 ± 1.4 | 0.07 ± 0.03 | 0.05 ± 0.03 | 0.15 ± 0.05 | 0.02 ± 0.01 |
| <i>Candida cantarelli</i> | | | | | | |
| IOEB0L416 | 2.8 ± 1.0 | 2.0 ± 0.8 | 1.5 ± 0.5 | 0.03 ± 0.01 | 0.09 ± 0.05 | 0.04 ± 0.01 |
| <i>Pichia anomala</i> | | | | | | |
| IOEBL0534 | 10.2 ± 0.8 | 19.6 ± 0.2 | 2.9 ± 0.8 | 1.3 ± 0.2 | 1.5 ± 0.3 | 0.07 ± 0.02 |

Abbreviations: AF, alcoholic fermentation; MLF, malolactic fermentation.

Volatile phenol production

All the *B. bruxellensis* strains produced volatile phenols, especially 4-ethylphenol and 4-ethylguaiacol. However, amounts of volatile phenols produced varied by over three-fold according to the strain: 330 µg/L 4-ethylphenol for the IOEBL 0469 strain and 1200 µg/L for IOEBL 0462). Moreover, the ratio of 4-ethylphenol and 4-ethylguaiacol produced varied from one strain to another. Under the test conditions, it was nearly two for most strains. However, certain strains, such as IOEB L0447, produced equal amounts of 4-ethylphenol and 4-ethylguaiacol.

The other yeast species tested also produced volatile phenols, but only compounds with vinyl functions: 4-vinylphenol and 4-vinylguaiacol. Among these species, *S. cerevisiae* and *C. cantarelli* produced the largest amounts. Finally, while the *B. bruxellensis* strains produced more 4-ethylphenol than 4-ethylguaiacol, the other yeast species produced more 4-vinylguaiacol than 4-vinylphenol, except *P. anomola*, which also produced little 4-ethylphenol.

Discussion

The inhibitory concentrations for the drugs tested were obviously too high to be used to prevent *B. bruxellensis* development in wine but the findings revealed physiological differences among *B. bruxellensis* strains in terms of sensitivity or resistance to toxic constraints. Even if *B. bruxellensis* strains were more resistant than the other species to the drugs tested, the MIC measurements revealed significant differences in

responses to Calcofluor white and vanillin. These compounds act on cell-wall and membrane organisation, resulting in different resistance to various stressors, notably alcohol, one of the most important in wine, and others, due to cross-tolerance phenomena.¹⁹ The high resistance of *B. bruxellensis* strains to toxic pressure on a laboratory scale was not sufficient to determine their capacity to resist in wine, but revealed that they were able to adapt their metabolism to constraining environmental conditions.

That was confirmed by their ability to grow after direct incubation in wine. The most resistant strain to Calcofluor white and vanillin had also tolerated incubation in wine with a higher ethanol level. In general, *B. bruxellensis* strains were more resistant to ethanol than the other species, while *S. cerevisiae* performed better at high sugar concentrations. This explains the prevalence of *S. cerevisiae* strains during the first phases of AF, when the medium has a high sugar content and osmotic pressure.²⁰ *S. cerevisiae* had a higher growth rate under these conditions than the other genera. In contrast, *B. bruxellensis* prospers at the end of sluggish fermentations, when the ethanol content reaches high values.^{4,21,22} Moreover *B. bruxellensis* is not very demanding from a nutritional point of view.²³ Certain strains, notably those which originated from fermented wines, established themselves more easily in must than others, and were more resistant to high ethanol and SO₂ levels. These findings, however, did not establish a direct correlation between the stage where the strains were isolated and their intrinsic capacity to develop, but simply indicated

Table 4 Volatile phenol production

| | | 4-ethylphenol (µg/L) | 4-ethylguaiacol (µg/L) | 4-vinylphenol (µg/L) | 4-vinylguaiacol (µg/L) |
|---------------------------------|-----------|----------------------|------------------------|----------------------|------------------------|
| <i>B. bruxellensis</i> | CLIB300 | 605 ± 60 | 390 ± 40 | 8 ± 2 | 22 ± 2 |
| | IOEBL0468 | 775 ± 35 | 345 ± 65 | 0 | 0 |
| | IOEBL0469 | 330 ± 20 | 180 ± 22 | 0 | 6 ± 2 |
| | IOEBL0447 | 950 ± 20 | 540 ± 40 | 12 ± 3 | 20 ± 2 |
| | IOEBL0407 | 780 ± 70 | 760 ± 50 | 16 ± 2 | 22 ± 6 |
| | IOEBL0462 | 1200 ± 90 | 690 ± 40 | 4 ± 2 | 26 ± 2 |
| | IOEBL0453 | 770 ± 30 | 380 ± 10 | 4 ± 2 | 2 ± 2 |
| | IOEBL0506 | 850 ± 68 | 520 ± 8 | 0 | 0 |
| | IOEBL0522 | 1060 ± 110 | 800 ± 20 | 0 | 0 |
| <i>S. cerevisiae</i> | 522 Davis | 0 | 0 | 60 ± 6 | 264 ± 12 |
| <i>Metschnikowia fructicola</i> | IOEBL0530 | 0 | 0 | 26 ± 4 | 120 ± 8 |
| <i>Rhodotorula mucilaginosa</i> | CLIB370 | 0 | 0 | 14 ± 3 | 40 ± 14 |
| <i>Candida cantarelli</i> | IOEBL0416 | 10 ± 3 | 0 | 100 ± 4 | 530 ± 20 |
| <i>Pichia anomala</i> | IOEBL0451 | 54 ± 4 | 0 | 226 ± 12 | 47 ± 3 |
| Control: sterile must | | 0 | 0 | 12 ± 4 | 22 ± 2 |

that they were preferentially selected during fermentation. Some strains were clearly more sensitive than others to variations in pH, ethanol content, and SO₂, resulting in the preferential selection of some strains and the elimination of others, resulting in the decrease in intraspecific heterogeneity during the winemaking process observed by Miot-Sertier.⁷

Volatile phenol production also varied according to the strain. Some strains are likely to be more prejudicial than others with regard to the problem of volatile phenols. Further work in molecular characterization, focused on identifying enzymes, genes, and the regulation patterns involved, is required to elucidate the volatile phenol production mechanism. Comparisons with other species that produce only vinyl derivatives will be made.

The *B. bruxellensis* strains collected in vineyards²⁴ differed from one estate to another,²⁵ so the physiological differences identified in this research may explain why certain cellars are more exposed to this problem than others and why nonobservance of hygiene recommendations is not the only relevant parameter. The prevalence and degree of spoilage may vary according to the ability of the strains present to develop under winemaking conditions and their capacity to produce volatile phenols.

Disclosure

The authors report no conflicts of interest in this work.

References

- Renouf V, Lonvaud-Funel A, Coulon J. The origin of *Brettanomyces bruxellensis* in wines: a review. *J Int Sc Vigne Vin*. 2007;41:161–173.
- Renouf V, Gindreau E, Claisse O, Lonvaud-Funel A. Microbial changes during malolactic fermentation in red wine elaboration. *J Int Sc Vigne Vin*. 2005;39:1–12.
- Renouf V, Perello MC, Strehaiano P, Lonvaud-Funel A. Global survey of the microbial ecosystem during alcoholic fermentation. *J Int Sc Vigne Vin*. 2006;40:101–116.
- Renouf V, Falcou M, Miot-Sertier C, Perello MC, de Revel G, Lonvaud-Funel A. Interactions between *Brettanomyces bruxellensis* and other yeasts species during the first steps of winemaking. *J Appl Microbiol*. 2006;100:1208–1219.
- Couto JA, Neves F, Campos F, Hogg T. Thermal inactivation of wine spoilage yeast *Dekkera/Brettanomyces*. *Int J Food Microbiol*. 2005;104:337–344.
- Delaherche A, Claisse O, Lonvaud-Funel A. Detection and quantification of *Brettanomyces bruxellensis* and ‘ropy’ *Pediococcus damnosus* strains in wine by real-time polymerase chain reaction. *J Appl Microbiol*. 2004;97:910–915.
- Miot-Sertier C, Lonvaud-Funel A. Development of a molecular method for the typing of *Brettanomyces bruxellensis* (*Dekkera bruxellensis*) at the strain level. *J App Microbiol*. 2007;102:555–562.
- Miot-Sertier C, Renouf V, Lonvaud-Funel A. REA-PFGE: a tool for traceability of *Brettanomyces bruxellensis* strains during the winemaking process. 30th SASEV Congress, Cape Town, South Africa.
- Conterno L, Lucy Joseph CM, Arvik TJ, Henick-Kling T, Bisson L. Genetic and physiological characterization of *Brettanomyces bruxellensis* strains isolated from wines. *Am J Enol Vitic*. 2006;57:139–147.
- Curtin CD, Bellon JR, Henschke PA, Godden PW, De Barros Lopes MA. Genetic diversity of *Dekkera bruxellensis* yeast isolated from Australian winery. *FEMS Yeast Res*. 2007;7:471–481.
- Ibeas JL, Lozano I, Perdogones F, Jimenez J. Detection of *Dekkera-Brettanomyces* strains in sherry by a nested PCR method. *Appl Environ Microbiol*. 1996;62:998–1003.
- Oelofse A, Pretorius IS, du Toit M. Significance of *Brettanomyces* and *Dekkera* during winemaking: a synoptic review. *S Afr J Enol Vitic*. 2008;29:128–144.
- Millet V, Lonvaud-Funel A. The viable but non-cultivable state of wine microorganisms during storage. *Lett Appl Microbiol*. 2000;30:136–141.
- Conner DE, Beuchat LR, Worthington RE, Hitchcock HL. Effects of essential oils and oleoresins of plants on ethanol production, respiration and sporulation of yeast. *Int J Food Microbiol*. 1984;1:63–74.
- Rico-Munoz E, Bargiota EE, Davidson PM. Effect of selected phenolic compounds on the membran-bound adenosine triphosphatase of *Staphylococcus aureus*. *Food Microbiol*. 1987;4:239–249.
- Elorza V, Rico H, Sentandreu R. Calcofluor white alters the assembly of chitin fibrils in *Saccharomyces cerevisiae* and *Candida albicans* cells. *J Gen Microbiol*. 1983;129:1577–1582.
- Roncero C, Duran A. Effect of Calcofluor white and congo red on fungal cell wall morphogenesis in vivo activation of chitin polymerisation. *J Bacteriol*. 1985;163:1180–1185.
- Chatonnet P, Boiron JN. Dosage des phénols volatils dans les vins par la chromatographie en phase gazeuse. *Sci Aliments*. 1988;8:479–488.
- Piper PW. The heat shock and ethanol stress responses of yeast exhibit extensive similarity and functional overlap. *FEMS Microbiol Lett*. 1995;134:121–127.
- Abbott DA, Hynes SH, Ingledew WN. Growth rates of *Dekkera/Brettanomyces* yeasts hinder their ability to compete with *Saccharomyces cerevisiae* in batch corn mash fermentation. *Appl Microbiol Biotechnol*. 2004;66:641–647.
- Medawar W, Strehaiano P, Delia ML. Yeast growth: lag phase modeling in alcoholic media. *Food Microbiol*. 2003;20:527–532.
- Silva P, Cardoso H, Geros H. Studies on the wine spoilage capacity of *Brettanomyces/Dekkera* spp. *Am J Enol Vitic*. 2004;55:65–72.
- Uscanga MG, Delia ML, Strehaiano P. Nutritional requirements of *Brettanomyces bruxellensis*: growth and physiology in batch chemostat cultures. *Can J Microbiol*. 2000;46:1046–1050.
- Renouf V, Miot-Sertier C, Strehaiano P, Lonvaud-Funel A. The wine microbial consortium: a real terroir characteristic. *J Int Sc Vigne Vin*. 2006;40:209–217.
- Renouf V, Lonvaud-Funel A. Development of an enrichment medium to detect *Dekkera/Brettanomyces bruxellensis*, a spoilage wine yeast, on the surface of grape berries. *Microbiol Res*. 2007;161:154–167.