

The association between COPD and outcomes of patients with advanced chronic kidney disease

Chih-Cheng Lai¹
 Che-Hsiung Wu^{2,3}
 Ya-Hui Wang⁴
 Cheng-Yi Wang⁴
 Vin-Cent Wu⁵
 Likwang Chen⁶

¹Department of Intensive Care Medicine, Chi Mei Medical Center, Liouying, Taiwan; ²Division of Nephrology, Taipei Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Taipei, Taiwan; ³School of Medicine, Tzu Chi University, Hualien, Zhunan, Taiwan; ⁴Medical Research Center, Cardinal Tien Hospital and School of Medicine, College of Medicine, Fu-Jen Catholic University, New Taipei City, Taiwan; ⁵Department of Internal Medicine, National Taiwan University Hospital and College of Medicine, National Taiwan University, Taipei, Taiwan; ⁶Institute of Population Health Sciences, National Health Research Institutes, Zhunan, Taiwan

Objective: The aim of this study was to investigate the impact of COPD on the outcomes of patients with advanced chronic kidney disease (CKD).

Patients and methods: All patients with advanced CKD from 2000 to 2010 were identified from the Taiwanese National Health Insurance Research Database. Associations between COPD and the risk of long-term dialysis and all-cause mortality were assessed.

Results: A total of 33,399 advanced CKD patients were enrolled, of whom 31,536 did not have COPD (non-COPD group) and 1,863 had COPD (COPD group). The incidence of end-stage renal disease (ESRD) was higher for those with COPD than those without COPD (744.2 per 1,000 person-years vs 724.6 per 1,000 person-years, adjusted HR [aHR] 1.04; 95% CI 0.96–1.12). The cumulative incidence rates of ESRD were similar between the COPD and non-COPD groups (log-rank test, $P=0.356$). Overall, the patients with COPD had a higher risk of death than those without COPD (151.7 per 1,000 person-years vs 125.5 per 1,000 person-years, aHR 1.22; 95% CI 1.11–1.33). The cumulative mortality rate was higher in the COPD group than in the non-COPD group (log-rank test, $P<0.001$).

Conclusion: COPD increased the risk of mortality among the advanced CKD patients in this study, especially the elderly and male patients. In contrast, COPD did not increase the risk of ESRD among the advanced CKD patients.

Keywords: COPD, CKD, ESRD, mortality

Introduction

COPD population is gradually increasing worldwide,^{1,2} and comorbidities including heart failure, ischemic heart disease, lung cancer, depression/anxiety, arrhythmia, sleep apnea, hypertension, and diabetes mellitus often coexist with COPD.^{3–5} In addition, a recent meta-analysis⁶ showed that COPD patients carry a higher risk of developing chronic kidney disease (CKD) (OR 2.20; 95% CI 1.83–2.65). Furthermore, a nationwide case cohort study⁷ using the Taiwan National Health Insurance Research Database (NHIRD) reported similar findings in which the overall incidence of CKD was higher among patients with COPD (470.9 per 10,000 person-years) than in those without COPD (287.52 per 10,000 person-years), with an adjusted HR (aHR) of 1.61 (95% CI 1.52–1.72). Conversely, a Korean population-based study⁸ reported that CKD could be one of the risk factors for COPD. Moreover, another study reported a 41% (95% CI 1.31–1.52) increased risk of mortality among COPD patients.⁹

Advanced CKD is the most severe form of CKD, which involves a severe reduction in glomerular filtration rate. Early detection and appropriate management of advanced CKD can help improve long-term morbidity and slow the deterioration of renal function. Despite the complicated association between COPD and CKD that has been reported, information regarding the impact of COPD on advanced CKD is

Correspondence: Ya-Hui Wang;
 Cheng-Yi Wang
 Medical Research Center, Cardinal Tien Hospital and School of Medicine, College of Medicine, Fu Jen Catholic University, No 362, Zhongzheng Road, Xindian District, New Taipei City 23148, Taiwan
 Tel +886 2 2219 3391x15433
 Fax +886 2 2219 0651
 Email yhwang531@gmail.com;
 cywang@mospital.com

limited. Thus, this study investigated the impact of COPD on two major outcomes: the risk of end-stage renal disease (ESRD) and mortality of advanced CKD patients.

Patients and methods

Data sources

The Taiwan NHIRD is established by the National Health Research Institute (NHRI) and includes detailed medical care records of more than 97% of the Taiwanese residents. NHIRD included all claims data for outpatient visits, prescriptions, inpatient care records, and disease. This study obtained ethical approval from the institutional review board (IRB) of National Taiwan University Hospital. The used data from NHIRD were de-identified; therefore, the IRB waived informed consent from the enrolled patients.

Study group

All patients aged ≥ 40 years who were diagnosed with CKD and received erythropoiesis-stimulating agent (ESA) treatment from January 1, 2000, to December 31, 2010, were identified from NHIRD. Due to the reimbursement regulations of the National Health Insurance (NHI), only anemic patients with advanced CKD with a hematocrit level of $\leq 28\%$ and a serum creatinine level of > 6 mg/dL can be prescribed ESAs. In this study, we defined advanced CKD as the presence of CKD coding and concomitant reimbursements for prescriptions of ESA like the previous study.¹⁰ The index date was defined as the date of the first ESA. Patients were excluded if they commenced dialysis before the index date, received dialysis, or died within 90 days after the index date, or if they received a renal transplant.

The COPD group comprised patients with advanced CKD and incident COPD within 1 year after the index hospitalization. To ensure accuracy, the diagnosis of COPD was validated based on one inpatient or three outpatient records of International Classification of Diseases (ICD), ninth revision, Clinical Modification (ICD-9-CM) codes 491, 492, and 496 and the prescription of at least one bronchodilator during the follow-up period.^{11–13} Patients without COPD and no history of asthma or COPD medications were included in the control (non-COPD) group.

Initially, 78,551 eligible patients with advanced CKD were identified between January 1, 2000, and December 31, 2010. Then, 41,361 patients who had a history of dialysis in the previous year, 2,040 patients aged < 40 years or > 100 years, 669 patients who received a kidney transplantation, and 1,082 patients who died within 3 months were excluded. The remaining 33,399 patients with advanced CKD were included for further analysis.

Research variables

Baseline underlying diseases/conditions including CKD were defined as at least three outpatient visits or one inpatient claim within 1 year preceding the first ESA prescription like previous studies.^{14,15} We collected the Charlson comorbidity index using baseline comorbidities as previously described.¹⁶ The prescriptions that might be associated with the risks of dialysis or all-cause mortality were identified if they were used within 30 days before the events, including alpha-blocker, beta-blocker, calcium channel blocker, diuretics, angiotensin-converting enzyme inhibitors (ACEIs), angiotensin II receptor blockers (ARBs), aspirin, clopidogrel, ticlopidine, dipyridamole, nitrate, statin, and proton pump inhibitor.

Outcome measures

The follow-up period started from the index date to the date of commencing long-term dialysis, death, or December 31, 2010, whichever occurred first. The primary end point was the long-term dialysis,¹⁷ and the secondary end point was all-cause mortality.

Statistical analysis

Continuous variables are presented as mean \pm SD, and categorical variables are described as counts and percentages. We used R software, version 2.8.1 (Free Software Foundation, Inc., Boston, MA, USA) for the statistical analysis. Given the differences in baseline characteristics and risk of long-term dialysis or all-cause mortality between the incident COPD and non-COPD groups, we matched the COPD patients with non-COPD patients using a greedy matching algorithm with a caliper width of 0.2 SD of the log of the odds of the estimated propensity score at a 1:1 ratio. Crude HRs with 95% CIs for the outcomes of interest were derived from Cox proportional hazards models, with matched individuals without COPD constituting the reference group. Because of the high mortality rate in patients with COPD and advanced CKD, competing risk regression was also performed using the Fine and Gray model considering the sub-distribution hazard.^{18,19}

Results

Patient characteristics

Among the 33,399 patients, 31,536 did not have COPD (non-COPD group) and 1,863 patients had COPD (COPD group). Significant differences were noted between the two groups in demographic characteristics and baseline comorbidities. In addition, we found significant differences between

the two groups in medications for hypertension and other diseases. Thus, pairwise matching (1:1) of the non-COPD and COPD groups was used and resulted in two similar subgroups with 1,820 patients in each. No significant differences were noted regarding age, gender, monthly income, hospital level, Charlson scores, underlying disease, and antihypertension and cardiovascular agents between these two groups (Table 1).

The incidence of ESRD and mortality

Table 2 summarizes the event rates of ESRD and mortality in the matched COPD and non-COPD cohorts. The incidence

of ESRD was 744.2 per 1,000 person-years in the COPD group and 724.6 per 1,000 person-years in the non-COPD group. However, this difference did not reach statistical significance (aHR 1.04; 95% CI 0.96–1.12; Table 2). A similar trend was noted in competing analysis in which we treated mortality as a competing risk. The mortality rate was 151.7 per 1,000 person-years in the COPD group and 125.5 per 1,000 person-years in the non-COPD group. Overall, the patients with COPD had a higher risk of mortality (aHR 1.22; 95% CI 1.11–1.33) than those without COPD (Table 2). The cumulative incidence rates of ESRD were similar between the COPD and non-COPD groups (log-rank test, $P=0.356$;

Table 1 Demographics and characteristics of COPD cohort and matched non-COPD cohort

Variables	Non-COPD cohort (n=1,820)	COPD cohort (n=1,820)	Standardized difference
Age (years, mean \pm SD)	71.53 \pm 10.86	71.56 \pm 10.88	-0.003
Male gender	1,145 (62.91%)	1,142 (62.75%)	-0.003
Monthly income (New Taiwan dollar), n (%)			
<19,100	991 (54.45%)	973 (53.46%)	0.010
19,100–41,999	749 (41.15%)	784 (43.08%)	-0.026
\geq 42,000	80 (4.40%)	63 (3.46%)	0.048
Hospital level, n (%)			
Urban	685 (37.64%)	697 (38.30%)	-0.008
Suburban	453 (24.89%)	455 (25.00%)	-0.002
Rural	682 (37.47%)	668 (36.70%)	0.014
Baseline comorbidities			
Charlson score	3.63 \pm 1.74	3.57 \pm 1.75	0.034
Myocardial infarction	36 (1.98%)	45 (2.47%)	0.034
Congestive heart failure	368 (20.22%)	368 (20.22%)	0.000
Peripheral vascular disease	31 (1.70%)	24 (1.32%)	-0.032
Cerebrovascular disease	201 (11.04%)	195 (10.71%)	-0.011
Dementia	42 (2.31%)	52 (2.86%)	0.035
Rheumatologic disease	10 (0.55%)	19 (1.04%)	0.056
Peptic ulcer disease	380 (20.88%)	374 (20.55%)	-0.008
Hemiplegia or paraplegia	9 (0.49%)	10 (0.55%)	0.008
Diabetes	829 (45.55%)	794 (43.63%)	-0.039
Moderate or severe liver disease	106 (5.82%)	82 (4.51%)	-0.060
Asthma	187 (10.27%)	190 (10.44%)	0.005
Medication for hypertension			
Alpha-blocker	399 (21.92%)	420 (23.08%)	0.028
Beta-blocker	818 (44.95%)	814 (44.73%)	-0.004
Calcium channel blocker	1,358 (74.62%)	1,360 (74.73%)	0.003
Diuretic	1,255 (68.96%)	1,258 (69.12%)	0.004
ACEI or ARB	1,051 (57.75%)	1,044 (57.36%)	-0.008
Other medications			
Aspirin	193 (10.60%)	197 (10.82%)	0.007
Clopidogrel	142 (7.80%)	155 (8.52%)	0.026
Ticlopidine	49 (2.69%)	57 (3.13%)	0.026
Dipyridamole	656 (36.04%)	666 (36.59%)	0.011
Nitrate	30 (1.65%)	36 (1.98%)	0.025
Statin	398 (21.87%)	407 (22.36%)	0.012
Proton pump inhibitor	264 (14.51%)	280 (15.38%)	0.025

Abbreviations: ACEI, angiotensin-converting enzyme inhibitor; ARB, angiotensin II receptor blocker.

Table 2 Cumulative incidence rates and HRs of ESRD and mortality in COPD and non-COPD patients

	COPD			Non-COPD			Crude HR (95% CI)	aHR ^a (95% CI)	Competing risk aHR ^a (95% CI)
	Event	Person-year	IR ^b	Event	Person-year	IR ^b			
ESRD	1,443	1,939.08	744.2	1,429	1,972.08	724.6	1.03 (0.96, 1.11)	1.04 (0.96, 1.12)	1.04 (0.96, 1.11)
Mortality	994	6,553.03	151.7	838	6,679.34	125.5	1.21 (1.10, 1.33)	1.22 (1.11, 1.33)	

Notes: ^aAdjusted for propensity score. ^bIR per 1,000 person-years.

Abbreviations: aHR, adjusted HR; ESRD, end-stage renal disease; IR, incidence rate.

Figure 1). In contrast, the mortality rate was higher in the COPD group than in the non-COPD group (log-rank test, $P < 0.001$; Figure 2).

Subgroup analysis

Although we performed further subgroup analysis, we did not find significant differences in terms of ESRD among the age and gender groups. However, a trend of a higher risk of ESRD with age was observed (Table 3). For the CKD patients aged 55–69 and ≥ 70 years, those with COPD had a significantly higher risk of mortality than those without COPD. In contrast, no significant difference was found in the CKD patients aged 40–54 years. A similar trend was observed in the male patients. For the female patients, those with COPD had a higher risk of mortality than those without COPD; however, none of the differences reached statistical significance (Table 4).

Discussion

In this study, we used a national database to assess the impact of COPD on the renal outcomes and mortality of advanced CKD patients. We found that the advanced CKD patients with COPD had a significantly higher risk of mortality than those without COPD (aHR 1.22; 95% CI 1.11–1.33). A similar finding was reported in the study based on CKD registry records at the Cleveland Clinic, in which COPD was

associated with a 41% increased risk (95% CI 1.31–1.52) of mortality among patients with stage 3 and 4 CKD.⁹ Another survey²⁰ of 3,358 patients who underwent vascular surgery revealed that both moderate and severe COPD were associated with increased mortality in CKD patients compared to patients without COPD (HR 1.27; 95% CI 1.03–1.56 and HR 1.61; 95% CI 1.10–2.35, respectively). Overall, our findings are in line with previous studies,^{9,20} suggesting that COPD can negatively impact mortality rates in patients with advanced CKD.

Navaneethan et al⁹ reported that significant associations between mortality and COPD were more pronounced in younger CKD patients and female patients with CKD. In contrast, the negative impact of COPD on the mortality rate of the advanced CKD patients in this study was noted only in patients aged older than 55 years and in male patients, but not in the female patients. Further studies are warranted to investigate this issue among specific groups.

Although the risk of ESRD was higher in the COPD group than in the non-COPD group in this study, the difference did not reach statistical significance (aHR 1.04; 95% CI 0.96–1.12). In the subgroup analysis according to age and gender, we still did not find any significant differences regarding the risk of ESRD between the COPD and non-COPD groups. Thus, our findings suggest that COPD does not affect the risk of ESRD among advanced CKD




Figure 1 Cumulative incidence of ESRD in patients with or without COPD.
Abbreviation: ESRD, end-stage renal disease.




Figure 2 Cumulative incidence of death in patients with or without COPD.

Table 3 Rates and risk ratio of cumulative incidence of ESRD in COPD and non-COPD patients by age and gender

	Cumulative incidence by age			
	All age	40–54 years	55–69 years	>70 years
COPD patients				
ESRD (n)	1,443	140	459	844
COPD patients (n)	1,820	153	525	1,142
Rate in COPD patients	0.79	0.92	0.87	0.74
Non-COPD patients				
ESRD (n)	1,429	146	474	809
Non-COPD patients (n)	1,820	153	531	1,136
Rate in non-COPD patients	0.79	0.95	0.89	0.71
Risk ratio (95% CI)	1.01 (0.98, 1.04)	0.959 (0.90, 1.02)	0.98 (0.94, 1.02)	1.04 (0.99, 1.09)
COPD male				
ESRD (n)	889	80	276	533
COPD male (n)	1,142	89	322	731
Rate in COPD male	0.78	0.90	0.86	0.73
Non-COPD male				
ESRD (n)	889	91	296	502
Non-COPD male (n)	1,145	95	330	720
Rate in non-COPD male	0.78	0.96	0.90	0.70
Risk ratio	1.00 (0.96, 1.05)	0.94 (0.87, 1.02)	0.96 (0.90, 1.01)	1.05 (0.98, 1.12)
COPD female				
ESRD (n)	554	60	183	311
COPD female (n)	678	64	203	411
Rate in COPD female	0.82	0.94	0.90	0.76
Non-COPD female				
ESRD (n)	540	55	178	307
Non-COPD female (n)	675	58	201	416
Rate in non-COPD female	0.80	0.95	0.89	0.74
Risk ratio	1.02 (0.97, 1.08)	0.99 (0.91, 1.08)	1.02 (0.95, 1.09)	1.03 (0.95, 1.11)

Abbreviation: ESRD, end-stage renal disease.

patients and also that COPD itself may not impact the renal outcomes of COPD patients.

In addition, the prevalence of COPD among the advanced CKD patients was about 5.6% (1,863/33,399) in this study. This result is consistent with the findings of one recent investigation in a large health care system among 5,960 patients with stage 3 and 4 CKD, of whom 4.7% (n=2,667) had underlying COPD.⁹ Another study of CKD and COPD among patients who had undergone vascular surgery reported that 4.9% (n=45) of 918 patients with a glomerular filtration rate of <60 mL/min/1.73 m² had COPD and that the prevalence of COPD was inversely associated with renal function.²⁰ Overall, these findings suggest that a significant proportion of CKD patients have underlying COPD and that clinicians should be alert to the presence of COPD among patients with CKD.

Because this study aimed to investigate the effect of COPD on the long-term risk of dialysis and mortality among patients with advanced CKD, we excluded patients who commenced dialysis before the index date and died within 90 days after the index date. However, CKD itself

can also affect the long-term outcome of COPD patients,²¹ and even acute kidney injury was associated with in-hospital mortality among patients with COPD exacerbation.²² Thus, the interaction between CKD and COPD could be bidirectional, and further study should be warranted to clarify their interaction.

This study had one major strength. We enrolled a large cohort of CKD patients with a long follow-up period. Therefore, we can include many known confounding factors to minimize their possible effects. However, this study had several limitations. Some confounding factors, such as lung function test and clinical symptoms/signs were not available from the NHIRD. Therefore, we cannot assess the severity of COPD. In addition, we can obtain only the data on all-cause mortality from NHIRD, so we cannot investigate specific causes of mortality. Moreover, data based on ICD-9-CM codes could not give information on pathology severity, functional status, or intensity of care given, including aggressive therapy and devices used. Diagnosis based on ICD-9-CM codes could be influenced from individual ability and codifying hospital procedures. Finally, we could

Table 4 Rates and risk ratio of cumulative incidence of death in COPD and non-COPD patients by age and gender

	Cumulative incidence by age			
	All age	40–54 years	55–69 years	>70 years
COPD patients				
Death (n)	994	35	258	701
COPD patients (n)	1,820	153	525	1,142
Rate in COPD patients	0.55	0.23	0.49	0.61
Non-COPD patients				
Death (n)	838	32	206	600
Non-COPD patients (n)	1,820	153	531	1,136
Rate in non-COPD patients	0.46	0.21	0.39	0.53
Risk ratio (95% CI)	1.19 (1.11, 1.27)	1.09 (0.72, 1.67)	1.27 (1.10, 1.45)	1.16 (1.08, 1.25)
COPD male				
Death (n)	652	22	176	454
COPD male (n)	1,142	89	322	731
Rate in COPD male	0.57	0.25	0.55	0.62
Non-COPD male				
Death (n)	523	24	132	367
Non-COPD male (n)	1,145	95	330	720
Rate in non-COPD male	0.46	0.25	0.40	0.51
Risk ratio	1.25 (1.15, 1.36)	0.98 (0.59, 1.62)	1.37 (1.16, 1.61)	1.22 (1.11, 1.34)
COPD female				
Death (n)	342	13	82	247
COPD female (n)	678	64	203	411
Rate in COPD female	0.50	0.20	0.40	0.60
Non-COPD female				
Death (n)	315	8	74	233
Non-COPD female (n)	675	58	201	416
Rate in non-COPD female	0.47	0.14	0.37	0.56
Risk ratio	1.08 (0.97, 1.21)	1.47 (0.66, 3.30)	1.10 (0.86, 1.40)	1.07 (0.96, 1.21)

not ascribe worsening renal function to any specific cause based on this database.

Conclusion

COPD increased the risk of mortality among the patients with advanced CKD in this study, especially the elderly and male patients. In contrast, COPD did not affect the risk of ESRD among the patients with advanced CKD.

Disclosure

The authors report no conflicts of interest in this work.

References

- Landis SH, Muellerova H, Mannino DM, et al. Continuing to Confront COPD International Patient Survey: methods, COPD prevalence, and disease burden in 2012–2013. *Int J Chron Obstruct Pulmon Dis.* 2014;9:597–611.
- Rycroft CE, Heyes A, Lanza L, Becker K. Epidemiology of chronic obstructive pulmonary disease: a literature review. *Int J Chron Obstruct Pulmon Dis.* 2012;7:457–494.
- Barnes PJ, Celli BR. Systemic manifestations and comorbidities of COPD. *Eur Respir J.* 2009;33(5):1165–1185.
- Miller J, Edwards LD, Agustí A, et al. Comorbidity, systemic inflammation and outcomes in the ECLIPSE cohort. *Respir Med.* 2013;107(9):1376–1384.
- Soriano JB, Visick GT, Muellerova H, Payvandi N, Hansell AL. Patterns of comorbidities in newly diagnosed COPD and asthma in primary care. *Chest.* 2005;128(4):2099–2107.
- Gaddam S, Gunukula SK, Lohr JW, Arora P. Prevalence of chronic kidney disease in patients with chronic obstructive pulmonary disease: a systematic review and meta-analysis. *BMC Pulm Med.* 2016;16(1):158.
- Chen CY, Liao KM. Chronic Obstructive Pulmonary Disease is associated with risk of Chronic Kidney Disease: A Nationwide Case-Cohort Study. *Sci Rep.* 2016;6:25855.
- Kim MY, Boo S, Yoo M, Lee J, Kang NR. Impact of chronic kidney disease among Korean adults with chronic obstructive pulmonary disease. *Int Urol Nephrol.* 2017;49(7):1225–1232.
- Navaneethan SD, Schold JD, Huang H, et al. Mortality Outcomes of Patients with Chronic Kidney Disease and Chronic Obstructive Pulmonary Disease. *Am J Nephrol.* 2016;43(1):39–46.
- Hsu TW, Liu JS, Hung SC, et al. Renoprotective effect of renin-angiotensin-aldosterone system blockade in patients with predialysis advanced chronic kidney disease, hypertension, and anemia. *JAMA Intern Med.* 2014;174(3):347–354.
- Li CH, Chen WC, Liao WC, et al. The association between chronic obstructive pulmonary disease and Parkinson's disease: a nationwide population-based retrospective cohort study. *QJM.* 2015;108(1):39–45.
- Huang JY, Jian ZH, Nfor ON, et al. The effects of pulmonary diseases on histologic types of lung cancer in both sexes: a population-based study in Taiwan. *BMC Cancer.* 2015;15:834.
- Liao KM, Chen CY. Incidence and risk factors of atrial fibrillation in Asian COPD patients. *Int J Chron Obstruct Pulmon Dis.* 2017;12:2523–2530.

14. Wu VC, Hu YH, Wu CH, et al. Administrative data on diagnosis and mineralocorticoid receptor antagonist prescription identified patients with primary aldosteronism in Taiwan. *J Clin Epidemiol*. 2014;67(10):1139–1149.
15. Cheng CL, Kao YH, Lin SJ, Lee CH, Lai ML. Validation of the National Health Insurance Research Database with ischemic stroke cases in Taiwan. *Pharmacoepidemiol Drug Saf*. 2011;20(3):236–242.
16. Charlson ME, Pompei P, Ales KL, Mackenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373–383.
17. Ou SM, Chen YT, Chao PW, et al. Nonsteroidal anti-inflammatory drug use is associated with cancer risk reduction in chronic dialysis patients. *Kidney Int*. 2013;84(1):198–205.
18. Wu VC, Chueh SJ, Chen L, et al. Risk of new-onset diabetes mellitus in primary aldosteronism: a population study over 5 years. *J Hypertens*. 2017;35(8):1698–1708.
19. Wu VC, Chang CH, Wang CY, et al. Risk of Fracture in Primary Aldosteronism: A Population-Based Cohort Study. *J Bone Miner Res*. 2017;32(4):743–752.
20. van Gestel YR, Chonchol M, Hoeks SE, et al. Association between chronic obstructive pulmonary disease and chronic kidney disease in vascular surgery patients. *Nephrol Dial Transplant*. 2009;24(9):2763–2767.
21. Fedeli U, de Giorgi A, Gennaro N, et al. Lung and kidney: a dangerous liaison? A population-based cohort study in COPD patients in Italy. *Int J Chron Obstruct Pulmon Dis*. 2017;12:443–450.
22. Fabbian F, de Giorgi A, Manfredini F, et al. Impact of renal dysfunction on in-hospital mortality of patients with severe chronic obstructive pulmonary disease: a single-center Italian study. *Int Urol Nephrol*. 2016;48(7):1121–1127.

International Journal of COPD

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>

This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress