

Use of transcutaneous electrical acupoint stimulation for prevention of postoperative delirium in geriatric patients with silent lacunar infarction

This article was published in the following Dove Medical Press journal:
Clinical Interventions in Aging

Hai-Jun Hou
Fu-Shan Xue
Rui-Juan Guo

Department of Anesthesiology, Beijing Friendship Hospital, Capital Medical University, Beijing, People's Republic of China

Dear editor

In a preliminary study, Gao et al¹ assessed the preventive effect of transcutaneous electrical acupoint stimulation (TEAS) on postoperative delirium (POD) in geriatric patients with silent lacunar infarction. They showed that TEAS could reduce the development of POD and might be related to attenuated neuroinflammation by reducing the permeability of the blood–brain barrier. Given that POD is a common postoperative complication associated with adverse events and outcomes including functional decline, and increased risks of morbidity and mortality in elderly surgical patients, their findings have the potential implications. To differentiate the real effect of one factor on primary endpoint in a randomized trial, however, all of other factors must be standardized for avoidance of potential bias. Other than the limitations described in the discussion, we noted several issues in this study that were not addressed well.

First, study subjects were elderly patients undergoing spine surgery. The study did not provide the baseline education level and preoperative serum albumin concentration of patients. In available literature, both lower baseline education level and preoperative albumin concentration have been identified as significant risk factor of POD in elderly surgical patients.^{2,3} Regarding intraoperative factors, moreover, only providing anesthesia and operation durations were insufficient. It has been shown that intraoperative major blood loss and blood transfusion are significantly associated with an increased risk of POD after noncardiac surgery in elderly patients.⁴ In addition, the authors did not observe and compare the incidences of postoperative adverse events and complications between groups. In fact, prolonged hospital and intensive care unit stay, postoperative complications including urinary tract infection, pneumonia, and cognitive impairment can increase the risk of POD after noncardiac surgery.⁵ We are concerned that any imbalance in the above unknown factors would have biased their findings.

Second, this study excluded patients with an increased risk of POD, such as those with mini-mental state examination score of <24 or dementia, preoperative delirium, history of neurological or mental illness, current use of tranquilizers or antidepressants.⁵ Thus, an important question that remains unanswered in this study is whether TEAS is an effective treatment in elderly patients with an increased risk of POD. Furthermore, this study only assessed the incidence of POD, but not the severity and duration of POD due to a short observed time. The available evidence indicates that both more severe POD and longer in-hospital POD duration are significantly associated with

Correspondence: Fu-Shan Xue
Department of Anesthesiology, Beijing Friendship Hospital, Capital Medical University, No 95 Yong-An Road, Xi-Cheng District, Beijing 100050, People's Republic of China
Tel +86 139 1117 7655
Fax +86 10 6313 8362
Email xuefushan@aliyun.com

worse postoperative outcomes.⁶ We believe that the results of this study would be more informative if the design had included these issues.

Finally, because of the small sample size, small inter-group mean differences and large standard deviations in this study, we questioned their results that serum levels of TNF- α and IL-6 were higher at T₂₋₃ and serum levels of MMP-9 and S100 β were higher at T₃ in group C compared with group TEAS ($P < 0.05$). Revalidation of these results with statistical software confirms our doubts.

Acknowledgment

All authors report no financial support.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Gao F, Zhang Q, Li Y, et al. Transcutaneous electrical acupoint stimulation for prevention of postoperative delirium in geriatric patients with silent lacunar infarction: a preliminary study. *Clin Interv Aging*. 2018;13:2127–2134.
2. Sprung J, Roberts RO, Weingarten TN, et al. Postoperative delirium in elderly patients is associated with subsequent cognitive impairment. *Br J Anaesth*. 2017;119(2):316–323.
3. Patti R, Saitta M, Cusumano G, Termine G, Di Vita G. Risk factors for postoperative delirium after colorectal surgery for carcinoma. *Eur J Oncol Nurs*. 2011;15(5):519–523.
4. Marcantonio ER, Goldman L, Orav EJ, Cook EF, Lee TH. The association of intraoperative factors with the development of postoperative delirium. *Am J Med*. 1998;105:380–384.
5. Smith TO, Cooper A, Peryer G, Griffiths R, Fox C, Cross J. Factors predicting incidence of post-operative delirium in older people following hip fracture surgery: a systematic review and meta-analysis. *Int J Geriatr Psychiatry*. 2017;32(4):386–396.
6. Hughes CG, Patel MB, Jackson JC, et al.; MIND-ICU, BRAIN-ICU Investigators. Surgery and anesthesia exposure is not a risk factor for cognitive impairment after major noncardiac surgery and critical illness. *Ann Surg*. 2017;265(6):1126–1133.

Authors' reply

Fang Gao,¹ Qi Zhang,² Yanan Li,¹ Yanlei Tai,¹ Xi Xin,³ Xiuli Wang,¹ Qiuju Wang¹

¹Department of Anesthesiology, The Third Hospital of Hebei Medical University, Shijiazhuang City, Hebei, People's Republic of China; ²Department of Anesthesiology, Children's Hospital of Hebei Province, Shijiazhuang City, Hebei, People's Republic of China; ³Department of Anesthesiology, Tianjin Third Central Hospital, Tianjin, People's Republic of China

Correspondence: Qiuju Wang
 Department of Anesthesiology, The Third Hospital of Hebei Medical University, No 139, Ziqiang Road, Shijiazhuang City, Hebei, People's Republic of China
 Tel/fax +86 311 8860 2072
 Email 13933178001@163.com

Dear editor

We thank Hou et al for their interest in our paper entitled "Transcutaneous electrical acupoint stimulation for prevention of postoperative delirium in geriatric patients with silent lacunar infarction: a preliminary study" and their valuable comments. Here we respond to their concerns.

Hou et al found that baseline education level, intraoperative major blood loss and blood transfusion, and postoperative complications were identified as significant risk factors of postoperative delirium (POD) in elderly surgical patients in the literature. However, some other previous studies have had conflicting results that these factors were not related to POD.¹⁻³ Further studies are needed to confirm these imparities.

For avoidance of potential bias, we set strict exclusion criteria. We would be pleased if our study provokes interest in further research into whether transcutaneous electrical acupoint stimulation (TEAS) is an effective treatment in elderly patients with an increased risk of POD. It will be useful to take into consideration the severity and duration of POD, but unfortunately we did not collect data on these variables due to the short observation time, and we have stated this in the discussion.

Our data were analyzed by SPSS, and the results are as follows.

	Levene's test for equality of variances		t-test for equality of means						
	F	Sig.	t	df	Sig. (2-tailed)	Mean difference	Std. error difference	95% confidence interval of the difference	
								Lower	Upper
MMP93									
Equal variances assumed	.069	.793	-2.084	62	.041	-.34469	.16536	-.67525	-.01413
Equal variances not assumed			-2.084	61.225	.041	-.34469	.16536	-.67533	-.01405
S100β3									
Equal variances assumed	.033	.857	-2.177	62	.033	-6.96437	3.19911	-13.35930	-.56945
Equal variances not assumed			-2.177	61.860	.033	-6.96437	3.19911	-13.35959	-.56916
TNF2									
Equal variances assumed	.217	.643	-2.606	62	.011	-3.66750	1.40735	-6.48075	-.85425
Equal variances not assumed			-2.606	61.986	.011	-3.66750	1.40735	-6.48076	-.85424
TNF3									
Equal variances assumed	.078	.781	-2.954	62	.004	-4.95875	1.67863	-8.31428	-1.60322
Equal variances not assumed			-2.954	61.270	.004	-4.95875	1.67863	-8.31507	-1.60243

(Continued)

	Levene's test for equality of variances		t-test for equality of means						
	F	Sig.	t	df	Sig. (2-tailed)	Mean difference	Std. error difference	95% confidence interval of the difference	
								Lower	Upper
IL62									
Equal variances assumed	.024	.877	-2.617	62	.011	-4.27750	1.63449	-7.54481	-1.01019
Equal variances not assumed			-2.617	61.998	.011	-4.27750	1.63449	-7.54481	-1.01019
IL63									
Equal variances assumed	.736	.394	-2.073	62	.042	-3.32375	1.60304	-6.52819	-1.1931
Equal variances not assumed			-2.073	61.566	.042	-3.32375	1.60304	-6.52864	-1.1886

Abbreviations: MMP9, the concentration of matrix metallo proteinase-9 at T₂; s100β3, the concentration of s100β3 at T₂; TNF2, the concentration of tumor necrosis factor-α at T₂; TNF3, the concentration of tumor necrosis factor-α at T₂; IL62, the concentration of interleukin-6 at T₂; IL63, the concentration of interleukin-6 at T₂.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Kobayashi K, Imagama S, Ando K, et al. Risk Factors for delirium after spine surgery in extremely elderly patients aged 80 years or older and review of the literature: Japan Association of Spine Surgeons with Ambition Multicenter Study. *Global Spine J.* 2017;7(6):560–566.
2. Brown CH 4th, LaFlam A, Max L, et al. Delirium after spine surgery in older adults: incidence, risk factors, and outcomes. *J Am Geriatr Soc.* 2016;64(10):2101–2108.
3. Jiang X, Chen D, Lou Y, et al. Risk factors for postoperative delirium after spine surgery in middle- and old-aged patients. *Aging Clin Exp Res.* 2017;29(5):1039–1044.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Interventions in Aging 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Interventions in Aging editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Clinical Interventions in Aging

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine,

CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>

Dovepress