

Association between the extent of coronary artery disease and prognosis in patients with diabetes mellitus [Letter]

This article was published in the following Dove Press journal:
Clinical Epidemiology

Huan Xu
Hui-Ju Bu
Jing-Zhao He

Intensive Care Unit, Dongguan Donghua Hospital, Dongguan, Guangdong, People's Republic of China

Dear editor

The correlation between the extent of coronary artery disease (CAD) and clinical outcomes among patients with diabetes mellitus (DM) was unclear. Recently, we read with great interest the study by Gyldenkerne et al.¹ The authors conducted a cohort study of patients with DM who underwent coronary angiography and were stratified according to the extent of coronary artery disease: 0-, 1-, 2- or 3-vessel disease or diffuse vessel disease, with myocardial infarction, all-cause death, and major adverse cardiovascular events (MACE) as endpoints. They concluded that the extent of coronary artery disease is a significant risk factor for myocardial infarction and death in patients with DM. The research appears informative clinically. Thus, we addressed some issues regarding this study.

First, low-density lipoprotein cholesterol (LDL) plays a vital role in the perpetuation and pathogenesis of atherosclerotic CVD.² The increased level of LDL is associated with an elevated risk of CVD events and decreasing of LDL is associated with a reduction in the event.³ Nevertheless, although the percentage of patients using statin was similar between the three groups, the LDL level was not described in the baseline characteristics. The conclusion may not be rigorous without considering this vital factor.

Second, the HbA1c level at baseline and during follow-up was associated with the risk of CVD.⁴ For patients with CAD and DM, the HbA1c was a routine test during the follow-up time. Unfortunately, the baseline HbA1c was not reported in the study by Gyldenkerne et al.

Third, although the coronary angiography (CAG) could not describe the lesion morphology and plaque vulnerability in the vessel wall, it could identify the lesion location and the number of lesions. Indeed, the lesion location was related to the prognosis for patients with CAD. For instance, patients with critical proximal left anterior descending may go on to develop massive anterior wall myocardial infarction. Additionally, one vessel may have more than one lesions. Therefore, the association between the extent of coronary artery disease and outcomes should be interpreted cautiously without reporting the lesion location and the number of lesions.

Fourth, more pieces of evidence showed that the major adverse cardiac events were different between the male and female in patients with CVD.⁵ In this study, the percentage of the male was different between the three groups. Did the authors perform the subgroup analysis based on gender?

Correspondence: Jing-Zhao He
Department of Intensive Care Unit,
Dongguan Donghua Hospital, No. 1
Dongcheng East Road, Dongguan,
Guangdong, People's Republic of China
Tel/Fax +86 769 2282 3333
Email dgkhhejingzhao@yeah.net

Fifth, treatment compliance was poor in patients with CAD and DM because there are many oral drugs in these patients. In this situation, the comparison between the patients with CVD and those without CVD may be influenced by the treatment compliance.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Gyldenkerne C, Olesen KKW, Madsen M, et al. Extent of coronary artery disease is associated with myocardial infarction and mortality in patients with diabetes mellitus. *Clinical Epidemiology*. 2019;2019(11):419–428. doi:10.2147/CLEP.S200173
2. National Cholesterol Education Program Expert Panel on Detection E, Treatment of High Blood Cholesterol in A. Third report of the National Cholesterol Education Program (NCEP) expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III) final report. *Circulation*. 2002;106(25):3143–3421.
3. Ference BA, Ginsberg HN, Graham I, et al. Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. Evidence from genetic, epidemiologic, and clinical studies. A consensus statement from the European Atherosclerosis Society Consensus Panel. *Eur Heart J*. 2017;38(32):2459–2472. doi:10.1093/eurheartj/ehx144
4. Zhao W, Katzmarzyk PT, Horswell R, Wang Y, Johnson J, Hu G. HbA1c and coronary heart disease risk among diabetic patients. *Diabetes Care*. 2014;37(2):428–435. doi:10.2337/dc13-1525
5. Hess CN, McCoy LA, Duggirala HJ, et al. Sex-based differences in outcomes after percutaneous coronary intervention for acute myocardial infarction: a report from TRANSLATE-ACS. *J Am Heart Assoc*. 2014;3(1):e000523. doi:10.1161/JAHA.114.000844

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Epidemiology 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Epidemiology editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Clinical Epidemiology

Dovepress

Publish your work in this journal

Clinical Epidemiology is an international, peer-reviewed, open access, online journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification,

systematic reviews, risk & safety of medical interventions, epidemiology & biostatistical methods, and evaluation of guidelines, translational medicine, health policies & economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.

Submit your manuscript here: <https://www.dovepress.com/clinical-epidemiology-journal>