

Antimicrobial Susceptibility of *Mycobacterium abscessus* Complex Clinical Isolates from a Chinese Tertiary Hospital

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Yinjuan Guo^{1,2}
Xingwei Cao³
Jingyi Yu⁴
Qing Zhan³
Jinghui Yang¹
Xiaocui Wu¹
Baoshan Wan¹
Yin Liu¹
Fangyou Yu^{1,2}

¹Department of Laboratory Medicine, Shanghai Pulmonary Hospital, Tongji University School of Medicine, Shanghai 200082, People's Republic of China;

²Shanghai Key Laboratory of Tuberculosis, Shanghai Pulmonary Hospital, Tongji University School of Medicine, Shanghai 200082, People's Republic of China; ³Jiangxi Provincial Key Laboratory of Medicine, Clinical Laboratory of the Second Affiliated Hospital of Nanchang University, Nanchang 330000, People's Republic of China; ⁴Department of Laboratory Medicine, Wenzhou Medical University, Wenzhou 325000, People's Republic of China

Introduction: *Mycobacterium abscessus* complex (MABC) is a group of important infectious agents that are highly associated with drug resistance, and antibiotic treatment is usually ineffective. This study investigated the characteristics of antimicrobial susceptibility of MABC isolates and the synergy between certain β -lactam combinations against MABC infection.

Methods: We collected 129 MABC isolates from patients with lower respiratory tract infections and categorized them into three subspecies. The minimum inhibitory concentrations (MICs) of 15 antimicrobials for the MABC isolates were determined using commercial Sensititre RAPMYCOI MIC plates and the broth microdilution method, as recommended in the CLSI (M24-A2). In addition, the MICs of imipenem, alone and with ceftazidime and/or avibactam, were assessed in vitro for all isolates. The *erm(41)* and *rrl* genes were also sequenced.

Results: The MABC isolates exhibited >80% resistance to 11 of the 15 antimicrobials. Regarding the remaining four antimicrobials, the isolates were least resistant to tigecycline (12.4%) and amikacin (3.9%), and only partially resistant to two ceftazidime (39.5%) and imipenem (40.3%). Compared with *M. massiliense* isolates, *M. abscessus* and *M. bolletii* isolates were more resistant to amikacin and imipenem, whereas *M. abscessus* was significantly less resistant to tigecycline relative to *M. massiliense* and *M. bolletii* isolates. The clarithromycin inducible resistance rate was 68.4% and 74.3% among *M. bolletii* and *M. abscessus* isolates. Furthermore, 88.7% of the *M. abscessus* isolates carried a T at position 28 of *erm(41)*, which is associated with inducible clarithromycin resistance. In addition, compared to imipenem with avibactam only, the MIC₅₀ and MIC₉₀ values of imipenem after adding ceftazidime plus avibactam were decreased fourfold.

Conclusion: The antimicrobial resistance rates and the characteristics of the *erm(41)* gene associated with inducible clarithromycin resistance were different among the three MABC subspecies. There was also synergy between imipenem and 100 μ g/mL ceftazidime against MABC isolates.

Keywords: *Mycobacterium abscessus* complex, resistance, *erm(41)*, synergy, dual β -lactam therapy

Introduction

Mycobacterium abscessus complex (MABC) is a group of pathogens that account for 80% of rapidly growing mycobacteria (RGM) isolates. MABC strains are ubiquitous environmental bacteria often found in dust, soil, and water.¹ They often cause pulmonary infections (which can occur in patients with cystic fibrosis), complicated infections of the skin and soft tissues, and disseminated infections, leading to high mortality rates.² MABC strains can be classified into three clearly divergent subspecies:

Correspondence: Fangyou Yu
Email wzxyfy@163.com

M. abscessus subsp. *Abscessus*(*M. abscessus*), *M. abscessus* subsp. *Massiliense*(*M. massiliense*), and *M. abscessus* subsp. *bolletii* (*M. bolletii*) based on genomic analysis, which has dramatically increased our knowledge regarding MABC.³ The high rates of antimicrobial resistance of MABC strains can render even combination antibiotic treatment ineffective.⁴ Consequently, identifying novel treatment approaches is imperative.

Combination drug regimens, which can involve clarithromycin (CLA), amikacin (AMK), azithromycin (AZM), the cephalosporin (a type of β -lactam) cefoxitin (FOX), and the carbapenem (another type of β -lactam) imipenem (IPM), are recommended by the American Thoracic Society and the Infectious Disease Society of America, but less than half of patients with MABC infection can be cured with these treatments.^{5,6} One report indicated that *M. abscessus* and *M. bolletii* isolates often exhibit inducible resistance to CLA, while *M. massiliense* isolates were mostly susceptible.⁷ However, another study reported that *M. abscessus* is less responsive to CLA.⁸ Because of the high rates of resistance, treatment for MABC infections is often time-consuming and costly.

Recently, several molecular mechanisms underlying CLA resistance were identified. The primary innate mechanism underlying CLA resistance in MABC involves an increase in expression of the erythromycin ribosomal methylase gene, *erm*(41). Induction of *erm*(41) is mediated by the transcription factor *whiB7*,⁹ and rifabutin suppresses inducible CLA resistance by preventing induction of *whiB7* and *erm*(41) expression.¹⁰ Inducible resistance to CLA was identified in a sequevar with an intact *erm*(41) but with a single-nucleotide polymorphism (C to T) at position 28¹¹ (C at position 28 in *M. abscessus* results in CLA susceptibility). Notably, *M. massiliense* strains with a CLA susceptible phenotype have a nonfunctional *erm*(41) gene.⁷ Acquired high-level resistance to CLA can be caused by a point mutation (A2058G or A2059G) in the *rrl* gene (which encodes domain V of 23S rRNA).¹¹

In addition, there are some mechanisms of resistance of MABC to carbapenems. Resistance to β -lactams (such as the carbapenem IPM) was found to be related to the initial characterization of the MABC β -lactamase (Bla_{Mab}) and transpeptidases.^{12,13} The development of new antimicrobials is important to overcome the emergence of carbapenem resistance;¹⁴ currently, there are several active pharmaceutical programs developing β -lactamase inhibitors. Previous studies reported that the β -lactamase inhibitor avibactam (AVI) could effectively inhibit Bla_{Mab} and

thus improve the effects of β -lactams in MABC infection.^{13,15} Recent evidence suggests that dual β -lactam therapy may be more effective than using a single β -lactam.¹⁶

In this study, MABC isolates were assessed for susceptibility to various antimicrobials, and the relationship between the *erm*(41) gene and inducible CLA resistance in each subspecies of MABC was explored. The results provide guidance for the empirical therapy of RGM infections. Additionally, we assessed the individual and combined effects of two β -lactams (imipenem [IPM] and ceftazidime [CAZ]) and a β -lactamase inhibitor (avibactam [AVI]), and the results provide further insights into the treatment options.

Materials and Methods

Collection of MABC Isolates, Culture Conditions, and Informed Consent

Between 2014 and 2018, 129 MABC isolates were randomly isolated from the sputum and bronchoalveolar fluids of patients with clinical signs of lower respiratory tract infections in three tertiary hospitals (Shanghai Pulmonary Hospital, the Second Affiliated Hospital of Nanchang University, and the Second Affiliated Hospital of Suzhou University). All isolates were cultured in Middlebrook 7H10broth (BD, France) supplemented with 10% (vol/vol) oleic acid-albumin-dextrose-catalase (Thermo Fisher Scientific, USA). The cultures were incubated at 37°C for 7 days.

The Ethics Committee of Shanghai Pulmonary Hospital, Tongji University School of Medicine, exempted this study from ethical review because the assessment of the bacteria was part of routine hospital laboratory procedures. Verbal informed consent was obtained from all participants.

Identification of Subspecies and Detection of *erm*(41) and *rrl* Mutations

Genomic DNA was extracted from the MABC cultures for identification based on the sequences of the genes 16S rRNA, *rpoB*, and *hsp65*. The 16S rRNA gene was amplified using primers F (5'-AGAGTTTGATCCTGGCTCAG-3') and R (5'-ACGGGCGGTGTCTACAA-3'), as previously described.¹⁷ A 723-bp fragment of the *rpoB* gene was amplified using primers *rpoBF* (5'-GGCAAGGTCACCCGAAGGG-3') and *rpoBR* (5'-AGCGGCTGCTGGGTGATCATC-3'), as previously described.¹⁸ The *hsp65* gene was amplified using

primers *hsp65F* (5'-ACCAACGATGGTGTGCCAT-3') and *hsp65R* (5'-CTTGTCGAACCGCATACCCT-3'), as previously described.¹⁹ The *erm(41)* gene (F:5'-TGGTATCCGCTCACTGATGA-3' and R:5'-GCGGTGGATGATGGA AAG-3') and *rrl* gene (F: 5'-CCTGCACGAATGGCG TAACG-3' and R: 5'-CACCAGAGGTTTCGTCCGTC-3') were amplified as described by Maurer et al.²⁰ The Basic Local Alignment Search Tool (BLAST) program (<https://blast.ncbi.nlm.nih.gov>) was used for gene sequence comparisons.

Antibiotics

The carbapenem β -lactam imipenem (IPM; Sigma), the cephalosporin β -lactam ceftazidime (CAZ; Sigma), and the β -lactamase inhibitor avibactam (AVI; IHMA Inc.) were used alone or in combination. AVI was kindly provided by Prof. Chen Liang from Newark University (USA). Stock solutions of IPM and AVI were prepared in sterile water, while a stock solution of CAZ was prepared in dimethyl sulfoxide (DMSO). The stock concentrations of AVI, IPM, and CAZ were 1, 10, and 10 mg/mL, respectively. These stock solutions were diluted to the desired working concentrations with 7H9 medium.

Antimicrobial Susceptibility Tests

The susceptibility of all isolates to the following 15 antimicrobials (Thermo Fisher Scientific) were assessed according to the manufacturer's instructions: amikacin (AMK), ciprofloxacin (CIP), moxifloxacin (MXF), trimethoprim-sulfamethoxazole (SXT), linezolid (LZD), ceftriaxone (CRO), cefepime (FEP), ceftazidime (CAZ), ceftazidime-avibactam (CAZ-AVI), ceftazidime-avibactam (CAZ-AVI), tobramycin (TOB), tigecycline (TGC), minocycline (MIN), doxycycline (DOX), amoxicillin/clavulanic acid (AMC), imipenem (IPM) and clarithromycin (CLA). The minimum inhibitory concentrations (MICs) of the 15 antimicrobials for the 129 MABC isolates were determined using Sensititre RAPMYCOI MIC plates (Thermo Fisher Scientific) and the broth microdilution method, as recommended in the Clinical and Laboratory Standards Institute (CLSI)(M24-A2). The MIC ranges tested and CLSI M24-A2²¹ breakpoints for the 15 antimicrobials are listed in Table 2. We assessed the MICs of the 15 antimicrobials after 72 h of incubation, while we assessed the MICs of CLA after 3 and 14 days of incubation, in order to assess inducible CLA resistance. *M. abscessus* ATCC19,977 was used as the reference strain to compare the MICs.

Synergistic Antimicrobial Susceptibility Tests

The MIC values of IPM, CAZ, and AVI were also assessed, alone and in combination (involving 100 μ g/mL CAZ and 4 μ g/mL AVI). The MICs were determined for each of the 129 MABC isolates using the broth microdilution method in 96-well plates after incubation at 37°C for 48 h, according to CLSI (M24-A2) and Chen et al.²²

Statistical Analysis

Differences between groups were compared using Pearson's χ^2 test. A two-tailed $P < 0.05$ was considered statistically significant. All computations were performed using Graph Pad Prism (version 7.0, La Jolla, CA, USA).

Results

MABC Subspecies Identification

M. abscessus, *M. bolletii* and *M. massiliense* accounted for 75.2% (97/129), 14.7% (19/129), and 10.1% (13/129) of the MABC isolates.

Antimicrobial Susceptibility Profiles

The antimicrobial susceptibilities of the 129 MABC isolates of the three subspecies are summarized in Table 1. The drug sensitivity results of each isolates were provided in the [supplementary data](#) (the file called 16_Apr_2020_16_Apr_2020_data.xlsx). In general, the antimicrobial resistance rates were very high. The MABC isolates exhibited >90% resistance to nine of the 15 antimicrobials tested (FEP, CRO AMC, IMI, MIN, CIP, MXF, SXT, and TOB), and the rates of resistance to DOX and LZD were >70%. But the isolates were most susceptible to TGC (87.6%) and AMK (65.1%).

The rate of FOX resistance was significantly lower than the rates of resistance to the two other cephalosporins (FEP and CRO) ($P < 0.05$). The rates of FOX resistance for *M. abscessus*, *M. massiliense*, and *M. bolletii* were 34.0%, 46.2%, and 63.2%, respectively ($P < 0.05$). Additionally, the rate of IPM resistance was significantly lower for *M. massiliense* (23.1%) than *M. abscessus* (41.2%) and *M. bolletii* (47.4%) ($P < 0.05$).

CLA Susceptibility Testing and *erm* Genotyping in MABC Isolates

Out of all the MABC isolates, 72 (74.2%) were susceptible to CLA on day 3 but resistant on day 14 (indicating inducible resistance). Although the CLA resistance rate

Table 1 Antimicrobial Susceptibility of 129 *Mycobacterium abscessus* Complex (MABC) Isolates

Subspecies/Antimicrobial Agents	MIC50 MIC90 Range			Susceptibility % (n)			erm(41) Sequevars (n)
	(µg/mL)			R	I	S	
<i>Mycobacterium abscessus</i> subsp. <i>abscessus</i> (n=97)							
Aminoglycosides							
Amikacin	16	32	2->64	4.1	30.9	63.9	T(28) 86; C(28) 10
Tobramycin	16	>16	2->16	91.8	6.2	2.1	
Cephalosporins							Truncated(1)
Cefoxitin	64	128	16->128	34.0	56.7	9.3	
Cefepime ^a	>32	>32	32->32	100	0	0	
Ceftriaxone ^a	>64	>64	32->64	99	1	0	
Carbapenem							
Imipenem	16	64	2->64	41.2	45.4	13.4	
Fluoroquinolones							
Ciprofloxacin	>4	>4	2->4	93.8	6.2	0	
Moxifloxacin	>8	>8	0.5->8	94.9	3.1	2.1	
Folate pathway inhibitor							
SXT	>8	>8	1->8	94.9	-	5.1	
Tetracyclines							
Tigecycline	1	4	0.12->4	11.3	-	88.7	
Minocycline ^a	>8	>8	4->8	94.9	3.1	2.1	
Doxycycline	>16	>16	0.5->16	87.6	8.3	4.1	
Macrolide							
Clarithromycin - 3days	1	16	0.06->16	14.4	0	85.6	
Clarithromycin - 14days	16	>16	0.06->16	88.7	0	11.3	
Oxazolidinone							
Linezolid	>32	>32	2->32	86.6	8.3	5.2	
β-Lactam/β-lactamase inhibitor combinations							
AMC ^a	>64	>64	32->64	99	1	0	
<i>Mycobacterium abscessus</i> subsp. <i>massiliense</i> (n=13)							Truncated(13)
Aminoglycosides							rrl(1)
Amikacin	16	32	8-32	0	38.5	61.5	
Tobramycin	>16	>16	4->16	92.3	7.7	0	
Cephalosporins							
Cefoxitin	64	128	32-128	46.1	53.9	0	
Cefepime ^a	>32	>32	16->32	92.3	7.7	0	
Ceftriaxone ^a	>64	>64	64->64	100	0	0	
Carbapenem							
Imipenem	16	32	2-32	23.1	69.2	7.7	
Fluoroquinolones							
Ciprofloxacin	>4	>4	2->4	92.3	7.7	0	
Moxifloxacin	>8	>8	2->8	84.6	15.4	0	

(Continued)

Table 1 (Continued).

Subspecies/Antimicrobial Agents	MIC50 MIC90 Range			Susceptibility % (n)			<i>erm(41)</i> Sequences (n)
	(µg/mL)			R	I	S	
Folate pathway inhibitor SXT	>8	>8	1→8	84.6	-	15.4	
Tetracyclines							
Tigecycline	2	4	0.5–4	15.4	-	84.6	
Minocycline ^a	>8	>8	8→8	100	0	0	
Doxycycline	>16	>16	4→16	84.6	15.4	0	
Macrolide							
Clarithromycin - 3days	0.5	2	0.06→16	7.7	0	92.3	
Clarithromycin - 14days	0.5	2	0.06→16	7.7	0	92.3	
Oxazolidinones							
Linezolid	>32	>32	16→32	76.9	15.4	7.7	
β-Lactam/β-lactamase inhibitor combinations							
AMC ^a	>64	>64	16→64	92.3	0	7.7	
<i>Mycobacterium abscessus</i> subsp. <i>bolletii</i> (n=19)							No mutations
Aminoglycosides							
Amikacin	16	32	2–64	5.3	21	73.7	
Tobramycin	16	>16	4→16	89.5	10.5	0	
Cephalosporins							
Cefoxitin	128	128	32→128	63.2	36.8	0	
Cefepime ^a	>32	>32	32→32	100	0	0	
Ceftriaxone ^a	>64	>64	64→64	100	0	0	
Carbapenem							
Imipenem	16	32	2–32	47.4	31.6	21	
Fluoroquinolones							
Ciprofloxacin	>4	>4	4→4	100	0	0	
Moxifloxacin	>8	>8	2→8	94.8	5.2	0	
Folate pathway inhibitor							
SXT	>8	>8	2→8	94.5	-	5.3	
Tetracyclines							
Tigecycline	2	4	0.25–4	15.8	-	84.2	
Minocycline ^a	>8	>8	8→8	100	0	0	
Doxycycline	>16	>16	2→16	89.5	10.5	0	
Macrolide							
Clarithromycin - 3days	1	16	0.5→16	26.3	0	73.7	
Clarithromycin - 14days	16	>16	2→16	94.7	0	5.3	
Oxazolidinone							
Linezolid	>32	>32	8→32	73.7	5.3	21	
β-Lactam/β-lactamase inhibitor combinations							
AMC ^a	>64	>64	64→64	100	0	0	

Note: ^aDrugs out of guidelines.

Abbreviations: R, resistance; I, intermediate; S, susceptible; No mutations, There were no *erm* or *rrl* mutations in the *M. bolletii* isolates; SXT, trimethoprim/sulfamethoxazole; AMC, amoxicillin/clavulanic acid.

Table 2 Antimicrobial Concentration Ranges for Drug Susceptibility Testing and MIC Breakpoints of Antimicrobial Agents

Antimicrobial Agents	Tested Concentration	Breakpoints ($\mu\text{g/mL}$)		
	Ranges ($\mu\text{g/mL}$)	R	I	S
Cephalosporins				
Cefoxitin	4–128	≥ 128	32–64	≤ 16
Cefepime	1–32	≥ 32	16	≤ 8
Ceftriaxone	4–64	≥ 64	32	≤ 16
Carbapenem				
Imipenem	2–64	≥ 32	8–16	≤ 4
Aminoglycosides				
Amikacin	1–64	≥ 64	32	≤ 16
Tobramycin	1–16	≥ 8	4	≤ 2
Oxazolidinone				
Linezolid	1–32	≥ 32	16	≤ 8
Fluoroquinolones				
Ciprofloxacin	0.12–16	≥ 4	2	≤ 1
Moxifloxacin	0.25–8	≥ 4	2	≤ 1
Tetracyclines				
Tigecycline	0.015–4	≥ 4	–	< 4
Minocycline	43,838	≥ 8	2–4	≤ 1
Doxycycline	0.12–16	≥ 8	2–4	≤ 1
Macrolide				
Clarithromycin	0.06–16	≥ 8	4	≤ 2
Folate pathway inhibitors				
SXT	0.25/4.75–8/152	$\geq 4/76$	–	$\leq 2/38$
β-Lactam/β-lactamase inhibitor combinations				
AMC	2/164/32	≥ 64	32	≤ 16

Abbreviations: R, resistance; I, intermediate; S, susceptible.

was higher on day 14 than on day 3 for *M. abscessus* (88.7% versus 14.4%), the corresponding rates were similar for *M. massiliense* (7.7% versus 7.7%), with none of the *M. massiliense* isolates exhibiting inducible resistance. Of the 19 *M. bolletii* isolates studied, 68.4% had inducible resistance, which could not be attributed to the presence of the *erm*(41) T28 polymorphism. In contrast, the majority of *M. abscessus* isolates (86 isolates) belonged to the *erm*(41) T28 sequevar. Of the 129 isolates, 24 (18.6%) were susceptible to CLA on day 14. Among these, 14 (10.9%) had an *erm*(41) deletion

(comprising all 13 *M. massiliense* isolates and one *M. abscessus* isolates) and the remaining 10 (7.8%) were *M. abscessus* isolates with the *erm*(41) C28 mutation. Among the 129 MABC isolates, an *rrl* point mutation was found in only one isolate (an *M. massiliense* isolate). There were no *erm*(41) or *rrl* mutations in the *M. bolletii* isolates.

Susceptibility of MABC Isolates to β -Lactams (IPM and CAZ) and a β -Lactamase Inhibitor (AVI)

All 129 MABC isolates underwent susceptibility testing regarding IPM, CAZ, and AVI used alone or in combination. The average MIC₅₀ and MIC₉₀ values are shown in Table 3. The drug sensitivity results of each isolates were provided in the [supplementary data](#) (the file called 16_Apr_2020_16_Apr_2020_data.xlsx). The MIC values for CAZ alone ranged from 128 to >1024 $\mu\text{g/mL}$, with average MIC₉₀ and MIC₅₀ values of 1024 and 512 $\mu\text{g/mL}$, respectively; after adding AVI, the MIC values for CAZ did not change. The MIC values for IPM alone ranged from 2 to >64 $\mu\text{g/mL}$, with average MIC₉₀ and MIC₅₀ values of 32 and 16 $\mu\text{g/mL}$, respectively; there were also no significant changes when IPM was combined with AVI. However, after adding CAZ (100 $\mu\text{g/mL}$) only to IPM, there was a 4-fold reduction in the MIC₅₀ and MIC₉₀ values for IPM compared to when IPM was used alone. Additionally, after adding both CAZ (100 $\mu\text{g/mL}$) and AVI (4 $\mu\text{g/mL}$), the MIC range of IPM for the MABC isolates decreased from 2–>64 to 0.5–16 $\mu\text{g/mL}$; there was a 4-fold reduction in the MIC₅₀ and MIC₉₀ values compared to when IPM was used with AVI only, which also occurred in the reference strain (*M. abscessus* ATCC 19977).

Discussion

In this study, we assessed the antimicrobial susceptibility of 129 MABC strains belonging to three subspecies and examined the association between the *erm*(41) gene and inducible CLA resistance. The MABC subspecies exhibited varied resistance rates to antimicrobials. When the antimicrobial resistance rates obtained in the present study were compared with the results of previous studies, we found that there were high rates of resistance to multiple antimicrobials among the three MABC subspecies. The resistance rates regarding fluoroquinolones (CIP and MXF) and other broad-spectrum antibiotics (such as DOX, SXT, and MIN) were similar (all >70%) to those

Table 3 Antimicrobial Activities Imipenem (IPM), Ceftazidime (CAZ), and Avibactam (AVI) Alone or in Combination, Against 129 *Mycobacterium abscessus* Complex (MABC) Isolates

Strain	MIC ($\mu\text{g/mL}$)					
	CAZ	CAZ+ AVI4	IPM	IPM+ AVI4	IPM + CAZ100	IPM+ CAZ100+ AVI4
Avg MIC ₅₀	512	512	16	16	4	4
Avg MIC ₉₀	1024	1024	32	32	8	8
<i>M. abscessus</i> ATCC 19,977	512	512	8	4	1	1

Abbreviations: Avg, average; CAZ, ceftazidime; AVI, avibactam; IPM, imipenem; CAZ100, 100 $\mu\text{g/mL}$ ceftazidim; AVI4, 4 $\mu\text{g/mL}$ avibactam.

previously reported for MABC isolates in Taiwan Province of China,²³ as well as in Japan,²⁴ Thailand,²⁵ the USA,²⁶ and France.²⁷ In contrast, the rate of IPM resistance varied significantly among different studies. It was 40.9% in our study, which was in agreement with the rates reported in Beijing²⁸ and Japan.²⁴ However, studies conducted in 2017 in Taiwan (61%),²³ 2015 in Australia (68%),²⁹ and 2018 in Shanghai (65%)³⁰ reported higher rates of IPM resistance. Notably, the *M. abscessus* isolates in our study had a higher resistance rate to FOX (34.02%) than the *M. abscessus* strains in a study in Japan (16.7%).³¹ In Korea³² and Japan,³³ the TOB resistance rate was 30–32%, while it was 90% in our study. These variations may be due to differences among the diverse studies in patient treatment histories or in the isolates.

In the early 2000s, when LZD was first used clinically, it was reported to be active against many species of RGM.³⁴ However, we found that the rate of resistance to LZD was high, similar to rates reported in Taiwan (70%)²³ and the UK (96%),³⁵ but higher than the 5% rate reported by a study in Korea.³⁶ Notably, there have been few reports of a rate of LZD resistance as high as that found in *M. bolletii* in our study (73.7%). With the exception of the resistance rate for LZD, the resistance rates of *M. bolletii* were higher than those of *M. abscessus* and *M. massiliense*. Therefore, precise differentiation between these subspecies is important for clinical purposes.

Bastian et al reported that MABC infections were usually poorly responsive to CLA because of acquired and/or inducible resistance.³⁷ Mutations in the *rhl* gene confer acquired CLA resistance, while a single-nucleotide polymorphism (T28) in *erm(41)* at position 28 leads to inducible CLA resistance.⁷ Notably, the acquired CLA resistance rate (on day 3) in the *M. abscessus* isolates of 14.4% (14/97) was similar to the rate reported in South Korea (15.84%), higher than those reported in France (9.09%) and the USA (2.51%),

and lower than that reported in China (33.95%).^{27,38,39} We hypothesize that the geographic diversity in the population structure of MABC may be a major reason why researchers from various regions observed contradictory results. In addition, the high rate of inducible resistance that we identified is consistent with those reported by other studies.^{38,40} This supports the need to modify the CLA susceptibility test recommended by the CLSI,²¹ by assessing the MIC of CLA after an additional longer period of incubation (eg, 14 days).

In our study, most of the mutations identified involved the *erm(41)* gene (75.1%) rather than the *rhl* gene (0.8%). We concluded that the *erm(41)* mutations may be associated with the treatment failure that occurs in many cases of MABC infection. Nash et al and Kim et al studied the molecular profiles of MABC isolates and found that the *erm(41)* C28 polymorphism was related to susceptibility.^{7,41} Of our 129 isolates, 23 (17.8%) were susceptible to CLA. Among these, 14 (10.9%) had an *erm(41)* deletion (comprising all 13 *M. massiliense* isolates and one *M. abscessus* isolate) and 10 (7.8%) were *M. abscessus* isolates with the *erm(41)* C28 mutation. Of the 19 *M. bolletii* isolates studied, 68.4% exhibited inducible CLA resistance, which could not be attributed to the presence of the *erm(41)* T28 polymorphism. This indicates that there is another mechanism of resistance to CLA in *M. bolletii*. In addition, it has been reported that acquisition of CLA resistance is 100% mediated by structural 50S ribosomal subunit mutations for the *M. abscessus* *erm(41)* C28 sequevar and *M. massiliense*, whereas it is less common for the *M. abscessus* *erm(41)* T28 sequevar and *M. bolletii* (other mechanisms may be responsible for CLA resistance in these strains).⁴² Hence, we hypothesize that detecting an *erm(41)* deletion may differentiate *M. massiliense* from *M. abscessus* and *M. bolletii*, but more extensive research is needed to accurately define the reliability of using *erm(41)* deletions to identify *M. massiliense*.

Given the increasing prevalence of multidrug-resistant MABC infections, the development of novel treatment regimens is imperative. Although knowledge regarding the efficacy of β -lactams and β -lactamase inhibitors in MABC remains limited, several studies have used the available data to develop synergistic treatment regimens for treating MABC infections.^{22,43} In this study, we evaluated the in vitro susceptibility of the 129 MABC isolates to IPM in the presence of CAZ and/or AVI. The MIC₅₀ and MIC₉₀ values of IPM after the addition of CAZ plus AVI, compared to the values after the addition of AVI only, decreased 4-fold to 4 and 8 μ g/mL, respectively. Similarly, the addition of 100 μ g/mL CAZ only led to 4-fold decreases in the MIC₅₀ and MIC₉₀ values of IPM (while there were no significant changes in the MIC of IPM when IPM was combined with 4 μ g/mL AVI only). The initial resistance or intermediate resistance to IPM changed to sensitivity. Interestingly, CAZ alone had poor activity against MABC but combining IPM and CAZ led to effective activity, indicating that the effect of triple therapy may be driven primarily by CAZ rather than AVI. In contrast, Lefebvre et al.¹³ reported that inhibition of the β -lactamase Bla_{Mab} by AVI improves the in vitro effects of IPM against MABC and Pandey et al.²² reported that 4 μ g/mL AVI enhanced the bactericidal activity of the β -lactam ceftazidime; it can be deduced that the β -lactamase of MABC strains probably hydrolyzes IPM (though this was not confirmed by our study). Future studies will need to identify effective measures for reducing exposure to these difficult-to-treat pathogens.

Conclusion

The MABC isolates exhibited varied resistance rates to antimicrobials. The antimicrobial susceptibility profile and the characteristics of the *erm*(41) gene associated with inducible CLA resistance were different among the three MABC subspecies. Additionally, there was synergy between the β -lactam IPM and 100 μ g/mL of the β -lactam CAZ against MABC infection.

Ethics Statement

As the assessment of the *Mycobacterium abscessus* complex (MABC) isolates in this study was part of routine hospital laboratory procedures, the study was exempted from ethics review by the Ethics Committee of Shanghai Pulmonary Hospital, Tongji University School of Medicine.

Acknowledgments

We thank Prof. Liang Chen (Newark University, USA) for providing the avibactam used in this study.

Author Contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Disclosure

The authors declare that they have no conflicts of interest.

References

- Griffith DE, Girard WM, Wallace RJ Jr. Clinical features of pulmonary disease caused by rapidly growing mycobacteria. An analysis of 154 patients. *Am Rev Respir Dis*. 1993;147:1271–1278. doi:10.1164/ajrccm/147.5.1271
- Lee MR, et al. Mycobacterium abscessus complex infections in humans. *Emerg Infect Dis*. 2015;21:1638–1646. doi:10.3201/2109.141634
- Leao SC, Tortoli E, Euzeby JP, Garcia MJ. Proposal that *Mycobacterium massiliense* and *Mycobacterium bolletii* be united and reclassified as *Mycobacterium abscessus* subsp. *bolletii* comb. nov., designation of *Mycobacterium abscessus* subsp. *abscessus* subsp. nov. and emended description of *Mycobacterium abscessus*. *Int J Syst Evol Microbiol*. 2011;61:2311–2313. doi:10.1099/ijs.0.023770-0
- Jarand J, Levin A, Zhang L, et al. Clinical and microbiologic outcomes in patients receiving treatment for *Mycobacterium abscessus* pulmonary disease. *Clin Infect Dis*. 2011;52:565–571. doi:10.1093/cid/ciq237
- Stout JE, Koh WJ, Yew WW. Update on pulmonary disease due to non-tuberculous mycobacteria. *Int J Infect Dis*. 2016;45:123–134. doi:10.1016/j.ijid.2016.03.006
- Griffith DE, et al. An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases. *Am J Respir Crit Care Med*. 2007;175:367–416. doi:10.1164/rccm.200604-571ST
- Nash KA, Brown-Elliott BA, Wallace RJ Jr. A novel gene, *erm*(41), confers inducible macrolide resistance to clinical isolates of *Mycobacterium abscessus* but is absent from *Mycobacterium chelonae*. *Antimicrob Agents Chemother*. 2009;53:1367–1376. doi:10.1128/AAC.01275-08
- Nessar R, Cambau E, Reytrat JM, Murray A, Gicquel B. *Mycobacterium abscessus*: a new antibiotic nightmare. *J Antimicrob Chemother*. 2012;67:810–818. doi:10.1093/jac/dkr578
- Richard M, Gutierrez AV, Kremer L. Dissecting *erm*(41)-mediated macrolide-inducible resistance in *Mycobacterium abscessus*. *Antimicrob Agents Chemother*. 2020;64. doi:10.1128/AAC.01879-19.
- Aziz DB, Go ML, Dick T. Rifabutin suppresses inducible clarithromycin resistance in *Mycobacterium abscessus* by blocking induction of *whiB7* and *erm41*. *Antibiotics*. 2020;9. doi:10.3390/antibiotics9020072.
- Wallace RJ Jr. et al. Genetic basis for clarithromycin resistance among isolates of *Mycobacterium chelonae* and *Mycobacterium abscessus*. *Antimicrob Agents Chemother*. 1996;40:1676–1681. doi:10.1128/AAC.40.7.1676

12. Soroka D, et al. Characterization of broad-spectrum Mycobacterium abscessus class A beta-lactamase. *J Antimicrob Chemother.* 2014;69:691–696. doi:10.1093/jac/dkt410
13. Lefebvre AL, et al. Inhibition of the beta-Lactamase BlaMab by Avibactam improves the in vitro and in vivo efficacy of imipenem against Mycobacterium abscessus. *Antimicrob Agents Chemother.* 2017;61. doi:10.1128/AAC.02440-16
14. Rahme C, Butterfield JM, Nicasio AM, Lodise TP. Dual beta-lactam therapy for serious Gram-negative infections: is it time to revisit? *Diagn Microbiol Infect Dis.* 2014;80:239–259. doi:10.1016/j.diagmicrobio.2014.07.007
15. Soroka D, et al. Inhibition of beta-lactamases of mycobacteria by avibactam and clavulanate. *J Antimicrob Chemother.* 2017;72:1081–1088. doi:10.1093/jac/dkw546
16. Story-Roller E, Maggioncalda EC, Cohen KA, Lamichhane G. Mycobacterium abscessus and beta-Lactams: emerging insights and potential opportunities. *Front Microbiol.* 2018;9:2273. doi:10.3389/fmicb.2018.02273
17. Yu XL, et al. Identification and characterization of non-tuberculous mycobacteria isolated from tuberculosis suspects in Southern-central China. *PLoS One.* 2014;9:e114353. doi:10.1371/journal.pone.0114353
18. Macheras E, et al. Inaccuracy of single-target sequencing for discriminating species of the Mycobacterium abscessus group. *J Clin Microbiol.* 2009;47:2596–2600. doi:10.1128/JCM.00037-09
19. Nie W, Duan H, Huang H, Lu Y, Chu N. Species identification and clarithromycin susceptibility testing of 278 clinical nontuberculosis Mycobacteria Isolates. *Biomed Res Int.* 2015;506598. doi:10.1155/2015/506598
20. Maurer FP, Ruegger V, Ritter C, Bloemberg GV, Bottger EC. Acquisition of clarithromycin resistance mutations in the 23S rRNA gene of Mycobacterium abscessus in the presence of inducible erm (41). *J Antimicrob Chemother.* 2012;67:2606–2611. doi:10.1093/jac/dks279
21. Institute CA, L S. Susceptibility testing of Mycobacteria, Nocardiae, and other aerobic actinomycetes second edition: approved standard M24-A2. *CLSI.* 2011. Wayne, PA, USA.
22. Pandey R, et al. Dual beta-Lactam combinations highly active against Mycobacterium abscessus Complex In Vitro. *mBio.* 2019;10. doi:10.1128/mBio.02895-18
23. Lee MC, et al. Antimicrobial resistance in Mycobacterium abscessus complex isolated from patients with skin and soft tissue infections at a tertiary teaching hospital in Taiwan. *J Antimicrob Chemother.* 2017;72:2782–2786. doi:10.1093/jac/dkx212
24. Kusuki M, et al. Determination of the antimicrobial susceptibility and molecular profile of clarithromycin resistance in the Mycobacterium abscessus complex in Japan by variable number tandem repeat analysis. *Diagn Microbiol Infect Dis.* 2018;91:256–259. doi:10.1016/j.diagmicrobio.2018.02.008
25. Ananta P, et al. Analysis of drug-susceptibility patterns and gene sequences associated with clarithromycin and amikacin resistance in serial Mycobacterium abscessus isolates from clinical specimens from Northeast Thailand. *PLoS One.* 2018;13:e0208053. doi:10.1371/journal.pone.0208053
26. Brown-Elliott BA, Wallace R, J Jr. Clinical and taxonomic status of pathogenic nonpigmented or late-pigmenting rapidly growing mycobacteria. *Clin Microbiol Rev.* 2002;15:716–746. doi:10.1128/cmr.15.4.716-746.2002
27. Mougari F, et al. Standardized interpretation of antibiotic susceptibility testing and resistance genotyping for Mycobacterium abscessus with regard to subspecies and erm41 sequevar. *J Antimicrob Chemother.* 2016;71:2208–2212. doi:10.1093/jac/dkw130
28. Li YM, et al. Prevalence and antimicrobial susceptibility of Mycobacterium abscessus in a general hospital, China. *Biomed Environ Sci.* 2016;29:85–90. doi:10.3967/bes2016.009
29. Chua KY, Bustamante A, Jelfs P, Chen SC, Sintchenko V. Antibiotic susceptibility of diverse Mycobacterium abscessus complex strains in New South Wales, Australia. *Pathology.* 2015;47:678–682. doi:10.1097/PAT.0000000000000327
30. Shen Y, et al. In vitro susceptibility of Mycobacterium abscessus and Mycobacterium fortuitum isolates to 30 antibiotics. *Biomed Res Int.* 2018;4902941. doi:10.1155/2018/4902941.
31. Aono A, et al. Antimicrobial susceptibility testing of Mycobacteroides (Mycobacterium) abscessus complex, Mycolicibacterium (Mycobacterium) fortuitum, and Mycobacteroides (Mycobacterium) chelonae. *J Infect Chemother.* 2019;25:117–123. doi:10.1016/j.jiac.2018.10.010
32. Park S, et al. In vitro antimicrobial susceptibility of Mycobacterium abscessus in Korea. *J Korean Med Sci.* 2008;23:49–52. doi:10.3346/jkms.2008.23.1.49
33. Yoshida S, et al. Further isolation of Mycobacterium abscessus subsp. abscessus and subsp. bolletii in different regions of Japan and susceptibility of these isolates to antimicrobial agents. *Int J Antimicrob Agents.* 2013;42:226–231. doi:10.1016/j.ijantimicag.2013.04.029
34. Wallace RJ Jr, Brown-Elliott BA, Ward SC, et al. Activities of linezolid against rapidly growing mycobacteria. *Antimicrob Agents Chemother.* 2001;45:764–767. doi:10.1128/AAC.45.3.764-767.2001
35. Cowman S, Burns K, Benson S, Wilson R, Loebinger MR. The antimicrobial susceptibility of non-tuberculous mycobacteria. *J Infect.* 2016;72:324–331. doi:10.1016/j.jinf.2015.12.007
36. Park J, Cho J, Lee CH, Han SK, Yim JJ. Progression and Treatment outcomes of lung disease caused by mycobacterium abscessus and Mycobacterium massiliense. *Clin Infect Dis.* 2017;64:301–308. doi:10.1093/cid/ciw723
37. Bastian S, et al. Assessment of clarithromycin susceptibility in strains belonging to the Mycobacterium abscessus group by erm(41) and rrl sequencing. *Antimicrob Agents Chemother.* 2011;55:775–781. doi:10.1128/AAC.00861-10
38. Lee SH, et al. The drug resistance profile of Mycobacterium abscessus group strains from Korea. *Ann Lab Med.* 2014;34:31–37. doi:10.3343/alm.2014.34.1.31
39. Li B, et al. Relationship between antibiotic susceptibility and genotype in Mycobacterium abscessus clinical isolates. *Front Microbiol.* 2017;8:1739. doi:10.3389/fmicb.2017.01739
40. Hanson KE, Slechta ES, Muir H, Barker AP. Rapid molecular detection of inducible macrolide resistance in Mycobacterium chelonae and M. abscessus strains: a replacement for 14-day susceptibility testing? *J Clin Microbiol.* 2014;52:1705–1707. doi:10.1128/JCM.03464-13
41. Kim HY, et al. Mycobacterium massiliense is differentiated from Mycobacterium abscessus and Mycobacterium bolletii by erythromycin ribosome methyltransferase gene (erm) and clarithromycin susceptibility patterns. *Microbiol Immunol.* 2010;54:347–353. doi:10.1111/j.1348-0421.2010.00221.x
42. Mougari F, et al. Selection of resistance to clarithromycin in Mycobacterium abscessus subspecies. *Antimicrob Agents Chemother.* 2017;61. doi:10.1128/AAC.00943-16
43. Kaushik A, Ammerman NC, Parrish NM, Nuermberger EL. New beta-Lactamase inhibitors nacubactam and zidebactam improve the in vitro activity of beta-Lactam antibiotics against Mycobacterium abscessus complex clinical isolates. *Antimicrob Agents Chemother.* 2019;63. doi:10.1128/AAC.00733-19.

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>