


Anti-Diabetic Effect of Telmisartan Through its Partial PPAR γ -Agonistic Activity

This article was published in the following Dove Press journal:
Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Muluken Altaye Ayza 
Kaleab Alemayehu Zewdie 
Bekalu Amare Tesfaye 
Selamawit Tesfamariam
Gebrekirstos 
Derbew Fikadu Berhe 

Department of Pharmacology and
Toxicology, School of Pharmacy, Mekelle
University, Mekelle, Ethiopia

Abstract: Telmisartan is an angiotensin II receptor antagonist, which selectively inhibits the angiotensin II type 1 receptor. Thus, it is widely used for hypertension management. Nowadays, telmisartan's effect on peroxisome proliferator-activated receptors (PPARs) is gaining wider attention. PPARs are ligand-activated transcription factors that belong to the nuclear hormone receptor superfamily. Telmisartan is reported to have a partial PPAR γ -agonistic effect while avoiding the safety concerns found with full PPAR γ agonists (thiazolidinediones). Telmisartan could be an alternative treatment option, with dual benefit for diabetes mellitus (DM) and hypertension. This review summarizes the anti-diabetic activity of telmisartan via its partial PPAR γ -agonistic activity.

Keywords: diabetes mellitus, hypertension, peroxisome proliferator-activated receptors, telmisartan

Introduction Telmisartan

Telmisartan is a selective angiotensin II type 1 receptor antagonist (angiotensin receptor blocker [ARB]) which does not affect the other receptor systems involved in cardiovascular regulation.^{1,2} It is a more lipophilic compound than other ARBs, which facilitates oral absorption and benefits tissue and cell penetration.³ The drug is used for the management of hypertension, either as monotherapy or as combination therapy with other antihypertensive agents.^{1,4} It is considered as a first-line drug in mild-to-moderate hypertension with an excellent safety profile, and is used for the treatment of hypertension-related cardiovascular end-organ damage.⁵ A number of studies have shown that telmisartan has partial PPAR γ -agonist activity, activating 25–30% of the receptor compared to the full PPAR γ agonists.^{6–8}

Peroxisome Proliferator-Activated Receptors (PPARs)

PPARs are ligand-activated transcription factors that belong to the nuclear hormone receptor superfamily.⁹ PPARs comprise three subtypes, namely PPAR α , PPAR γ , and PPAR β/δ .¹⁰

Among the three subtypes, PPAR γ is the most studied nuclear receptor and is involved in the control of energy balance, glucose and lipid homeostasis.¹¹ Thiazolidinediones, including rosiglitazone and pioglitazone, are anti-diabetic drugs acting on PPAR γ .¹² Thiazolidinediones promote insulin sensitization and improve dyslipidemia in diabetic patients.¹³ It has been reported that PPAR γ agonists may have various therapeutic advantages in addition to anti-diabetic activity. Rosiglitazone decreased arterial stiffness and clinical inflammatory responses in diabetic patients with coronary artery disease.^{14,15}

Correspondence: Muluken Altaye Ayza
Tel +251-916707913
Email Mulealto1@gmail.com

Despite the beneficial effects of thiazolidinediones on the heart, they are also associated with several adverse effects, including cardiac dysfunction and ventricular fibrillation.^{16,17}

Challenges with Full-PPAR γ Agonists

Full PPAR γ agonists are associated with a number of side effects, including weight gain, fluid retention (edema), and cardiac toxicity.¹⁸ The first approved drug from the thiazolidinedione class was troglitazone, which was withdrawn from the market in 2000 owing to its fatal hepatotoxicity.¹⁹ This idiosyncratic hepatotoxicity is mediated through oxidative stress and involves mitochondrial dysfunction.^{20,21} Other thiazolidinediones are also associated with adverse effects, including weight gain (adiposity), increased rate of bone fracture, anemia, and hepatic damage.²²

Rosiglitazone is associated with granulomatous hepatitis.^{23,24} It has also been reported to increase cardiovascular risk.²⁵ Hypertension, myocardial ischemia, heart failure, upper respiratory tract infection, and diarrhea were also reported with rosiglitazone use.²⁶ Owing to its cardiovascular side effects and adverse effects on the lipid profile, rosiglitazone was banned in India in 2010.²⁷

Clinical use of pioglitazone is also limited by undesirable adverse effects, including weight gain, precipitation of heart failure, peripheral edema, and an increase in bone fractures.^{28,29} Pioglitazone is also reported to induce hepatic damage.^{30,31} Although the severity depends on the dose administered, subchronic administration of pioglitazone in mice resulted in nephrotoxicity, hepatotoxicity, cardiotoxicity, and hematological disorders, which were supported by biochemical abnormalities and histopathological changes.³² For this reason, new PPAR γ ligands with increased therapeutic efficacy and reduced adverse effects are needed. Partial PPAR γ modulators are one of the promising classes in this regard.³³

Advantages of Partial PPAR γ Agonists

PPAR γ partial agonists selectively modify the expression of genes needed only for insulin sensitization without activating the genes responsible for weight gain and edema.³⁴ This makes partial PPAR γ agonists preferable to full agonists.³⁵ Adverse effects including weight gain and edema may lead to heart failure, and other adverse effects may also be avoided by selective modulation of the PPAR γ receptor.³⁶

The partial PPAR γ agonist balaglitazone showed fewer untoward effects, including fluid and fat accumulation, compared to the full PPAR γ agonist pioglitazone.³⁷ Another selective partial activator of PPAR γ (PAR-1622) showed anti-diabetic activity with a broader safety margin against fluid retention in comparison with rosiglitazone.³⁸ PAM-1616 is also a selective modulator of PPAR γ with anti-hyperglycemic properties and an improved side effect profile compared to rosiglitazone.³⁹ KR-62980 is a selective PPAR γ modulator with anti-diabetic and anti-adipogenic activity with an improved side effect profile.^{40,41} CMHX008, a partial PPAR γ modulator, showed comparable insulin-sensitizing effects with lower adipogenic capacity and lower risk of bone loss compared to rosiglitazone, and it could be an effective agent for the management of DM and other metabolic disorders.^{42,43} GQ-16 and L312, PPAR γ partial agonists, also showed a promising anti-diabetic effect with an improved side effect profile.^{44,45} There are several partial PPAR γ agonists under investigation, with an insulin sensitization action and fewer adverse effects, including a reduced tendency to cause weight gain, adipose tissue gain, and fluid retention.⁴⁶

PPAR γ partial activation may have an important endothelial defensive action through activating eNOS, generating nitric oxide (NO) and enhancing NO bioavailability, reducing oxidative stress, and preventing the adhesion cascade and vascular inflammation.⁴⁷ Partial PPAR γ agonists may offer beneficial effects for the management of type II DM (T2DM) with attenuated adverse effects such as fluid retention and cardiac hypertrophy.³⁴

Patients with hypertension and T2DM may obtain a dual benefit from drugs such as telmisartan with multiple targets (AT1R and PPAR γ). That would potentially reduce the pill burden and side effects of multiple medication use, and hence improve medication adherence. This review focuses on the anti-diabetic effects of telmisartan mediated through PPAR γ .

Telmisartan for Diabetes Mellitus

Current management of DM is associated with major hurdles, comprising polypharmacy, striving for glycemic targets, patient adherence, and clinical inertia.⁴⁸ In spite of many anti-diabetic drugs being available, adequate control of diabetic hyperglycemia often remains difficult to achieve.⁴⁹ In T2DM, intense control of glycemia generally requires the addition of insulin to oral agents, which is consistently accompanied by weight gain. Furthermore, increased hypoglycemia carries a risk of cerebral damage, mental disorders and stroke, cognitive damage,

hypertension, and renal disease.⁵⁰ The adverse effects associated with currently available drugs include GI adverse effects and lactic acidosis with metformin,^{51,52} weight gain and hypoglycemia with sulfonylureas, and fluid retention and weight gain with thiazolidinediones.⁵³ Since the available agents are associated with various adverse effects, it is important to discover an alternative agent that is more effective and safer to use in the treatment of T2DM.^{53,54}

As telmisartan is reported to have partial PPAR γ -agonistic effect, the drug can regulate glucose and lipid metabolism, and improve insulin resistance.⁵⁵ PPAR γ phosphorylation and its downstream gene expression may be regulated by telmisartan, promoting glucose uptake and acting as an insulin-sensitizing agent in adipocytes.⁵⁶ Telmisartan augments glucose transporter-4 protein expression and 2-deoxyglucose uptake in the basal and insulin-stimulated states of adipocytes.⁵⁷ It further increases glucose transporter-4 localization to the plasma membrane and enhances glucose uptake through PPAR γ in adipocytes. Thus, telmisartan could be an effective therapy for metabolic syndrome.⁵⁸

Different mechanisms have been postulated for the glucose-lowering potential of telmisartan. It may occur through the suppression of reactive oxygen species and an agonistic effect on PPAR γ .⁵⁹ Telmisartan decreased malondialdehyde and increased glutathione, catalase, and superoxide dismutase levels in STZ-induced type 2 diabetic rats, and prevented cardiovascular complications.⁶⁰ Furthermore, telmisartan was shown to increase serum adiponectin and improved both oxidative stress and insulin resistance in type 2 diabetic patients.⁶¹ Also, in diabetic patients with hypertension, telmisartan increased high-molecular-weight adiponectin levels and improved insulin resistance, through partial PPAR γ activation.⁶² Adiponectin, through binding to its receptors, mediates the insulin-sensitizing effect, which is important in DM patients.⁶³

In addition, interruption of the renin–angiotensin–aldosterone system (RAAS) may improve glycemic control in T2DM, since the RAAS is implicated in the development of T2DM.⁶⁴ Owing to the ability of telmisartan, partial activation of PPAR γ and ARB blockade may have additional importance not merely in the management of metabolic syndrome and prevention of T2DM but also in the prevention and treatment of atherosclerotic cardiovascular disease.⁶⁵

Telmisartan improves insulin sensitivity in hypertensive patients with insulin resistance or T2DM.⁶⁶ Telmisartan treatment of hypertensive patients with T2DM not only

improved blood pressure, glucose and lipid metabolism, but also improved endothelial function for the prevention of atherosclerosis.⁶⁷ Through activating PPAR γ , telmisartan reduces cardiac fibrosis and hypertrophy, thus preventing unfavorable cardiac remodeling.⁷ In addition to PPAR γ activation, telmisartan improves hyperglycemia-induced cardiac fibrosis through PPAR δ activation.^{68,69} Telmisartan may be important in preventing diabetic cardiomyopathy and it could be used to treat both hemodynamic and metabolic aberrations.^{70,71} These cardioprotective effects of telmisartan may be due to improvement of endothelial function, which is associated with PPAR γ -eNOS, oxidative stress, and the Rho-kinase pathway.⁷² Targeting both angiotensin type 1 receptor and PPAR γ may be important in treating hyperlipidemia, insulin resistance, hypertension, and stroke, and ultimately mitigating the burden of cerebrovascular and cardiovascular disease.⁷³

Patients with uncomplicated hypertension and mild dyslipidemia can be effectively treated with telmisartan.⁷⁴ Both animal and human studies showed improvements in the lipid profile with telmisartan treatment. In high-fat diet-induced dyslipidemic guinea pigs, telmisartan markedly decreased triglyceride and slightly increased high-density lipoprotein cholesterol levels.⁷⁵ In a prospective open-label study, telmisartan significantly increased high-density lipoprotein and decreased triglyceride and total cholesterol in hypertensive patients with dyslipidemia.⁷⁶ Telmisartan may have hepatic partial PPAR α agonist activity and it increases lipoprotein lipase expression via a PPAR α -dependent pathway, which could explain telmisartan's anti-dyslipidemic actions.^{77,78}

Telmisartan has the additional benefit of renoprotection in patients diagnosed with diabetes and hypertension. In a randomized, multicenter, double-blind, placebo-controlled trial, telmisartan reduced the transition from incipient to overt nephropathy and induced remission of albuminuria in Japanese patients with T2DM; telmisartan achieved superior renoprotection.⁷⁹ In addition, in a multicenter, prospective, double-blind, forced-titration, randomized study, telmisartan and ramipril both increased nitric oxide activity of the renal endothelium significantly, which in turn may support the preservation of cardiovascular and renal function in patients with T2DM.⁸⁰

A number of preclinical and clinical studies have shown a decrease in blood glucose and improvement in diabetic complications after treatment with telmisartan (Table 1).

Table 1 Effects of Telmisartan in Diabetes Mellitus and Associated Complications

Reference	Study Design	Model	Method and Intervention	Major Outcome(s)
81	Experimental study	Streptozotocin (STZ)-induced DM in Wistar rats	DM was induced by the administration of STZ. Animals were administered with telmisartan for 28 days	Capillary blood glucose level was lower at all intervals Telmisartan improved the lipid profile. Hypoglycemic activity and improved lipid profile were comparable with glibenclamide
82	Experimental study	STZ-induced DM in rats	Diabetic rats were treated with/without telmisartan for 2 weeks	Telmisartan decreased SGLT2 expression in the PCT Insulin resistance was ameliorated in diabetic rats
[83]	Experimental study	STZ-induced DM in rats	Diabetic rats were challenged with nicotinamide to induce diabetic nephropathy	The addition of telmisartan to vildagliptin demonstrated the best control over blood pressure, glycemia, and diabetic nephropathy markers, renal structural changes, and improvement of renal function
84	Experimental study	STZ-induced diabetic nephropathy in rats	After grouping, rats were treated with telmisartan (10 mg/kg), ramipril (5 mg/kg), or vehicle for 16 weeks	Telmisartan pretreatment reduced blood glucose level after 8 weeks Telmisartan significantly altered other parameters towards normal levels
85	Experimental study	HFFD + STZ-induced diabetes and insulin resistant	Animals were treated with telmisartan and/or metformin	Addition of telmisartan to metformin successfully restored serum glucose back to normal levels and improved serum total cholesterol (TC), triglycerides (TG), and adiponectin
86	Experimental study	STZ-induced DM	Diabetic rats were treated with telmisartan or gliclazide	Telmisartan or gliclazide produced beneficial effects on serum glucose, Hb, and HbA _{1c} . Restored tissue GSH and SOD with a fall in tissue thiobarbituric acid reactive substances
87	Clinical study	Five-year, double-blind, double-dummy, randomized study	250 patients with hypertension and early type 2 diabetic nephropathy were included to compare the effect of telmisartan with enalapril	Telmisartan was not inferior to enalapril in reducing the decline in GFR None of the patients developed serum creatinine >200 µmol/L, and none of them required dialysis
88	Clinical study	Randomized controlled trial	3488 adults at high risk for CVD, but free from diabetes, were involved	21.8% of participants treated with telmisartan and 22.4% of those on placebo developed diabetes Participants originally diagnosed with IFG and/or IGT were equally likely to regress to normoglycemia (26.9% vs 24.5%)

(Continued)

Table I (Continued).

Reference	Study Design	Model	Method and Intervention	Major Outcome(s)
89	Clinical study	Homeostatic model assessment–insulin resistance (HOMA-IR) in hypertensive subjects with MS	42 patients were involved in the study. Cytokines and metabolic parameters were analyzed before and after treatment with telmisartan	Treatment with telmisartan diminished 35% of HOMA-IR Telmisartan had more favorable effects on glucose homeostasis in patients with MS and low levels of serum cytokines
90	Clinical study	Randomized, parallel-group, open-label, crossover, prospective study	Patients were on 20 mg of lisinopril or 80 mg of telmisartan once a day for 24 weeks. Then, patients were randomized to continuing treatment with the respective monotherapy or with lisinopril plus telmisartan for a further 28 weeks	Dual blockade may provide a new approach to prevention of diabetic nephropathy in patients with type 2 diabetes, hypertension, and microalbuminuria
91	Clinical study	Randomized, placebo-controlled, double-blind, crossover study	20 volunteers with insulin resistance and abdominal overweight were involved	Compared to placebo, telmisartan resulted in a reduction in glucose area under the curve during intravenous glucose tolerance There was an increase in the insulinogenic index, indicating improved beta-cell function
92	Clinical study	Non-comparative, open-label study	34 patients with diabetic nephropathy were involved. Telmisartan (40 mg) was added to ACEI treatment and the patients were observed for 12 weeks	The addition of telmisartan resulted in a significant reduction in albuminuria
93	Clinical study	Randomized, placebo-controlled, double-blind, multicenter study	Patients with type 2 diabetes and incipient nephropathy were involved. Normotensive type 2 diabetic patients treated with telmisartan or placebo over 52 weeks	Patients treated with telmisartan showed lower transition rates from microalbuminuria to overt nephropathy In addition, more patients on telmisartan reverted to normoalbuminuria. Telmisartan prevented the progression of microalbuminuria in normotensive patients with type 2 diabetes
94	Clinical study	Randomized controlled comparative trial	82 individuals with DM and HTN were involved. They were on telmisartan or lisinopril or placebo control for 1 month	Significant improvement in both serum malondialdehyde (MDA) and total antioxidants status (TAS) parameters. Both lisinopril and telmisartan were able to subside oxidative stress in diabetic hypertensive patients
95	Clinical study	Double-blind, parallel-group, randomized study	Patients were treated with once-daily doses of telmisartan or losartan for 3 months	Telmisartan significantly reduced free plasma glucose, free plasma insulin, HOMA-IR, and HbA _{1c} . Plasma glucose and insulin were reduced during the oral glucose tolerance test by telmisartan

(Continued)

Table I (Continued).

Reference	Study Design	Model	Method and Intervention	Major Outcome(s)
96	Clinical trial	Double-blind, parallel-group, randomized study	65 overweight and obese patients with mild to moderate hypertension were involved. Patients were randomized to telmisartan or olmesartan for 3 months	Telmisartan improved blood pressure, glucose, insulin, HOMA-IR, and leptin in hypertensive diabetic patients

Abbreviations: ACEI, angiotensin-converting enzyme inhibitor; CVD, cardiovascular disease; DM, diabetes mellitus; GFR, glomerular filtration rate; GSH, glutathione; Hb, hemoglobin; HbA_{1c}, hemoglobin A_{1c}; HFFD, high-fat, high-fructose diet; HOMA-IR, homeostatic model assessment–insulin resistance; HTN, hypertension; IFG, impaired fasting glucose; IGT, impaired glucose tolerance; MS, metabolic syndrome; PCT, proximal convoluted tubule; SGLT2, sodium–glucose co-transporter-2; SOD, superoxide dismutase; STZ, streptozotocin; TC, total cholesterol; TG, triglyceride.

Conclusion

Telmisartan is a well-known anti-hypertensive drug, which is currently in clinical use. The partial PPAR γ -agonistic activity and angiotensin receptor blockade activity of telmisartan have been shown to have multiple clinical benefits, including anti-diabetic and cardiovascular effects. In addition, telmisartan is reported to have PPAR α and PPAR δ agonist activity. Telmisartan would be an ideal alternative dual-purpose medication for patients with T2DM, hypertension and other cardiovascular disorders. However, this requires further detailed understanding of the therapeutic efficacy and molecular mechanism of telmisartan and other partial PPAR γ -agonist drugs.

Abbreviations

ARB, angiotensin receptor blocker; DM, diabetes mellitus; GC, glibenclamide; GSH, glutathione; HbA_{1c}, hemoglobin A_{1c}; HOMA-IR, homeostatic model assessment–insulin resistance; MDA, malondialdehyde; PCT, proximal convoluted tubule; RAAS, renin–angiotensin–aldosterone system; SOD, superoxide dismutase; STZ, streptozotocin; T2DM, type 2 diabetes mellitus; TAS, total antioxidant status.

Author Contributions

All authors contributed to data analysis, drafting or revising the article, have agreed on the journal to which the article will be submitted, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Wiene W, Entzeroth M, van Meel JC, et al. A review on telmisartan: a novel, long-acting angiotensin II-receptor antagonist. *Cardiovasc Drug Rev.* 2000;18(2):127–154. doi:10.1111/j.1527-3466.2000.tb00039.x
2. Sharpe M, Jarvis B, Goa KL. Telmisartan. *Drugs.* 2001;61(10):1501–1529. doi:10.2165/00003495-200161100-00009
3. Galzerano D, Capogrosso C, Di Michele S, et al. New standards in hypertension and cardiovascular risk management: focus on telmisartan. *Vasc Health Risk Manag.* 2010;6:113–133. doi:10.2147/VHRM.S7857
4. Battershill AJ, Scott LJ. Telmisartan. *Drugs.* 2006;66(1):51–83. doi:10.2165/00003495-200666010-00004
5. Borém LMA, Neto JFR, Brandi IV, Lelis DF, Santos SHS. The role of the angiotensin II type I receptor blocker telmisartan in the treatment of non-alcoholic fatty liver disease: a brief review. *Hypertens Res.* 2018;41(6):394–405. doi:10.1038/s41440-018-0040-6
6. Haraguchi T, Iwasaki K, Takasaki K, et al. Telmisartan, a partial agonist of peroxisome proliferator-activated receptor γ , improves impairment of spatial memory and hippocampal apoptosis in rats treated with repeated cerebral ischemia. *Brain Res.* 2010;1353:125–132. doi:10.1016/j.brainres.2010.07.017
7. Maejima Y, Okada H, Haraguchi G, et al. Telmisartan, a unique ARB, improves left ventricular remodeling of infarcted heart by activating PPAR gamma. *Lab Invest.* 2011;91(6):932–944. doi:10.1038/labinvest.2011.45
8. Amano Y, Yamaguchi T, Ohno K, et al. Structural basis for telmisartan-mediated partial activation of PPAR gamma. *Hypertens Res.* 2012;35(7):715–719. doi:10.1038/hr.2012.17
9. Rakhshandehroo M, Knoch B, Müller M, Kersten S. Peroxisome proliferator-activated receptor alpha target genes. *PPAR Res.* 2010;2010:612089. doi:10.1155/2010/612089
10. Tyagi S, Gupta P, Saini AS, Kaushal C, Sharma S. The peroxisome proliferator-activated receptor: A family of nuclear receptors role in various diseases. *J Adv Pharm Technol Res.* 2011;2(4):236. doi:10.4103/2231-4040.90879
11. Zieleniak A, Wójcik M, Woźniak LA. Structure and physiological functions of the human peroxisome proliferator-activated receptor γ . *Arch Immunol Ther Exp.* 2008;56(5):331. doi:10.1007/s00005-008-0037-y
12. Duan SZ, Usher MG, Mortensen RM. Peroxisome proliferator-activated receptor-gamma-mediated effects in the vasculature. *Circ Res.* 2008;102(3):283–294. doi:10.1161/CIRCRESAHA.107.164384
13. Fogo AB. Potential for peroxisome proliferator-activated receptor-gamma agonists in progression: beyond metabolism. *Curr Opin Nephrol Hypertens.* 2008;17(3):282–285. doi:10.1097/MNH.0b013e3282f9b1c0

14. Wang G, Wei J, Guan Y, Jin N, Mao J, Wang X. Peroxisome proliferator-activated receptor-gamma agonist rosiglitazone reduces clinical inflammatory responses in type 2 diabetes with coronary artery disease after coronary angioplasty. *Metabolism*. 2005;54(5):590–597. doi:10.1016/j.metabol.2004.11.017
15. Yu J, Jin N, Wang G, Zhang F, Mao J, Wang X. Peroxisome proliferator-activated receptor gamma agonist improves arterial stiffness in patients with type 2 diabetes mellitus and coronary artery disease. *Metabolism*. 2007;56(10):1396–1401. doi:10.1016/j.metabol.2007.05.011
16. Psaty BM, Furberg CD. Rosiglitazone and cardiovascular risk. *Mass Med Soc*. 2007.
17. Palee S, Weerateerangkul P, Chinda K, Chattipakorn SC, Chattipakorn N. Mechanisms responsible for beneficial and adverse effects of rosiglitazone in a rat model of acute cardiac ischaemia-reperfusion. *Exp Physiol*. 2013;98(5):1028–1037. doi:10.1113/expphysiol.2012.070433
18. Liberato MV, Nascimento AS, Ayers SD, et al. Medium chain fatty acids are selective peroxisome proliferator activated receptor (PPAR) γ activators and pan-PPAR partial agonists. *PLoS One*. 2012;7(5):e36297–e. doi:10.1371/journal.pone.0036297
19. Wang L, Waltenberger B, Pferschy-Wenzig E-M, et al. Natural product agonists of peroxisome proliferator-activated receptor gamma (PPAR γ): a review. *Biochem Pharmacol*. 2014;92(1):73–89. doi:10.1016/j.bcp.2014.07.018
20. Julie NL, Julie IM, Kende AI, Wilson GL. Mitochondrial dysfunction and delayed hepatotoxicity: another lesson from troglitazone. *Diabetologia*. 2008;51(11):2108–2116.
21. Yokoi T. Troglitazone. In: Uetrecht J, editor. *Adverse Drug Reactions*. Berlin: Springer; 2010:419–435.
22. Bhalla K, Hwang BJ, Choi JH, et al. N-Acetylfarnesylcysteine is a novel class of peroxisome proliferator-activated receptor γ ligand with partial and full agonist activity in vitro and in vivo. *J Biol Chem*. 2011;286(48):41626–41635. doi:10.1074/jbc.M111.257915
23. Dhawan M, Agrawal R, Ravi J, et al. Rosiglitazone-induced granulomatous hepatitis. *J Clin Gastroenterol*. 2002;34(5):582–584. doi:10.1097/00004836-200205000-00021
24. Bonkovsky HL, Azar R, Bird S, Szabo G, Banner B. Severe cholestatic hepatitis caused by thiazolidinediones: risks associated with substituting rosiglitazone for troglitazone. *Dig Dis Sci*. 2002;47(7):1632–1637. doi:10.1023/A:1015895925374
25. Wallach JD, Wang K, Zhang AD, et al. Updating insights into rosiglitazone and cardiovascular risk through shared data: individual patient and summary level meta-analyses. *BMJ*. 2020;368:l7078. doi:10.1136/bmj.l7078
26. Rodriguez BSQ, Correa R. *Rosiglitazone*. StatPearls [Internet]. StatPearls Publishing; 2019.
27. Chinnam P, Mohsin M, Shafee LM. Evaluation of acute toxicity of pioglitazone in mice. *Toxicol Int*. 2012;19(3):250–254. doi:10.4103/0971-6580.103660
28. Shah P, Mudaliar S. Pioglitazone: side effect and safety profile. *Expert Opin Drug Saf*. 2010;9(2):347–354. doi:10.1517/14740331003623218
29. Zou C, Hu H. Use of pioglitazone in the treatment of diabetes: effect on cardiovascular risk. *Vasc Health Risk Manag*. 2013;9:429.
30. Belcher G, Matthews D. Safety and tolerability of pioglitazone. *Exp Clin Endocrinol Diabetes*. 2000;108(Sup.2):267–273. doi:10.1055/s-2000-8529
31. Nagasaka S, Abe T, Kawakami A, et al. Pioglitazone-induced hepatic injury in a patient previously receiving troglitazone with success. *Diabet Med*. 2002;19(4):347–348. doi:10.1046/j.1464-5491.2002.00712_4.x
32. Elshama SS, El-Kenawy AE-M, Osman H-EH. Toxicological evaluation of subchronic use of pioglitazone in mice. *Iran J Basic Med Sci*. 2016;19(7):712–719.
33. Guasch L, Sala E, Castell-Auvi A, et al. Identification of PPARgamma partial agonists of natural origin (I): development of a virtual screening procedure and in vitro validation. *PLoS One*. 2012;7(11):e50816. doi:10.1371/journal.pone.0050816
34. Chang CH, McNamara LA, Wu MS, et al. A novel selective peroxisome proliferator-activator receptor- γ modulator—SPPAR γ M5 improves insulin sensitivity with diminished adverse cardiovascular effects. *Eur J Pharmacol*. 2008;584(1):192–201. doi:10.1016/j.ejphar.2007.12.036
35. Agrawal R, Jain P, N Dikshit S. Balaglitazone: a second generation peroxisome proliferator-activated receptor (PPAR) gamma (γ) agonist. *Mini Rev Med Chem*. 2012;12(2):87–97. doi:10.2174/138955712798995048
36. Burgermeister E, Schnoebelen A, Flament A, et al. A novel partial agonist of peroxisome proliferator-activated receptor- γ (PPAR γ) Recruits PPAR γ -Coactivator-1 α , prevents triglyceride accumulation, and potentiates insulin signaling in vitro. *Mol Endocrinol*. 2006;20(4):809–830. doi:10.1210/me.2005-0171
37. Henriksen K, Byrjalsen I, Qvist P, et al. Efficacy and safety of the PPARgamma partial agonist balaglitazone compared with pioglitazone and placebo: a Phase III, randomized, parallel-group study in patients with type 2 diabetes on stable insulin therapy. *Diabetes Metab Res Rev*. 2011;27(4):392–401. doi:10.1002/dmrr.1187
38. Kim MK, Chae YN, Kim HS, et al. PAR-1622 is a selective peroxisome proliferator-activated receptor γ partial activator with preserved antidiabetic efficacy and broader safety profile for fluid retention. *Arch Pharm Res*. 2009;32(5):721–727. doi:10.1007/s12272-009-1511-8
39. Kim MK, Chae YN, Choi SH, et al. PAM-1616, a selective peroxisome proliferator-activated receptor γ modulator with preserved antidiabetic efficacy and reduced adverse effects. *Eur J Pharmacol*. 2011;650(2–3):673–681. doi:10.1016/j.ejphar.2010.10.044
40. Kim KR, Lee JH, Kim SJ, et al. KR-62980: a novel peroxisome proliferator-activated receptor gamma agonist with weak adipogenic effects. *Biochem Pharmacol*. 2006;72(4):446–454. doi:10.1016/j.bcp.2006.05.005
41. Jung H, Lee MS, Jang EJ, et al. Augmentation of PPARgamma-TAZ interaction contributes to the anti-adipogenic activity of KR62980. *Biochem Pharmacol*. 2009;78(10):1323–1329. doi:10.1016/j.bcp.2009.07.001
42. Ming Y, Hu X, Song Y, et al. CMHX008, a novel peroxisome proliferator-activated receptor γ partial agonist, enhances insulin sensitivity in vitro and in vivo. *PLoS One*. 2014;9(7):e102102–e. doi:10.1371/journal.pone.0102102
43. Hou Y, Cao X, Hu X, et al. CMHX008, a PPAR γ partial agonist, enhances insulin sensitivity with minor influences on bone loss. *Genes Dis*. 2018;5(3):290–299. doi:10.1016/j.gendis.2018.05.004
44. Amato AA, Rajagopalan S, Lin JZ, et al. GQ-16, a novel peroxisome proliferator-activated receptor γ (PPAR γ) ligand, promotes insulin sensitization without weight gain. *J Biol Chem*. 2012;287(33):28169–28179. doi:10.1074/jbc.M111.332106
45. Xie X, Zhou X, Chen W, et al. L312, a novel PPAR γ ligand with potent anti-diabetic activity by selective regulation. *BBA - Gen Subjects*. 2015;1850(1):62–72. doi:10.1016/j.bbagen.2014.09.027
46. Chigurupati S, Dhanaraj SA, Balakumar P. A step ahead of PPAR γ full agonists to PPAR γ partial agonists: therapeutic perspectives in the management of diabetic insulin resistance. *Eur J Pharmacol*. 2015;755:50–57. doi:10.1016/j.ejphar.2015.02.043
47. Balakumar P, Kathuria S. Submaximal PPAR γ activation and endothelial dysfunction: new perspectives for the management of cardiovascular disorders. *Br J Pharmacol*. 2012;166(7):1981–1992.
48. Khan NM, Ahmad A, Tiwari K, A Kamal M, Mushtaq G, M Ashraf G. Current challenges to overcome in the management of type 2 diabetes mellitus and associated neurological disorders. *CNS Neurol Disord Drug Targets*. 2014;13(8):1440–1457.

49. Opie L. Sodium glucose co-transporter 2 (SGLT2) inhibitors: new among antidiabetic drugs. *Cardiovasc Drugs Ther.* 2014;28(4):331–334. doi:10.1007/s10557-014-6522-0
50. Opie LH, Yellon DM, Gersh BJ. Controversies in the cardiovascular management of type 2 diabetes. *Heart.* 2011;97(1):6–14. doi:10.1136/hrt.2010.214031
51. Shurrah NT, Arafa E-SA. Metformin: A review of its therapeutic efficacy and adverse effects. *Obes Med.* 2020;17:100186. doi:10.1016/j.obmed.2020.100186
52. Alam F, Islam MA, Kamal MA, Gan SH. Updates on managing type 2 diabetes mellitus with natural products: towards antidiabetic drug development. *Curr Med Chem.* 2018;25(39):5395–5431. doi:10.2174/0929867323666160813222436
53. Bonora E. Antidiabetic medications in overweight/obese patients with type 2 diabetes: drawbacks of current drugs and potential advantages of incretin-based treatment on body weight. *Int J Clin Pract.* 2007;61:19–28. doi:10.1111/j.1742-1241.2007.01441.x
54. Chakraborti CK. Exenatide: a new promising antidiabetic agent. *Indian J Pharm Sci.* 2010;72(1):1–11. doi:10.4103/0250-474X.62228
55. Wang J, Pang T, Hafko R, Benicky J, Sanchez-Lemus E, Saavedra JM. Telmisartan ameliorates glutamate-induced neurotoxicity: roles of AT1 receptor blockade and PPAR γ activation. *Neuropharmacology.* 2014;79:249–261. doi:10.1016/j.neuropharm.2013.11.022
56. Fang T, Di Y, Li G, et al. Effects of telmisartan on TNF α induced PPAR γ phosphorylation and insulin resistance in adipocytes. *Biochem Biophys Res Commun.* 2018;503(4):3044–3049. doi:10.1016/j.bbrc.2018.08.091
57. Fujimoto M, Masuzaki H, Tanaka T, et al. An angiotensin II AT1 receptor antagonist, telmisartan augments glucose uptake and GLUT4 protein expression in 3T3-L1 adipocytes. *FEBS Lett.* 2004;576(3):492–497. doi:10.1016/j.febslet.2004.09.027
58. Furukawa H, Mawatari K, Koyama K, et al. Telmisartan increases localization of glucose transporter 4 to the plasma membrane and increases glucose uptake via peroxisome proliferator-activated receptor gamma in 3T3-L1 adipocytes. *Eur J Pharmacol.* 2011;660(2–3):485–491. doi:10.1016/j.ejphar.2011.04.008
59. Zhao S, Chan LKY, Chen L, Cheng TW, Klein T, Leung PS. Combination of telmisartan and linagliptin preserves pancreatic islet cell function and morphology in db/db mice. *Pancreas.* 2016;45(4):584–592. doi:10.1097/MPA.0000000000000505
60. Goyal BR, Parmar K, Goyal RK, Mehta AA. Beneficial role of telmisartan on cardiovascular complications associated with STZ-induced type 2 diabetes in rats. *Pharmacol Rep.* 2011;63(4):956–966. doi:10.1016/S1734-1140(11)70611-9
61. Watanabe M, Inukai K, Sumita T, et al. Effects of telmisartan on insulin resistance in Japanese type 2 diabetic patients. *Internal Med.* 2010;49(17):1843–1847. doi:10.2169/internalmedicine.49.3189
62. Mori H, Okada Y, Arao T, Nishida K, Tanaka Y. Telmisartan at 80 mg/Day Increases High-Molecular-Weight Adiponectin Levels and Improves Insulin Resistance in Diabetic Patients. *Adv Ther.* 2012;29(7):635–644. doi:10.1007/s12325-012-0032-x
63. Yadav A, Kataria MA, Saini V, Yadav A. Role of leptin and adiponectin in insulin resistance. *Clin Chim Acta.* 2013;417:80–84. doi:10.1016/j.cca.2012.12.007
64. Khan A, Khan BT, Qayyum A. Comparative study of telmisartan with pioglitazone on insulin resistance in type 2 diabetic mice. *Pak Armed Forces Med J.* 2017;67(1):31–36.
65. Kurtz TW. New treatment strategies for patients with hypertension and insulin resistance. *Am J Med.* 2006;119(5 Suppl 1):011. doi:10.1016/j.amjmed.2006.01.011
66. Sukomboon N, Poolsup N, Prasit T. Systematic review of the effect of telmisartan on insulin sensitivity in hypertensive patients with insulin resistance or diabetes. *J Clin Pharm Ther.* 2012;37(3):319–327. doi:10.1111/j.1365-2710.2011.01295.x
67. Wago T, Yoshimoto T, Akaza I, et al. Improvement of endothelial function in patients with hypertension and type 2 diabetes after treatment with telmisartan. *Hypertens Res.* 2010;33(8):796–801. doi:10.1038/hr.2010.107
68. Chang W-T, Cheng J-T, Chen Z-C. Telmisartan improves cardiac fibrosis in diabetes through peroxisome proliferator activated receptor δ (PPAR δ): from bedside to bench. *Cardiovasc Diabetol.* 2016;15(1):113. doi:10.1186/s12933-016-0430-5
69. Mikami D, Kimura H, Kamiyama K, et al. Telmisartan activates endogenous peroxisome proliferator-activated receptor- δ and may have anti-fibrotic effects in human mesangial cells. *Hypertens Res.* 2014;37(5):422–431. doi:10.1038/hr.2013.157
70. Tuck ML. Angiotensin-receptor blocking agents and the peroxisome proliferator-activated receptor-gamma system. *Curr Hypertens Rep.* 2005;7(4):240–243. doi:10.1007/s11906-005-0019-y
71. Malek V, Gaikwad AB. Telmisartan and thiorphan combination treatment attenuates fibrosis and apoptosis in preventing diabetic cardiomyopathy. *Cardiovasc Res.* 2019;115(2):373–384. doi:10.1093/cvr/cvy226
72. Kobayashi N, Ohno T, Yoshida K, et al. Cardioprotective mechanism of telmisartan via PPAR-gamma-eNOS pathway in dahl salt-sensitive hypertensive rats. *Am J Hypertens.* 2008;21(5):576–581. doi:10.1038/ajh.2008.27
73. Towfighi A, Ovbiagele B. Partial peroxisome proliferator-activated receptor agonist angiotensin receptor blockers. *Cerebrovasc Dis.* 2008;26(2):106–112. doi:10.1159/000139656
74. Vanitha M, Vijayar R. Effect of telmisartan on serum lipid profile in patients with hypertension and dyslipidemia. *Int J Med Res Health Sci.* 2013;2(4):745–749. doi:10.5958/j.2319-5886.2.4.119
75. Xu C, Fang D, Chen X, et al. Effect of telmisartan on the therapeutic efficacy of pitavastatin in high-fat diet induced dyslipidemic guinea pigs. *Eur J Pharmacol.* 2015;762:364–371. doi:10.1016/j.ejphar.2015.06.005
76. Jayapriya B, Thamilarasi S, Shanthi M, Jafrin AL. Effect of telmisartan on blood pressure and lipid profile in hypertensive patients with dyslipidemia. *Int J Pharm Life Sci.* 2013;4(10).
77. Clemenz M, Frost N, Schupp M, et al. Liver-specific peroxisome proliferator-activated receptor α target gene regulation by the angiotensin type 1 receptor blocker telmisartan. *Diabetes.* 2008;57(5):1405–1413. doi:10.2337/db07-0839
78. Yin SN, Liu M, Jing DQ, Mu YM, Lu JM, Pan CY. Telmisartan increases lipoprotein lipase expression via peroxisome proliferator-activated receptor- α in HepG2 cells. *Endocr Res.* 2014;39(2):67–73. doi:10.3109/07435800.2013.828741
79. Makino H, Haneda M, Babazono T, et al. Prevention of transition from incipient to overt nephropathy with telmisartan in patients with type 2 diabetes. *Diabetes Care.* 2007;30(6):1577–1578. doi:10.2337/dc06-1998
80. Schmieder RE, Delles C, Mimran A, Fauvel JP, Ruilope LM. Impact of telmisartan versus ramipril on renal endothelial function in patients with hypertension and type 2 diabetes. *Diabetes Care.* 2007;30(6):1351–1356. doi:10.2337/dc06-1551
81. Amoghmath S, Suresha R. Effect of telmisartan on blood glucose levels and blood lipid levels in streptozotocin induced diabetic rats. *Biomed Pharmacol J.* 2019;12(3):1303–1308. doi:10.13005/bpj/1758
82. Tojo A, Hatakeyama S, Kinugasa S, Nangaku M. Angiotensin receptor blocker telmisartan suppresses renal gluconeogenesis during starvation. *Diabetes Metab Syndr Obes.* 2015;8:103–113. doi:10.2147/DMSO.S78771
83. Sharma AK, Kanawat DS, Mishra A, et al. Dual therapy of vildagliptin and telmisartan on diabetic nephropathy in experimentally induced type 2 diabetes mellitus rats. *J Renin-Angio-Aldo S.* 2014;15(4):410–418.
84. Senapaty S, Rath B. Biochemical effects of telmisartan versus ramipril in experimental diabetic nephropathy. *Int J Health Sci Res.* 2015;5(5):195–202.

85. Salama RM, Schaalan MF, Ibrahim ME, Khalifa AE, Elkoussi AA. Effectiveness of telmisartan as an adjunct to metformin in treating type II diabetes mellitus in rats. *Open J Endocr Metab Dis.* 2013;3:186–196. doi:10.4236/ojemd.2013.33026
86. Hamed AA, Malek HA. Effect of telmisartan in experimentally induced diabetes mellitus in rats. *Int J Health Sci.* 2007;1(2):249.
87. Barnett A. Preventing renal complications in type 2 diabetes: results of the diabetics exposed to telmisartan and enalapril trial. *J Am Soc Nephrol.* 2006;17(4 suppl 2):S132–S5. doi:10.1681/ASN.2005121326
88. Barzilay JI, Gao P, Rydén L, et al. Effects of telmisartan on glucose levels in people at high risk for cardiovascular disease but free from diabetes: the TRANSCEND study. *Diabetes Care.* 2011;34(9):1902–1907. doi:10.2337/dc11-0545
89. Muñoz-Torrero JFS, Rivas MD, Costo A, et al. Telmisartan improves insulin resistance in patients with low cytokine levels. *J Investig Med.* 2011;59(3):602–605. doi:10.2310/JIM.0b013e31820bf26b
90. Sengul AM, Altuntas Y, Kürklü A, Aydın L. Beneficial effect of lisinopril plus telmisartan in patients with type 2 diabetes, microalbuminuria and hypertension. *Diabetes Res Clin Pract.* 2006;71(2):210–219. doi:10.1016/j.diabres.2005.06.010
91. Nagel JM, Tietz AB, Göke B, Parhofer KG. The effect of telmisartan on glucose and lipid metabolism in nondiabetic, insulin-resistant subjects. *Metabolism.* 2006;55(9):1149–1154. doi:10.1016/j.metabol.2006.04.011
92. Mazerska M, Myśliwiec M. Telmisartan lowers albuminuria in type 2 diabetic patients treated with angiotensin enzyme inhibitors. *Adv Med Sci.* 2009;54(1). doi:10.2478/v10039-009-0015-6
93. Makino H, Haneda M, Babazono T, et al. Microalbuminuria reduction with telmisartan in normotensive and hypertensive Japanese patients with type 2 diabetes: a post-hoc analysis of The Incipient to Overt: angiotensin II Blocker, Telmisartan, Investigation on Type 2 Diabetic Nephropathy (INNOVATION) study. *Hypertens Res.* 2008;31(4):657–664.
94. Al-Hammami FA, Hasan OA. Comparative study of lisinopril versus telmisartan effects on oxidative stress in diabetic type 2 hypertensive patients. *Raf J Sci.* 2012;23(2E):34–41.
95. Vitale C, Mercurio G, Castiglioni C, et al. Metabolic effect of telmisartan and losartan in hypertensive patients with metabolic syndrome. *Cardiovasc Diabetol.* 2005;4(1):6. doi:10.1186/1475-2840-4-6
96. De Luis D, Conde R, González-Sagrado M, et al. Effects of telmisartan vs olmesartan on metabolic parameters, insulin resistance and adipocytokines in hypertensive obese patients. *Nutr Hosp.* 2010;25(2):275–279.

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion

and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-targets-and-therapy-journal>