


Reply to: “Intraclass Difference in Pneumonia Risk with Fluticasone and Budesonide in COPD: A Systematic Review of Evidence from Direct-Comparison Studies” [Letter]

Neil C Barnes ^{1,2}

Dave Singh³

David A Lipson ^{4,5}

¹Respiratory Therapy Area, GlaxoSmithKline, Middlesex, UK; ²William Harvey Institute, Bart's and the London School of Medicine and Dentistry, London, UK; ³Centre for Respiratory Medicine and Allergy, Institute of Inflammation and Repair, Manchester Academic Health Science Centre, The University of Manchester, Manchester University NHS Foundation Trust, Manchester, UK; ⁴Clinical Sciences, GlaxoSmithKline, Collegeville, PA, USA; ⁵Pulmonary, Allergy and Critical Care Division, Department of Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

Dear editor

We read with interest the article by Lodise et al.¹ The analysis is fundamentally flawed. The authors conclude that there is a greater risk of pneumonia with fluticasone compared with budesonide (BUD). There is no such drug as fluticasone; there is fluticasone propionate (FP) and fluticasone furoate (FF). These are different compounds with different molecular structures, potency and dosages. Of the six fluticasone studies discussed,¹ five investigated FP, whereas one (Lipson et al) investigated FF. These drugs need to be analysed separately.

A major problem with analyses that compare pneumonia rates is that multiple risk factors for ICS-related pneumonia exist.² Such analyses require valid comparison groups, which can only be done reliably by comparing pneumonia rates between ICS- and non-ICS-containing treatments within the same trial. This was done in a systematic review³ not cited by Lodise et al. It showed no difference between fluticasone- versus BUD-containing treatments with regards to serious or fatal pneumonia events; there was a significantly higher risk of any pneumonia event with fluticasone-containing treatments but the authors advised caution due to differences in pneumonia assessment between studies and the lack of head-to-head trials.³

Unfortunately, there are few direct comparisons between FF and BUD. Lodise et al do quote Lipson et al but only partially. At 24 weeks there was a higher rate of pneumonia with FF/umeclidinium/vilanterol (FF/UMEC/VI) compared with BUD/formoterol (BUD/FOR), but at 52 weeks there was no difference, which the authors fail to mention.¹

When Lodise et al was published they will have been aware of the results of the ETHOS study.⁴ The ETHOS and IMPACT studies recruited similar groups of patients with symptomatic COPD and a history of exacerbations, but defined pneumonia in different ways, which may result in different absolute pneumonia rates.^{4,5} A within-study comparison of pneumonia events shows very similar pneumonia rate ratios between triple therapy and non-ICS-containing treatments (1.7 in ETHOS for BUD 320µg/glycopyrronium [GLY]/FOR vs GLY/FOR; 1.6 in IMPACT for FF/UMEC/VI vs UMEC/VI).^{4,5}

Correspondence: Neil C Barnes
GlaxoSmithKline, 980 Great West Road,
London, TW8 9GS, UK
Email neil.c.barnes@gsk.com

The other studies referenced are observational, and the increased pneumonia risk identified in this systematic review is highly dependent on the PATHOS study by Janson et al. This was an observational study in Sweden for which Lodise et al¹ failed to adequately identify and assess risk of bias, an essential element of systematic literature reviews. Propensity matching was used in an attempt to balance patients using the two ICS/LABA treatments in PATHOS, but this did not include factors known to be important risk predictors for ICS-related pneumonia in COPD.^{2,6} Lung function data were included where available in the propensity score, but it is unclear how many patients had missing data. Contemporaneous data suggest that spirometry is only conducted in 45–52% of patients with COPD in primary care in Sweden,^{7,8} suggesting that many patients in PATHOS may have lung function unaccounted for. Janson et al was a secondary analysis of a study whose stated primary aim was to describe the evolution of COPD management in Sweden over an 11-year period.⁹ Unusually, the pneumonia analysis was published before the paper describing the protocol and primary purpose of the study meaning that referees and readers of the pneumonia analysis could not fully understand the potential biases. The primary study results showed significant improvements in outcomes in Sweden over the 11-year period and a follow-up paper showed that clinics with a trained respiratory nurse had better outcomes than those that did not,^{9,10} but neither the index date nor these differences were accounted for in propensity matching in the PATHOS pneumonia analysis. Lastly, and perhaps most importantly, although death due to pneumonia was more common in the patients prescribed FP by their physicians, overall mortality was the same, implying that there were more deaths from other causes with BUD compared with FP, this is not discussed.

Finally the European Medicines Agency, an independent body with a responsibility for patient safety, has twice examined whether there is any difference in pneumonia rates between different ICSs and concluded that there is no intraclass difference.¹¹

Acknowledgments

The authors would like to thank Paul Jones, Kieran J. Rothnie and Mark T. Dransfield for their contribution to the data analysis and interpretation. Editorial support (in the form of collating author comments, grammatical editing and referencing) was provided by Rachel Edwards, PhD, at Fishawack Indicia Ltd, part of Fishawack Health,

and was funded by GSK. Dave Singh is supported by the National Institute for Health Research (NIHR) Manchester Biomedical Research Centre (BRC).

Disclosure

N.C. Barnes and D.A. Lipson are employees of GSK and hold stocks and shares in GSK. D. Singh has received personal fees from GSK, Cipla, Genentech and Peptinnovate, and personal fees and grant support from AstraZeneca, Boehringer Ingelheim, Chiesi, Glenmark, Menarini, Mundipharma, Novartis, Pfizer, Pulmatrix, Theravance and Verona. The authors report no other conflicts of interest in this communication.

References

- Lodise TP, Li J, Gandhi HN, O'Brien G, Sethi S. Intraclass difference in pneumonia risk with fluticasone and budesonide in COPD: a systematic review of evidence from direct-comparison studies. *Int J Chron Obstruct Pulmon Dis.* 2020;15:2889–2900. doi:10.2147/COPD.S269637
- Hartley BF, Barnes N, Lettis S, Compton CH, Papi A, Jones P. Risk factors for exacerbations and pneumonia in patients with chronic obstructive pulmonary disease: a pooled analysis. *Respir Res.* 2020;21(1):5. doi:10.1186/s12931-019-1262-0
- Kew KM, Seniukovich A. Inhaled steroids and risk of pneumonia for chronic obstructive pulmonary disease. *Cochrane Database Syst Rev.* 2014;3:Cd010115.
- Rabe KF, Martinez FJ, Ferguson GT, et al. Triple inhaled therapy at two glucocorticoid doses in moderate-to-very-severe COPD. *N Engl J Med.* 2020;383:35–48. doi:10.1056/NEJMoa1916046
- Lipson DA, Barnhart F, Brealey N, et al. Once-daily single-inhaler triple versus dual therapy in patients with COPD. *N Engl J Med.* 2018;378(18):1671–1680. doi:10.1056/NEJMoa1713901
- Crim C, Calverley PM, Anderson JA, et al. Pneumonia risk in COPD patients receiving inhaled corticosteroids alone or in combination: TORCH study results. *Eur Respir J.* 2009;34(3):641–647. doi:10.1183/09031936.00193908
- Lofdahl CG, Tilling B, Ekstrom T, Jorgensen L, Johansson G, Larsson K. COPD health care in Sweden - A study in primary and secondary care. *Respir Med.* 2010;104(3):404–411. doi:10.1016/j.rmed.2009.10.007
- Weidinger P, Nilsson JL, Lindblad U. Adherence to diagnostic guidelines and quality indicators in asthma and COPD in Swedish primary care. *Pharmacoepidemiol Drug Saf.* 2009;18(5):393–400. doi:10.1002/pds.1734
- Ställberg B, Janson C, Johansson G, et al. Management, morbidity and mortality of COPD during an 11-year period: an observational retrospective epidemiological register study in Sweden (PATHOS). *Prim Care Respir J.* 2014;23(1):38–45. doi:10.4104/pcrj.2013.00106
- Lisspers K, Johansson G, Jansson C, et al. Improvement in COPD management by access to asthma/COPD clinics in primary care: data from the observational PATHOS study. *Respir Med.* 2014;108(9):1345–1354. doi:10.1016/j.rmed.2014.06.002
- European Medicines Agency (EMA). Pharmacovigilance Risk Assessment Committee (PRAC), recommendation 2016. Inhaled corticosteroids (ICS) containing medicinal products indicated in the treatment of chronic obstructive pulmonary disease (COPD). 2016. Available from: <https://www.ema.europa.eu/medicines/human/referals/inhaled-corticosteroids-containing-medicinal-products-indicated-treatment-chronic-obstructive>. Accessed November, 2020.

Dove Medical Press encourages responsible, free and frank academic debate. The content of the International Journal of Chronic Obstructive Pulmonary Disease 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the International Journal of Chronic Obstructive Pulmonary Disease editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

International Journal of Chronic Obstructive Pulmonary Disease

Dovepress

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management

protocols. This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>