

Adherence to Adjuvant Hormonal Therapy and Associated Factors Among Women with Breast Cancer [Response To Letter]

Zerko Wako ¹
Daniel Mengistu ²
Negalign Getahun Dinegde ²

¹Department of Medical Nursing, School of Nursing, College of Medicine and Health Science, University of Gondar, Gondar, Ethiopia; ²School of Nursing and Midwifery, College of Health Science, Addis Ababa University, Addis Ababa, Ethiopia

Dear editor

We appreciate the Editor-in-Chief for allowing us to respond to comments from Bekele et al. We also express our appreciation to Bekele et al for the attention on our study entitled ‘Adherence to adjuvant hormonal therapy and associated factors among women with breast cancer attending the Tikur Anbessa Specialized Hospital, Addis Ababa, Ethiopia’. Hereby, we provide a point-by-point response to the comments raised by Bekele et al from the following aspects.

For the first suggestion of Bekele et al as they were a clinical pharmacy specialist by profession will make them to enrich their study by addressing our suggestion and limitation as well. They appreciated our study on the importance of assessing the adherence to adjuvant hormonal therapy in developing countries like Ethiopia.¹ They should acknowledge that it is the first study in Ethiopia regarding adherence to adjuvant hormonal therapy.

For the second suggestion of Bekele et al on the need of using different methodological approaches, following the authors’ recommendation, as we also recommended on the use of different methods including follow up study and as adherence study is multifaceted there are so many factors that can affect the patients’ adherence to adjuvant hormonal therapy that need to be identified.

Appropriate medication adherence could result in a reduction of the recurrence and mortality rates among women diagnosed with breast cancer.² Despite this, many studies reported poor adherence to hormonal therapy.^{3,4}

For the third suggestion of Bekele et al listed that the authors only assess the rate of medication adherence. We would like enlighten Bekele et al that we have not only tried to assess the rate of adherence but also tried to identify factors affecting it. On the other hand, I agree with Bekele et al that we have not studied the outcome or impact of adherence on breast cancer recurrence and mortality, we also strongly recommend others to consider it.

We also believed that the impact of non-adherence on the treatment outcomes of breast cancer should be studied at the large. We support the fact that

Correspondence: Zerko Wako
Tel +251918728268
Email zerwabek13@gmail.com

Bekele et al studied using retrospectively reviewing the patients' charts as per our recommendation. Therefore, I believe and expect a huge number of non-adherences if conducted prospectively. We agree on their recommendation that a future researchers should assess the adherences prospectively by using a validated tool. Even though the design we used was cross-sectional, but we had checked patient chart review retrospectively.

About the fourth suggestion of Bekele et al, on the authors identifying the therapeutic communication and side effects of the drug as predictors of adherence. I strongly support the idea that a clinical pharmacy intervention in the study area to tackle any medication-related problems. Furthermore, effective patient health-care provider communication was significantly associated with adherence to Tamoxifen and Aromatase inhibitors.⁵ On the other hand, we the authors only assessed whether the patients had experienced side effects or not but not the predictors for the occurrences of side effects as these variables need a follow-up study.

According to a study done by Brito et al found that the rate of adherence to hormonal therapy was lower among alcohol drinkers.⁶ We the authors of the paper tried to assess the magnitudes of social drug uses like smoking a cigarette and drinking alcohol. However, we left the variable because it was not fit for binary logistic regression analysis as it violates cross-tabulation. Besides, the nature of the variables was so unwelcoming to our culture and respondents were not willingly responded to it.

As per the suggestion of providing quality pharmaceutical care service globally could satisfy the demands of the patients that might increase the adherences.⁷ Hence, the effect of increased patient satisfaction on treatment adherence should be studied at large. We strongly suggest Bekele et al to use more recent data because ref number 7 was studied in 2001. However, we believed that patient satisfaction increases with increasing adherence by reducing morbidity and mortality among breast cancer patients.⁸

As per comorbidity, the authors believe that those having comorbid illness likely to be non-adhered. We authors also support Bekele et al idea

that multiple drugs prescription (polypharmacy) and the presence of comorbidity can reduce the patient's adherence.⁹ On contrary, study conducted by Liu et al revealed that those with at least one comorbid illness were having odds of increasing adherence.⁵ Therefore, we the authors' should recommend to use Charlson Comorbidity Index (CCI) to assess the presence of comorbidity and we should also pay special attention to those patients with multiple comorbidity.

Lastly, we the authors recommended specifically for health care workers to distribute the leaflets, the use of alarm as an alert for the elderly, and creation of adherence counselling to scale up adherence. Furthermore, Bekele et al recommended the establishment of the drug information centre at the hospital to address issues that affect patient adherence to the medication.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Wako Z, Mengistu D, Dinegde NG, Asefa T, Wassie M. Adherence to adjuvant hormonal therapy and associated factors among women with breast cancer attending the Tikur Anbessa Specialized Hospital, Addis Ababa Ethiopia, 2019: a cross-sectional study. *Breast Cancer*. 2021;13:383.
2. Felder TM, Heiney SP, Hebert JR, et al. Improving adherence to adjuvant hormonal therapy among disadvantaged women diagnosed with breast cancer in South Carolina: proposal for a multimethod study. *JMIR Res Protoc*. 2020;9(9):e17742. doi:10.2196/17742
3. Sella T, Chodick G. Adherence and persistence to adjuvant hormonal therapy in early-stage breast cancer patients: a population-based retrospective cohort study in Israel. *Breast Care*. 2020;15(1):45–54. doi:10.1159/000500318
4. Farias AJ, Hansen RN, Zeliadt SB, Ornelas IJ, Li CI, Thompson B. Factors associated with adherence to adjuvant endocrine therapy among privately insured and newly diagnosed breast cancer patients: a quantile regression analysis. *J Manag Care Spec Pharm*. 2016;22(8):969–978. doi:10.18553/jmcp.2016.22.8.969
5. Liu Y, Malin JL, Diamant AL, Thind A, Maly RC. Adherence to adjuvant hormone therapy in low-income women with breast cancer: the role of provider-patient communication. *Breast Cancer Res Treat*. 2013;137(3):829–836. doi:10.1007/s10549-012-2387-8
6. Brito C, Portela MC, de Vasconcellos MT. Adherence to hormone therapy among women with breast cancer. *BMC Cancer*. 2014;14:397. doi:10.1186/1471-2407-14-397
7. Andaleeb SS. Service quality perceptions and patient satisfaction: a study of hospitals in a developing country. *Soc Sci Med*. 2001;52(9):1359–1370. doi:10.1016/S0277-9536(00)00235-5

8. Murphy CC, Bartholomew LK, Carpentier MY, Bluethmann SM, Vernon SW. Adherence to adjuvant hormonal therapy among breast cancer survivors in clinical practice: a systematic review. *Breast Cancer Res Treat.* 2012;134(2):459–478. doi:10.1007/s10549-012-2114-5
9. Bekele F, Fekadu G, Bekele K, Dugassa D, Sori J. Drug-related problems among patients with infectious disease admitted to medical wards of Wollega University Referral Hospital: prospective observational study. *SAGE Open Med.* 2021;9:2050312121989625. doi:10.1177/2050312121989625

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Patient Preference and Adherence 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Patient Preference and Adherence editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Patient Preference and Adherence

Dovepress

Publish your work in this journal

Patient Preference and Adherence is an international, peer-reviewed, open access journal that focusing on the growing importance of patient preference and adherence throughout the therapeutic continuum. Patient satisfaction, acceptability, quality of life, compliance, persistence and their role in developing new therapeutic modalities and compounds to optimize clinical outcomes for existing disease

states are major areas of interest for the journal. This journal has been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/patient-preference-and-adherence-journal>