

Potential Impact of Active *H. pylori* Infection-Related Metabolic Syndrome on Preeclampsia Among Pregnant Women [Letter]

Maria Tziritidou-Chatzopoulou^{1,2}, Dimitra Gialamprinou³, Jannis Kountouras¹

¹Department of Medicine, Second Medical Clinic, Aristotle University of Thessaloniki, Ippokraton Hospital, Thessaloniki, Macedonia, Greece; ²School of Healthcare Sciences, Midwifery Department, University of West Macedonia, Kozani, Macedonia, 50100, Greece; ³Second Neonatal Department and NICU, Aristotle University of Thessaloniki, Papageorgiou General Hospital, Thessaloniki, Macedonia, 56403, Greece

Correspondence: Jannis Kountouras, Professor of Medicine, Gastroenterologist, 8 Fanariou St, Byzantio, Thessaloniki, Macedonia, 551 33, Greece, Tel +30-2310-892238, Fax +30-2310-892744, Email jannis@auth.gr; ancoratus2010@gmail.com

Dear editor

In their case control study, Tsegaye et al¹ by introducing the stool antigen test (SAT) for the diagnosis of *Helicobacter* infection (*Hp-I*) concluded that *Hp-I* is linked with preeclamptic pregnant women in Ethiopia associated with metabolic syndrome (MetS) parameters such as body mass index, diabetes mellitus, dyslipidemia and arterial hypertension.

Although the SAT has been introduced broadly in Africa, the limitation of this study, as also mentioned by others² is that only the SAT was used. By introducing several diagnostic approaches increases the detection of *Hp-I*.²

Specifically, despite the advantages (easy to perform and rapid for *Hp* detection), SATs display several disadvantages. Because the principle of SAT is an antigen-antibody reaction, differences in antigens used for SAT in different geographic regions may cause heterogeneity of results; the diagnostic performances of different SATs are heterogeneous, and this could relate to the designs of the test like enzyme immunoassay and immunochromatographic assay (ICA) and for the selection of antibody, such as monoclonal antibody and polyclonal antibody. SAT shows a high rate of false positivity. Moreover, SAT negative results may not signify the absence of *Hp-I* because the low gastric colonization of bacteria results in low concentration of *Hp* antigen in the sample. Furthermore, concerning ICA-based methods, subjective interpretation of the results, especially for the tracing line, makes it problematic to diagnostic accuracy. The accuracy of SAT is influenced by several factors, including antibiotics, proton pump inhibitors, N-acetylcysteine, bowel movement and upper gastrointestinal tract bleeding. Preservation of the specimen, like temperature and transport time before testing, and cut-off valve also have impacts on the SAT diagnostic accuracy; if study results are not compared to the gold standard examinations for the diagnosis of active *Hp-I* (histology, rapid urease test, culture or urea breath test), positive or negative results are still controversial.

In this regard, important to note that, only active *Hp-I* induces humoral and cellular immune responses that, owing to the sharing of homologous epitopes (molecular mimicry), cross-react with components of the host, thereby contributing and possibly perpetuating *Hp*-related systemic disorders^{3,4} including preeclampsia.⁵ In this respect, maternal inflammatory immune reactions appear to be involved in the pathogenesis of preeclampsia; several studies of both patients and animal models have established that changes in the immune system (cellular plus humoral immunity and/or adaptive-innate immunity) could contribute to the pathophysiology of preeclampsia.

Hp-I, a worldwide burden with a mean global prevalence of 58% and accounting of about 4.4 billion of *Hp* infected individuals, is also related with MetS-systemic systemic pathologies such as preeclampsia, cardio-cerebrovascular, and neurodegenerative diseases, the end outcomes of MetS;³⁻⁵ and it appears to be a critical risk factor of MetS in pregnancy and influences the occurrence of many opposing pregnancy outcomes⁵ such as gestational diabetes mellitus, arterial hypertension, hyperemesis of pregnancy, preeclampsia in obese or overweight pregnant women, fetal growth restriction,

premature delivery, abortion, *Hp-I* of newborn, and neural tube defects. Moreover, eradicating *Hp-I* may reduce or delay such systemic pathologies.³

Therefore, active *Hp-I*-related MetS may affect pregnancy critical outcomes, *Hp* eradication may benefit pregnant women and their neonates and thus further studies are needed.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Tsegaye N, Diriba R, Araya S. Magnitude of *H. pylori* and its association with preeclampsia among pregnant women in Ethiopia: a Case Control Study. *Int J Womens Health*. 2022;14:635–642. doi:10.2147/IJWH.S363137
2. Yisak H, Belete D, Mahtsentu Y. *Helicobacter pylori* infection and related factors among pregnant women at Debre Tabor General Hospital, Northwest Ethiopia, 2021. *Anemia Highly Related H Pylori Womens Health*. 2022;18:17455057221092266. doi:10.1177/17455057221092266
3. Kountouras J, Doulberis M, Polyzos SA, et al. Impact of *Helicobacter pylori* and/or *Helicobacter pylori*-related metabolic syndrome on incidence of all-cause and Alzheimer's dementia. *Alzheimers Dement*. 2019;15(5):723–725. doi:10.1016/j.jalz.2019.01.008
4. Kountouras J, Papaefthymiou A, Polyzos SA, et al. Impact of *Helicobacter pylori*-Related metabolic syndrome parameters on arterial hypertension. *Microorganisms*. 2021;9(11):2351. doi:10.3390/microorganisms9112351
5. Zhou B, Wang F. Research progress in relation of *Helicobacter pylori* infection with pregnancy-related diseases and adverse pregnancy outcomes. *Zhong Nan Da Xue Xue Bao Yi Xue Ban*. 2020;45(3):338–344. doi:10.11817/j.issn.1672-7347.2020.190032

Dove Medical Press encourages responsible, free and frank academic debate. The content of the International Journal of Women's Health 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the International Journal of Women's Health editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

International Journal of Women's Health

Dovepress

Publish your work in this journal

The International Journal of Women's Health is an international, peer-reviewed open-access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of women's healthcare including gynecology, obstetrics, and breast cancer. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-womens-health-journal>

<https://doi.org/10.2147/IJWH.S378482>