

The Doctors' Effect on Patients' Physical Health Outcomes Beyond the Intervention: A Methodological Review

Christoph Schnelle , Mark A Jones

Institute of Evidence-Based Healthcare, Bond University, Robina, QLD, 4226, Australia

Correspondence: Christoph Schnelle, Institute of Evidence-Based Healthcare, Bond University, 14 University Drive, Robina, QLD, 4226, Australia, Email christoph.schnelle@student.bond.edu.au



Background: Previous research suggests that when a treatment is delivered, patients' outcomes may vary systematically by medical practitioner.

Objective: To conduct a methodological review of studies reporting on the effect of doctors on patients' physical health outcomes and to provide recommendations on how this effect could be measured and reported in a consistent and appropriate way.

Methods: The data source was 79 included studies and randomized controlled trials from a systematic review of doctors' effects on patients' physical health. We qualitatively assessed the studies and summarized how the doctors' effect was measured and reported.

Results: The doctors' effects on patients' physical health outcomes were reported as fixed effects, identifying high and low outliers, or random effects, which estimate the variation in patient health outcomes due to the doctor after accounting for all available variables via the intra-class correlation coefficient. Multivariable multilevel regression is commonly used to adjust for patient risk, doctor experience and other demographics, and also to account for the clustering effect of hospitals in estimating both fixed and random effects.

Conclusion: This methodological review identified inconsistencies in how the doctor's effect on patients' physical health outcomes is measured and reported. For grading doctors from worst to best performances and estimating random effects, specific recommendations are given along with the specific data points to report.

Keywords: methodological study, meta-epidemiology, meta-epidemiological review, research methods, doctors' effect

Introduction

A fundamental question in medical research is whether medical practitioners have an effect on patients' health beyond the intervention, patient risk, and hospital variables. Previous research has revealed that when a treatment is delivered by a doctor (ie surgeon or medical physician), patient outcomes may vary systematically by medical practitioner.^{1,2} It is well known that hospitals can have an influence on patients' health outcomes, with wide variation between hospitals.³⁻⁷ Such outcomes include adverse events,⁴ prescribing errors,⁴ hospital readmission,^{5,6} and mortality.⁷⁻⁹ Comparing hospitals requires a sound methodology and reliable estimates that take into account the multiple variables involved.^{8,10} In contrast to the substantial research on hospital effects, there is minimal research on the effect of doctors.

The influence of doctor-patient communication has been investigated as a "doctor effect" on patients' health outcomes,^{1,11,12} including symptoms,^{13,14} readmission rates in the emergency department,^{13,15} health-related quality of life,¹⁶ and improved diabetes control.¹⁷

Research on the therapist effect in psychotherapy has shown significant effects of therapists on patient outcomes beyond the therapy technique or modality applied.^{18,19} This wide variation among practitioners has been acknowledged and incorporated into the training material for psychotherapists.^{20,21} In surgery, outcomes associated with procedure volume, seniority, level of experience, or doctor specialty, include mortality rate,²² length of hospital stay,^{23,24} post-operative complications,²⁵ and readmission.^{26,27} While research on the doctors' effect in non-surgical specialties is limited, there is evidence from studies in primary care,^{1,28} intensive care,²⁹ acute care,³⁰ and obstetrics,³¹ where medical practitioners had an effect on patients' health outcomes.

Given the significant therapist effect in psychotherapy, and the known wide variation in patient outcomes across hospitals, but unclear effect of individual doctors on patient outcomes, we conducted a systematic review of the effect of doctors on patients' physical outcomes. We aimed to assess whether doctor effects vary with specificity, outcome and intervention. However, in conducting the review, we found substantial variation in the way a doctor effect is measured and reported, therefore making data synthesis challenging and meta-analysis impossible. This has led to the present study where we have conducted a methodological review of studies that measure and report on doctors' effect on physical patient outcomes. The focus of the methodological review is on the method of measurement of the doctors' effect as well as how it is reported. The data source for the review is the included studies from our systematic review.³²

Objective

To conduct a methodological review of studies reporting on the effect of doctors on patients' physical health outcomes and to provide recommendations on how this effect could be measured and reported in a consistent and appropriate way.

Materials and Methods

Design

The present study is a methodological review where the focus is on statistical analysis and reporting.³³ The search strategy, data collection, and extraction are explained in detail in a previous report of a systematic review of the surgeons' effect on patients' physical health outcomes.³²

Search Strategy

Three databases were searched initially: PubMed, Embase, and PsycINFO; and over 10,000 publications were screened. For each of the studies identified that met the inclusion criteria, a citation analysis on Scopus was conducted to identify further eligible studies. The full search strategy and keywords can be found in the [Supplementary Material](#).

Study Selection and Eligibility Criteria

The studies selected in the initial electronic search and the studies added through the citation analysis were independently reviewed by two researchers with a third reviewer acting as an arbitrator if required. This process resulted in 79 included studies, all of which are included in the present study. Any physical patient health-related outcome was eligible for inclusion. Studies that fulfilled any of the following criteria were excluded: (1) studies that only described a doctors' effect on particular doctor-related variables (such as specialty of doctor), (2) studies with fewer than 15 doctors, (3) cross-sectional studies, and (4) studies that mention fixed or random effects but did not list them either graphically or in numerical form.

Data Extraction and Quality Assessment

CS extracted the relevant information for assessing doctor effects from each included study, and the extracted data was then reviewed by a second researcher. The data items extracted can be found in [Table 1](#). For quality assessment, the Newcastle-Ottawa Scale (NOS) was used, with the majority of studies scoring between 8 and 9 (9 being the maximum total).³⁴⁻³⁶

Table 1 Data Items Extracted

Data Item	Comment
Publication	First author, year
Surgeon or Other Medical Specialty	Surgeon, Other
Practitioner Type	Surgeon, GP, Cardiologist, etc.
Medical Specialty of Doctor	
Detailed Intervention	
General Outcome	
Specific Outcome	Often same as General Outcome
Type of Study	Cohort or Randomized Controlled Trial
Newcastle Ottawa Scale Score	0–9
Count of Doctors in Study	
Count of Patients	
Count of Institutions	
Doctor ICC	Intra-class correlation coefficient, here a measure of the strength of the effect on patients' physical health
Multivariate Data Analysis used	Y/N
Percentage of Doctors that are Outliers	Positive and Negative Outliers
Country of dataset analyzed	

Methodological Review

We planned to describe the methods used to estimate and report the doctors' effect on patients' physical outcomes including the statistical model used, types of confounding variables adjusted for (patient variables, hospital/institution variables, doctor variables), and the method of reporting the doctor effect.

Results

Of the 79 included studies, 62 used a multivariable multilevel regression model to estimate the doctors' effect, 72 studies included patient variables in their model, 41 studies included hospital or institution variables in the model, 60 studies included doctors' volume, and 24 studies included other doctor variables. There were two different ways that the doctors' effect was reported: fixed effects and random effects,^{37,38} with 54 studies reporting fixed effects and 34 studies reporting random effects.

Table 2 provides details for each included study, presenting in part the wide variety of statistical methods used.

Fixed Effects – Grading Doctors by Their Effect

Fixed effects are represented by the range of patient outcomes that doctors are responsible for after all available confounding variables have been accounted for. They are shown visually using a caterpillar plot, which ranks doctors by outcomes from lowest to highest, or a funnel plot, which shows each doctor as a dot and indicates whether doctors are outside a 95% or 99% confidence interval. For example, Papachristofi et al³⁹ showed caterpillar graphs with an ICC of 4.0% (surgeons) and an ICC of 0.25% (anesthetists) (Figure 1), while Kunadian et al⁴⁰ showed a funnel plot with an ICC of 6.5% (Figure 2), redone at a higher resolution by the authors (Figure 3) and the same data as a caterpillar plot (Figure 4). Measuring fixed effects allows

Table 2 Detailed Results for Each Study

Publication	Doctors	Patients/ Procedures	Institutions	ICC %	Neg Outlier %	Pos Outlier %	Country	MLR*	MV**	Statistical Analysis	PV^	HV^^	DVo#	ODV###	Confidence Interval Calculation
Anderson, 2016 ²²	NS	2880	35		Other	Other	US	Y		"Gaussian Kernel Densities were constructed to show the relative distributions of the effects of individual institutions and surgeons"	Y	Y	Y	N	None
Aquina, 2015 ⁵¹ pg e163	NS	158,596	NS		Other	Other	US	Y		"Mixed Effects Multivariable Logistic Regression", conference abstract	Y	Y	Y	N	95% CI given, but not method
Aquina, 2015 ⁵²	223	14,875	99		13.0	28.0	US	Y		"Bivariate and hierarchical logistic regression with further multivariable analysis" R 3.1 SAS 9.3	Y	Y	Y	N	95% CI given, but not method
Aquina, 2016 ⁵³	3481	125,160	210	24.3	Other	Other	US	Y		"Three-level mixed-effects logistic regression analyses were performed" R 3.2.0 SAS 9.4	Y	Y	Y	Y	None
Aquina, 2017 ⁵⁴	1572/ 2012	124,416/ 78,267	260/256	40.5/ 14			US	Y		"Mixed-effects Cox proportional hazards analyses" R 3.2.1 SAS 9.4	Y	Y	Y	Y	95% CI given, but not method
Arvidsson, 2005 ⁵⁵	25	1068	7		Other	Other	Sweden	N	Y	SAS 8.2 NL Mixed model	Y	Y	N	N	None
Becerra, 2017 ⁵⁶	1503/ 814	12,332	187	7.9			US	Y		"Multilevel logistic regression", "multilevel competing-risks Cox models" SAS 9.3 R	Y	Y	Y	Y	95% CI given, but not method
Beckett, 2018 ³⁰	22	21,570	1				UK	N	N	Analysis based on r-square	N	N	Y	N	None
Begg, 2002 ⁵⁷	159	10,737	72		8/13/9	3/14/3	US	Y		Correlation-adjusted and GEE logistic regression	Y	Y	Y	N	None
Bianco, 2005 ⁵⁸	159	5238	NS		7.5	2.5	US	Y		Logistic regression, binomial distribution, histograms, extra-binomial variation	Y	Y	Y	N	None
Bianco, 2010 ⁵⁹	54	7725	4		9.3	13.0	US	Y		"[M]ultivariable, parametric random-effects regression survival-time model, using a log-logistic survival distribution to model hazard over time" Stata 9.2	Y	N	Y	N	95% CI given, but not method

Bolling, 2010 ⁶⁰	1088	28,507	639		6.6	7.4	US	Y		GEE logistic regression SAS 9.2 GENMOD	Y	N	Y	N	Funnel plot with 95% CI
Bridgewater, 2003 ⁶¹	23	8572	4		0.0	0.0	UK	Y		Unspecified, using SAS	Y	N	Y	N	95% CI given, but not method
Bridgewater, 2005 ⁶²	25	1097/9066	4		0.0	0.0	UK	Y		Unspecified, using SAS	Y	N	Y	N	Clopper-Pearson 95% CI
Brown, 2016 ⁶³	133	14,033	84		6.0	6.8	US	Y		Bayesian hierarchical logistic regression	Y	N	Y	Y	95% CI given, but not method, f did better
Burns, 2011 ⁶⁴	1557	246,469	156		0.7	4.5	UK	N	Y	Logistic regression	Y	Y	Y	N	“We constructed funnel plots using exact Poisson control limits by means of the web tool available at www.erpho.org.uk/topics/tools/funnel.aspx .”
Cirillo, 2020 ⁶⁵	32	19,824	1		3.1	0.0	Italy	Y		Logistic regression, random effects meta-analysis	Y	N	Y	N	95% CI given, but not method
Cromwell, 2013 ⁶⁶	490	1194	126/129		0.0	0.0	UK	N	Y	Stata Funnel plot, Wilcoxon extended by Cuzick	Y	Y	Y	N	Binomial distribution 95% CI
Dagenais, 2019 ⁶⁷	19	1461	1	14.4	10.5	10.5	US	Y		Hierarchical logistic regression	Y	N	Y	Y	95% CI given, but not method
Davenport, 2020 ⁶⁸	55	25,596	1				US	Y		SAS 9.4 inference testing	Y	N	Y	Y	95% CI given, but not method, though not relevant for mortality
Duclos, 2012 ⁶⁹	28	2357/2904	5	10/32			France	Y		Mixed effects logistic regression	Y	Y	Y	Y	Binomial distribution 95% CI
Eastham, 2003 ⁷⁰	44	4629	2		Other	Other	US	Y		Logistic mixed model	Y	Y	Y	N	None
Eijkenaar, 2013 ⁷¹	447/537	26,684/37,832	N/A	2.5/0.6			Netherlands	Y		Generalized Linear Multilevel Models using SAS 9.2 GLIMMIX	Y	N/A	Y	N	95% CI given, but not method
Eklund, 2009 ⁷²	48	1275	>1		2.1		Sweden	N	Y	RCT Pearson Chi2, Fisher's exact, Cox regression, “Z-test for heterogeneity”	Y	N	Y	Y	None
Faschinger, 2011 ⁷³	17	36,329	1		Other	Other	Austria	N	Not specified	Correlations calculated	Y	N/A	Y	N	None
Fountain, 2004 ⁷⁴	43	876/504	28	7.4			UK	Y		SAS NLMIXED, dealing with convergence issues	Y	N	N	N	Standard error calculated

(Continued)

Table 2 (Continued).

Publication	Doctors	Patients/ Procedures	Institutions	ICC %	Neg Outlier %	Pos Outlier %	Country	MLR*	MV**	Statistical Analysis	PV^	HV^^	DVo#	ODV##	Confidence Interval Calculation
Gani, 2015 ²⁶	56	22,559	1	2.8			US	Y		"[M]ultilevel multivariable logistic regression" Stata 12.1 GLLAM	Y	N/A	Y	Y	95% CI given, but not method
Glance, 2006 ⁷⁵	138	51,750	33	5.9	5.1	8.7	US	Y		Stata 8.2 SAS GLIMMIX	Y	Y	Y	Y	"Quality outliers were identified using 1) the ratio of observed-to-expected mortality rates (O/E ratio) and confidence intervals (CIs) calculated using both parametric (Poisson distribution) and nonparametric (bootstrapping) techniques; and 2) shrinkage estimators."
Glance, 2016 ⁴⁵	420/241	55,436	40	0.5/ 1.8	0.0/3.3	0.0/1.7	US	Y		Hierarchical logistic regression	Y	Y	N	N	95% CI given, but not method
Goodwin, 2013 ⁴²	1099	131,710	268	0.75	0.6	1.5	US	Y		"[H]ierarchical general linear model"	Y	Y	N	N	95% CI given, but not method
Gossl, 2013 ⁷⁶	21	8187	3		0.0	4.8	US	N	Y	Logistic regression	Y	N/A	N	N	Deviation from normal distribution
Grant, 2008 ⁷⁷	31	14,637	4		0.0	0.0	UK	N	Y	SAS 8.2 Logistic regression	Y	N	Y	N	95% CI given, but not method
Gutacker, 2018 ⁴³	212– 3760	24,505– 405,671	30–152	0.4– 12.7			UK	Y		"Three-level hierarchical generalised linear mixed models"	Y	Y	Y	N	None
Hannan, 2017 ⁷⁸	403	27,560	60	12.0	18.6	12.7	US	Y		Hierarchical logistic regression	Y	N	Y	Y	95% CI given, but not method
Harley, 2005 ³¹	143	NS	Multiple		6.3	2.1	UK	N	Y	Multivariate Analysis	N	N	N	N	95% CI given, but not method
Healy, 2017 ²⁵	97	3118/2078	46		10.3/9.3	7.2/4.1	US	Y		"Multi-level mixed-effects logistic regression" Stata 13	Y	N	Y	N	95% CI given, but not method
Hermanek, 1999 ⁷⁹	43	1121	7		9.3	16.3	Germany	N	Y	"Multiple logistic regression analyses"	N	N	N	N	95% CI given, but not method

Hermann, 2002 ⁸⁰	20	16,443	1		Other	Other	Austria	N	N	Chi-square, Brandt and Snedecor contingency tables for binomial distributions	N	N	N	N	None
Hofer, 1999 ⁸¹	232	3642	3	1.0			US	Y		"hierarchical regression for general linear models"	Y	N	N/A	N	None
Hoffman, 2017 ⁸²	1128	183,283	601	6.2			US	Y		"Generalized linear mixed effects models"	Y	Y	Y	Y	Conference abstract ICC CI not specified how
Holmboe, 2010 ⁸³	236	22,526	13 states	12.0			US	Y		SAS 9.1.3 NLMIXED	Y	N	Y	Y	Delta method for 95% CI
Huesch, 2009 ⁸⁴	398	221,327	75		1.2	Other	US	N	Y	Using SEMA by SEMATECH	Y	Y	N	N	Binomial distribution 95% CI
Hyder, 2013 ⁸⁵	575	1488	298	0.3			US	Y		Multilevel Models SAS 9.3	Y	Y	Y	N	95% CI given, but not method
Jemt, 2016 ⁸⁶	23	8808	1		8.7	Other	Sweden	N	N	Chi-square	N	N/A	N	N	None
Johnston, 2010 ⁸⁷	404	55,515	12		Other	Other	UK	N	N	Funnel plots	N	N	Y	N	None
Justiniano, 2019 ⁸⁸	345	1251	118		Other	Other	US	Y		Bayesian hierarchical regression	Y	Y	Y	Y	95% CI given, but not method
Kaczmarek, 2019 ⁸⁹	5337	291,065	NS		17.5	3.7	US	Y		Hierarchical logistic regression SAS 7.1	Y	N	Y	Y	95% CI given, but not method
Kaplan, 2009 ⁹⁰	210	7574		33.0/ 30.6	27.6	43.8	US	Y		Binary mixed models SAS NLMIXED	Y	N	N	N	Standard error calculated
Kerlin, 2018 ⁹¹	345	11,268	104	1.8	22.9	25.2	US	Y		Bayesian hierarchical regression Stata 14.2	Y	Y	Y	Y	Bayesian 95% credible intervals of odds ratios
Kissenberth, 2018 ⁹²	57	1703	NS	44.0			US	N	Y	Linear regression	Y	N	N	N	Conference abstract, no CI
Krein, 2002 ⁹³	258	12,110	9	1.0			US	Y		Multilevel analysis MLwiN 2000	Y	Y	N	N	None
Kunadian, 2009 ⁹⁰	261	149,888	48		1.6	1.1	US	Y		Multivariate Logistic Regression	Y	Y	Y	N	Binomial distribution 95% CI
Landercasper, 2019 ⁹⁴	71	3954	NS		5.7	4.3	US	Y		Mixed effects multivariate model SAS 9.4	Y	N	Y	Y	95% CI given, but not method
LaPar, 2014 ⁹⁵	93	4194	17		Other	Other	US	N	Y	[M]Multivariable, mortality risk-adjusted models with restricted cubic splines	Y	Y	Y	N	None

(Continued)

Table 2 (Continued).

Publication	Doctors	Patients/ Procedures	Institutions	ICC %	Neg Outlier %	Pos Outlier %	Country	MLR*	MV**	Statistical Analysis	PV^	HV^^	DVo#	ODV###	Confidence Interval Calculation
Likosky, 2012 ⁹⁶	32	11,838	8		Other	Other	US	N	Y	Multivariate Logistic Regression	Y	N	N	N	None
Luan, 2019 ⁹⁷	38	1277	21		2.6	15.8	US	Y		Multivariate Mixed Effects Logistic regression Stata 15	Y	Y	Y	N	Bonferroni corrected 95% CI, no further details
Martin, 2013 ⁹⁸	298	6091	43	2.5	Graph too small	Graph too small	US	Y		Logistic regression	Y	Y	Y	N	Bayesian 95% coverage intervals, surgeon performance assumed normally distributed
McCahill, 2012 ⁹⁹	54	2206	4		11.1	31.5	US	Y		Logistic regression	Y	Y	Y	N	95% CI given, but not method
Navar-Boggan, 2012 ¹⁰⁰	47	5979	1		6.4	12.8	US	Y		"Multilevel multivariable random-effects logistic regression" Stata 9	Y	N/A	Y	Y	95% CI given, but not method
O'Connor, 2008 ¹⁰¹	120	2589	18	0.8			US	Y		"Multivariate hierarchical models" MLwiN	Y	Y	Y	Y	None
Orueta, 2015 ¹⁰²	1479	2,207,175	130	4.2			Spain	Y		"Four-level mixed effect models" inc district SAS 9.2 GLIMMIX	Y	Y	Y	Y	95% CI given, but not method
Papachristofi, 2014 ¹⁰³	24/18	18,426	1	0.1/ 2.8	0.0/16.7	0.0/0.0	UK	Y		"Logistic random effects regression" with random effects	Y	N/A	Y	N	95% CI given, but not method
Papachristofi, 2016 ³⁹	190/127	110,769	10	0.25/ 4.0	0.0/15.0	0.0/6.3	UK	Y		"[L]ogistic random-effects regression analysis" using R 3.01	Y	Y	Y	N	95% CI given, but not method for practitioners, comment why no 95% CI for ICC
Papachristofi, 2017 ²³	190/127	107,038	10	0.19/ 2.8	2.1/11.8	0.5/14.2	UK	Y		"Logistic mixed effects models" using R 3.2.2	Y	Y	Y	N	95% CI given, but not method
Quinn, 2018 ¹⁰⁴	2724	123,141	51	2.2	0.2	0.2	US	Y		"3-level crossed random effects logistic regression models" Stata MP 14.2, SAS 9.4	Y	Y	Y	N	"Ninety-five percent CIs were calculated according to Agency for Healthcare Research and Quality methods for risk-adjusted rates."
Rudmik, 2017 ¹⁰⁵	43	2168	Multiple		16.3	4.7	Canada	Y		Logistic regression	Y	N	Y	N	Binomial distribution 95% CI

Selby, 2010 ¹⁰⁶	1005/ 1,049	169,156/ 232,053	35	1.9/ 1.9			US	Y		"Multilevel linear and logistic regression"	Y	Y	N	N	Standard error calculated
Shih, 2015 ¹⁰⁷	345	5033	24	14.0			US	Y		"Hierarchical logistic regression", Stata 12.0	Y	N	Y	N	None
Singh, 2015 ¹⁵	525	48,883	143	15.0	12.8	12.5	US	Y		"[M]ultilevel, multi-variable models"	Y	Y	Y	Y	95% CI given, but not method
Singh, 2018 ²⁴	3987	39,884	NS		10.0/0.1	7.2/0.0	US	Y		Mixed models, SAS GLMM	Y	Y	Y	Y	95% CI given, but not method
Singh, 2019 ²⁸	4230	565,579			0.0	0.1	US	Y		"Multilevel logistic regression" SAS 9.4 GENMOD, GLIMMIX, Stata 15.1 margins	Y	N	Y	N	Formula for 95% CI given and bootstrapping
Thigpen, 2018 ²⁷	34	995	I		5.9	8.8	US	N	Y	"Linear regression model"	Y	N	Y	N	Efron's bootstrap for 95% CI
Tuerk, 2008 ¹⁰⁸	42	1381	I		2.0		US	Y		"Hierarchical linear models" HLM6	Y	N	N	N	ICC as per Bryk Raudenbusch, 95% CI not calculated
Udyavar, 2018 ¹⁰⁹	2149	569,767	224		2.3		US	Y		"Multilevel random effects modelling" Stata 14 MELOGIT	Y	Y	Y	Y	95% CI given, but not method
Udyavar, 2018 ¹¹⁰	175	65,706	31		8.7		US	Y		"[M]ultilevel random effects models" Stata 14	Y	Y	Y	Y	ICC 95% CI not calculated
Udyavar, 2019 ¹¹¹	5816	215,745	198		27.3		US	Y		"[M]ultilevel mixed effects modeling"	Y	Y	Y	Y	Odds ratio 95% CI given, but not method
Verma, 2020 ¹¹²	135	103,085	7		18.5	14.8	Canada	Y		Six different multivariable regression analyses R 3.5	Y	Y	Y	N	95% CI given, but not method
Xu, 2016 ¹¹³	276	2525	44		3.3	0.0	US	Y		"Logistic regression and post-estimation"	Y	Y	Y	Y	None
Xu, 2019 ¹¹⁴	14,598	1,884,842			Other	Other	US	Y		"Multivariable logistic regressions" Stata MP 14	Y	N	Y	Y	95% CI given, but not method

Abbreviations: *MLR, Multi-level regression; **MV, If no MLR, was multivariate regression used? ^PV, Patient variables; ^^HV, Hospital variables; #DVo, Doctors' volume of procedures used; ###ODV, Other doctor variables than volume used.




Figure 1 Estimated probability of in-hospital death within three months of surgery for a patient with average Euro-SCORE risk: (a) surgeons adjusted for centre and anaesthetist; (c) anaesthetists adjusted for centre and surgeon. The horizontal line is average probability (1.8%) for the study cohort. Error bars = 95% CI.

Notes: Reproduced from: Papachristofi O, Sharples LD, Mackay JH, Nashef SAM, Fletcher SN, Klein AA. The contribution of the anaesthetist to risk-adjusted mortality after cardiac surgery. *Anaesthesia*. 2016;71(2):138–146. doi:10.1111/anae.13291.³⁹ © 2015 The Authors. Anaesthesia published by John Wiley & Sons Ltd on behalf of Association of Anaesthetists of Great Britain and Ireland. Creative Commons CC BY (<https://creativecommons.org/about/cclicenses/>).




Figure 2 A funnel plot with each cardiologist represented by a black dot with 95% and 99% confidence intervals. The grey horizontal line is the average mortality for percutaneous coronary intervention (PCI) in New York State 2002–2004.

Notes: Reproduced/used with permission of John Wiley & Sons - Books, from: Kunadian B, Dunning J, Roberts AP, Morley R, de Belder MA. Funnel plots for comparing the performance of PCI performing hospitals and cardiologists: demonstration of utility using the New York hospital mortality data. *Catheter Cardiovasc Interv*. 2009;73(5):589–94. doi:10.1002/ccd.21893.⁴⁰ Copyright © 2009 Wiley-Liss, Inc. Permission conveyed through Copyright Clearance Center, Inc.




Figure 3 This figure was created by the authors and is a higher resolution version of [Figure 2](#) using the same data. It is a funnel plot with each cardiologist represented by a dot with 95% and 99% confidence intervals. Cardiologists whose mortality confidence interval is above the 95% line are marked in red, those below marked in green. **Notes:** Adapted/used with permission of John Wiley & Sons - Books, from: Kunadian B, Dunning J, Roberts AP, Morley R, de Belder MA. Funnel plots for comparing the performance of PCI performing hospitals and cardiologists: demonstration of utility using the New York hospital mortality data. *Catheter Cardiovasc Interv.* 2009;73(5):589–94. doi:10.1002/ccd.21893.⁴⁰ Copyright © 2009 Wiley-Liss, Inc. Permission conveyed through Copyright Clearance Center, Inc.

Example of distribution of doctor performance with ICC of 6.54% and mortality of .63%




Figure 4 A caterpillar plot created by the authors. It uses the same data as [Figures 2 and 3](#). Beige (on left) and brown (on right) confidence intervals have an upper limit above 10%. Green confidence intervals are wholly below average mortality, red confidence intervals wholly above. **Notes:** Data from this publicly available source¹¹⁷ which is the same one as used by Kunadian et al.⁴⁰

identification of high and low outliers and how heterogeneously doctors perform. They also show whether variation in performance is consistent with chance or whether the variation is more significant than that. Fixed effects are calculated through “modelling fixed provider effects”.⁴¹

Random Effects – Estimating the Variation Due to the Doctor

Random effects represent a percentage of the total variation in outcomes between patients that the doctors are responsible for. They are estimated via the intra-class correlation coefficient (ICC), which is the proportion of the total variation in the patient outcome attributed to doctors. There are many different ways to describe this effect.³⁷ The ICCs measured and reported in the studies ranged from 0% to 47% with a median of 3%.

Discussion

This methodological review of studies that report a doctors’ effect on a patient’s physical outcomes has identified wide variations in how researchers measure and report a doctors’ effect. However, there were 2 broad methods identified: fixed effects that allow doctors to be ranked; and random effects where the proportion of variance attributed to unexplained differences between doctors is estimated. The most common statistical model used in the analyses was a multivariable multilevel regression where the types of confounding variables adjusted for included those assessing patient risk, known doctor attributes, and, to a lesser degree, differences between hospitals or institutions.

Glance et al³⁸ discuss in some detail three approaches of provider profiling for binary outcomes, namely conventional logistic regression, hierarchical logistic regression, and fixed-effects logistic regression. They conclude that hierarchical logistic regression is generally preferred, except in the case where providers have low case volume, where hierarchical logistic regression understates the provider effect. We agree that hierarchical logistic regression is an acceptable method for provider profiling as it allows measurement of the strength of the providers’ effect on physical patient health.

This review identified substantial heterogeneity in how the percentage of the variation due to the doctors is reported. For example, Goodwin et al⁴² reported the percentage of the variation for the null model as the “ICC” and the variation calculated after taking all available information into account as “partitioned variance”. It is helpful to calculate the variation of the null model as, if there is negligible or no variation, there is no need to include additional levels in the analysis. In both cases, the null and adjusted models, the ICC was calculated. In contrast, Gutacker et al⁴³ referred to the random effect measure as the “variance partition coefficient”.

A crucial element of reporting fixed effects is the calculation of the confidence intervals of each doctors’ performance. Glance et al^{38,44,45} provide a detailed technical discussion of the respective advantages of using fixed (grading doctors from worst to best) and random effects (calculating the percentage of outcome variation due to the doctor). One pertinent issue discussed is that the smaller the cluster is, ie the fewer patients the doctor has, the greater the shrinkage towards the mean,^{46,47} reducing the calculated ICC, and leading to an underestimate of the difference in performance between doctors.

Interpreting the Doctor’s Effect

The clinical importance of the findings from the studies assessed in this methodological review depends on how common the outcome assessed is and how varied the doctors’ effect is among practitioners. The more common and the more varied, the more the finding is clinically important. The choice between a doctor with an above or below average effect will have implications for the patients’ health outcomes at different levels of how common the outcome is and how strong the doctors’ effect is. The stronger the doctors’ effect and the more important the outcome, the more the choice of doctor matters for the individual. The more common the outcome is, the more the choice of doctor matters for population health.

Table 3 by Baldwin et al,²¹ originally from Wampold et al,⁴⁸ and augmented by Kraemer et al,⁴⁹ shows effect sizes for different ICCs. The intra-class correlation coefficient (ICC) can measure the percentage of the variation in patients’ physical health outcomes due to each component of a medical interaction,²¹ which is typically the patient, the doctor, the hospital, and the intervention. Table 3 shows a scenario where 50% of the patients recover from an intervention when there is no doctors’ effect, ie for an ICC of zero. However, an ICC of 5.9% is reported to produce a medium-sized effect

Table 3 Relationship Between ICC, Cohen's d, Success Rates and NNT

ICC	Cohen's d ⁵⁰	Proportion of Untreated Controls Below Mean of Treated Persons	Success Rate of Untreated Persons	Success Rate of Treated Persons	NNT – Numbers Needed to Treat ⁴⁹
Small					
0.0%	0.0	0.500	0.500	0.500	∞
0.2%	0.1	0.540	0.475	0.525	17.7
1.0%	0.2	0.579	0.450	0.550	8.9
Medium					
2.2%	0.3	0.618	0.426	0.574	6.0
3.8%	0.4	0.655	0.402	0.598	4.5
5.9%	0.5	0.691	0.379	0.621	3.6
8.3%	0.6	0.726	0.356	0.644	3.0
10.9%	0.7	0.758	0.335	0.665	2.6
Large					
13.8%	0.8	0.788	0.314	0.686	2.3

Notes: Cohen's d's aim is to describe the magnitude of response to treatments between two groups, for example, a treatment and a control group. More technically, "The difference between the Treatment and Control group means, divided by the within-group standard deviation".⁵⁰ The Number Needed to Treat (NNT) is defined as the number of patients one would expect to treat with Treatment to have one more success (or one less failure) than if the same number were treated with Control.⁴⁹

(Cohen's d) with a Number Needed to Treat (NNT) of 3.6. Under such circumstances, an ICC of 5.9% can mean that doctors have a clinically significant effect that is greater than many interventions.

Recommendations

How to Report a Doctors' Effect

If researchers wish to report a doctors' effect that has been estimated, we recommend the following:

- Including "doctors" effect' or "physicians" effect' in the keywords and optionally in the title or abstract
- Using multivariable multilevel regression for the analysis with adjustments for patient risk, doctor experience, hospital effects, and any other potential confounding variable
- For describing fixed effects – grading doctors from worst to best, showing individual results for each doctor in a Table or a Figure
- For describing random effects, calculation of the intra-class correlation coefficient (ICC), describing the variation with 95% confidence interval and whether the outcome is a binary or continuous variable
- Making the dataset used for the analysis available for other researchers to conduct their own analyses.

What to Report

Observational Studies

We recommend reporting doctor effects after all available confounding variables have been taken into account, either by (a) the percentage of variation in the patient outcomes which is attributed to the doctor but unexplained by known attributes such as their experience, or (b) the ordering of doctors by their patients' physical health outcomes, or (c) ideally both.

Reporting this data offers the potential to identify both low and high outliers among doctors, as well as how much of an unexplained doctors' effect there is on patient outcomes.

Data Points to Report

Table 4 lists the data points that are recommended to report. Table 5 shows a specific example of those reported data points employing the dataset used in Kunadian et al.⁴⁰

Table 4 Data Points to Report

Data Points to Report	Description
1. Intervention	Type of intervention
2. Type of study	We do not recommend using cross-sectional studies (surveys), as response rates can introduce a selection bias. This does not concern patient-reported data recorded by the doctor, like levels of pain or mobility.
3. Count of doctors	Count of doctors overall. For randomized controlled trials, the count of doctors for each arm.
4. Count of patients or procedures	If available, both patients and procedures. <i>Randomized controlled trials:</i> In addition to the above: number of arms to the study. If the trial is not too large, a matrix showing how many patients of each arm were served by each doctor.
5. Count of higher aggregation, if any – hospital, practices, counties, states	If there are more than two levels, ie not just patients/procedures and doctors, but also hospital, or medical practice, or county, or state, reporting their number could be useful. As there is a well-known hospital effect, distinguishing between hospital and doctors' effects will be useful.
6. Outcome type	The patients' physical health outcomes measured, for example mortality, length of stay, complications, pregnancies, blood pressure or HbA1c levels under control/ not under control. Definitions for each outcome. For example, with mortality, whether it is in-hospital, 30-days, or five years. Whether the outcome is binary, ordinal, or continuous. If feasible, all 30-days, in-hospital, and longer times, if they are available.
7. Percentage of patients/procedures with this outcome	For binary outcomes, the percentage of patients by doctor with that outcome – lowest percentage, highest, mean, and median. For ordinal or continuous outcomes, lowest, highest, average, mean, and median outcome by doctor.
8. Multivariate analysis (Y/N)	Has there been a multivariate analysis, and which variables were considered for exclusion in the analysis, and which were included in the final analysis?
9. Volume effect Y/N/NS (NS="not stated")	Was the number of patients/procedures per doctor included in the analysis? Was the effect, if any, reported as being substantial, not substantial, or not stated?
10. Observed vs expected recorded Y/N/NS	Were investigations done to identify low and high outliers among the doctors, and their count, or proportion recorded? Were funnel plot(s) provided, pointing out 95% and optionally, 90% and or 99% outliers? Alternatively, a caterpillar plot, ie a fixed effect chart showing the patient outcome for each doctor, together with the individual doctor's 95% CI, sorted by patient outcome, showing outliers among doctors. <i>Confidence interval options:</i> ¹¹⁵ Binomial (normal distribution in patient outcomes) Delta method – what are the details, and how is it done? Other – bootstrap, simulation ¹¹⁶
11. Percentage variation number/NS	The variation due to the doctors in the patients' physical health outcome as a percentage of the total variance of all investigated levels, with 95% confidence levels. Optionally, absolute variance and total variance as well.
12. ICC calculated during multilevel, multivariate analysis	As the percentage of the total variance of all investigated levels is the definition of the ICC, reporting of the ICC (intra-class correlation coefficient) as such with 95% confidence intervals as a more detailed alternative to reporting only the variation.
13. Pre-shrinkage ICC calculated through simulation	The ICC calculated in multilevel analysis is often reported as lower than it really is due to shrinkage. ^{46,47} In order to find the pre-shrinkage ICC, the following approach can be taken: Simulated datasets that have the same distribution as the doctor/patient clusters in the data investigated can be generated using increasing ICCs until a generating ICC is found that has the same post-shrinkage ICC as the dataset investigated. Reporting this pre-shrinkage ICC can be valuable, as it can be much larger than the post-shrinkage ICC when, for example, the patients' physical effect is not common (under 10%).

Table 5 Data Points Reported by Kunadian et al

Data Points to Report	Kunadian et al: ⁴⁰ an Example*
1. Type of intervention	Percutaneous Coronary Interventions (PCI) in New York State 2002–2004, also known as angioplasty.
2. Type of study	Cohort study from medical records.
3. Count of doctors	261
4. Count of patients or procedures	149,888 patients, procedures not stated.
5. Count of higher aggregation, if any – hospital, practices, counties, states	48 hospitals
6. Outcome type	30-day and 3-year mortality following PCI.
7. Percentage of patients/procedures with this outcome	Overall, 944 deaths out of 149,888 PCI procedures. After excluding patients listed as “All Other doctors in this hospital”, 912 deaths in 146,781 procedures.
8. Multivariate analysis (Y/N)	Yes. Risk-adjusted mortality rate.
9. Volume effect Y/N/NS (NS=‘not stated’)	Yes. Neither the downloadable paper nor Kunadian state whether there is a volume effect for cardiologists. Kunadian states there is no significant relationship between hospital volume and risk of in-hospital death from these data.
10. Observed vs expected recorded Y/N/NS	Yes.
Were funnel plot(s) provided?	Yes, provided in Kunadian as Figure 2 .
Were caterpillar plots provided?	Not by Kunadian et al ⁴⁰ . See Figure 4 as provided by authors.
Were confidence intervals calculated?	Neither the downloadable document nor Kunadian state how the confidence interval was calculated.
11. Percentage variation Number/NS	NS
12. ICC calculated during multilevel, multivariate analysis	ICC was calculated by the authors of this paper to be 6.54%, 95% CI (4.32%, 9.79%).
13. Pre-shrinkage ICC calculated through simulation	Using simulated data with the same number of doctors, cases per doctor, and deaths per doctor, resulted in an average ICC of 6.48%, 95% CI (4.47%, 9.32%) after 550 simulations. Therefore, there is no substantial shrinkage at work, which is not unexpected as the mean number of cases per doctor is high at 558.

Notes: *Kunadian et al's 2009 paper⁴⁰ refers to a version of the original dataset¹¹⁷ that can be freely downloaded and is sufficiently detailed for our purposes.

Strengths and Limitations

This is the first methodological review on the reporting of doctors' effect on patient outcomes. The clarity and simplicity of how doctors' and surgeons' effects are described here and the suggested standardization of such reporting should allow meta-analysis to be conducted, allow robust identification of outliers, and make the re-analysis of much existing data feasible. However, a limitation is that, as all of the included studies were conducted in North America or Europe, it is unclear whether the findings can be generalized to other regions, particularly in developing nations.

Conclusion

A doctors' effect on patients' physical health can be measured and reported in two ways:

Firstly, by calculating the percentage of variation in patients' physical health outcomes due to the doctor in the form of the intra-class correlation coefficient (ICC). Secondly, by grading doctors from best to worst patients' physical health outcomes, assigning a confidence interval to those outcomes, and reporting how many doctors' confidence intervals fall wholly above or below the overall average. Ideally, both should be reported.

Ethical Approval

As this is a methodological review, no ethical approval was required.

Acknowledgments

The authors thank Dr Aya Ashraf Ali and Tulia Gonzalez Flores for their excellent editorial contributions.

The authors thank Dr Rachel Mascord for her support during the systematic review.

Author Contributions

Both authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Disclosure

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare no support from any organization for the submitted work, no financial relationships with any organizations that might have an interest in the submitted work in the previous three years, and no other relationships or activities that could appear to have influenced the submitted work. The lead author affirms that the manuscript is an honest, accurate, and transparent account of the study being reported, that no important aspects of the study have been omitted, and that any discrepancies from the study as originally planned (and, if relevant, registered) have been explained.

References

1. Moreau A, Boussageon R, Girier P, Figon S. Efficacité thérapeutique de “l’effet médecin” en soins primaires. *Presse Med.* 2006;35(6):967–973. doi:10.1016/S0755-4982(06)74729-7
2. Cook JA, Bruckner T, MacLennan GS, Seiler CM. Clustering in surgical trials - database of intracluster correlations. *Trials.* 2012;132. doi:10.1186/1745-6215-13-2
3. De Vries EN, Ramrattan MA, Smorenburg SM, Gouma DJ, Boermeester MA. The incidence and nature of in-hospital adverse events: a systematic review. *Qual Saf Health Care.* 2008;17(3):216–223. doi:10.1136/qshc.2007.023622
4. Tam VC, Knowles SR, Cornish PL, Fine N, Marchesano R, Etchells EE. Frequency, type and clinical importance of medication history errors at admission to hospital: a systematic review. *CMAJ.* 2005;173(5):510–515. doi:10.1503/cmaj.045311
5. Van Walraven C, Bennett C, Jennings A, Austin PC, Forster AJ. Proportion of hospital readmissions deemed avoidable: a systematic review. *CMAJ.* 2011;183(7):E391–E402. doi:10.1503/cmaj.101860
6. Leppin AL, Gionfriddo MR, Kessler M, et al. Preventing 30-day hospital readmissions: a systematic review and meta-analysis of randomized trials. *JAMA Intern Med.* 2014;174(7):1095–1107. doi:10.1001/jamainternmed.2014.1608
7. Tjarda Van Heek N, Kuhlmann KFD, Scholten RJ, et al. Hospital volume and mortality after pancreatic resection: a systematic review and an evaluation of intervention in The Netherlands. Conference Paper. *Ann Surg.* 2005;242(6):781–790. doi:10.1097/01.sla.0000188462.00249.36
8. Kristensen PK, Merlo J, Ghith N, Leckie G, Johnsen SP. Hospital differences in mortality rates after Hip fracture surgery in Denmark. *Clin Epidemiol.* 2019;11:605–614. doi:10.2147/CLEP.S213898
9. Nilsen SM, Bjørngaard JH, Carlsen F, et al. Hospitals’ discharge tendency and risk of death-an analysis of 60,000 Norwegian Hip fracture patients. *Clin Epidemiol.* 2020;12:173–182. doi:10.2147/CLEP.S237060
10. Krell RW, Girotti ME, Dimick JB. Extended length of stay after surgery: complications, inefficient practice, or sick patients? *JAMA Surg.* 2014;149(8):815–820. doi:10.1001/jamasurg.2014.629
11. Riedl D, Schübler G. The influence of doctor-patient communication on health outcomes: a systematic review. *Z Psychosom Med Psychother.* 2017;63(2):131–150. doi:10.13109/zptm.2017.63.2.131
12. Di Blasi Z, Harkness E, Ernst E, Georgiou A, Kleijnen J. Influence of context effects on health outcomes: a systematic review. *Lancet.* 2001;357(9258):757–762. doi:10.1016/S0140-6736(00
13. Cabana MD, Slish KK, Evans D, et al. Impact of Physician Asthma Care Education on Patient Outcomes. *Health Educ Behav.* 2014;41(5):509–517. doi:10.1177/1090198114547510
14. Thomas KB. General practice consultations: is there any point in being positive? *Br Med J.* 1987;294(6581):1200–1202. doi:10.1136/bmj.294.6581.1200
15. Singh S, Lin YL, Nattinger AB, Kuo YF, Goodwin JS. Variation in readmission rates by emergency departments and emergency department providers caring for patients after discharge. *J Hospital Med.* 2015;10(11):705–710. doi:10.1002/jhm.2407

16. Alruthia Y, Sales I, Almalag H, et al. The relationship between health-related quality of life and trust in primary care physicians among patients with diabetes. *Clin Epidemiol.* 2020;12:143–151. doi:10.2147/CLEP.S236952
17. Sperl-Hillen J, Beaton S, Fernandes O, et al. Comparative effectiveness of patient education methods for type 2 diabetes: a randomized controlled trial. *Arch Intern Med.* 2011;171(22):2001–2010. doi:10.1001/archinternmed.2011.507
18. Walwyn R, Roberts C. Therapist variation within randomised trials of psychotherapy: implications for precision, internal and external validity. *Stat Methods Med Res.* 2010;19(3):291–315. doi:10.1177/0962280209105017
19. Walwyn R, Roberts C. Meta-analysis of absolute mean differences from randomised trials with treatment-related clustering associated with care providers. *Stat Med.* 2015;34(6):966–983. doi:10.1002/sim.6379
20. Wampold BE, Imel ZE. *The Great Psychotherapy Debate: The Evidence for What Makes Psychotherapy Work: Second Edition.* Taylor and Francis Inc; 2015:1–323.
21. Baldwin SA, Imel Z. Therapist effects: findings and methods. *Bergin Garfield's Handbook Psychother Behavior Change.* 2013;6:258–297.
22. Anderson BR, Ciarleglio AJ, Cohen DJ, et al. The Norwood operation: relative effects of surgeon and institutional volumes on outcomes and resource utilization. *Cardiol Young.* 2016;26(4):683–692. doi:10.1017/s1047951115001031
23. Papachristofi O, Klein AA, Mackay J, Nashef S, Fletcher N, Sharples LD. Effect of individual patient risk, centre, surgeon and anaesthetist on length of stay in hospital after cardiac surgery: association of Cardiothoracic Anaesthesia and Critical Care (ACTACC) consecutive cases series study of 10 UK specialist centres. *BMJ Open.* 2017;7(9):e016947. doi:10.1136/bmjopen-2017-016947
24. Singh S, Sparapani R, Wang MC. Variations in 30-day readmissions and length of stay among spine surgeons: a national study of elective spine surgery among US Medicare beneficiaries. *J Neurosurg Spine.* 2018;29(3):286–291. doi:10.3171/2018.1.Spine171064
25. Healy MA, Regenbogen SE, Kanters AE, et al. Surgeon variation in complications with minimally invasive and open colectomy: results from the Michigan surgical quality collaborative. *JAMA Surg.* 2017;152(9):860–867. doi:10.1001/jamasurg.2017.1527
26. Gani F, Lucas DJ, Kim Y, Schneider EB, Pawlik TM. Understanding Variation in 30-Day Surgical Readmission in the Era of Accountable Care: effect of the Patient, Surgeon, and Surgical Subspecialties. *JAMA Surg.* 2015;150(11):1042–1049. doi:10.1001/jamasurg.2015.2215
27. Thigpen CA, Floyd SB, Chapman C, et al. Comparison of surgeon performance of rotator cuff repair: risk adjustment toward a more accurate performance measure. *J Bone Joint Surg Am.* 2018;100(24):2110–2117. doi:10.2106/jbjs.18.00211
28. Singh S, Goodwin JS, Zhou J, Kuo YF, Nattinger AB. Variation among primary care physicians in 30-day readmissions. *Ann Intern Med.* 2019;170(11):749–755. doi:10.7326/m18-2526
29. Kerlin MP, Weissman GE, Wonneberger KA, et al. Validation of administrative definitions of invasive mechanical ventilation across 30 intensive care units. *Am J Respir Crit Care Med.* 2016;194(12):1548–1552. doi:10.1164/rccm.201605-0953LE
30. Beckett DJ, Spears M, Thomson E. Reliable consultant level data from an Acute Medical Unit: a powerful tool for improvement. *J R Coll Physicians Edinb.* 2018;48(2):108–113. doi:10.4997/jrcpe.2018.202
31. Harley M, Mohammed MA, Hussain S, Yates J, Almasri A. Was Rodney Ledward a statistical outlier? Retrospective analysis using routine hospital data to identify gynaecologists' performance. *Br Med J.* 2005;330(7497):929–932. doi:10.1136/bmj.38377.675440.8F
32. Schnelle C, Clark J, Mascord R, Jones M. Is there a surgeons' effect on patients' physical health, beyond the intervention, that requires further investigation?. *Ther Clin Risk Manag.* 2022;1(18):467–490. doi:10.2147/TCRM.S357934
33. Mbuagbaw L, Lawson DO, Puljak L, Allison DB, Thabane L. A tutorial on methodological studies: the what, when, how and why. *BMC Med Res Methodol.* 2020;20(1):226. doi:10.1186/s12874-020-01107-7
34. Higgins JPT TJ, Chandler J, Cumpston M, Li T, Page MJ, Welch VA Cochrane Handbook for Systematic Reviews of Interventions version 6.1 (updated September 2020); 2020. Available from: <https://training.cochrane.org/cochrane-handbook-systematic-reviews-interventions>. Accessed January 17, 2021.
35. Wells GA, Shea B, O'Connell D, et al. *The Newcastle-Ottawa Scale (NOS) for Assessing the Quality of Nonrandomised Studies in Meta-Analyses.* Oxford; 2000.
36. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol.* 2010;25(9):603–605. doi:10.1007/s10654-010-9491-z
37. Allison PD. *Fixed Effects Regression Models. Vol 160. (Quantitative Applications in the Social Sciences.* SAGE publications; 2009.
38. Glance LG, Dick AW. In Response. *Anesth Analg.* 2016;122(5):1722–1727. doi:10.1213/ANE.0000000000001194
39. Papachristofi O, Sharples LD, Mackay JH, Nashef SAM, Fletcher SN, Klein AA. The contribution of the anaesthetist to risk-adjusted mortality after cardiac surgery. *Anaesthesia.* 2016;71(2):138–146. doi:10.1111/anae.13291
40. Kunadian B, Dunning J, Roberts AP, Morley R, de Belder MA. Funnel plots for comparing performance of PCI performing hospitals and cardiologists: demonstration of utility using the New York hospital mortality data. *Catheter Cardiovasc Interv.* 2009;73(5):589–594. doi:10.1002/ccd.21893
41. DeLong ER, Peterson ED, DeLong DM, Muhlbaier LH, Hackett S, Mark DB. Comparing risk-adjustment methods for provider profiling. *Stat Med.* 1997;16(23):2645–2664. doi:10.1002/(SICI)1097-0258(19971215)16:23<2645::
42. Goodwin JS, Lin YL, Singh S, Kuo YF. Variation in length of stay and outcomes among hospitalized patients attributable to hospitals and hospitalists. *J Gen Intern Med.* 2013;28(3):370–376. doi:10.1007/s11606-012-2255-6
43. Gutacker N, Bloor K, Bojke C, Walshe K. Should interventions to reduce variation in care quality target doctors or hospitals? *Health Policy.* 2018;122(6):660–666. doi:10.1016/j.healthpol.2018.04.004
44. Glance LG, Kellermann AL, Hannan EL, et al. RETRACTED The impact of anesthesiologists on coronary artery bypass graft surgery outcomes. *Anesth Analg.* 2015;120(3):526–533. doi:10.1213/ANE.0000000000000522
45. Glance LG, Hannan EL, Fleisher LA, et al. Feasibility of Report Cards for Measuring Anesthesiologist Quality for Cardiac Surgery. *Anesth Analg.* 2016;122(5):1603–1613. doi:10.1213/ane.0000000000001252
46. Hox JJ. *Multilevel Analysis: Techniques and Applications: Second Edition.* Routledge Taylor & Francis Group; 2010:1–382.
47. Raudenbush SW, Bryk AS. *Hierarchical Linear Models: Applications and Data Analysis Methods.* Vol. 1. sage; 2002.
48. Wampold BE. *The Great Psychotherapy Debate: Models, Methods, and Findings.* Lawrence Erlbaum Associates, Inc; 2001.
49. Kraemer HC, Kupfer DJ. Size of treatment effects and their importance to clinical research and practice. *Biol Psychiatry.* 2006;59(11):990–996. doi:10.1016/j.biopsych.2005.09.014
50. Cohen J. *Statistical Power Analysis for the Behavioral Sciences.* 2nd ed. Hillsdale, NJ: Lawrence Earlbam Associates; 1988.

51. Aquina C, Probst C, Hensley B, et al. High variability in nosocomial clostridium difficile infection rates among both surgeons and hospitals following colorectal resection. Conference Abstract. *Dis Colon Rectum*. 2015;58(5):e163. doi:10.1097/01.dcr.0000464773.42498.34
52. Aquina CT, Blumberg N, Probst CP, et al. Significant variation in blood transfusion practice persists following upper gi cancer resection. *J Gastrointest Surg*. 2015;19(11):1927–1937. doi:10.1007/s11605-015-2903-3
53. Aquina CT, Blumberg N, Probst CP, et al. Large variation in blood transfusion use after colorectal resection: a call to action. *Dis Colon Rectum*. 2016;59(5):411–418. doi:10.1097/dcr.0000000000000588
54. Aquina CT, Fleming FJ, Becerra AZ, et al. Explaining variation in ventral and inguinal hernia repair outcomes: a population-based analysis. *Surgery*. 2017;162(3):628–639. doi:10.1016/j.surg.2017.03.013
55. Arvidsson D, Berndsen FH, Larsson LG, et al. Randomized clinical trial comparing 5-year recurrence rate after laparoscopic versus Shouldice repair of primary inguinal hernia. *Br J Surg*. 2005;92(9):1085–1091. doi:10.1002/bjs.5137
56. Becerra AZ, Aquina CT, Berho M, et al. Surgeon-, pathologist-, and hospital-level variation in suboptimal lymph node examination after colectomy: compartmentalizing quality improvement strategies. *Surgery*. 2017;161(5):1299–1306. doi:10.1016/j.surg.2016.11.029
57. Begg CB, Riedel ER, Bach PB, et al. Variations in morbidity after radical prostatectomy. *N Eng J Med*. 2002;346(15):1138–1144. doi:10.1056/NEJMsa011788
58. Bianco FJ, Riedel ER, Begg CB, Kattan MW, Scardino PT. Variations among high volume surgeons in the rate of complications after radical prostatectomy: further evidence that technique matters. *J Urol*. 2005;173(6):2099–2103. doi:10.1097/01.ju.0000158163.21079.66
59. Bianco FJ, Vickers AJ, Cronin AM, et al. Variations among experienced surgeons in cancer control after open radical prostatectomy. *J Urol*. 2010;183(3):977–983. doi:10.1016/j.juro.2009.11.015
60. Bolling SF, Li S, O'Brien SM, Brennan JM, Prager RL, Gammie JS. Predictors of mitral valve repair: clinical and surgeon factors. *Ann Thoracic Surgery*. 2010;90(6):1904–1911. doi:10.1016/j.athoracsur.2010.07.062
61. Bridgewater B, Grayson AD, Jackson M, et al. Surgeon specific mortality in adult cardiac surgery: comparison between crude and risk stratified data. *BMJ*. 2003;327(7405):13–17. doi:10.1136/bmj.327.7405.13
62. Bridgewater B. Mortality data in adult cardiac surgery for named surgeons: retrospective examination of prospectively collected data on coronary artery surgery and aortic valve replacement. *Br Med J*. 2005;330(7490):506–510. doi:10.1136/bmj.330.7490.506
63. Brown EC, Robicsek A, Billings LK, et al. Evaluating Primary Care Physician Performance in Diabetes Glucose Control. *Am J Med Qual*. 2016;31(5):392–399. doi:10.1177/1062860615585138
64. Burns EM, Bottle A, Aylin P, Darzi A, John Nicholls R, Faiz O. Variation in reoperation after colorectal surgery in England as an indicator of surgical performance: retrospective analysis of Hospital Episode Statistics. *BMJ*. 2011;343(7820):d4836. doi:10.1136/bmj.d4836
65. Cirillo F, Patrizio P, Baccini M, et al. The human factor: does the operator performing the embryo transfer significantly impact the cycle outcome? *Human Reproduction*. 2020;35(2):275–282. doi:10.1093/humrep/dez290
66. Cromwell D, Hilton P. Retrospective cohort study on patterns of care and outcomes of surgical treatment for lower urinary-genital tract fistula among English National Health Service hospitals between 2000 and 2009. *BJU Int*. 2013;111(4 B):E257–E262. doi:10.1111/j.1464-410X.2012.11483.x
67. Dagenais J, Bertolo R, Garisto J, et al. Variability in Partial Nephrectomy Outcomes: does Your Surgeon Matter? *Eur Urol*. 2019;75(4):628–634. doi:10.1016/j.eururo.2018.10.046
68. Davenport MS, Khalatbari S, Keshavarzi N, et al. Differences in Outcomes Associated With Individual Radiologists for Emergency Department Patients With Headache Imaged With CT: a Retrospective Cohort Study of 25,596 Patients. *AJR Am J Roentgenol*. 2020:1–8. doi:10.2214/ajr.19.22189
69. Duclos A, Peix JL, Colin C, et al. Influence of experience on performance of individual surgeons in thyroid surgery: prospective cross sectional multicentre study. *BMJ*. 2012;344(7843):d8041. doi:10.1136/bmj.d8041
70. Eastham JA, Kattan MW, Riedel E, et al. Variations among individual surgeons in the rate of positive surgical margins in radical prostatectomy specimens. *J Urol*. 2003;170(6):2292–2295. doi:10.1097/01.ju.0000091100.83725.51
71. Eijkenaar F, van Vliet RC. Profiling individual physicians using administrative data from a single insurer: variance components, reliability, and implications for performance improvement efforts. *Med Care*. 2013;51(8):731–739. doi:10.1097/MLR.0b013e3182992bc1
72. Eklund AS, Montgomery AK, Rasmussen IC, Sandbue RP, Bergkvist LÅ, Rudberg CR. Low Recurrence Rate After Laparoscopic (TEP) and Open (Lichtenstein) Inguinal Hernia Repair: a Randomized, Multicenter Trial With 5-Year Follow-Up. *Ann Surg*. 2009;249(1):33–38. doi:10.1097/SLA.0b013e31819255d0
73. Faschinger C. Quality assessment of cataract surgery of the Department of Ophthalmology, Medical University of Graz. *Spektrum Augenheilkd*. 2011;25(3):215–219. doi:10.1007/s00717-011-0013-5
74. Fountain J, Gallagher J, Brown J. A practical approach to a multi-level analysis with a sparse binary outcome within a large surgical trial. *J Eval Clin Pract*. 2004;10(2):323–327. doi:10.1111/j.1365-2753.2003.00462.x
75. Glance LG, Dick A, Osler TM, Li Y, Mukamel DB. Impact of changing the statistical methodology on hospital and surgeon ranking: the case of the New York State cardiac surgery report card. *Med Care*. 2006;44(4):311–319. doi:10.1097/01.mlr.0000204106.64619.2a
76. Gossli M, Rihal CS, Lennon RJ, Singh M. Assessment of individual operator performance using a risk-adjustment model for percutaneous coronary interventions. *Mayo Clin Proc*. 2013;88(11):1250–1258. doi:10.1016/j.mayocp.2013.07.017
77. Grant SW, Grayson AD, Jackson M, et al. Does the choice of risk-adjustment model influence the outcome of surgeon-specific mortality analysis? A retrospective analysis of 14 637 patients under 31 surgeons. *Heart*. 2008;94(8):1044–1049. doi:10.1136/hrt.2006.110478
78. Hannan EL, Zhong Y, Jacobs AK, et al. Incomplete revascularization for percutaneous coronary interventions: variation among operators, and association with operator and hospital characteristics. *Am Heart J*. 2017;186:118–126. doi:10.1016/j.ahj.2017.01.015
79. Hermanek P. Impact of surgeon's technique on outcome after treatment of rectal carcinoma. *Dis Colon Rectum*. 1999;42(5):559–562. doi:10.1007/bf02234128
80. Hermann M, Alk G, Roka R, Glaser K, Freissmuth M. Laryngeal recurrent nerve injury in surgery for benign thyroid diseases: effect of nerve dissection and impact of individual surgeon in more than 27,000 nerves at risk. *Ann Surg*. 2002;235(2):261–268. doi:10.1097/0000658-200202000-00015
81. Hofer TP, Hayward RA, Greenfield S, Wagner EH, Kaplan SH, Manning WG. The unreliability of individual physician 'report cards' for assessing the costs and quality of care of a chronic disease. *J Am Med Assoc*. 1999;281(22):2098–2105. doi:10.1001/jama.281.22.2098

82. Hoffman RL, Kelz RR, Wirtalla CJ, et al. Variations in surgical outcomes: is it the residency program, the surgeon or the practice venue? Conference Abstract. *J Am Coll Surg.* 2017;225(4):S185.
83. Holmboe ES, Weng W, Arnold GK, et al. The comprehensive care project: measuring physician performance in ambulatory practice. *Health Serv Res.* 2010;45(6 Pt 2):1912–1933. doi:10.1111/j.1475-6773.2010.01160.x
84. Huesch MD. Can managed care plans reliably infer the quality of cardiac surgeons' outcomes? *Am J Manag Care.* Dec. 2009;15(12):890–896.
85. Hyder O, Dodson RM, Nathan H, et al. Influence of patient, physician, and hospital factors on 30-day readmission following pancreatoduodenectomy in the United States. *JAMA Surg.* 2013;148(12):1095–1102. doi:10.1001/jamasurg.2013.2509
86. Jemt T, Olsson M, Renouard F, Stenport V, Friberg B. Early Implant Failures Related to Individual Surgeons: an Analysis Covering 11,074 Operations Performed during 28 Years. *Clin Implant Dent Relat Res.* 2016;18(5):861–872. doi:10.1111/cid.12379
87. Johnston RL, Taylor H, Smith R, Sparrow JM. The Cataract National Dataset electronic multi-centre audit of 55,567 operations: variation in posterior capsule rupture rates between surgeons. *Eye.* 2010;24(5):888–893. doi:10.1038/eye.2009.195
88. Justiniano CF, Aquina CT, Fleming FJ, et al. Hospital and surgeon variation in positive circumferential resection margin among rectal cancer patients. *Am J Surg.* 2019;218(5):881–886. doi:10.1016/j.amjsurg.2019.02.029
89. Kaczmarek K, Wang P, Gilmore R, et al. Surgeon re-excision rates after breast-conserving surgery: a measure of low-value care. *J Am Coll Surg.* 2019;228(4):504–512.e2. doi:10.1016/j.jamcollsurg.2018.12.043
90. Kaplan SH, Griffith JL, Price LL, Pawlson LG, Greenfield S. Improving the reliability of physician performance assessment: identifying the “physician effect” on quality and creating composite measures. *Med Care.* 2009;47(4):378–387. doi:10.1097/MLR.0b013e31818dce07
91. Prasad-Kerlin MP, Epstein A, Kahn JM, et al. Physician-level variation in outcomes of mechanically ventilated patients. *Ann Am Thorac Soc.* 2018;15(3):371–379. doi:10.1513/AnnalsATS.201711-867OC
92. Kissenberth M, Thigpen C, Brooks J, Floyd S, Hawkins RJ, Tokish JM. Comparing surgeon performance of rotator cuff repair: risk adjustment toward a fair performance measure. *Arthroscopy J Arthroscopic Related Surgery.* 2018;34(12):e3. doi:10.1016/j.arthro.2018.10.022
93. Krein SL, Hofer TP, Kerr EA, Hayward RA. Whom should we profile? Examining diabetes care practice variation among primary care providers, provider groups, and health care facilities. *Health Serv Res.* 2002;37(5):1159–1180. doi:10.1111/1475-6773.01102
94. Landercasper J, Borgert AJ, Fayanju OM, et al. Factors associated with reoperation in breast-conserving surgery for cancer: a prospective study of American society of breast surgeon members. *Ann Surg Oncol.* 2019;26(10):3321–3336. doi:10.1245/s10434-019-07547-w
95. LaPar DJ, Ailawadi G, Isbell JM, et al. Mitral valve repair rates correlate with surgeon and institutional experience. *J Thorac Cardiovasc Surg.* 2014;148(3):995–1003. doi:10.1016/j.jtcvs.2014.06.039
96. Likosky DS, Goldberg JB, DiScipio AW, et al. Variability in surgeons' perioperative practices may influence the incidence of low-output failure after coronary artery bypass grafting surgery. *Circ Cardiovasc Qual Outcomes.* 2012;5(5):638–644. doi:10.1161/circoutcomes.112.967091
97. Luan WP, Leroux TC, Olsen C, Robb D, Skinner JS, Richard P. Variation in bariatric surgery costs and complication rates in the military health system. *Mil Med.* 2019. doi:10.1093/milmed/usz454
98. Martin BI, Mirza SK, Franklin GM, Lurie JD, MacKenzie TA, Deyo RA. Hospital and surgeon variation in complications and repeat surgery following incident lumbar fusion for common degenerative diagnoses. *Health Serv Res.* 2013;48(1):1–25. doi:10.1111/j.1475-6773.2012.01434.x
99. McCahill LE, Single RM, Aiello Bowles EJ, et al. Variability in reexcision following breast conservation surgery. *JAMA.* 2012;307(5):467–475. doi:10.1001/jama.2012.43
100. Navar-Boggan AM, Boggan JC, Stafford JA, Muhlbaier LH, McCarver C, Peterson ED. Hypertension control among patients followed by cardiologists. *Circ Cardiovasc Qual Outcomes.* 2012;5(3):352–357. doi:10.1161/circoutcomes.111.963488
101. O'Connor PJ, Rush WA, Davidson G, et al. Variation in quality of diabetes care at the levels of patient, physician, and clinic. *Prev Chronic Dis.* 2008;5(1):A15.
102. Orueta JF, Garcia-Alvarez A, Grandes G, Nuno-Solinis R. The origin of variation in primary care process and outcome indicators: patients, professionals, centers, and health districts. *Medicine.* 2015;94(31):e1314. doi:10.1097/md.0000000000001314
103. Papachristofi O, Mackay JH, Powell SJ, Nashif SAM, Sharples L. Impact of the anesthesiologist and surgeon on cardiac surgical outcomes. *J Cardiothorac Vasc Anesth.* 2014;28(1):103–109. doi:10.1053/j.jvca.2013.07.004
104. Quinn CM, Bilimoria KY, Chung JW, Ko CY, Cohen ME, Stulberg JJ. Creating individual surgeon performance assessments in a statewide hospital surgical quality improvement collaborative. *J Am Coll Surg.* 2018;227(3):303–312.e3. doi:10.1016/j.jamcollsurg.2018.06.002
105. Rudmik L, Xu Y, Alt JA, et al. Evaluating surgeon-specific performance for endoscopic sinus surgery. *JAMA Otolaryngol Head Neck Surg.* 2017;143(9):891–898. doi:10.1001/jamaoto.2017.0752
106. Selby JV, Schmittiel JA, Lee J, et al. Meaningful variation in performance: what does variation in quality tell us about improving quality? *Med Care.* 2010;48(2):133–139. doi:10.1097/MLR.0b013e3181c15a6e
107. Shih T, Cole AI, Al-Attar PM, et al. Reliability of surgeon-specific reporting of complications after colectomy. *Ann Surg.* 2015;261(5):920–925. doi:10.1097/sla.0000000000001032
108. Tuerk PW, Mueller M, Egede LE. Estimating physician effects on glycemic control in the treatment of diabetes. *Diabetes Care.* 2008;31(5):869–873. doi:10.2337/dc07-1662
109. Udyavar R, Cornwell EE, Havens JM, et al. Surgeon-driven variability in emergency general surgery outcomes: does it matter who is on call? *Surgery.* 2018;164(5):1109–1116. doi:10.1016/j.surg.2018.07.008
110. Udyavar NR, Salim A, Havens JM, et al. The impact of individual physicians on outcomes after trauma: is it the system or the surgeon? *J Surg Res.* 2018;229:51–57. doi:10.1016/j.jss.2018.02.051
111. Udyavar NR, Salim A, Cornwell EE, et al. Racial differences in complication risk following emergency general surgery: who your surgeon is may matter. *J Surg Res.* 2019;235:424–431. doi:10.1016/j.jss.2018.05.086
112. Verma AA, Guo Y, Jung HY, et al. Physician-level variation in clinical outcomes and resource use in inpatient general internal medicine: an observational study. *BMJ Qual Saf.* 2020. doi:10.1136/bmjqs-2019-010425
113. Xu T, Makary MA, Al Kazzi E, Zhou M, Pawlik TM, Hutfless SM. Surgeon-level variation in postoperative complications. *J Gastrointest Surg.* 2016;20(7):1393–1399. doi:10.1007/s11605-016-3139-6

114. Xu T, Mehta A, Park A, Makary MA, Price DW. Association Between Board Certification, Maintenance of Certification, and Surgical Complications in the United States. *Am J Med Qual.* 2019;34(6):545–552. doi:10.1177/1062860618822752
115. Ukoumunne OC. A comparison of confidence interval methods for the intraclass correlation coefficient in cluster randomized trials. *Stat Med.* 2002;21(24):3757–3774. doi:10.1002/sim.1330
116. Carpenter J, Bithell J. Bootstrap confidence intervals: when, which, what? A practical guide for medical statisticians. *Stat Med.* 2000;19(9):1141–1164. doi:10.1002/(SICI)1097-0258(20000515)19:9<1141::
117. KI Shine. *Percutaneous Coronary Interventions (PCI) in New York State 2002-2004*. New York State Department of Health; 2006.

Clinical Epidemiology

Dovepress

Publish your work in this journal

Clinical Epidemiology is an international, peer-reviewed, open access, online journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification, systematic reviews, risk & safety of medical interventions, epidemiology & biostatistical methods, and evaluation of guidelines, translational medicine, health policies & economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.

Submit your manuscript here: <https://www.dovepress.com/clinical-epidemiology-journal>