

Genome Analysis of STI *Bartonella henselae*, a Zoonotic Pathogen Causing Endocarditis in an Elderly Patient in China

Xinli Mu^{1,*}, Jianghong Liang^{2,*}, Linyan Qian³, Bing Zhou⁴, Xuehan Zou⁵, Ying Fu⁶, Yongze Zhu², Xi Li², Jiana Shi⁷

¹Department of Infectious Diseases, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang, People's Republic of China; ²Laboratory Medicine Center, Department of Clinical Laboratory, Zhejiang Provincial People's Hospital (Affiliated People's Hospital, Hangzhou Medical College), Hangzhou, Zhejiang, People's Republic of China; ³Heart Center, Department of Cardiovascular Medicine, Zhejiang Provincial People's Hospital (Affiliated People's Hospital, Hangzhou Medical College), Hangzhou, Zhejiang, People's Republic of China; ⁴Heart Center, Department of Cardiovascular Surgery, Zhejiang Provincial People's Hospital (Affiliated People's Hospital, Hangzhou Medical College), Hangzhou, Zhejiang, People's Republic of China; ⁵Center for General Practice Medicine, Department of Infectious Diseases, Zhejiang Provincial People's Hospital, Affiliated People's Hospital, Hangzhou Medical College, Hangzhou, Zhejiang, People's Republic of China; ⁶Department of Clinical Laboratory, Sir Run Run Shaw Hospital, Zhejiang University, Hangzhou, Zhejiang, People's Republic of China; ⁷Center for Clinical Pharmacy, Cancer Center, Department of Pharmacy, Zhejiang Provincial People's Hospital (Affiliated People's Hospital, Hangzhou Medical College), Hangzhou, Zhejiang, People's Republic of China

*These authors contributed equally to this work

Correspondence: Xi Li, Laboratory Medicine Center, Department of Clinical Laboratory, Zhejiang Provincial People's Hospital, Affiliated People's Hospital, Hangzhou Medical College, Hangzhou, Zhejiang, 310014, People's Republic of China, Tel/Fax +86-571-8589-3267, Email lixl_0611@163.com; Jiana Shi, Center for Clinical Pharmacy, Cancer Center, Department of Pharmacy, Zhejiang Provincial People's Hospital, Affiliated People's Hospital, Hangzhou Medical College, Hangzhou, Zhejiang, 310014, People's Republic of China, Tel +86-571-8589-3899, Email shijinn@126.com

Abstract: Infective endocarditis (IE) is a rare disease but with high associated mortality. Currently, the mainstays of diagnosis are still echocardiography and blood cultures. Here, we reported a case of infective endocarditis with negative blood cultures, and blood and aortic valve tissue metagenomic next-generation sequencing (mNGS) results suggested *Bartonella henselae*. In addition, we obtained the whole genomic sequence of *B. henselae* ZJBH strain. To our knowledge, this is the first report of *B. henselae* genomic analysis isolated from clinic in China. Furthermore, we described the whole genome sequencing (WGS) data incorporating all *B. henselae* from diverse sources worldwide and shed light on underlying risk of *B. henselae* transmitted between cats and humans.

Keywords: infective endocarditis, *Bartonella henselae*, metagenomic next-generation sequencing

Introduction

Infective endocarditis (IE), as a disease first recognized in 1885, is defined by infection of a native or prosthetic heart valve, the endocardial surface, or an indwelling cardiac device.^{1–3} Despite advances in diagnostic capabilities and treatment options, IE remains a rare condition but with high associated mortality. And epidemiological studies have shown that 1-year mortality in IE has not improved over the past 20 years.^{4–6} One of the key reasons is that the diagnosis of pathogens is difficult. Current diagnosis is still based on echocardiography and blood culture.⁷ Imaging examinations have irreplaceable value for the diagnosis of IE. Correspondingly, etiological examination also plays an important role in the treatment of IE, especially the formulation of antibiotic regimens.⁸ The commonly described IE-associated pathogens are *Staphylococci*, *Streptococci*, and *Enterococci*, with *Staphylococcus aureus* being the most frequently diagnosed species.⁹ These microorganisms were mostly identified in blood cultures.^{9,10} Notably, blood-culture negative endocarditis (BCNE) is reported to be up to 31% of all cases of IE in the current guidelines.¹¹ For patients with suspected infective endocarditis with negative blood cultures, traditional investigations include serological studies, polymerase chain reaction (PCR) assays of heart valves, and histopathology. In addition, metagenomic next-generation sequencing

(mNGS) is a promising technology that has been widely used in the detection of pathogens in IE in recent years.^{12–16} mNGS is a technology capable of sequencing extremely large numbers of DNA fragments (thousands to millions) simultaneously. A chief advantage of mNGS is unbiased sampling, which enables broad identification of known as well as unexpected pathogens or even the discovery of new organisms.


Bartonella emerged in the form of trench fever in World War I, but the disease it causes has been less well understood. Until the early 1990s, in the form of opportunistic infections in AIDS patients and homeless patients.¹⁷ *Bartonella* species are gram-negative and intracellular pathogens with a unique erythrocytic intracellular lifestyle.¹⁸ They are often found in the gut of obligate blood-feeding arthropod carriers and in the blood of mammalian hosts.¹⁹ It has been reported that its animal host range has steadily expanded in recent years, such as cats, sheep, rodents, humans, and Marine mammals.^{20–23} *Bartonella* endocarditis is a serious disease that can cause clinical complications and has a high mortality rate.^{18,24} *Bartonella henselae* accounts for about one-fourth of *Bartonella* endocarditis cases.²⁵

Here, we report a case of infective endocarditis with negative blood cultures, and blood and aortic valve tissue mNGS results suggested *B. henselae*. In addition, the whole-genome sequence of *B. henselae* were obtained. To our knowledge, this is the first report of *B. henselae* genomic analysis isolated from clinic in China.

Case Presentation

A 64-year-old man was admitted to Zhejiang Provincial People's Hospital from a community (28°70' 66' N; 118°31' 98' E), Yushan county of Jiangxi Province. Fifteen days before presentation, the patient developed chest tightness, shortness of breath, and fatigue during activities, which improved with rest. During the period accompanied by fever, cough, headache, and expectoration. He had a history of smoking and drinking for more than 40 years.

On admission, his vital signs were as follows: blood pressure, 106/51 mmHg; heart rate 81 beats/min; body temperature, 37.7 °C; respiratory rate, 20/min. Physical examination revealed a diastolic murmur in the patient's aortic valve auscultation area. Laboratory test results revealed normal white blood cell count, neutrophil count, C-reactive protein and procalcitonin levels, decreased hemoglobin (103 g/L, normal 130–175 g/L) and albumin (30.8 g/L, normal 40.0–55.0 g/L), increased cardiac troponin I (TnI 0.580 ug/L, normal ≤0.050 ug/L) and brain natriuretic peptide (BNP 1405.4 pg/L, normal ≤119.0 pg/L). Blood cultures were drawn on admission. The next day, echocardiography revealed severe regurgitation of aortic valve with vegetation formation (Figure 1). The patient was diagnosed with infective endocarditis and acute heart failure. The most common pathogen of infective endocarditis outside the hospital is gram-positive bacteria. Ceftriaxone combined with daptomycin was used for anti-infection treatment, and blood cultures were collected at the same time.



A Echocardiography of aortic valve (before operation)

B Echocardiography of aortic valve (after operation)

Figure 1 Echocardiography of aortic valve. (A) Echocardiography of aortic valve (before operation). Yellow circle: aortic coronal valve with a slightly elevated echogenic mass, with a cross-sectional size of approximately 7mm×15mm. (B) Echocardiography of aortic valve (after operation). Normal position and function of the biological valve are, no obvious perivalve leakage.

After 1 week of treatment with the above-mentioned antibiotic regimen, the patient's symptoms of infection, such as low-grade fever, remained unrelieved. In addition, multiple blood/sputum cultures and smears were negative on the first, third, and seventh days of admission. Due to the need for an effective antibiotic regimen, the peripheral blood mNGS examination was performed on the sixth day of admission to clarify the pathogen, which showed: *B. henselae*. Afterward, the patient's contact history was carefully inquired again, and it was found that the patient lived with a juvenile for a long time, which is the most common source of *Bartonella* infection.²⁶ He recalled pet scratches and bites.

Subsequently, the antibiotic regimen was changed to amikacin combined with doxycycline, and aortic valve prosthetic valve replacement was performed. The pathological examination of the aortic valve tissue after surgery showed that the valve tissue had fibrinous necrosis, calcification, with hyaline degeneration, and local infiltration of a large number of inflammatory cells, which was in line with the pathology of infective endocarditis (Figure 2). Concurrently, mNGS of aortic valve tissue was still suggestive of *B. henselae*. After surgery, the patient continued to receive targeted anti-infective treatment with the antibiotic regimen in addition to ceftriaxone treatment, intravenous amikacin (0.2 g/12 hrs) and doxycycline (0.1 g/12 hrs). His symptoms improved quickly, and he finally recovered and was discharged from hospital. At the 4-week and 2-month follow-up, the patient had no sign of heart failure or ongoing infection.

Genomic Analysis of ST I *Bartonella henselae*

PCR-free library preparation and metagenomic sequencing were performed according to the previous study.²⁷ Shotgun sequencing was carried out on illumina Nextseq platform (Illumina Inc., San Diego, CA). The raw reads of *B. henselae* ZJBH were assembled into draft genomes using CLC Genomics Workbench v.10.0 software (QIAGEN, Hilden, Germany; <https://www.qiagenbioinformatics.com/products/clc-genomics-work-bench>). The assembled *B. henselae* ZJBH strain genome, with a size of 1.73 Mb, is similar to the generally observed genome sizes of other *Bartonellae*.

In order to shed some light on the molecular epidemiology and evolutionary relatedness of *B. henselae* around the world, we used whole-genome phylogenetic analyses to examine the phylogenetic relatedness between *B. henselae* and other 38 *B. henselae* strains from the National Center of Biotechnology Information (NCBI) GenBank database (as of 09 October 2022). The basic information of *B. henselae* strains was downloaded from NCBI GenBank in [Supplementary Table 1](#). All *B. henselae* strains obtained from at least six countries (China, France, Denmark, Switzerland, Germany and the USA) could be assigned into five distinct clades. It was noted that *B. henselae* ZJBH strain was the first genome sequence of *B. henselae* isolated from human in China (Figure 3). *B. henselae* ZJBH strain showed close relationship with human and felis catus-derived strains detected in three countries (Denmark, Germany and USA) in the same clade. Furthermore, these *B. henselae* strains belonged to several diverse ST types. A previous report described that different STs may be associated with different clinical manifestations, such as bacillary




Figure 2 Vegetation of aortic valve (HE staining). **(A)** Vegetation of aortic valve, HE staining (×200). Vitreous degeneration, cellulose like necrosis, calcification, and extensive infiltration of inflammatory cells. **(B)** Vegetation of aortic valve, HE staining (×400). Inflammatory cells with a large number of lobulated nuclei (neutrophils) and a small number of plasma cells.




Figure 3 The evolutionary tree of *B. henselae*.
Abbreviations: CSD, cat-scratch disease; NA, not available.

angiomatosis and endocarditis.¹⁶ Multilocus sequence typing (MLST) analysis indicated that the *B. henselae* ZJBH belonged to ST1 using the bioinformatics tools MLST v.2.0 from the Center for Genomic Epidemiology (<http://www.genomicepidemiology.org/>), which is the most frequently found sequence type in China by molecular detection. The other three strains (A20, A71 and F1) collected from cats also belonged to ST1, but they were rarely assigned to the same clade. These findings implicated that *Bartonella* infection among cats could pose a potential risk to human.

Discussion

Bartonella species are facultative intracellular, small fastidious Gram-negative bacilli responsible for one of the rare causes of IE. The genus *Bartonella* contains aerobic or microaerophilic, fastidious, Gram-negative bacilli, and belongs to the alpha-2 subgroup of the class *Proteobacteria*. Of the 13 species known to be associated with human diseases, at least 8 of them can cause infective endocarditis, the most important of which are *B. henselae* and *B. quintana*. The majority, but not all, of *B. henselae* endocarditis cases have a history of contact or interaction with a cat.¹⁸ *Felis catus* plays a vital role since it is the major reservoir of *B. henselae*, and the cat fleas are considered the principal vector for transmission between cats and humans.²⁸ Like our case, IE caused by *B. henselae* may be associated with cats or cat fleas because the patient lived with a cat for a long time and recalled pet scratches and bites. Furthermore, *Bartonella* species have been found in ticks from China,²⁹ and tick-borne *Bartonellosis* in our case is a possibility.

Notably, a recent study described at least two diverse STs of *B. henselae* isolates from cats samples in Brazil by MLST and microbiological analyses.³⁰ Even though *B. henselae* has been extensively studied by serological and molecular detection in humans and cats around the world, the genomic analyses of *B. henselae* and evolutionary trajectory of this zoonotic pathogen among humans and mammals have yet to be elucidated.^{31,32} To our knowledge, this is the first genomic report of *B. henselae* in patient from China. MLST results showed the *B. henselae* ZJBH strain belonged to ST1, which is commonly identified in cats from other countries. Furthermore, we described whole genome sequencing (WGS) data incorporating all *B. henselae* from diverse sources worldwide and shed light on underlying risk of *B. henselae* transmitted between cats and humans. In addition, many virulence factors could contribute to host-specific pathogenesis of *B. henselae* ZJBH strain, mainly including bacterial type IV secretion system (T4SS) proteins VirB and Trw. The VirB/T4SS system in *B. henselae* strain mediates invasion, proinflammatory activation, and anti-apoptotic protection of endothelial cells. Trw type T4SS mediates adhesion to erythrocytes and diversifies the host specificity in *B. henselae*.³³ Overall, our results suggest that the presence of these virulence factors may be an important determinant to the pathogenicity of *B. henselae* strains.

Historically, patients with *Bartonella* IE had a high mortality rate despite surgery,³⁴ so early diagnosis and timely treatment were very necessary. Nowadays, the rapidly developing mNGS technology can rapidly detect unknown pathogenic microorganisms, breaking through the limitations of traditional microbial detection, showing broad prospects in the field of clinical microbiology. Taking the case reported above as an example, the rapid and accurate identification of pathogenic microorganisms through mNGS detection helped us formulate a targeted antibiotic regimen. We also highlight the major role of zoonotic agents and the underestimated role of infective diseases in BCNE. Clinicians should consider *Bartonella* serology, echocardiography and infectious disease consultation when patients present unwell with a history of body lice infestation.

Conclusions

This is the first genomic report of *B. henselae* in patient from China, to the best of our knowledge. Early diagnosis of *Bartonella* spp. infectious endocarditis, is challenging, especially for patients with preexisting valvular heart disease. The epidemiology and management of infective endocarditis are continually changing and many uncertainties remain. Traditional etiological testing has not worked and blood-culture is negative, mNGS examination may be a good option for BCNE. Prompt diagnosis and effective treatment can improve the prognosis.

Abbreviations

IE, Infective endocarditis; mNGS, metagenomic next-generation sequencing; BCNE, blood-culture negative endocarditis; PCR, polymerase chain reaction; NCBI, National Center of Biotechnology Information; MLST, Multilocus sequence typing; WGS, whole genome sequencing.

Data Sharing Statement

The complete genome sequences of the *B. henselae* ZJBH reported here have been deposited at DDBJ/ENA/GenBank under accession no JARVXB000000000. The version described in this paper is the first version JARVXB000000000.

Ethics Approval and Consent for Publication

This case report was approved by Zhejiang Provincial People's Hospital Research Ethics Board (QT2022429). This patient provided consent for publication of the clinical details, and written informed consent was obtained.

Funding

The work was supported by the Research Fund for Applications of High-throughput Sequencing Technology in Infectious Diseases (No. 2024645138), Zhejiang Provincial CONBA Hospital Management Soft Science Research Project (No. 2019ZHA-KEB313) and Zhejiang Provincial Natural Science Foundation of China (LQ21H310004).

Disclosure

The authors declare that they have no competing interests in this work.

References

1. Cahill TJ, Prendergast BD. Infective endocarditis. *Lancet*. 2016;387(10021):882–893. doi:10.1016/S0140-6736(15)00067-7
2. Osler W. The gulstonian lectures, on malignant endocarditis. *Br Med J*. 1885;1(1262):467–470. doi:10.1136/bmj.1.1262.467
3. Shmueli H, Thomas F, Flint N, Setia G, Janjic A, Siegel RJ. Right-sided infective endocarditis 2020: challenges and updates in diagnosis and treatment. *J Am Heart Assoc*. 2020;9(15):e017293. doi:10.1161/JAHA.120.017293
4. Baddour LM, Wilson WR, Bayer AS, et al. Infective Endocarditis In Adults: Diagnosis, Antimicrobial Therapy, And Management Of Complications: A Scientific Statement For Healthcare Professionals From the American Heart Association. *Circulation*. 2015;132(15):1435–1486. doi:10.1161/CIR.0000000000000296
5. Thuny F, Grisoli D, Collart F, Habib G, Raoult D. Management of infective endocarditis: challenges and perspectives. *Lancet*. 2012;379(9819):965–975. doi:10.1016/S0140-6736(11)60755-1
6. Rajani R, Klein JL. Infective endocarditis: a contemporary update. *Clin Med*. 2020;20(1):31–35. doi:10.7861/clinmed.cme.20.1.1
7. Cahill TJ, Baddour LM, Habib G, et al. Challenges in infective endocarditis. *J Am Coll Cardiol*. 2017;69(3):325–344. doi:10.1016/j.jacc.2016.10.066
8. Liesman RM, Pritt BS, Maleszewski JJ, Patel R, Kraft CS. Laboratory diagnosis of infective endocarditis. *J Clin Microbiol*. 2017;55(9):2599–2608. doi:10.1128/JCM.00635-17

9. Habib G, Lancellotti P, Antunes MJ, et al. 2015 ESC Guidelines for the management of infective endocarditis: the task force for the management of infective endocarditis of the European Society of Cardiology (ESC). Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM). *Eur Heart J*. 2015;36(44):3075–3128. doi:10.1093/eurheartj/ehv319
10. Rodríguez-Pastor R, Shafran Y, Knossow N, et al. A road map for in vivo evolution experiments with blood-borne parasitic microbes. *Mol Ecol Resour*. 2022;22(8):2843–2859. doi:10.1111/1755-0998.13649
11. Shi S, Huang Q, Xiao K. Letter to the Editor: rare pathogen of infective endocarditis: *Bartonella henselae*. *Surg Infect*. 2023;24(2):203. doi:10.1089/sur.2022.305
12. Guo Y, Li H, Chen H, et al. Metagenomic next-generation sequencing to identify pathogens and cancer in lung biopsy tissue. *EBioMedicine*. 2021;73. doi:10.1016/j.ebiom.2021.103639
13. Li M, Yan K, Jia P, Wei E, Wang H. Metagenomic next-generation sequencing may assist diagnosis of cat-scratch disease. *Front Cell Infect Microbiol*. 2022;12:946849. doi:10.3389/fcimb.2022.946849
14. Drummond MR, Lania BG, Diniz PPVDP, et al. Improvement of *Bartonella henselae* DNA detection in cat blood samples by combining molecular and culture methods. *J Clin Microbiol*. 2018;56(5):e01732–17. doi:10.1128/JCM.01732-17
15. Next-generation sequencing combined with routine methods to detect the pathogens of encephalitis/meningitis from a Chinese tertiary pediatric neurology center - PubMed. Available from: <https://pubmed.ncbi.nlm.nih.gov/30797792/>. Accessed August 14, 2023.
16. Clinical metagenomic next-generation sequencing for pathogen detection - PubMed. Available from: <https://pubmed.ncbi.nlm.nih.gov/30355154/>. Accessed August 14, 2023.
17. The centenary of the discovery of trench fever, an emerging infectious disease of World War 1 - PubMed. Available from: <https://pubmed.ncbi.nlm.nih.gov/27375211/>. Accessed August 14, 2023.
18. Okaro U, Addisu A, Casanas B, Anderson B. *Bartonella* species, an emerging cause of blood-culture-negative endocarditis. *Clin Microbiol Rev*. 2017;30(3):709–746. doi:10.1128/CMR.00013-17
19. Roden JA, Wells DH, Chomel BB, Kasten RW, Koehler JE, McCormick BA. Hemin binding protein C is found in outer membrane vesicles and protects *Bartonella henselae* against toxic concentrations of hemin. *Infect Immun*. 2012;80(3):929–942. doi:10.1128/IAI.05769-11
20. Ereqat S, Nasereddin A, Vayssier-Taussat M, et al. Molecular evidence of *Bartonella* species in ixodid ticks and domestic animals in Palestine. *Front Microbiol*. 2016;7:1217. doi:10.3389/fmicb.2016.01217
21. Maggi RG, Raverty SA, Lester SJ, et al. *Bartonella henselae* in captive and hunter-harvested beluga (Delphinapterus leucas). *J Wildl Dis*. 2008;44(4):871–877. doi:10.7589/0090-3558-44.4.871
22. Molecular evidence of *Bartonella* species in ixodid ticks and domestic animals in Palestine - PubMed. Available from: <https://pubmed.ncbi.nlm.nih.gov/27540374/>. Accessed August 14, 2023.
23. Hertig MP. Phlebotomus and Carrion's disease. *Am J Trop Med*. 1942;22(Suppl):1–80.
24. Angelakis E, Raoult D. Pathogenicity and treatment of *Bartonella* infections. *Int J Antimicrob Agents*. 2014;44(1):16–25. doi:10.1016/j.ijantimicag.2014.04.006
25. Shapira L, Rasis M, Binsky Ehrenreich I, et al. Laboratory diagnosis of 37 cases of *Bartonella* endocarditis based on enzyme immunoassay and real-time PCR. *J Clin Microbiol*. 2021;59(6):e02217–20. doi:10.1128/JCM.02217-20
26. Brown LD, Maness R, Greer K. Detection of *Bartonella* spp. and Rickettsia spp. in cat fleas (*Ctenocephalides felis*) collected from free-roaming domestic cats in southeastern Georgia, USA. *Vet Parasitol Reg Stud Rep*. 2022;32:100743. doi:10.1016/j.vprsr.2022.100743
27. Luan Y, Hu H, Liu C, et al. A proof-of-concept study of an automated solution for clinical metagenomic next-generation sequencing. *J Appl Microbiol*. 2021;131(2):1007–1016. doi:10.1111/jam.15003
28. Furquim MEC, Do Amaral R, Dias CM, et al. Genetic diversity and multilocus sequence typing analysis of *Bartonella henselae* in domestic cats from Southeastern Brazil. *Acta Trop*. 2021;222:106037. doi:10.1016/j.actatropica.2021.106037
29. Cao XQ, Gu XL, Zhang L, Xu J, Han HJ, Yu XJ. Molecular detection of *Rickettsia*, *Anaplasma*, and *Bartonella* in ticks from free-ranging sheep in Gansu Province, China. *Ticks Tick Borne Dis*. 2023;14(3):102137. doi:10.1016/j.ttbdis.2023.102137
30. Dias CM, Do Amaral RB, Perles L, et al. Multi-locus Sequencing Typing of *Bartonella henselae* isolates reveals coinfection with different variants in domestic cats from Midwestern Brazil. *Acta Trop*. 2023;237:106742. doi:10.1016/j.actatropica.2022.106742
31. Segers FH, Kešnerová L, Kosoy M, Engel P. Genomic changes associated with the evolutionary transition of an insect gut symbiont into a blood-borne pathogen. *ISME J*. 2017;11(5):1232–1244. doi:10.1038/ismej.2016.201
32. Nystedt B, Frank AC, Tholleson M, Andersson SGE. Diversifying selection and concerted evolution of a type IV secretion system in *Bartonella*. *Mol Biol Evol*. 2008;25(2):287–300. doi:10.1093/molbev/msm252
33. Mullins KE, Hang J, Clifford RJ, et al. Whole-genome analysis of *bartonella* ancashensis, a novel pathogen causing Verruga Peruana, Rural Ancash Region, Peru. *Emerg Infect Dis*. 2017;23(3):430–438. doi:10.3201/eid2303.161476
34. Raoult D, Fournier PE, Drancourt M, et al. Diagnosis of 22 new cases of *Bartonella* endocarditis. *Ann Intern Med*. 1996;125(8):646–652. doi:10.7326/0003-4819-125-8-199610150-00004

Infection and Drug Resistance

Dovepress

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>