

Unhealthy Eating Habits and Insomnia Symptoms are Associated with Internet Addiction in Chinese Left-Behind Children: The Gender Difference

Liqing Yao ^{1,2}, Kaixin Liang ^{3,4}, Qiu Zhang⁵, Xinli Chi^{3,4}

¹School of Pharmacy, Guangdong Pharmaceutical University, Guangzhou, People's Republic of China; ²Faculty of Medicine, Medical Sciences Division, Macau University of Science and Technology, Macau, People's Republic of China; ³School of Psychology, Shenzhen University, Shenzhen, People's Republic of China; ⁴The Shenzhen Humanities & Social Sciences Key Research Bases of the Center for Mental Health, Shenzhen University, Shenzhen, People's Republic of China; ⁵The Center for Health Economics and Health Promotion Research, Guangdong Pharmaceutical University, Guangzhou, People's Republic of China

Correspondence: Qiu Zhang, The Center for Health Economics and Health Promotion Research, Guangdong Pharmaceutical University, Guangzhou, 510006, People's Republic of China, Email zhangqiu@gdpu.edu.cn; Xinli Chi, School of Psychology, Shenzhen University, Shenzhen, 518060, People's Republic of China, Email xinlich@szu.edu.cn

Purpose: This study aimed to evaluate the associations between unhealthy eating habits and insomnia symptoms with Internet addiction (IA) among Chinese left-behind children (LBC) based on potential gender difference.

Methods: This cross-sectional study was conducted in junior high schools from an economically disadvantaged area of China in May 2020. Participants were asked to complete the information and scales of socio-demographic characteristics, eating habits (the frequency of eating carbohydrates, fruits, vegetables, protein, and breakfast), insomnia severity and IA. A total of 3156 LBCs (43.9% males and 56.1% females, aged 14.2 years in average) were used for final analyses. Associations between eating habits and insomnia symptoms with IA were analyzed by generalized linear model with binary logistic regression.

Results: The study found that left-behind children (LBCs) had unhealthy eating habits, particularly among females. Females reported more insomnia symptoms than males. The prevalence of IA was similar across genders. Consuming fewer fruits and vegetables and skipping breakfast were linked to increased IA risk, especially in females. Insomnia was significantly associated with higher IA risk in both genders.

Conclusion: In LBCs, unhealthy eating and sleep were notably linked to IA, especially in females. Addressing these factors might reduce the prevalence of IA among LBCs.

Keywords: eating habits, insomnia, Internet addiction, left-behind children, gender difference

Introduction

With the rapid development of science and technology, the Internet has become an increasingly important aspect of human daily life due to its convenience, accessibility and powerful functions.^{1,2} According to the 50th Statistical Report on China's Internet development released by the China Internet Network Information Center (CNNIC) in 2022,³ as of June 2022, the number of the young netizens who were aged 10 to 19 years in China reached 142 million, accounting for 13.5% of the total netizens. However, in the process of experiencing the convenience brought by the Internet, negative effects cannot be ignored, for example, Internet addiction (IA) among adolescents, which has strongly demonstrated a worrisome situation. IA refers to excessive time spent on Internet activities, resulting in behavioral impairment and psychological dysfunction in daily life.⁴ Regarding IA problem among adolescent groups, left-behind children (LBC) may need special academic attention. LBCs refer to the children and adolescents under 18 who are left behind in rural communities and whose one or both parents have been working outside the home for more than six consecutive months.⁵⁻⁷ In general, due to poverty, low socioeconomic status and other risky social situations of the family in original residence, parents have to work remotely.⁸ Hence, these parents occasionally learn about their children's growth through mobile devices (e.g., telephone), which is difficult for them to give timely guidance when their children encounter

problems in the process of growth.⁸ Additionally, LBCs are usually taken care of by their grandparents, who may care less about their psychological needs compared to their parents.⁵ Living in unstable family environments, having inadequate parental care, receiving less mental health education from schools, facing financial difficulties and social discrimination contribute to high risk of LBCs' mental health disorders and problem behaviors.^{5,9,10} Regarding behavioral problem of IA, LBCs are at higher risk for IA than non-LBCs, and shorter time spent at home by parents each year was correlated with a higher rate of IA among LBCs.¹¹ IA could have severe negative effects on LBCs' educational outcome, psychological well-being, and behavioral development.¹² A previous study showed that the proportion of IA was significantly higher among LBCs (20.0%) than that of non-LBCs (12.2%), and another study concluded that IA might be associated with an increased risk of depression in LBCs.^{9,13} Due to the high prevalence of IA in LBCs, it is necessary to pay more attention to the current situation of IA for LBCs and put forward more effective and feasible improvement strategies. The initial step is to explore the factors correlating with IA of LBCs.

Healthy behaviors, which are essential to be promoted at an early age, are possible determinants of IA as healthy behaviors can have a significant impact on the overall health outcomes.¹⁴ Based on healthy behaviors perspective, previous studies have demonstrated that eating habits and sleep status played crucial roles in IA in adolescents. In terms of eating habits, several studies have shown that the presence of disordered eating attitudes and behavior were found to be the strongest predictor variables of IA.¹⁵ A study from France reported that IA was correlated with disordered eating among young participants.¹⁶ Furthermore, a Korean study found that the favorite snacks of high-risk Internet users were confectionery and fast food, which are nutritionally poor; this study also reported a high incidence of meal skipping in individuals with IA symptoms.¹⁷ Regarding the sleep problems, a longitudinal study among children and adolescents from Taiwan indicated that IA was associated with decreased sleep duration during the night-time, increased sleep need and prospectively predicted disturbed circadian rhythm; IA was also predicted by dyssomnias within the follow-up time frame.¹⁸ The evidence from a systematic review reported that participants who had experienced insomnia were about 1.5 times as likely to be a problematic Internet user in comparison to those without sleep problems.¹⁹ As such, unhealthy eating habits and sleep problems (especially insomnia symptoms) may be correlated with the levels of IA among LBCs.

Gender difference may exist in the associations between unhealthy eating habits and sleep problems with IA. A previous study found that LBC boys were at higher risk of skipping breakfast and IA.²⁰ Another previous study indicated that the prevalence of insomnia among LBC girls from rural areas was higher than boys due to the gender discrimination.²¹ However, there is a lack of evidence related to the associations between unhealthy eating habits and insomnia symptoms with IA in Chinese LBCs, and associated studies did not typically compare strengths of these relationships within samples of female vs samples of male. Investigating gender difference is beneficial to analyze targeted associations from more perspectives. Thus, based on the existing studies, this study has two main purposes that address literature gaps related to IA literature in Chinese LBCs: (1) to explore the gender-specific prevalence of IA among Chinese LBCs; (2) to explore the gender-specific associations between unhealthy eating habits and insomnia with IA in Chinese LBCs. Based on the above content, we point out two hypotheses: (1) the gender-specific prevalence of IA was significant in Chinese LBCs; (2) the gender-specific associations between unhealthy eating habits and insomnia with IA were also significant in Chinese LBCs. LBCs are an important vulnerable group in China, exploring the psychological and behavioral problems in LBCs may be a key point to establish guidelines for them to develop their healthy lifestyles, improve their mental health and quality of life. Hence, if both of our hypotheses were supported, this study would provide preliminary evidence that improving diet and sleep quality would be a promising way for reducing the IA problems among Chinese LBCs.

Materials and Methods

Participants and Procedure

The cross-sectional study was conducted in Hechi city, Guangxi province in May 2020. Hechi is one of the most economically underdeveloped places in China with a large number of LBCs.^{7,22} The survey was conducted on junior high-school students in Yizhou district of Hechi, the permission from the local educational department was asked. During the period of the research, there were 30 public junior high schools located in Yizhou with 38,407 students.²³ We randomly recruited 13 local junior high schools (from Grade 7 to Grade 9) to participate in the survey. There were 8939 students providing data. In this study, LBCs were those who reported "Yes" to the question "Whether one or both of your

parents had been absent for more than six consecutive months in the past year since worked in other cities?"^{24–27} After eliminating incomplete and missing responses, 3156 LBCs were successfully matched in our study for final analyses.

This study was a part of annual mental health assessment from the local junior high schools. All students completed our questionnaires via a commonly used online survey platform in China (<https://www.wjx.cn/>). Before filling the questionnaires, the graduate students majoring in Psychology and mental health teachers explained to all students and parents about the purpose of the study, and also asked for their informed consents. The study was approved by the school administrative committees and the Human Research Ethics Committee of Shenzhen University. This study also complied with the Declaration of Helsinki.

Measures

Sociodemographic Characteristics

Information including gender, grade, residence, siblings status, migrants status, family structure, parental education and family income was collected. Previous studies indicated that most of these sociodemographic characteristics included are associated with IA and needed to be adjusted in statistical analyses.^{28–30}

Internet Addiction (IA)

The Chinese version of Young's 10-item Internet Addiction Test, which was developed by Shek et al,³¹ was used to measure severity of Internet addiction which is adjusted specifically for Chinese adolescents. Students participating in the survey selected the options of "Yes" or "No" based on their reality. If the participants expressed 4 or more IA behaviors, they were classified as addicted to the Internet.³¹ The classification of IA in our study referred to 1 = Non-addicted (<4), 2 = Addicted (≥ 4). Confirmatory factor analysis (CFA) with the values of Comparative Fit Index (CFI), Tucker-Lewis Index (TLI), Root Mean Square Error of Approximation (RMSEA), and Standardized Root Mean Square Residual (SRMR) showed good model fits in this sample: CFI = 0.995, TLI = 0.993, RMSEA = 0.056, SRMR = 0.047, and p of chi-square test <0.001, indicating good structural validity. Kuder-Richardson 20 coefficient of 0.89 indicated a great reliability.

Eating Habits

Data on eating habits were collected by 5 items from the sub-scale of the Chinese version of the Health Promoting Lifestyle Profile-II developed by Cao et al.³² Specifically, these items asked about the frequency of eating carbohydrates, fruit, vegetables, protein, and breakfast. All of the dietary contents in this scale are necessities in daily life. All items were scored according to a 4-point Likert scale (1 = Never to 4 = Routinely). CFA indicated good structural validity, with good model fits in this sample: CFI = 0.999, TLI = 0.999, RMSEA = 0.035, SRMR = 0.014, and p of chi-square test <0.001. Cronbach's α coefficient of 0.84 indicated a good reliability.

Insomnia

Insomnia was evaluated using the Chinese version of the Youth Self-Rating Insomnia Scale developed by Liu et al,³³ which consisted of 8 items with each item that is rated on a 5-point scale. A total score that ranged from 8 to 40 can be formed. The recommended grades of insomnia severity refer to 1 = Normal (<22), 2 = Mild (22–25), 3 = Moderate (26–29) and 4 = Severe (≥ 30).³³ CFA showed good model fits in this sample: CFI = 0.979, TLI = 0.970, RMSEA = 0.214, SRMR = 0.106, and p of chi-square test <0.001, indicating good structural validity. Cronbach's α coefficient of 0.90 indicated an excellent reliability.

Statistical Analyses

Analyses were mainly stratified by gender. Descriptive results of sociodemographic characteristics, eating habits, insomnia symptoms and IA were presented as counts (n) and percentages (%). Gender difference in eating habits, insomnia symptoms and IA were firstly examined using chi-square test since all variables were categorical. Partial correlation analysis was used to preliminarily analyze the relationships between eating habits, insomnia symptoms and IA, adjusting for age, grade, sibling status, migrant status, family structure, parental education and family income. Generalized linear model with binary logistic regression was performed for further analysis, adjusted for the covariates same as the partial correlation analyses. The odds ratio (OR) and the 95% confidence interval (CI) were reported. Specifically, we established three models: model 1 included the frequency of eating carbohydrates, fruits, vegetables, and

protein, model 2 included the frequency of eating breakfast, and model 3 included the insomnia severity. Statistical analyses above were performed using JASP 0.16.4 and SPSS 26, $p < 0.05$ was statistically significant.

Results

Participants' Sociodemographic Information

The average age of the LBCs included was 14.2 years old, and there were 1384 (43.9%) males and 1772 (56.1%) females in the sample. More details of the sociodemographic information are summarized in [Table 1](#).

Table 1 Participants' Sociodemographic Information

| Variables | n (%) |
|--------------------------------|-------------|
| Gender | |
| Males | 1384 (43.9) |
| Females | 1772 (56.1) |
| Grade | |
| 7th graders | 1288 (40.8) |
| 8th graders | 1032 (32.7) |
| 9th graders | 836 (26.5) |
| Residence | |
| Urban | 389 (12.3) |
| Rural | 2767 (87.7) |
| Siblings status | |
| Only child | 595 (18.9) |
| Non-only child | 2561 (81.1) |
| Migrants status | |
| Yes | 213 (6.7) |
| No | 2943 (93.3) |
| Family structure | |
| Full | 2480 (78.6) |
| Divorced | 450 (14.3) |
| Other | 226 (7.2) |
| Paternal education | |
| Junior middle school or below | 2590 (82.1) |
| High school or equivalent | 436 (13.8) |
| Bachelor | 86 (2.7) |
| Master or above | 44 (1.4) |
| Maternal education | |
| Junior middle school or below | 2649 (83.9) |
| High school or equivalent | 381 (12.1) |
| Bachelor | 71 (2.2) |
| Master or above | 55 (1.7) |
| Household income (yuan) | |
| Less than 1000 | 405 (12.8) |
| 1000–1999 | 626 (19.8) |
| 2000–2999 | 655 (20.8) |
| 3000–3999 | 699 (22.1) |
| 4000–4999 | 282 (8.9) |
| 5000–5999 | 162 (5.1) |
| More than 6000 | 327 (10.4) |

Gender Difference of Eating Habits, Insomnia Severity, and IA

In Table 2, for eating habits, most participants chose “Sometimes” on all items of eating habits (males: 36.4% (vegetables) to 51.4% (fruits), females: 40.7% (vegetables) to 59.3% (fruits)). The proportion of “Routinely” frequencies that participants reported on these eating habits was low (males: 11.7% (fruits) to 24.1% (breakfast), females: 8.7% (fruits) to 19.6% (carbohydrates)). The proportion of “Routinely” that females reported on all eating habits items was significantly lower compared to males (all $p < 0.01$). For insomnia severity, the proportion of insomnia with moderate severity and above in males and females was 18.3% and 28.2%, respectively. Compared to males, the prevalence of insomnia was significantly higher in females ($p < 0.001$). For the classification of IA, the proportion of “Addicted” participants in males and females was 39.7% and 39.3%, respectively. There is no significant difference of IA between males and females.

Table 2 Gender Difference of Eating Habits, Insomnia Severity and IA

| Variables | Males | | Females | | p |
|-----------------------------|-------|------|---------|------|--------|
| | n | % | n | % | |
| Carbohydrates | | | | | 0.002 |
| Never | 159 | 11.5 | 145 | 8.2 | |
| Sometimes | 510 | 36.8 | 740 | 41.8 | |
| Often | 415 | 30.0 | 540 | 30.5 | |
| Routinely | 300 | 21.7 | 347 | 19.6 | |
| Fruits | | | | | <0.001 |
| Never | 190 | 13.7 | 195 | 11.0 | |
| Sometimes | 712 | 51.4 | 1051 | 59.3 | |
| Often | 320 | 23.1 | 371 | 20.9 | |
| Routinely | 162 | 11.7 | 155 | 8.7 | |
| Vegetables | | | | | 0.008 |
| Never | 128 | 9.2 | 115 | 6.5 | |
| Sometimes | 504 | 36.4 | 722 | 40.7 | |
| Often | 486 | 35.1 | 612 | 34.5 | |
| Routinely | 266 | 19.2 | 323 | 18.2 | |
| Protein | | | | | <0.001 |
| Never | 133 | 9.6 | 109 | 6.2 | |
| Sometimes | 561 | 40.5 | 981 | 55.4 | |
| Often | 457 | 33.0 | 476 | 26.9 | |
| Routinely | 233 | 16.8 | 206 | 11.6 | |
| Breakfast | | | | | <0.001 |
| Never | 146 | 10.5 | 122 | 6.9 | |
| Sometimes | 511 | 36.9 | 873 | 49.3 | |
| Often | 394 | 28.5 | 454 | 25.6 | |
| Routinely | 333 | 24.1 | 323 | 18.2 | |
| Insomnia severity | | | | | <0.001 |
| Normal | 939 | 67.8 | 963 | 54.3 | |
| Mild | 191 | 13.8 | 309 | 17.4 | |
| Moderate | 129 | 9.3 | 245 | 13.8 | |
| Severe | 125 | 9.0 | 255 | 14.4 | |
| Classification of IA | | | | | 0.817 |
| Non-addicted | 834 | 60.3 | 1075 | 60.7 | |
| Addicted | 550 | 39.7 | 697 | 39.3 | |

Abbreviation: IA, Internet addiction.

Correlation Analysis

In Table 3, after adjusting for covariates, the frequencies of eating carbohydrates and protein were significantly negatively correlated to the severity of IA in females. The frequencies of eating fruits, vegetables and breakfast were significantly negatively correlated to the severity of IA in both males and females. Additionally, the severity of insomnia symptoms provided a significantly positive correlation for the severity of IA in both males and females.

Associations Between Eating Habits and Insomnia Severity with IA

In Table 4, the results of the regression model are described with the references of “Routinely” for eating habits groups and “Normal” for insomnia group. The covariates were adjusted when analyzing the data. In males of model 1, the frequency of never eating fruits (OR = 2.252; 95% CI: 1.182, 4.291; $p = 0.014$) and the frequency of sometimes eating

Table 3 Partial Correlations Between Eating Habits and Insomnia Severity with IA

| Variables | IA | |
|-------------------|-----------|-----------|
| | Males | Females |
| Carbohydrates | -0.046 | -0.073** |
| Fruits | -0.085** | -0.151*** |
| Vegetables | -0.096*** | -0.129*** |
| Protein | -0.045 | -0.049* |
| Breakfast | -0.129*** | -0.174*** |
| Insomnia severity | 0.258*** | 0.294*** |

Notes: Adjusted for age, grade, sibling status, migrants status, family structure, parental education and family income. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Abbreviation: IA, Internet addiction.

Table 4 Regression Model of the Associations Between Eating Habits and Insomnia Severity with IA

| Variables | Males | | | | Females | | | |
|---|--------|-------|-------|-------|---------|-------|-------|--------|
| | 95% CI | | | | 95% CI | | | |
| | OR | LL | UL | p | OR | LL | UL | p |
| Model 1 | | | | | | | | |
| Carbohydrates (reference: routinely) | | | | | | | | |
| Never | 0.671 | 0.356 | 1.265 | 0.218 | 1.141 | 0.701 | 1.856 | 0.596 |
| Sometimes | 0.835 | 0.550 | 1.267 | 0.396 | 0.975 | 0.697 | 1.362 | 0.880 |
| Often | 0.735 | 0.487 | 1.110 | 0.143 | 0.896 | 0.640 | 1.256 | 0.525 |
| Fruits (reference: routinely) | | | | | | | | |
| Never | 2.252 | 1.182 | 4.291 | 0.014 | 2.834 | 1.596 | 5.034 | <0.001 |
| Sometimes | 1.150 | 0.696 | 1.900 | 0.585 | 1.855 | 1.170 | 2.941 | 0.009 |
| Often | 1.247 | 0.722 | 2.153 | 0.429 | 1.197 | 0.727 | 1.971 | 0.481 |
| Vegetables (reference: routinely) | | | | | | | | |
| Never | 2.012 | 0.912 | 4.436 | 0.083 | 1.996 | 1.105 | 3.606 | 0.022 |
| Sometimes | 1.729 | 1.086 | 2.755 | 0.021 | 1.504 | 1.028 | 2.202 | 0.036 |
| Often | 1.434 | 0.897 | 2.292 | 0.132 | 1.071 | 0.743 | 1.544 | 0.712 |

(Continued)

Table 4 (Continued).

| Variables | Males | | | | Females | | | |
|--|--------|-------|-------|--------|---------|-------|-------|--------|
| | 95% CI | | | | 95% CI | | | |
| | OR | LL | UL | p | OR | LL | UL | p |
| Protein (reference: routinely) | | | | | | | | |
| Never | 0.638 | 0.321 | 1.269 | 0.200 | 0.553 | 0.288 | 1.061 | 0.075 |
| Sometimes | 1.009 | 0.658 | 1.547 | 0.967 | 0.630 | 0.412 | 0.964 | 0.033 |
| Often | 1.042 | 0.676 | 1.607 | 0.851 | 0.781 | 0.511 | 1.193 | 0.253 |
| Model 2 | | | | | | | | |
| Breakfast (reference: routinely) | | | | | | | | |
| Never | 2.046 | 1.360 | 3.078 | 0.001 | 3.729 | 2.395 | 5.807 | <0.001 |
| Sometimes | 2.136 | 1.586 | 2.876 | <0.001 | 1.914 | 1.452 | 2.523 | <0.001 |
| Often | 1.603 | 1.170 | 2.195 | 0.003 | 0.991 | 0.725 | 1.355 | 0.956 |
| Model 3 | | | | | | | | |
| Insomnia severity (reference: normal) | | | | | | | | |
| Mild | 2.028 | 1.476 | 2.787 | <0.001 | 2.494 | 1.910 | 3.257 | <0.001 |
| Moderate | 3.188 | 2.181 | 4.658 | <0.001 | 3.910 | 2.912 | 5.249 | <0.001 |
| Severe | 4.821 | 3.192 | 7.281 | <0.001 | 4.497 | 3.361 | 6.018 | <0.001 |

Notes: Model 1 included age, grade, sibling status, migrants status, family structure, parental education, family income, and the frequency of eating carbohydrates, fruit, vegetables, and protein. Model 2 included age, grade, sibling status, migrants status, family structure, parental education, family income, and the frequency of eating breakfast. Model 3 included age, grade, sibling status, migrants status, family structure, parental education, family income, and the insomnia severity. The reference category is “routinely” frequency for model 1 and 2, is “normal” for model 3.

Abbreviations: IA, Internet addiction; OR, odds ratio; LL, lower limit; UL, upper limit.

vegetables (OR = 1.729; 95% CI: 1.086, 2.755; $p = 0.021$) were significantly associated with a higher risk of IA. As for females, the frequencies of never and sometimes eating fruits (never: OR = 2.834; 95% CI: 1.596, 5.034; $p < 0.001$) (sometimes: OR = 1.855; 95% CI: 1.170, 2.941; $p = 0.009$), together with the frequencies of never and sometimes eating vegetables (never: OR = 1.996; 95% CI: 1.105, 3.606; $p = 0.022$) (sometimes: OR = 1.504; 95% CI: 1.028, 2.202; $p = 0.036$) were significantly associated with a higher risk of IA. However, the frequency of sometimes eating protein (OR = 0.630; 95% CI: 0.412, 0.964; $p = 0.033$) was significantly associated with a lower risk of IA in females. In males of model 2, the frequencies of never, sometimes and often eating breakfast (never: OR = 2.046; 95% CI: 1.360, 3.078; $p = 0.001$) (sometimes: OR = 2.136; 95% CI: 1.586, 2.876; $p < 0.001$) (often: OR = 1.603; 95% CI: 1.170, 2.195; $p = 0.003$) were all significantly associated with a higher risk of IA compared to the “Routinely” frequencies. As for females, the frequencies of never and sometimes eating breakfast (never: OR = 3.729; 95% CI: 2.395, 5.807; $p < 0.001$) (sometimes: OR = 1.914; 95% CI: 1.452, 2.523; $p < 0.001$) were both associated with a higher risk of IA compared to the “Routinely” frequencies significantly. Lastly, in model 3, severe insomnia symptoms were more strongly associated with higher risk of IA significantly in both genders (all p values < 0.001).

Discussion

This is one of the few studies to determine the associations between eating habits and insomnia symptoms with IA in Chinese LBC sample. It was found that both non-optimal eating habits and insomnia were significantly associated with high risk of IA among LBCs, confirming our hypotheses. Further discussion of our findings is presented below.

In the present study, we found that the dietary patterns among LBCs were undesirable as the proportion of routinely adopting healthy eating habits were low, especially in fruits intake. These results also supported the previous publications that LBCs were at risk of inappropriate food intake and malnutrition as disadvantaged economic conditions, and this may be due to the lack of nutritional knowledge among elderly caregivers compared to younger parents.^{34,35} In line with other

studies, our present research also shows that more female participants reported a higher level of insomnia compared with males.^{36,37} The gender difference in insomnia among LBCs is possibly due to higher vulnerability to stressful life events in females than males.³⁸ It was also explained by the fact that females appear to report poorer sleep quality during periods of hormonal flux such as the luteal phase of the menstrual cycle in adolescents.³⁹ These explanations highlight the necessity to put more effort into improving the sleep quality of female LBCs. Additionally, the prevalence of IA among LBCs is generally high, and more efforts need to be put into guiding LBCs to access the Internet in a healthy way. The prevalence of IA in females (39.3%) was similar to that in males (39.7%), which was similar with a previous study on Chinese general adolescents.⁴⁰

LBCs with deficiencies in fruit and vegetable intakes presented a substantial risk of IA, and the associations were more pronounced in females. A possible explanation is that fruit and vegetable intakes may indirectly affect IA through the associations with mental health problems. Previous studies have proved that having inadequate intakes of fruits and vegetables were associated with higher risk of depression severity, because the deficiencies in minerals and vitamins which were rich in fruits and vegetables may increase the risk of depression.^{7,41} Simultaneously, certain psychological risk factors including depression and anxiety may increase the risk of IA, and persistent depressive symptoms may have long-lasting influence on IA among adolescents especially LBCs.⁴²⁻⁴⁴ In this study, females were more likely to be influenced by nutritional factors in our results. As the associations between fruit/vegetable intakes and depression were stronger in females,⁷ the associations between eating fruits/vegetables and IA were more significant among females compared with males. Moreover, it was found that healthful eating behaviors like eating fruits and vegetables could significantly prevent eating disorders, and the prevalence of IA in students without eating disorders was lower than those with eating disorders.⁴⁵⁻⁴⁷ Therefore, developing healthy eating habits by improving fruit and vegetable consumption may prevent IA through reducing the risk of developing eating disorders.

The current study also noted that moderate intake of protein might be beneficial to the prevention for IA among females. This may be hypothesized that moderate dietary protein could maintain body mass index (BMI) in normal level and prevent obesity, which may be significantly protective factors for IA. Previous studies have reported that individuals with overweight or obesity were at greater risk of IA.^{45,48} This may be explained that individuals (especially females) with overweight or obesity had lower average levels of body appreciation than did individuals with normal BMI, and low levels of body appreciation were related to significantly high severities of depression, anxiety or other mental health problems, which may lead to IA.⁴⁹ However, gender difference in the associations needs more other studies analyzing the relationship between nutrient intakes and IA for further explanations.

Compared with LBCs with keeping eating breakfast, individuals with skipping breakfast may be more prone to IA. This result is consistent with a previous study which conducted in university freshmen group from Taiwan and showed that the habit of skipping breakfast was one of the risk factors of IA.⁵⁰ Nevertheless, there are no exact or common explanations on the mechanisms that the reasons why skipping breakfast is harmful to the prevention for IA or pathological Internet use.^{50,51} Whole grains, dairy products and eggs regularly contained in breakfast are rich in magnesium, calcium, tryptophan and choline, which has been proven to be essential for endogenous serotonin synthesis (a hormone involved in mood regulation) and producing neurotransmitters (an organic base involved in affecting mental health).⁵² It is possible to conjecture that skipping breakfast may reduce intake of these essential nutrients and negatively impact mental health. Besides, skipping breakfast may increase the risk of obesity then cause mental health problems.⁵² Similar to the above findings, mental health problems caused by skipping breakfast may be a risk factor for the problem of IA in LBCs. It is essential to supervise LBCs to develop the good habit of routine eating breakfast, which has a vital role in reducing the prevalence of IA in LBCs.

Insomnia may be a significant pathway to IA among LBCs. LBCs who have low-quality parent-child attachments, are disadvantaged when solving individual problems related to physical development, peer interaction, school adjustment, studying and so on in daily life,⁵³ and these may cause a series of mental health problems including insomnia. Therefore, the possible explanation of this result is that LBCs who are suffering from insomnia symptoms may seek out Internet as a coping mechanism to help regulate their negative emotions (e.g., depressive and anxious emotions from dyssomnia or loneliness), which is similar to another study among other participants.⁵⁴ LBCs will become addicted to Internet unless to

be controlled over time. Conversely, IA also leads to irregular sleep patterns and insomnia, which may form a vicious cycle.¹⁸ Therefore, it is essential to develop monitoring rules about Internet use and bedtimes for LBCs.

This study has several strengths and implications. First, few study has simultaneously examined the associations between eating habits and insomnia symptoms with IA in Chinese LBCs based on the gender difference and explained the possible factors. Our study provides insights on the quality of life of this population, therefore pointing out the need for improvements. Second, this study had a large sample size with the number of 3156, which was helpful to reduce errors and interpret the research findings. Additionally, the study adjusted for possible confounders such as several important sociodemographic factors, allowing for a more accurate interpretation of the main findings. More importantly, our findings suggest that healthy lifestyle behaviors (e.g., healthy diet and good sleep) should be promoted as an important strategy to prevent Chinese LBCs from IA and make sure of their healthy growth. The reasons for parents having to leave their children should be also further investigated and relevant coping strategies should be suggested, which is also beneficial to improve LBCs' physical and mental health.

Several limitations should be considered in our research. First, this is a cross-sectional study and no definite conclusion can be drawn. Therefore, future prospective studies with higher levels of inferential power are needed to provide more insights regarding the hypotheses raised by this study. Second, the data were only collected from Guangxi province, more future data can be collected and analyzed in different provinces to improve the representativeness of the sample. Third, since self-reported measures were used in this study, there might be some response bias and shared method variance. Future studies can consider combining subjective indicators and objective ones to make the results more accurate. For example, wearable devices (e.g., smart watch or bracelet) can be used to assist in monitoring sleep indicators. Fourth, we only measured the frequency of nutrient intake with a lack of more detailed dietary materials such as the quantitative data, which could not rule out the potential confounding factors from overall diet quality and the actual diagnosis may be unapparent. Hence, more accurate and comprehensive dietary measurements, such as semi-quantitative food frequency questionnaires and the method of 24-hour dietary recalls, are recommended to obtain more precise information in future studies. Fifth, the present study only reported the findings of LBCs instead of non-LBCs, which may limit the extensivities of research perspectives. Further studies need to be prepared for supplementing information of non-LBCs in data processing, in order to enhance the richness of similar research findings. Sixth, other possible confounders such as physical health conditions (e.g., congenital and hereditary diseases) and emotional problems (e.g., depression or anxiety) were not considered or controlled in our study. Additionally, no addiction medical history (e.g., drug abuse and eating disorders) that may constitute a vulnerability factor to develop IA has been asked to participants, so that adjustment for these covariates has not been made, either. Finally, although we have some information regarding the parents of these LBCs, more information on the caregivers was not collected. The parameter related to caregivers may constitute another variable to adjust. Indeed, lifestyle behaviors at home routinely depend on caregivers (e.g., grandparents), which may be associated with dietary and sleeping habits, and Internet use. Hence, these possible confounders above are suggested to be considered in future studies to provide a more complete understanding of links among gender, lifestyle, and IA among LBCs.

Conclusion

Lower frequencies of fruit and vegetable intake and breakfast consumption, and higher level of insomnia symptoms were significantly associated with higher risk of IA problem of LBCs in both genders. The associations between unhealthy eating habits and IA were more significant among females than among males. Further studies are still required to confirm the internal mechanisms. Developing healthy lifestyles, for example, improving diet quality (e.g., avoiding skipping breakfast and improving fruit/vegetable intakes) and relieving the symptoms of insomnia, may be beneficial to prevent IA or reduce the prevalence rate among LBCs.

Acknowledgments

Preparation of this article was supported by Guangdong Higher Education Teaching Reform and Construction Project: Research on a New Teaching Model Based on the Internet Environment – A case Study of the Course “Sexual Health Education for College Students” (506), and Humanities and Social Science Research Planning Foundation of the

Ministry of Education: Research on the mechanism and intervention of inhibitory control on the occurrence of short video addiction in adolescents (23YJA190002). We are grateful to all participants in this study sincerely.

Disclosure

The authors report no conflicts of interest in this work.

References

- Cui G, Yin Y, Li S, et al. Longitudinal relationships among problematic mobile phone use, bedtime procrastination, sleep quality and depressive symptoms in Chinese college students: a cross-lagged panel analysis. *BMC Psychiatry*. 2021;21(1):449. doi:10.1186/s12888-021-03451-4
- Wang J, Hao QH, Peng W, et al. Relationship between smartphone addiction and eating disorders and lifestyle among Chinese college students. *Front Public Health*. 2023;11:1111477. doi:10.3389/fpubh.2023.1111477
- China Internet Network Information Center. 50th statistical report on China's internet development; 2022. Available from: <https://www.cnnic.cn/n4/2022/0914/c88-10226.html>. Accessed March 1, 2023.
- Zhou M, Zhu W, Sun X, et al. Internet addiction and child physical and mental health: evidence from panel dataset in China. *J Affect Disord*. 2022;309:52–62. doi:10.1016/j.jad.2022.04.115
- Wu W, Qu G, Wang L, et al. Meta-analysis of the mental health status of left-behind children in China. *J Paediatr Child Health*. 2019;55(3):260–270. doi:10.1111/jpc.14349
- Wang Y, Liu W, Wang W, et al. Left-behind children's social adjustment and relationship with parental coping with children's negative emotions during the COVID-19 pandemic in China. *Int J Psychol*. 2021;56(4):512–521. doi:10.1002/ijop.12754
- Liang K, Chen S, Chi X. Care their diet and mind: association between eating habits and mental health in Chinese left-behind children. *Nutrients*. 2022;14(3):524. doi:10.3390/nu14030524
- Ding L, Yuen LW, Buhs ES, et al. Depression among Chinese left-behind children: a systematic review and meta-analysis. *Child Care Health Dev*. 2019;45(2):189–197. doi:10.1111/cch.12642
- Cai J, Wang Y, Wang F, et al. The association of parent-child communication with internet addiction in left-behind children in China: a Cross-Sectional Study. *Int J Public Health*. 2021;66(630070). doi:10.3389/ijph.2021.630700
- Wong FK, Chang YL, He XS. Correlates of psychological wellbeing of children of migrant workers in Shanghai, China. *Soc Psychiatry Psychiatr Epidemiol*. 2009;44(10):815–824. doi:10.1007/s00127-009-0003-y
- Ren Y, Yang J, Liu L. Social anxiety and internet addiction among rural left-behind children: the mediating effect of loneliness. *Iran J Public Health*. 2017;46(12):1659–1668.
- Fan X. Unpacking the association between family functionality and psychological distress among Chinese left-behind children: the mediating role of social support and internet addiction. *Int J Environ Res Public Health*. 2022;19(20):13327. doi:10.3390/ijerph192013327
- Guo J, Chen L, Wang X, et al. The relationship between internet addiction and depression among migrant children and left-behind children in China. *Cyberpsychol Behav Soc Netw*. 2012;15(11):585–590. doi:10.1089/cyber.2012.0261
- Marques A, Loureiro N, Avelar-Rosa B, et al. Adolescents' healthy lifestyle. *J Pediatr*. 2020;96(2):217–224. doi:10.1016/j.jpmed.2018.09.002
- Alpaslan AH, Koçak U, Avci K, et al. The association between internet addiction and disordered eating attitudes among Turkish high school students. *Eat Weight Disord*. 2015;20(4):441–448. doi:10.1007/s40519-015-0197-9
- Rodgers RF, Melioli T, Laconi S, et al. Internet addiction symptoms, disordered eating, and body image avoidance. *Cyberpsychol Behav Soc Netw*. 2013;16(1):56–60. doi:10.1089/cyber.2012.1570
- Kim Y, Park JY, Kim SB, et al. The effects of internet addiction on the lifestyle and dietary behavior of Korean adolescents. *Nutr Res Pract*. 2010;4(1):51–57. doi:10.4162/nrp.2010.4.1.51
- Chen YL, Gau SS. Sleep problems and internet addiction among children and adolescents: a longitudinal study. *J Sleep Res*. 2016;25(4):13327–13365. doi:10.1111/jsr.12388
- Lam LT. Internet gaming addiction, problematic use of the internet, and sleep problems: a systematic review. *Curr Psychiatry Rep*. 2014;16(4):444. doi:10.1007/s11920-014-0444-1
- Gao Y, Li LP, Kim JH, et al. The impact of parental migration on health status and health behaviours among left behind adolescent school children in China. *BMC Public Health*. 2010;10(1):56. doi:10.1186/1471-2458-10-56
- Tang D, Choi WI, Deng L, et al. Health status of children left behind in rural areas of Sichuan Province of China: a cross-sectional study. *BMC Int Health Hum Rights*. 2019;19(1):4. doi:10.1186/s12914-019-0191-9
- Lu C, Chi X, Liang K, et al. Moving more and sitting less as healthy lifestyle behaviors are protective factors for insomnia, depression, and anxiety among adolescents during the COVID-19 pandemic. *Psychol Res Behav Manag*. 2020;13:1223–1233. doi:10.2147/PRBM.S284103
- Statistics Bureau of Yizhou District, Hechi City. Statistical bulletin of national economic and social development in Yichou District of Hechi City in 2019; 2020.
- Tang W, Wang G, Hu T, et al. Mental health and psychosocial problems among Chinese left-behind children: a cross-sectional comparative study. *J Affect Disord*. 2018;241:133–141. doi:10.1016/j.jad.2018.08.017
- Tang W, Dai Q, Wang G, et al. Impact of parental absence on insomnia and nightmares in Chinese left-behind adolescents: a structural equation modeling analysis. *Child Youth Serv Rev*. 2020;114:105076. doi:10.1016/j.childyouth.2020.105076
- Li X, Coid JW, Tang W, et al. Sustained effects of left-behind experience during childhood on mental health in Chinese university undergraduates. *Eur Child Adolesc Psychiatry*. 2021;30(12):1949–1957. doi:10.1007/s00787-020-01666-6
- Hou T, Xie Y, Mao X, et al. The mediating role of loneliness between social support and depressive symptoms among Chinese rural adolescents during COVID-19 outbreak: a Comparative Study between left-behind and non-left-behind students. *Front Psychiatry*. 2021;12:740094. doi:10.3389/fpsy.2021.740094
- Ibrahim AK, Fouad I, Kelly SJ, et al. Prevalence and determinants of internet addiction among medical students and its association with depression. *J Affect Disord*. 2022;314:94–102. doi:10.1016/j.jad.2022.07.007

29. Martins MV, Formiga A, Santos C, et al. Adolescent internet addiction - role of parental control and adolescent behaviours. *Int J Pediatr Adolesc Med.* 2020;7(3):116–120. doi:10.1016/j.ijpam.2019.12.003
30. Chi X, Hong X, Chen X. Profiles and sociodemographic correlates of internet addiction in early adolescents in southern China. *Addict Behav.* 2020;106:106385. doi:10.1016/j.addbeh.2020.106385
31. Shek DT, Tang VM, Lo CY. Internet addiction in Chinese adolescents in Hong Kong: assessment, profiles, and psychosocial correlates. *ScientificWorldJournal.* 2008;8:776–787. doi:10.1100/tsw.2008.104
32. Cao W, Guo Y, Ping W, et al. Development and psychometric tests of a Chinese version of the HPLP-II scales. *Chin J Dis Control Prev.* 2016;20(03):286–289.
33. Liu X, Yang Y, Liu -Z-Z, et al. Psychometric properties of Youth Self-Rating Insomnia Scale (YSIS) in Chinese adolescents. *Sleep Biol Rhythms.* 2019;17(3):339–348. doi:10.1007/s41105-019-00222-3
34. Bel-Serrat S, von der Schulenburg A, Marques-Previ M, et al. What are the determinants of vegetable intake among adolescents from socioeconomically disadvantaged urban areas? A systematic review of qualitative studies. *Int J Behav Nutr Phys Act.* 2022;19(1):158. doi:10.1186/s12966-022-01396-9
35. Mo X, Xu L, Luo H, et al. Do different parenting patterns impact the health and physical growth of ‘left-behind’ preschool-aged children? A cross-sectional study in rural China. *Eur J Public Health.* 2016;26(1):18–23. doi:10.1093/eurpub/ckv181
36. Chi X, Liang K, Chen ST, et al. Mental health problems among Chinese adolescents during the COVID-19: the importance of nutrition and physical activity. *Int J Clin Health Psychol.* 2021;21(3):100218. doi:10.1016/j.ijchp.2020.100218
37. Brodar KE, La Greca AM, Hysing M, et al. Stressors, repetitive negative thinking, and insomnia symptoms in adolescents beginning high school. *J Pediatr Psychol.* 2020;45(9):1027–1038. doi:10.1093/jpepsy/jsaa064
38. Suh S, Cho N, Zhang J. Sex differences in insomnia: from epidemiology and etiology to intervention. *Curr Psychiatry Rep.* 2018;20(9):69. doi:10.1007/s11920-018-0940-9
39. Li SH, Graham BM, Werner-Seidler A. Gender differences in adolescent sleep disturbance and treatment response to smartphone app-delivered cognitive behavioral therapy for insomnia: Exploratory Study. *JMIR Form Res.* 2021;5(3):e22498. doi:10.2196/22498
40. Shen Y, Wang L, Huang C, et al. Sex differences in prevalence, risk factors and clinical correlates of internet addiction among Chinese college students. *J Affect Disord.* 2021;279:680–686. doi:10.1016/j.jad.2020.10.054
41. Liu MW, Chen QT, Towne SD Jr, et al. Fruit and vegetable intake in relation to depressive and anxiety symptoms among adolescents in 25 low- and middle-income countries. *J Affect Disord.* 2020;261:172–180. doi:10.1016/j.jad.2019.10.007
42. Veisani Y, Jalilian Z, Mohamadian F. Relationship between internet addiction and mental health in adolescents. *J Educ Health Promot.* 2020;9:303. doi:10.4103/jehp.jehp_362_20
43. Bu H, Chi X, Qu D. Prevalence and predictors of the persistence and incidence of adolescent internet addiction in Mainland China: a two-year longitudinal study. *Addict Behav.* 2021;122:107039. doi:10.1016/j.addbeh.2021.107039
44. Gao T, Qin Z, Hu Y, et al. Trajectories of depression and anxiety in Chinese high school freshmen: associations with Internet addiction. *J Affect Disord.* 2021;286:180–186. doi:10.1016/j.jad.2021.02.074
45. Tayhan Kartal F, Yabancı Ayhan N. Relationship between eating disorders and internet and smartphone addiction in college students. *Eat Weight Disord.* 2021;26(6):1853–1862. doi:10.1007/s40519-020-01027-x
46. Chung A, Vieira D, Donley T, et al. Adolescent peer influence on eating behaviors via social media: scoping review. *J Med Internet Res.* 2021;23(6):e19697. doi:10.2196/19697
47. Yu Z, Tan M. Disordered eating behaviors and food addiction among nutrition major college students. *Nutrients.* 2016;8(11):673. doi:10.3390/nu8110673
48. Eliacik K, Bolat N, Koçyiğit C, et al. Internet addiction, sleep and health-related life quality among obese individuals: a comparison study of the growing problems in adolescent health. *Eat Weight Disord.* 2016;21(4):709–717. doi:10.1007/s40519-016-0327-z
49. Liu ZH, Cai H, Bai W, et al. Gender differences in body appreciation and its associations with psychiatric symptoms among Chinese college students: a nationwide survey. *Front Psychiatry.* 2022;13:771398. doi:10.3389/fpsy.2022.771398
50. Tsai HF, Cheng SH, Yeh TL, et al. The risk factors of internet addiction—a survey of university freshmen. *Psychiatry Res.* 2009;167(3):294–299. doi:10.1016/j.psychres.2008.01.015
51. Yamada M, Sekine M, Tatsuse T, et al. Prevalence and associated factors of pathological internet use and online risky behaviors among Japanese elementary school children. *J Epidemiol.* 2021;31(10):537–544. doi:10.2188/jea.JE20200214
52. Zahedi H, Djalalinia S, Sadeghi O, et al. Breakfast consumption and mental health: a systematic review and meta-analysis of observational studies. *Nutr Neurosci.* 2022;25(6):1250–1264. doi:10.1080/1028415X.2020.1853411
53. Shuang M, Yiqing W, Ling J, et al. Relationship between parent-child attachment and depression among migrant children and left-behind children in China. *Public Health.* 2022;204:1–8. doi:10.1016/j.puhe.2021.12.015
54. Tavernier R, Willoughby T. Sleep problems: predictor or outcome of media use among emerging adults at university? *J Sleep Res.* 2014;23(4):389–396. doi:10.1111/jsr.12132

Psychology Research and Behavior Management

Dovepress

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior modification and management; Clinical applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/psychology-research-and-behavior-management-journal>