

Prevalence of Risk Factors for Hypertension Among Faculty at an Urban University in Uganda [Letter]

Elanda Fikri ^{1,2}

¹Department of Environmental Health, Poltekkes Kemenkes Bandung, Bandung, Indonesia; ²Center of Excellence on Utilization of Local Material for Health Improvement, Bandung Health Polytechnic, Bandung, Indonesia

Correspondence: Elanda Fikri, Department of Environmental Health, Poltekkes Kemenkes Bandung, Jl. Pajajaran 56, Bandung, Jawa Barat, Indonesia, Email elandafikri@yahoo.com

Dear editor

We have read a research article entitled “Prevalence of Risk Factors for Hypertension Among Faculty at an Urban University in Uganda” by Alinaitwe et al,¹ recently published in Integrated Blood Pressure Control Journal. We congratulate the authors on this successful article and make some contributions. There are four strengths of this study: 1) This is the first study conducted at a university in Uganda to assess the risk profile of hypertension among teaching staff, thus making an important contribution to existing knowledge, 2) Involved 141 teaching staff from five of the ten faculties at Makerere University, thus covering a diverse range of academic positions, 3) Data were collected using a semi-structured questionnaire developed based on the STEP-wise approach to NCD surveys by WHO, indicating that the study used a standardized tool, 4) Provides implications for the development of hypertension risk screening tools in the formal sector of work, as well as emphasizing the importance of prevention and health promotion to reduce the risk burden of hypertension.

However, we have also discovered several limitations that need to be corrected in the future, namely: 1) Most of the data collected on participants' behavior was based on self-reported assessments, which can be prone to response bias, 2) Some other indicators of increased risk factors for hypertension, such as diabetes and blood cholesterol levels, were not evaluated in this study, thus limiting the understanding of the total risk profile, 3). This study used blood pressure measurements taken on only one day, which may not be definitive enough to diagnose hypertension, 4) The specific study population, ie university teaching staff, may not be able to directly generalize these findings to the general population in Uganda.

To obtain better results, we recommend that further research be carried out by 1) There is a need to develop and test hypertension risk screening tools in the formal work sector to promote primary and secondary prevention of hypertension,² 2) Further studies could expand the scope of the study to include additional risk indicators such as diabetes³ and blood cholesterol levels, thus providing a more comprehensive understanding of the risk profile, 3) Further evaluation of factors inhibiting physical activity among teaching staff and design of targeted interventions to promote physical exercise as a preventive measure for hypertension,⁴ 4) Consider blood pressure measurements taken over a wider time span to gain a more comprehensive understanding of hypertension status.

In conclusion, this study is one of the first to demonstrate the prevalence of hypertension risk factors among university teaching staff in Uganda, as well as confirm the association between modifiable risk factors such as obesity, physical activity, and consumption of added salt with hypertension. The study also highlights the need for routine screening efforts and health promotion campaigns to reduce the risk burden of hypertension among this population.

Disclosure

There is no conflict of interest related to this communication.

References

1. Alinaitwe B, Amany C, Muwanguzi PA, Ngabirano TD. Prevalence of risk factors for hypertension among faculty at an urban university in Uganda. *Integr Blood Press Control*. 2024;17:1–11. doi:10.2147/IBPC.S440972
2. Schmidt BM, Duraõ S, Toews I, et al. Screening strategies for hypertension. *Cochrane Database Syst Rev*. 2020;5(5):CD013212. doi:10.1002/14651858.cd013212.pub2
3. Petrie JR, Guzik TJ, Touyz RM. Diabetes, hypertension, and cardiovascular disease: clinical insights and vascular mechanisms. *Can J Cardiol*. 2018;34(5):575–584. doi:10.1016/j.cjca.2017.12.005
4. Diaz KM, Shimbo D. Physical activity and the prevention of hypertension. *Curr Hypertens Rep*. 2013;15(6):659–668. doi:10.1007/s11906-013-0386-8

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Integrated Blood Pressure Control 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Integrated Blood Pressure Control editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Integrated Blood Pressure Control

Dovepress

Publish your work in this journal

Integrated Blood Pressure Control is an international, peer-reviewed open-access journal focusing on the integrated approach to managing hypertension and risk reduction. Treating the patient and comorbidities together with diet and lifestyle modification and optimizing healthcare resources through a multidisciplinary team approach constitute key features of the journal. This journal is indexed on American Chemical Society's Chemical Abstracts Service (CAS). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/integrated-blood-pressure-control-journal>

<https://doi.org/10.2147/IBPC.S458680>