

# Statin-Associated Liver Dysfunction and Muscle Injury: epidemiology, Mechanisms, and Management Strategies

Ming-Ming Ma<sup>1,\*</sup>, Yao-Yao Xu<sup>2,\*</sup>, Li-Hua Sun<sup>1,\*</sup>, Wen-Jie Cui<sup>1</sup>, Miao Fan<sup>3</sup>, Su Zhang<sup>4</sup>, Lei Liu<sup>5</sup>, Ling-Zhi Wu<sup>6</sup>, Liu-Cheng Li<sup>7</sup>

<sup>1</sup>Department of Diagnostic Ultrasound & Echocardiography, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, 310016, People's Republic of China; <sup>2</sup>Department of Pharmacy, The People's Hospital of Pingyang, Wenzhou, 325400, People's Republic of China; <sup>3</sup>Department of Pharmacy, The First Affiliated Hospital of Ningbo University, Ningbo, 315010, People's Republic of China; <sup>4</sup>Center for Clinical Pharmacy, Cancer Center, Department of Pharmacy, Zhejiang Provincial People's Hospital, Affiliated People's Hospital, Hangzhou Medical College, Hangzhou, 310014, People's Republic of China; <sup>5</sup>Department of Orthopaedics, Shaoxing Hospital of Traditional Chinese Medicine, Shaoxing, 312000, People's Republic of China; <sup>6</sup>Department of Pharmacy, The Affiliated Hospital of Jiaxing University (The First Hospital of Jiaxing), Jiaxing, 314000, People's Republic of China; <sup>7</sup>Department of Pharmacy, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, 310016, People's Republic of China

\*These authors contributed equally to this work

Correspondence: Liu-Cheng Li, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, No. 3 Qingchun East Road, Hangzhou, 310016, People's Republic of China, Email 3415116@zju.edu.cn; Ling-Zhi Wu, The Affiliated Hospital of Jiaxing University (The First Hospital of Jiaxing), No. 1882 South Central Road, Jiaxing, 314000, People's Republic of China, Email wulingzhi1992@163.com

**Abstract:** Surveillance of drug safety is an important aspect in the routine medical care. Adverse events caused by real-world drug utilization has become one of the leading causes of death and an urgent issue in the field of toxicology. Cardiovascular disease is now the leading cause of fatal diseases in most countries, especially in the elderly population who often suffer from multiple diseases and need long-term multidrug therapy. Among which, statins have been widely used to lower bad cholesterol and regress coronary plaque mainly in patients with hyperlipidemia and atherosclerotic cardiovascular diseases (ASCVD). Although the real-world benefits of statins are significant, different degrees and types of adverse drug reactions (ADR) such as liver dysfunction and muscle injury, have a great impact on the original treatment regimens as well as the quality of life. This review describes the epidemiology, mechanisms, early identification and post-intervention of statin-associated liver dysfunction and muscle injury based on the updated clinical evidence. It provides systematic and comprehensive guidance and necessary supplement for the clinical safety of statin use in cardiovascular diseases.

**Keywords:** statin, cardiovascular diseases, liver dysfunction, muscle injury, countermeasures

## Introduction

Drug safety, an urgent problem in clinical pharmacology and toxicology, is one of the primary health programs.<sup>1,2</sup> Drug exposure inevitably produces negative effects such as adverse drug reaction (ADR), which is related to many influencing factors such as age, drug categories, and route of administration.<sup>3</sup> ADR could damage patients' health, even delay the disease process, or lead to fatal consequences. The drug safety issues have become increasingly prominent.

In recent years, the disease burden from cardiovascular diseases consistently ranks the primary contributor among all diseases.<sup>4</sup> There are many types of cardiovascular diseases with complex treatment strategies which include lipid-lowering drugs such as statins or its combination with other drugs. Recent pairwise, network, and dose-response meta-analyses of 47 randomized controlled trials showed that statins decreased the risk of myocardial infarction (OR = 0.66, 95% CI: 0.61–0.71, P < 0.001), stroke (OR = 0.78, 95% CI: 0.72–0.84, P < 0.001), death from cardiovascular diseases (OR = 0.77, 95% CI: 0.72–0.83, P < 0.001) and all-cause death (OR = 0.83, 95% CI: 0.79–0.88, P < 0.001).<sup>5</sup> Statin combined with other lipid-lowering drug therapy could also significantly improve the anatomical and physiological

function of the coronary arteries and down-regulate the re-hospitalization rate due to unstable angina in ST elevation myocardial infarction patients with non-infarct-related artery.<sup>6</sup> A retrospective study found that statins reduced the intubation, ICU admission, and mortality in COVID-19 patients.<sup>7</sup> These evidence reveal the importance of statins in lowering cardiovascular diseases risk and potential additional benefits of comorbidities.

However, the issues of ADR-related drug safety of statins have also aroused great concerns.<sup>8</sup> ADR not only causes additional harm to patients, but also greatly interferes with clinical routine treatment. It is necessary to put on the agenda to systematically manage the use of statins in clinical practice. This study describes the typical toxic consequences and epidemiological characteristics of statin-associated liver dysfunction and muscle injury in clinical practice. Besides, detailed countermeasures accordingly are presented to minimize the risks of statins use and ensure timely medication schedule adjustments for patients.

## Epidemiology of Statins Use

Statins are 3-hydroxy-3-methylglutaryl-Coenzyme A (HMG-CoA) reductase inhibitors that reduce bad lipid levels by inhibiting the biosynthesis of cholesterol.<sup>9</sup> Several drugs including simvastatin, pravastatin, atorvastatin, and rosuvastatin, are currently on the market to reduce the bad cholesterol such as low-density lipoprotein cholesterol (LDL-C). Clinically, it is mainly applicable to atherosclerotic cardiovascular diseases (ASCVD) and hyperlipidemia. Statins have also become the main drug class for the primary and secondary prevention and treatment of cardiovascular and cerebrovascular diseases.

In recent years, statins have been increasingly used. It is reported that the use of statins in the primary prevention of cardiovascular disease in patients over 75 years old was as high as 62.6% in USA.<sup>10</sup> In a retrospective analysis of more than 125,000 patients receiving post-carotid endarterectomy in USA, statin use gradually increased from 36.2% to 62% during 2013 to 2021.<sup>11</sup> New statin prescriptions were associated with higher rates of diabetes, coronary heart disease, stroke, and transient ischemic attack.<sup>11</sup> A 12-month prospective cross-sectional study of 356 patients with acute coronary syndrome demonstrated that the most prescribed drug was statins (93%), followed by antiplatelet drugs and anticoagulants in Saudi Arabia.<sup>12</sup> In a database from Australia's Pharmaceutical Benefits Scheme, the prevalence of long-term statins users over the age of 65 had increased 35.6% with ten years.<sup>13</sup> Meanwhile, atorvastatin is the most commonly used statin in each year and the new use of high-intensity statins had also increased from 23.6% to 30.5% in a 10-year trend analysis.<sup>13</sup> In China, the overall rate of statins prescription for ASCVD patients was as high as 58.8%.<sup>14</sup>

These studies demonstrate that the reported use of statins has been rising in different countries and diverse indications in recent years. However, such scale of real-world use also exposes the risk of of statin-associated liver dysfunction and myopathy, suggesting that these patients need to be more closely concerned and monitored.<sup>15</sup>

## Liver Dysfunction Caused by Statins

Liver dysfunction had attracted considerable attentions in a large number of drug types such as statins. The auxiliary examination methods for diagnosis of liver injury including B-mode ultrasound, hepatic arteriography, pathology, and hepatic enzyme examination. The deteriorated liver function, especially the increase of serum alanine aminotransferase (ALT) and/or aspartate aminotransferase (AST), is one of the most common side effects of statins. It is of great significance to correctly understand and evaluate the liver dysfunction caused by statins during the use of statins.

In vitro experiments have found that the damage of liver mitochondria caused by statins is time-dependent and dose-dependent.<sup>16</sup> Statins can result in a decrease in membrane potential and a significant increase in superoxides in mitochondria, which then lead to serious mitochondrial damage.<sup>17</sup> Some statins (atorvastatin, simvastatin, and lovastatin) are metabolized by the enzyme CYP3A4 in the liver. When it is used in large doses or in combination with CYP3A4 inhibitors (such as macrolides), both of the drug concentration and the liver load increase significantly, which can directly cause liver damage.<sup>15</sup> In addition, gene polymorphisms of statin metabolizing enzymes are also associated with liver dysfunction.<sup>18</sup>

## Manifestations and Epidemiology of Liver Dysfunction Caused by Statins

The reported mechanisms of hepatotoxicity caused by statins are constantly being updated. Although statin-induced liver dysfunction (mainly liver transaminase abnormalities) is common in clinical practice, the incidence of statins-induced liver injury is 1.9% to 5.5%.<sup>19</sup> Among which the incidence rates of ALT  $\geq$  3 times upper limit of normal (ULN) were fluvastatin 1.0–2.0%, atorvastatin 0.9–1.3%, pitavastatin 0.9%, simvastatin 0.7%, and rosuvastatin  $<$ 0.4%, respectively.<sup>20</sup> The types of liver dysfunction caused by different statins were also different. Most of the liver injury caused by simvastatin was hepatocellular liver injury (85%), while most of the liver injury caused by atorvastatin, fluvastatin, lovastatin and pravastatin was cholestatic or mixed liver injury.<sup>21</sup> A meta-analysis indicated that atorvastatin showed a higher risk of transaminase elevations than non-statin control (OR = 4.0, 95% CI: 2.2–7.6), pravastatin (OR = 3.49, 95% CI: 1.77–6.92) and simvastatin (OR = 2.77, 95% CI: 1.31–5.09), respectively.<sup>5</sup> Besides, an Emax dose–response relationship was identified for the effect of atorvastatin on transaminase elevations.<sup>5</sup> Hepatocellular, cholestasis, and autoimmune hepatitis may infrequently appear months to years after the initiation of statins.<sup>22</sup>

## Countermeasures of Liver Dysfunction Caused by Statins

Firstly, statin-induced liver dysfunction is usually mild to moderate (ALT and/or AST  $<$  3ULN), among whom the elevated ALT and/or AST may voluntarily decrease. If the statins users are in the absence of other relevant clinical manifestations of liver damage, it is not necessary to reduce or stop statins. However, it is recommended to recheck the liver function every 4 to 8 weeks. Secondly, patients with elevated liver enzymes  $\geq$ 3ULN or combined with elevated total bilirubin should reduce or discontinue statins, or switch to cholesterol absorption inhibitors or PCSK9 inhibitors.<sup>23</sup> An appropriate combination of liver protectant (intravenous N-acetylcysteine, magnesium isoglycyrhizinate, or bicyclic alcohol) may be needed, but the liver function should be rechecked weekly until normal.<sup>24</sup> Due to the important benefits of statins in coronary plaque regression,<sup>25</sup> restarting low-dose statin therapy or switching to other statins such as pravastatin is still encouraged for patients at high and very high risk of ASCVD. The liver damage may occur again after switching to another statin, so the hepatotoxicity should also be carefully monitored. However, it has been reported that prospective monitoring of statin-induced liver injury is not justified unless the patient would otherwise benefit from such monitoring.<sup>22</sup>

Thirdly, the lowest effective dose of statins should be recommended, and the combination with CYP3A4 inhibitors or other drugs that may aggravate liver damage should be avoided, eg macrolide antimicrobials, acetaminophen, fibrates. Besides, in patients who used both vitamin K antagonists and statins, the risk of ADR observed was lower than that of concomitant treatment with non-vitamin K antagonist oral anticoagulants and statins in an observational cross-sectional study.<sup>26</sup> Fourthly, a history of liver and biliary diseases should be checked before using statins. If the liver transaminase is elevated, differential diagnosis and treatment should be performed, and statins should be started after the transaminase has returned to normal.

Fifthly, statins are not recommended for patients with active chronic liver disease (alcoholic liver disease, autoimmune liver disease, etc.) and decompensated cirrhosis, which is due to that decreased CYP3A4 activity in such patients not only worsens liver damage, but also significantly increases the risk of rhabdomyolysis due to significant increased blood concentrations of statins.<sup>22</sup> The last but not least, the patients with stable chronic liver disease and liver cirrhosis compensatory stage can start with a small dose of statins and monitor changes in liver function. Statins should be discontinued and liver protection therapy should be performed when ALT and/or AST  $\geq$  3ULN.<sup>22</sup>

## Muscle Injury Caused by Statins

Myopathy/rhabdomyolysis is a disorder characterized by muscle dysfunction that could be linked to genetics, inflammation, or a serious side effect of medications such as statins.<sup>27</sup> The auxiliary examination methods for diagnosis of muscle injury including B-mode ultrasound, magnetic resonance imaging, and creatine kinase (CK) examination. Clinical features of different statin-associated muscle symptoms including myalgia, myopathy, myositis, myonecrosis, and rhabdomyolysis, are complex.<sup>28</sup> The mechanism of muscle injury caused by statins was reported related to the over-expression of HMG-CoA reductase in various tissues and cell types, and then activate T cell receptors to induce autoimmunity, which ultimately leads to muscle fiber atrophy and degeneration.<sup>29</sup>

## Manifestations and Epidemiology of Muscle Injury Caused by Statins


There are appropriate 10% to 25% of patients treated with statins report statin-related muscle symptoms, while the real incidence of myopathy was 1.5% to 5%.<sup>30</sup> Myalgia can manifest as aches, tenderness, and cramps with normal CK level, while myopathy can be characterized by muscle weakness, but the CK level may not be elevated.<sup>28</sup> Pathological staining of statin-induced myopathy showed atrophy, degeneration, and regeneration of muscle fibers, but without obvious lymphocyte inflammation.<sup>29</sup> If the muscle injury develops into myositis, it may manifest as pain, redness, and elevated temperature with histopathological features of macrophages and inflammatory T and B cells. Rhabdomyolysis is the most severe muscle injury state with dead tissue, manifested by myonecrosis with varying degrees of elevated CK levels, as well as the increase of serum creatinine  $\geq 0.5$  mg/dL from baseline.<sup>28</sup>

Statin-associated muscle symptoms were higher in female, younger subjects (age < 65 years), and those who reported physical activity.<sup>31</sup> However, rhabdomyolysis was reported more common in elder patients  $\geq 75$  years of age than in younger patients and in male than female among a 33 million report in VigiBase®.<sup>32</sup> Besides, the most commonly co-reported drugs were gemfibrozil, cyclosporine, ezetimibe, clarithromycin, amiodarone, and fenofibrate, indicating a great risk of interacting with these drugs. The meta-analyses of 47 randomized controlled trials showed that simvastatin was associated with a lower risk of muscle problems (OR = 0.70, 95% CI: 0.55–0.90), while rosuvastatin showed a higher risk (OR = 1.75, 95% CI: 1.17–2.61) when compared with atorvastatin.<sup>5</sup> However, simvastatin was associated with the highest incidence of rhabdomyolysis, followed by atorvastatin, rosuvastatin, fluvastatin, pitavastatin, pravastatin, and lovastatin.<sup>32</sup>

## Risk Factors and Countermeasures of Muscle Injury Caused by Statins

Many risk factors seem to increase the potential of muscle injury caused by statins, and avoiding which can early identify the risk (Figure 1A). From the patient perspective, elderly, kidney disease, muscle disease, excessive physical activity, and genetic variants of CYP isoenzymes lead to increased likelihood of muscle injury caused by statins.<sup>30,33</sup>

It has been reported that lipophilic statins (atorvastatin, simvastatin, fluvastatin, lovastatin), compared with hydrophilic statins (rosuvastatin, pravastatin), are able to cross the cell membrane of extrahepatic tissue in a non-selective manner, and thus increasing muscular toxicity.<sup>30</sup> A meta-analysis of randomized controlled trials found that lipophilic



**Figure 1** Early identification of risk factors (A) and intervention strategies (B) for statin-related muscle injury.

statins were associated with a greater risk of muscle injury caused by statins.<sup>34</sup> It may reveal a lower risk of muscle injury when using hydrophilic statins than that of lipophilic statins.

Change of CYP450 enzyme activity is also a key factor affecting the adverse reactions of statins.<sup>35</sup> Most statins such as lovastatin, simvastatin, and atorvastatin are mainly metabolized by CYP3A4 enzymes, while fluvastatin, rosuvastatin, and pravastatin are mainly metabolized by CYP2C9 enzymes. A systematic review showed that the drug–drug interactions between CYP3A4 inhibitors (eg clarithromycin, amiodarone, ritonavir) and statins would increase statin-related muscle injury.<sup>35</sup> Thus, it is suggested to avoid CYP3A4 inhibitors in combination with simvastatin, lovastatin, or atorvastatin, reduce the dose of pitavastatin or pravastatin, but can continue to use fluvastatin or rosuvastatin. Additionally, a systematic review showed that the adverse effects in muscle in the elderly showed an increased risk of 16% with high-intensity statins comparing with low/moderate-intensity statins.<sup>34</sup> Adverse herb–drug interaction based on CYP450 may also affect the metabolism of statins. It proved that Jiangzhi Recipe, a traditional Chinese medicine compound, can inhibit the activity of CYP3A4, enhance the blood concentration and bioavailability of atorvastatin, which could cause or aggravate the side effect of statins (eg muscular toxicity).<sup>36</sup> It has been reported that the SCLO1B1 genotype can inform the risk of statin-related muscle symptoms, and new users of statins in particular may benefit from pharmacogenomic testing of the SCLO1B1 gene.<sup>37</sup>

It has also been demonstrated that the food like Grapefruit and Seville oranges contain furanocoumarins, and the mechanisms by which these furanocoumarins would cause food–drug interactions focus on inhibiting CYP3A4.<sup>38</sup> Of the current statins, simvastatin, lovastatin, and atorvastatin can be significantly enhanced by interaction with grapefruit and its juice, while pravastatin, fluvastatin and rosuvastatin seem least likely to interact, and these interactions can produce serious adverse effects such as rhabdomyolysis, to which the elderly may be most susceptible.<sup>38,39</sup> Because food–drug interactions vary from person to person and are difficult to predict, at the very least, interactions that are known to help prevent ADR should be avoided, and thus requiring careful daily dietary advice for patients.


From the management perspective, muscle injury caused by statins includes avoiding or reducing above risk factors, and switching to different statins and/or alternative lipid-lowering drug (eg ezetimibe, PCSK9 inhibitors, inclisiran). It is also important to consider weighing the costs and benefits of statins versus other medications to reduce long-term atherosclerotic cardiovascular disease events.

## Discussion and Future Perspectives

ADR is a natural element during drug use, which can occur at any age and has attracted great concerns from doctors and pharmacists.<sup>40</sup> In face of the successful benefits and high proportion use of statins in the prevention of adverse cardiovascular events for more than 40 years, the statin-related liver dysfunction and muscle injury have been health care challenge.<sup>41</sup> It is certainly far-sighted to identify the balance of benefits and risks before using statins, and to focus on the countermeasures once those risks occur.

Statins are typically used by elderly patients whose physiological functions such as immune capacity, rehabilitation function, and drug metabolism are deteriorating, resulting in damaged organ and decreased ability to respond to ADR. In addition, the cardiovascular diseases of statin users often require multiple drugs, which increases the incidence of ADR caused by drug interactions to some extent. Therefore, in the actual clinical application of statin, medical staff should pay attention to the monitoring of the elderly population, reduce the incidence of adverse reactions, and improve the treatment effect and quality of life. A cross-sectional study enrolling 752 statins users in Jordan showed that only half of the patients (49.7%) received statin treatment at the intensity recommended by the guidelines.<sup>42</sup> Besides, 40.7% cases were undertreated with statins, and the management of dyslipidemia was not accompanied by appropriate follow-up.<sup>42</sup> Statin users who are older, have been on statins for a longer period of time, and have a history of angina or stroke, should be given sufficient follow-up to determine patient compliance and treatment response including the potential ADR.<sup>42</sup>

The precautionary principles of statin-related ADR are to ensure correct and personalized medication, avoid unnecessary or easy to cause serious risks of combined medication, and closely identify the patients' initial drug response. The countermeasures for statin-related muscle injury (Figure 1B) and liver dysfunction (Figure 2) including stopping the suspected drug; reducing statin dosage; changing to another lipid-lowering agent; and taking effective treatment measures using liver protectors and hydration treatments, for instance. It is worth mentioning that the adverse



**Figure 2** Countermeasures of liver dysfunction caused by statins.

interactions between herbs and drugs are often ignored. Even in the USA, 40% of herbal users reported potential adverse herb–drug interactions.<sup>43</sup> There is evidence that traditional Chinese medicine flavonoids can raise plasma levels of statins and increase the risk of statin induced myopathy by interacting with the liver CYP450 enzyme, OATP uptake, or HMG-CoA reductase.<sup>44</sup>

A large body of evidence shows that the benefits of statin therapy far outweigh any actual or perceived risks.<sup>45</sup> Whether the adverse effects of statins are caused by the effects of statins or placebo is still debated. Recent study included 3027 statin users showed that high-intensity statin was statistically significant in resulting in an LDL-C level < 40 mg/dL when compared with low- and moderate-intensity statin, but the rate of ADR related to different intensity statin showed no statistical significance.<sup>46</sup> A retrospective cohort study showed that high-intensity atorvastatin was associated with an increased incidence of ADR.<sup>47</sup> These studies have yielded conflicting conclusions about whether different intensity doses of statins are associated with an increased incidence of adverse reactions. Therefore, detailed studies on the adverse risks of statins are needed in the future.

Firstly, it is necessary to accurately extract, clean, and standardize the flow of adverse reaction records involving statins and their combination drugs (Chinese medicine and Western medicine). Then the gender, age, route, appropriate statin intensity dose, time of administration, symptoms, and severity of the patients should be analyzed via multivariate analysis of variance. Secondly, the real-world discovery may provide an accurate basis for the identification of drug use risk.<sup>1,48,49</sup> Thirdly, too low LDL-C level may lead to unexpected adverse events, but there was evidence showing that the lipid-lowering drugs such as PCSK9 inhibitors have been shown not only to be harmless but also to produce additional cardiovascular benefits.<sup>50</sup> Whether there is an optimal minimum safe level of LDL-C after statin treatment for different types of patients needs further study.

## Conclusions

Statins have become the main lipid-lowering drugs for the primary and secondary prevention and treatment of ASCVD. Meanwhile, the issues of drug safety of statins have aroused increasing concerns. This review describes the epidemiology

and mechanisms of statin-associated liver dysfunction and muscle injury. Besides, detailed management strategies including early identification and post-intervention are presented aiming to minimize the risks of statins use and ensure timely medication schedule adjustments for patients.

## Abbreviations

ADR, adverse drug reaction; ALT, alanine aminotransferase; ASCVD, atherosclerotic cardiovascular diseases; AST, aspartate aminotransferase; CK, creatine kinase; HMG-CoA, 3-hydroxy-3-methylglutaryl-Coenzyme A; LDL-C, low density lipoprotein cholesterol; ULN, upper limit of normal.

## Data Sharing Statement

No data was used for the research described in the article.

## Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, review design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

## Funding

This work was supported by Medical Health Science and Technology Project of Zhejiang Provincial Health Commission (No. 2023KY1075), Jiaxing Key Discipline of Medicine–Clinical Pharmacy (2023-ZC-008), Public Technology Application Research Program of Zhejiang Province of China (NO. LGD21H310003), Science and Technology Project of Jiaxing of China (NO. 2021AD10023), National Natural Science Foundation of China (Nos. 81974470, 82272004), Zhejiang Provincial Natural Science Foundation of China (Nos. LYY19H280006, LQ21H310004, LY18H180001, LGF22H180008), and Zhejiang Provincial Science Technology Projects of Traditional Chinese Medicine (Nos. 2021ZB174, 2021ZB301).

## Disclosure

All authors declare no conflicts of interest in this work.

## References

1. Sun L, Zhao Q, Wang Y, et al. Efficacy and safety of anlotinib-containing regimens in advanced non-small cell lung cancer: a real-world study. *Int J Gen Med.* 2023;16:4165–4179. doi:10.2147/IJGM.S424777
2. Tchio-Nighie KH, Mpoh MM, Tchokomeni H, et al. Drug safety monitoring in health programs of Cameroon. *J Pharmacovigil.* 2022;10(9):10000385.
3. Zhang XJ, Zhou JG, Pan M, Yuan W, Gao B. Analysis of adverse drug reaction reports from a public hospital in Shanxi Province in 2022. *Risk Manag Healthc Policy.* 2023;16:1391–1401. doi:10.2147/RMHP.S418386
4. Liu S, Li Y, Zeng X, et al. Burden of cardiovascular diseases in China, 1990–2016: findings from the 2016 global burden of disease study. *JAMA Cardiol.* 2019;4(4):342–352. doi:10.1001/jamacardio.2019.0295
5. Wang X, Li J, Wang T, et al. Associations between statins and adverse events in secondary prevention of cardiovascular disease: pairwise, network, and dose-response meta-analyses of 47 randomized controlled trials. *Front Cardiovasc Med.* 2022;9:929020. doi:10.3389/fcvm.2022.929020
6. Zhao Q, Sun S, Zhou F, Yue J, Luo X, Qu X. The inhibition of evolocumab on non-infarct-related artery disease in patients with ST-elevation myocardial infarction. *Int J Gen Med.* 2023;16:2771–2781. doi:10.2147/IJGM.S417481
7. Kouhpeikar H, Khosaravizade Tabasi H, Khazir Z, et al. Statin use in COVID-19 hospitalized patients and outcomes: a retrospective study. *Front Cardiovasc Med.* 2022;9:820260. doi:10.3389/fcvm.2022.820260
8. Tuleutayeva RY, Makhatova AR, Rakhzyhanova SO, Zhazykbayeva LK, Kozhakhmetova DK. Candidate genes for prediction of efficacy and safety of statin therapy in the Kazakh Population. *Twin Res Hum Genet.* 2023;1–7. doi:10.1017/thg.2023.28
9. Vohra LI, Rizwan K, Saeed E, Hamza MSA, Ochani S. Inclisiran adjuvant therapy to statins for the use of hypercholesterolemia: a commentary. *Egypt Heart J.* 2023;75(1):60. doi:10.1186/s43044-023-00389-8
10. Nanna MG, Navar AM, Wang TY, et al. Statin use and adverse effects among adults >75 years of age: insights from the patient and Provider Assessment of Lipid Management (PALM) registry. *J Am Heart Assoc.* 2018;7(10):e008546. doi:10.1161/JAHA.118.008546
11. Eppler M, Singh N, Ding L, Magee G, Garg P. Discharge prescription patterns for antiplatelet and statin therapy following carotid endarterectomy: an analysis of the vascular quality initiative. *BMJ Open.* 2023;13(7):e071550. doi:10.1136/bmjopen-2022-071550

12. Balaha MF, Alamer AA, Kabel AM, Aldosari SA, Fatani S. A prospective cross-sectional study of acute coronary syndrome patients' quality of life and drug prescription patterns at Riyadh Region Hospitals, Saudi Arabia. *Healthcare*. 2023;11(13):1973. doi:10.3390/healthcare11131973
13. Ofori-Asenso R, Ilomäki J, Zomer E, Curtis AJ, Zoungas S, Liew D. A 10-year trend in statin use among older adults in Australia: an analysis using national pharmacy claims data. *Cardiovasc Drugs Ther*. 2018;32(3):265–272. doi:10.1007/s10557-018-6794-x
14. Huang BT, Peng Y, Huang FY, et al. Trends in prescribing rate of statins at discharge and modifiable factors in patients with atherosclerotic cardiovascular disease. *Intern Emerg Med*. 2017;12(8):1121–1129. doi:10.1007/s11739-017-1694-9
15. Bellosta S, Corsini A. Statin drug interactions and related adverse reactions: an update. *Expert Opin Drug Saf*. 2018;17(1):25–37. doi:10.1080/14740338.2018.1394455
16. Abdoli N, Azarmi Y, Eghbal MA. Protective effects of N-acetylcysteine against the statins cytotoxicity in freshly isolated rat hepatocytes. *Adv Pharm Bull*. 2014;4(3):249–254. doi:10.5681/apb.2014.036
17. Karahalil B, Hare E, Koç G, Uslu İ, Şentürk K, Özkan Y. Hepatotoxicity associated with statins. *Arh Hig Rada Toksikol*. 2017;68(4):254–260. doi:10.1515/aiht-2017-68-2994
18. Qu -K-K, Zhang C-N, Dong L-X, Wang -S-S, Zhang Z-D, Zhang L. Association of ABCB1 polymorphisms with lipid homeostasis and liver injury response to atorvastatin in the Chinese population. *Can J Physiol Pharmacol*. 2020;98(1):15–22. doi:10.1139/cjpp-2019-0339
19. Björnsson ES. Hepatotoxicity of statins and other lipid-lowering agents. *Liver Int*. 2017;37(2):173–178. doi:10.1111/liv.13308
20. Zhang XY, Zheng SB. Liver dysfunction caused by statins and its treatment strategy. *Chin J New Drugs Clin Remedies*. 2021;40(4):246–250. Chinese.
21. Perdices EV, Medina-Cáliz I, Hernando S, et al. Hepatotoxicity associated with statin use: analysis of the cases included in the Spanish hepatotoxicity registry. *Rev Esp Enferm Dig*. 2014;106(4):246–254.
22. Russo MW, Hoofnagle JH, Gu J, et al. Spectrum of statin hepatotoxicity: experience of the drug-induced liver injury network. *Hepatology*. 2014;60(2):679–686. doi:10.1002/hep.27157
23. Joint Committee on the Chinese Guidelines for Lipid Management. Chinese guidelines for lipid management(2023). *Chin Circ J*. 2023;38(3):237–271. Chinese.
24. Technology Committee on DILI Prevention and Management, Chinese Medical Biotechnology Association; Study Group of Drug-Induced Liver Disease, Chinese Medical Association for the Study of Liver Diseases. Chinese guideline for diagnosis and management of drug-induced liver injury(2023 Version). *Chin J Gastroenterol*. 2023;28(7):397–431. Chinese.
25. Räber L, Ueki Y, Otsuka T, et al. Effect of alirocumab added to high-intensity statin therapy on coronary atherosclerosis in patients with acute myocardial infarction: the PACMAN-AMI randomized clinical trial. *JAMA*. 2022;327(18):1771–1781. doi:10.1001/jama.2022.5218
26. Jimenez-Serrania M-I, Treceño-Lobato C. Influence of concomitant treatments under anticoagulants and statins in detecting signals of adverse drug reactions. *Semin Thromb Hemost*. 2019;45(8):837–845. doi:10.1055/s-0039-1695734
27. Awad M, Al-hussaniy HA, Alburghaif AH, Tawfeeq KT. The role of COVID-19 in myopathy: incidence, causes, treatment, and prevention. *J Med Life*. 2022;15(12):1458–1463. doi:10.25122/jml-2022-0167
28. Gupta R, Alcantara R, Popli T, et al. Myopathy associated with statins and SGLT2 - A review of literature. *Curr Probl Cardiol*. 2021;46(4):100765. doi:10.1016/j.cpcardiol.2020.100765
29. Mohassel P, Mammen AL. Anti-HMGCR Myopathy. *J Neuromuscul Dis*. 2018;5(1):11–20. doi:10.3233/JND-170282
30. Vinci P, Panizon E, Tosoni LM, et al. Statin-associated myopathy: emphasis on mechanisms and targeted therapy. *Int J Mol Sci*. 2021;22(21):11687. doi:10.3390/ijms222111687
31. Casula M, Gazzotti M, Bonaiti F, et al. Reported muscle symptoms during statin treatment amongst Italian dyslipidaemic patients in the real-life setting: the PROSISA Study. *J Intern Med*. 2021;290(1):116–128. doi:10.1111/joim.13219
32. Montastruc JL. Rhabdomyolysis and statins: a pharmacovigilance comparative study between statins. *Br J Clin Pharmacol*. 2023;89(8):2636–2638. doi:10.1111/bcp.15757
33. Vrablik M, Zlatohlavek L, Stulc T, et al. Statin-associated myopathy: from genetic predisposition to clinical management. *Physiol Res*. 2014;63(Suppl 3):S327–S334. doi:10.33549/physiolres.932865
34. Iwere RB, Hewitt J. Myopathy in older people receiving statin therapy: a systematic review and meta-analysis. *Br J Clin Pharmacol*. 2015;80(3):363–371. doi:10.1111/bcp.12687
35. Hougaard Christensen MM, Bruun Haastrup M, Øhlenschlaeger T, et al. Interaction potential between clarithromycin and individual statins-A systematic review. *Basic Clin Pharmacol Toxicol*. 2020;126(4):307–317. doi:10.1111/bcpt.13343
36. Wang XL. Potential herb-drug interaction in the prevention of cardiovascular diseases during integrated traditional and Western medicine treatment. *Chin J Integr Med*. 2015;21(1):3–9. doi:10.1007/s11655-014-1892-5
37. Tomcsanyi KM, Tran KA, Bates J, et al. Veterans health administration: implementation of pharmacogenomic clinical decision support with statin medications and the SLCO1B1 gene as an exemplar. *Am J Health Syst Pharm*. 2023;80(16):1082–1089. doi:10.1093/ajhp/zxad111
38. Straughan JL. Grapefruit-drug interactions. *Cardiovasc J S Afr*. 2007;18(1):39–40.
39. Dahan A, Altman H. Food-drug interaction: grapefruit juice augments drug bioavailability--mechanism, extent and relevance. *Eur J Clin Nutr*. 2004;58(1):1–9. doi:10.1038/sj.ejcn.1601736
40. Balasubramanian R, Maideen NMP. HMG-CoA reductase inhibitors (Statins) and their drug interactions involving CYP enzymes, P-glycoprotein and OATP transporters-an overview. *Curr Drug Metab*. 2021;22(5):328–341. doi:10.2174/1389200222666210114122729
41. Strandberg TE. Role of statin therapy in primary prevention of cardiovascular disease in elderly patients. *Curr Atheroscler Rep*. 2019;21(8):28. doi:10.1007/s11883-019-0793-7
42. Gharaibeh L, Al Zoubi S, Sartawi H, Ayyad D, Al-Hawamdeh M, Alrashdan R. The appropriateness of the use of statins for the secondary and primary prevention of atherosclerotic cardiovascular disease: a cross-sectional study from Jordan. *Eur Rev Med Pharmacol Sci*. 2023;27(12):5480–5492. doi:10.26355/eurrev\_202306\_32785
43. Bush TM, Rayburn KS, Holloway SW, et al. Adverse interactions between herbal and dietary substances and prescription medications: a clinical survey. *Altern Ther Health Med*. 2007;13(2):30–35.
44. Zechner J, Britza SM, Farrington R, Byard RW, Musgrave IF. Flavonoid-statin interactions causing myopathy and the possible significance of OATP transport, CYP450 metabolism and mevalonate synthesis. *Life Sci*. 2022;291:119975. doi:10.1016/j.lfs.2021.119975
45. Adhyaru BB, Jacobson TA. Safety and efficacy of statin therapy. *Nat Rev Cardiol*. 2018;15(12):757–769. doi:10.1038/s41569-018-0098-5

46. Plummer S, Wright M, Brown JM. How low is too low? A retrospective analysis of very low LDL-C levels in veterans. *Fed Pract.* 2022;39(Suppl 5).
47. Stein B, Ward T, Hale G, Lyver E. Safety of high-intensity statins in the veteran population: atorvastatin 40 to 80 mg compared with rosuvastatin 20 to 40 mg. *Ann Pharmacother.* 2020;54(5):405–413. doi:10.1177/1060028019888487
48. Li L-C, Sun W, Lv X-Q, Xu -Y-Y, Hu Y, Shi J-N. Escitalopram-induced sinus bradycardia in coronary heart disease combined with depression: a case report and review of literature. *Front Cardiovasc Med.* 2024;10:1133662. doi:10.3389/fcvm.2023.1133662
49. Qian Q, Wang YZ, Kan LD, et al. Real-world prescribing patterns for hypertensive children in China from 2018 to 2021: a cross-sectional multicenter study. *Risk Manag Healthc Policy.* 2023;16:287–299. doi:10.2147/RMHP.S392224
50. Copping C, Movahed MR, Azemawah V, Peyton L, Gregory J, Hashemzadeh M. A comprehensive review of PCSK9 inhibitors. *J Cardiovasc Pharmacol Ther.* 2022;27:10742484221100107. doi:10.1177/10742484221100107

International Journal of General Medicine

Dovepress

## Publish your work in this journal

The International Journal of General Medicine is an international, peer-reviewed open-access journal that focuses on general and internal medicine, pathogenesis, epidemiology, diagnosis, monitoring and treatment protocols. The journal is characterized by the rapid reporting of reviews, original research and clinical studies across all disease areas. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-general-medicine-journal>