

# Exploring Thermal Dynamics in Wound Healing: The Impact of Temperature and Microenvironment

Jun Huang<sup>1,2</sup>, Chunjie Fan<sup>2</sup>, Yindong Ma<sup>2</sup>, Guobao Huang<sup>1,2</sup> 

<sup>1</sup>Department of Clinical Medicine, Shandong Second Medical University (Weifang Medical University), Weifang, 261000, People's Republic of China;

<sup>2</sup>Department of Burns and Reconstructive Surgery, Central Hospital Affiliated to Shandong First Medical University, Jinan, 250013, People's Republic of China

Correspondence: Guobao Huang, Department of Burns and Reconstructive Surgery, Central Hospital Affiliated to Shandong First Medical University, No. 105, Jiefang Road, Lixia District, Jinan, Shandong Province, 250013, People's Republic of China, Tel +86 531-55865707, Email huanguobao@sdu.edu.cn

**Abstract:** Exploring the critical role of thermal dynamics in wound healing, this manuscript navigates through the complex biological responses initiated upon wound infliction and how temperature variations influence the healing trajectory. Integrating biothermal physics, clinical medicine, and biomedical engineering, it highlights the significance of thermal management in wound care, emphasizing the wound microenvironment's division into internal and external domains and their collaborative impact on tissue repair. Innovations in real-time wound temperature monitoring, especially through intelligent wireless sensor dressings, are spotlighted as transformative, enabling precise wound condition management. The text underscores the necessity for further research to elucidate thermal regulation's molecular and cellular mechanisms on healing processes. It advocates for standardized protocols for localized heating treatments, integrating them into personalized wound care strategies to enhance therapeutic outcomes, improve patient well-being, and achieve cost-effective healthcare practices. This work presents a forward-looking perspective on refining wound management through sophisticated, evidence-based interventions, emphasizing the interplay between thermal dynamics and wound healing.

**Keywords:** wound healing, thermal dynamics, wound microenvironment, intelligent dressings, localized heating treatment

## Introduction

The disruption of epidermal integrity instigates a multifaceted biological response within the dermal stratum. This response encompasses immune activation,<sup>1</sup> vasodilation mediated by pro-inflammatory cytokines,<sup>2</sup> and an intensification of metabolic processes within the afflicted tissues.<sup>3</sup> Collectively, these phenomena modify the thermal profiles at both the lesion and its adjacent epidermal areas.

A seminal study conducted in 2015 elucidated the complex architecture of the wound microenvironment by dichotomizing it into external and internal components.<sup>4</sup> The external wound microenvironment is defined as the area immediately adjacent to the lesion's surface, interfacing directly with the ambient environment. In contrast, the internal wound microenvironment comprises the subepidermal layers, rich in cellular constituents and the extracellular matrix (ECM).<sup>5</sup> This bifurcation is crucial for understanding the dynamic interactions that dictate the course of wound healing.<sup>6</sup> Clinical interventions are strategically designed to modulate these microenvironments to optimize conditions conducive to reparative processes. Pertinent external variables affecting both acute and chronic wound contexts include thermal fluctuations,<sup>7</sup> mechanical pressure,<sup>8</sup> moisture levels,<sup>9</sup> and microbial colonization.<sup>10</sup> An intricate comprehension of both the synergistic and antagonistic interactions among these factors, as well as their aggregate impact on the healing trajectory, is paramount for the advancement of therapeutic approaches.

Within the discourse of wound management, the imperative of localized surveillance and modulation of the external wound environment is unequivocal. Current clinical guidelines advocate for the manipulation of pressure, humidity, and microbial load as fundamental elements of localized intervention protocols. Nonetheless, the regulation of wound

temperature has not been adequately integrated into standard therapeutic regimens.<sup>11,12</sup> The nuanced interplay between thermal conditions and reparative processes in wounds represents a relatively unexplored field, necessitating enhanced empirical examination. This study endeavors to provide a comprehensive exploration and critical evaluation of the correlation between thermal parameters within the external wound microenvironment and the holistic process of wound healing, thereby contributing to the expansion of both theoretical and practical paradigms in clinical wound management.

## Comprehensive Modeling of Human Biothermal Transfer Dynamics

The constitution of human tissues, encompassing a tripartite composition of solids, liquids, and gases, demonstrates a labyrinthine array of thermal transfer phenomena that diverge markedly from conventional mechanisms.<sup>13</sup> Embedded within the biological milieu, these tissues engage in metabolic activities that are distinct in nature from rudimentary chemical reactions, anchored by an intrinsically sophisticated apparatus for thermoregulation.<sup>14</sup> The spectrum of biothermal transfer within the human organism is delineated through a confluence of conductive, convective, and radiative modalities, rendering the narrative of thermal energy conveyance notably complex.<sup>15</sup> Scholarly endeavors have been directed towards the elucidation of these complexities, yielding theoretical constructs and algebraic expressions that encapsulate the quintessential principles of biothermal transfer in biological matrices. This intellectual journey has led to the creation of analytical models that allow for a detailed examination of thermal phenomena.

A watershed moment in this scholarly trajectory was realized in 1948 with the conceptualization of the human arm as a cylindrical model, leading to the articulation of the seminal bioheat transfer equation, colloquially known as the “Pennes equation”.<sup>16</sup> This theoretical innovation provided a substantive framework for the elucidation of thermal distribution patterns within biological tissues. Progressing beyond this foundational milestone, research further meticulously charted the intricacies of the human body’s thermoregulatory functions, deploying mathematical equations to quantitatively delineate this mechanism.<sup>17</sup> The findings from this inquiry resonated with a high degree of fidelity to established physiological paradigms. Recent scholarly initiatives have introduced a cutting-edge model, meticulously tailored for the demographic contours of the Chinese populace.<sup>18</sup> This model is predicated on the objective of precisely prognosticating instantaneous average skin temperatures under the ambience of warm environmental conditions (26.0 ~ 33.8°C). This analytical tool extends its utility beyond mere temperature estimation to encompass the delineation of evaporative sweat regulation efficiency, the dynamics of skin blood flow, and the quantification of total heat dissipation ratios. Critically, it unveils the gender-specific thermal response disparities, augmenting the granularity of thermal physiological understanding.


In light of the criticality of temperature as an arbiter of physiological equilibrium, its quantification and interpretive analysis have ascended to a position of paramount importance within the diagnostic lexicon of medical science. The quest for precision in mapping and modeling the temperature landscapes across biological tissues has emerged as a vibrant focal point of inquiry within the interdisciplinary domains of biothermal transfer and biomedical engineering.<sup>19,20</sup> This pursuit is emblematic of a broader scientific commitment to harness computational innovations for the real-time simulation and intricate reconstruction of biotemperature fields, thereby advancing the frontiers of knowledge in biothermal phenomena and refining the precision of diagnostic methodologies.

## Wound Temperature Measurement

### Single-Point Thermometric Evaluation

The archetypical modalities for quantifying thermal variations predominantly rely on single-point detection mechanisms, incorporating instruments such as thermocouple thermometers,<sup>21</sup> infrared thermometers,<sup>22</sup> and infrared thermographic cameras (Figure 1).<sup>23</sup> In the landscape of empirical investigation, a preference is discernible towards non-contact measurement devices, attributed to their non-intrusive nature and the convenience they offer in operational deployment.<sup>24</sup> Nevertheless, a discernible fraction of investigative efforts integrates contact-based thermometric tools, including liquid crystal thermometry and thermocouple probes, delineating a methodological diversity.<sup>25</sup> The deployment of infrared thermometers, characterized by their portability, ease of manipulation, and economic viability, engenders a spectrum of viewpoints regarding their utility in elucidating the correlation between thermal dynamics at the wound site

## Single-Point Thermometric Evaluation



**Figure 1** Overview of thermometric assessment tools. The left panel displays single-point measurement tools including thermocouple thermometers, probes, infrared thermometers, and thermographic cameras for diverse temperature assessments. The right panel introduces a real-time thermometric surveillance system with a smart wireless sensor dressing connected to digital platforms for continuous monitoring.

and the consequent healing trajectory.<sup>26</sup> Detractors highlight the susceptibility of infrared thermometric readings to perturbations from a multitude of exogenous and endogenous factors, casting aspersions on their reliability within the domain of wound diagnostic research.<sup>26</sup> In contrast, proponents assert that, under a regime of controlled environmental parameters, infrared thermometry emerges as a credible modality for the assessment of wound temperature, thereby facilitating an analytical lens into the state of wound convalescence.<sup>27</sup>

The fidelity of single-point thermal monitoring methodologies is contingent upon a complex interplay of factors, encompassing ambient thermal conditions, the systemic temperature of the patient, circulatory dynamics, and the presence of exudative fluids, which collectively precipitate variances in the resultant measurements.<sup>28,29</sup> The paucity of a globally endorsed standard for executing these measurements amplifies the challenges associated with the universal adoption and implementation of single-point wound temperature monitoring within both clinical and investigative contexts. This scenario underscores an imperative for methodological refinement and the formulation of standardized procedural frameworks to bolster the reliability and generalizability of these techniques in the broader discourse of wound management and scholarly inquiry.

## Pioneering Developments in Real-Time Thermometric Surveillance

Recent scholarly discourse and investigative endeavors have heralded the advent of advanced intelligent wireless sensor dressings.<sup>30–32</sup> These novel interventions represent a confluence of biocompatible materials and micro-engineered temperature sensors, epitomizing the fusion of biomedical engineering and sensor technology. Utilizing sophisticated mechanisms such as radio-frequency identification (RFID) systems or platinum-based sensors, these sensor-integrated dressings facilitate uninterrupted and meticulous monitoring of thermal variations at the wound site.<sup>33,34</sup> This innovation surmounts the limitations endemic to conventional single-point thermometric approaches, notably their diminished stability and pronounced vulnerability to a broad spectrum of environmental and physiological perturbations.

The utility of intelligent wireless sensor dressings extends beyond mere thermal measurement, enabling comprehensive real-time analysis of fluctuating parameters critical to the wound healing milieu.<sup>35,36</sup> This integrative monitoring capacity permits a detailed evaluation of the wound microenvironment's suitability for healing, embodying a proactive approach to wound management.<sup>37</sup> Moreover, these advanced dressings are endowed with the capability to wirelessly transmit the amassed data to contemporary digital devices, including smartphones and tablets, via Bluetooth technology (Figure 1).<sup>38,39</sup> This facilitates the prompt dissemination of vital health metrics to medical practitioners and integrates seamlessly with cloud-based infrastructures, ensuring the longitudinal conservation of wound-related data. Such integration heralds a significant leap forward in remote healthcare provision, aligning with the principles of personalized medicine. This digital convergence in wound care management underscores a paradigmatic shift towards customizing

therapeutic interventions to meet the unique physiological and clinical profiles of individual patients, thereby enhancing the efficacy and precision of medical care.

## The Relationship Between Wound Temperature Monitoring and Wound Healing Status

Temperature has been acknowledged as a critical factor influencing wound healing.<sup>40</sup> In the distinct stages following the formation of acute and chronic wounds, variations in wound temperature exhibit specific trends. These temperature changes are influenced by environmental conditions and local blood flow and can modulate vascular active molecules, facilitating vasoconstriction and dilation.

### Acute Wounds

#### Surgical Wounds

Studies show that normal surgical wounds can exhibit localized temperature increases in the initial days post-operation, returning to baseline within two weeks.<sup>41</sup> Zheng et al analyzed the impact of preoperative warming on reducing the risk of surgical site infections (SSI), finding that using mixed warming methods before surgery significantly lowers the incidence of SSI, recommending their adoption in surgical protocols.<sup>42</sup> Another study also highlights that maintaining normothermia with active warming techniques to keep body temperature above 36°C during surgery is a crucial strategy among others to significantly reduce the risk of surgical site infections.<sup>43</sup>

#### Early-Stage Burn Wounds

Determining the temperature of burn wounds assists in diagnosing the depth of burns. Studies employing contact-type rapid-response thermocouple probes to measure wound temperature within 10 hours post-burn reveal that, compared to adjacent normal skin, elevated wound temperatures indicate shallower burn depth or shorter healing times.<sup>44</sup> Such thermometric methods achieve a 78% accuracy rate in assessing burn depth, surpassing the 60% accuracy of evaluations based on clinical experience. However, the correlation between wound temperature and injury depth is not evident in hands, faces, feet, or beyond 10 hours post-burn.<sup>44</sup> Other research indicates that the optimal timeframe for using infrared thermal imaging to diagnose burn wound depth is within 3 days post-injury, as the correlation between wound temperature and burn depth diminishes after this period.<sup>45</sup> Furthermore, Ganon et al validated the use of the Flir One Thermal Imager<sup>®</sup> for early burn wound assessment in pediatrics, demonstrating its ability to distinguish between superficial and deep burns by measuring skin temperature differences, thereby aiding in the early decision-making process for treatments such as skin grafting.<sup>46</sup> Carrière et al highlighted the validity of using infrared thermography with a new, more sensitive thermal imager for assessing burn wound healing potential, showing its effectiveness in discriminating between different healing time frames when compared to Laser Doppler Imaging, marking it as a promising tool for burn wound triage and evaluation.<sup>47</sup>

### Chronic Wounds

In chronic wounds, a sudden increase in local temperature is a typical sign of infection and inflammation. Most chronic wounds, often occurring in the lower limbs, exhibit temperatures approximately 5°C lower than core body temperature due to compromised blood supply and oxygenation.<sup>48</sup>

#### Chronic Venous Leg Ulcers

Studies utilizing handheld thermometers to monitor the temperature of chronic venous leg ulcers show that infected wound sites exhibit skin temperatures more than 2°C higher than corresponding healthy skin on the opposite limb, indicating a strong correlation between elevated periwound skin temperature and infection.<sup>48</sup> The infrared thermography can identify distinct temperature patterns in the lower legs of patients with chronic venous diseases (CVD), showing potential for early detection and classification based on the severity of CVD, including the presence of florid ulcers.<sup>49</sup>

## Diabetic Foot Ulcers

In clinical trials employing infrared thermometers to measure the temperature of diabetic foot ulcers and surrounding skin, results indicate that smaller temperature differences between the wound and intact skin on the opposite foot predict a positive wound status and a tendency towards healing, suggesting that wound temperature effectively reflects inflammatory changes and healing trends in diabetic foot ulcers.<sup>50</sup> These studies collectively reveal that wound temperatures can significantly rise during the initial stages of injury and in cases of abnormal healing, but tend to decrease back to normal as the healing process normalizes.<sup>51</sup> Research combining temperature measurement with image evaluation has shown diagnostic sensitivities >60% and specificities >79% for detecting wound infections, whereas temperature measurement alone has shown a diagnostic sensitivity of up to 90% but a specificity of <25%.<sup>52</sup> However, a systematic analysis of existing literature suggests a lack of sufficient evidence to recommend wound temperature as a predictive indicator for wound healing.<sup>53</sup> Another meta-analysis also revealed that daily foot temperature monitoring at six points can significantly reduce the incidence of foot ulcers in patients with diabetes by prompting increased preventive measures in response to temperature differences greater than 2.2°C between the left and right corresponding sites.<sup>54</sup> Future research, if it further clarifies and confirms the role of wound temperature variations in early diagnosis and predicting infections and healing outcomes, could establish wound temperature measurement as an effective non-invasive method for early assessment of wound status and prognosis.

## Pressure Injury

Clinical research demonstrates that measuring wound temperature can accurately predict the anatomical locations of ulcers before the formation of pressure injuries, proving more accurate than the commonly used Braden scale for pressure injury risk assessment.<sup>55,56</sup> This technique can monitor healing progress and improve management and prevention of ulcers.<sup>27</sup> Another study indicates that, three weeks post-pressure injury formation, wound temperatures significantly exceed those of surrounding skin, suggesting the presence of infection or other healing-compromising factors.<sup>57</sup>

## The Impact of Wound Temperature Treatment on Wound Healing

For both acute and chronic wounds, it is imperative to ameliorate systemic factors and provide an optimal wound microenvironment to facilitate timely and orderly healing and functional recovery. Beyond maintaining a moist microenvironment, alleviating hypothermic states and sustaining normothermic conditions within the wound vicinity can also expedite wound healing. During the healing process, the temperature of the wound exerts a significant influence on various repair-related effector cells. Studies have demonstrated that early-stage wound temperatures falling below core body temperature can hinder collagen deposition and reduce the presence of late-stage inflammatory cells and fibroblasts, thereby delaying the healing process.<sup>58</sup> Especially, 33°C represents a critical threshold temperature for the occurrence of various biological changes within the wound; temperatures below this threshold lead to decreased activity of neutrophils, fibroblasts, and keratinocytes.<sup>59,60</sup> Moreover, investigations revealed that during the replacement of dressings on acute traumatic wounds, the immediate post-removal average wound temperature is measured at 32.6°C, slightly below the minimum requisite of 33°C for cellular activity.<sup>61</sup> Prior to applying a new dressing, the average temperature drops to 29.9°C, indicating a persistently hypothermic state throughout the dressing change process.<sup>62</sup>

Observations suggest that elevating the temperature of pressure injury wounds to between 36–38°C significantly reduces wound area.<sup>63</sup> Thermal radiation dressings, as a therapeutic device, have been proven to increase capillary blood flow perfusion and blood oxygen partial pressure, thereby promoting the healing of pressure injuries.<sup>64</sup> Various modalities exist for heating treatment. A study utilizing thermal radiation dressings to heat both normal skin on human thighs and surgical wounds (set at 38°C for 2 hours of heating followed by 2 hours of rest, over a period of 7 days) maintained wound temperatures between 36.0–37.5°C.<sup>65</sup> This demonstrated that local heating can enhance microvascular blood flow and subcutaneous oxygen content in both normal and injured skin, potentially offering resistance to infection without increasing collagen deposition at the wound site. In addition, thermal radiation dressings (set at 38°C, three times a day for 1 hour) were used for the treatment of Stage I–II uninfected pressure injuries over 6 weeks or until healing showed that, compared to standard treatment with alginate dressings, this method significantly improved wound healing rates.<sup>66</sup> Subsequent research using heated dressing systems for intermittent heating of wounds

for 5 hours (set at 38°C, heating for 3 hours followed by a 2-hour rest without the system) found that, compared to a control group using standard dressings, the proportion of CD3+ T lymphocytes in the wound increased after 24 and 48 hours of dressing use, while neutrophil and macrophage counts showed no significant difference.<sup>67</sup> This suggests lymphocyte infiltration into the local wound microenvironment, thereby enhancing the innate immune response within the wound healing milieu.

## Conclusion

The healing of various acute and chronic wounds in clinical settings is influenced by numerous factors pertaining to the external microenvironment of the wound. Wound temperature, as a pivotal indicator of the wound's external microenvironment, has been identified in research as having potential for predicting infection and assessing healing outcomes, though further validation is necessary. Clarification of the interrelationship between wound temperature and wound healing could offer novel insights and foundational support for clinical treatment methodologies and the development of smart dressings.

By employing intelligent wireless sensor dressings, it is possible to observe comprehensively the variations in wound temperature within the external microenvironment, enabling real-time assessment of wound status. Furthermore, these smart dressings facilitate modulation and adjustment of the wound microenvironment, reduce the frequency of dressing changes, and promote wound healing. Significantly, this approach has the potential to alleviate patient discomfort and decrease hospitalization costs.

In addressing hypothermic wound states, localized heating treatments have been shown to enhance capillary blood flow perfusion and blood oxygen partial pressure, increase the activity and collagen deposition of neutrophils, fibroblasts, and keratinocytes, and elevate the proportion of lymphocytes at the wound site. This, in turn, strengthens the innate immune response within the wound healing microenvironment, thereby improving the wound condition and fostering repair. However, further research is necessary to elucidate the precise mechanisms by which localized heating modulates the proliferation and function of wound repair-related effector cells and to validate its efficacy in clinical intervention for wound infection. The optimal constant temperature required for the wound microenvironment and the duration of heating treatments remain to be established. Nonetheless, localized heating treatment represents an innovative approach for regulating the local wound microenvironment and the associated healing processes.

## Funding

This work was supported by the Youth Science Fund Cultivation Funding Program of Shandong First Medical University [202201-130] and Rong-Xiang Regenerative Medicine Fund of Shandong University [2019SDRX-08].

## Disclosure

The authors declare no conflicts of interest in this work.

## References

1. Sawada Y, Gallo RL. Role of epigenetics in the regulation of immune functions of the skin. *J Invest Dermatol*. 2021;141(5):1157–1166. doi:10.1016/j.jid.2020.10.012
2. Jahanshahi M, Hamdi D, Godau B, et al. An engineered infected epidermis model for in vitro study of the skin's pro-inflammatory response. *Micromachines*. 2020;11(2):227. doi:10.3390/mi11020227
3. Trompette A, Pernot J, Perdijk O, et al. Gut-derived short-chain fatty acids modulate skin barrier integrity by promoting keratinocyte metabolism and differentiation. *Mucosal Immunol*. 2022;15(5):908–926. doi:10.1038/s41385-022-00524-9
4. Kruse CR, Nuutila K, Lee CCY, et al. The external microenvironment of healing skin wounds. *Wound Repair Regen*. 2015;23(4):456–464. doi:10.1111/wrr.12303
5. Sorg H, Sorg CGG. Skin wound healing: of players, patterns, and processes. *Eur Surg Res*. 2023;64(2):141–157. doi:10.1159/000528271
6. Hu C, Chu C, Liu L, et al. Dissecting the microenvironment around biosynthetic scaffolds in murine skin wound healing. *Sci Adv*. 2021;7(22). doi:10.1126/sciadv.abf0787
7. Monshipouri M, Aliahmad B, Ogrin R, et al. Thermal imaging potential and limitations to predict healing of venous leg ulcers. *Sci Rep*. 2021;11(1):13239. doi:10.1038/s41598-021-92828-2
8. Kimura S, Tsuji T. Mechanical and immunological regulation in wound healing and skin reconstruction. *Int J Mol Sci*. 2021;22(11):5474. doi:10.3390/ijms22115474

9. Li Y, Zhang Y, Wang Y, et al. Regulating wound moisture for accelerated healing: a strategy for the continuous drainage of wound exudates by mimicking plant transpiration. *Chem Eng J.* 2022;429:131964. doi:10.1016/j.cej.2021.131964
10. Wolcott R. Disrupting the biofilm matrix improves wound healing outcomes. *J Wound Care.* 2015;24(8):366–371. doi:10.12968/jowc.2015.24.8.366
11. Eriksson E, Liu PY, Schultz GS, et al. Chronic wounds: treatment consensus. *Wound Repair Regen.* 2022;30(2):156–171. doi:10.1111/wrr.12994
12. Rayman G, Vas P, Dhataria K, et al. Guidelines on use of interventions to enhance healing of chronic foot ulcers in diabetes (IWGDF 2019 update). *Diabetes Metab Res Rev.* 2020;36(Suppl 1):e3283. doi:10.1002/dmrr.3283
13. Coccarelli A, Boileau E, Parthimos D, Nithiarasu P. An advanced computational bioheat transfer model for a human body with an embedded systemic circulation. *Biomech Model Mechanobiol.* 2016;15(5):1173–1190. doi:10.1007/s10237-015-0751-4
14. Namisnak LH, Haghayegh S, Khoshnevis S, Diller KR. Bioheat transfer basis of human thermoregulation: principles and applications. *J Heat Transfer.* 2022;144(3):031203. doi:10.1115/1.4053195
15. Park G, Kim J, Woo S, et al. Modeling heat transfer in humans for body heat harvesting and personal thermal management. *Appl Energy.* 2022;323:119609. doi:10.1016/j.apenergy.2022.119609
16. Pennes HH. Analysis of tissue and arterial blood temperatures in the resting human forearm. *J Appl Physiol.* 1948;1(2):93–122. doi:10.1152/jappl.1948.1.2.93
17. Santee WR, Reardon MJ, Pandolf KB. Modeling the physiological and medical effects of exposure to environmental extremes. *Military Quant Physiol.* 2012;2012:39–72.
18. Li B, Yang Y, Yao R, Liu H, Li Y. A simplified thermoregulation model of the human body in warm conditions. *Appl Ergon.* 2017;59(Pt A):387–400. doi:10.1016/j.apergo.2016.09.010
19. Zahra S, Róbert K, Péter V. Lagging heat models in thermodynamics and bioheat transfer: a critical review. *Cont Mech.* 2022;34(3):637–679. doi:10.1007/s00161-022-01096-6
20. Dinda A, Acharya J, Bhanja D, Nath S. Local thermal non-equilibrium bioheat transfer model for interstitial hyperthermia treatment of tumour cell: a numerical approach. *J Therm Biol.* 2022;110:103368. doi:10.1016/j.jtherbio.2022.103368
21. Dolibog P, Pietrzyk B, Kierszniok K, Pawlicki K. Comparative analysis of human body temperatures measured with noncontact and contact thermometers. *Healthcare.* 2022;10(2):331. doi:10.3390/healthcare10020331
22. Pecoraro V, Petri D, Costantino G, et al. The diagnostic accuracy of digital, infrared and mercury-in-glass thermometers in measuring body temperature: a systematic review and network meta-analysis. *Intern Emerg Med.* 2021;16(4):1071–1083. doi:10.1007/s11739-020-02556-0
23. Wang Q, Zhou Y, Ghassemi P, McBride D, Casamento JP, Pfefer TJ. Infrared thermography for measuring elevated body temperature: clinical accuracy, calibration, and evaluation. *Sensors.* 2021;22(1):215. doi:10.3390/s22010215
24. Yang B, Li X, Hou Y, Meier A, Cheng X, Choi JH. Non-invasive (non-contact) measurements of human thermal physiology signals and thermal comfort/discomfort poses—a review. *Energy.* 2020;224:110261.
25. Zhao Y, Bergmann JHM. Non-contact infrared thermometers and thermal scanners for human body temperature monitoring: a systematic review. *Sensors.* 2023;23(17). doi:10.3390/s23177439
26. Ramirez-GarciaLuna JL, Bartlett R, Arriaga-Caballero JE, Fraser RDJ, Saiko G. Infrared thermography in wound care, surgery, and sports medicine: a review. *Front Physiol.* 2022;13:838528. doi:10.3389/fphys.2022.838528
27. Lin YH, Chen YC, Cheng KS, Yu PJ, Wang JL, Ko NY. Higher periwound temperature associated with wound healing of pressure ulcers detected by infrared thermography. *J Clin Med Res.* 2021;10(13). doi:10.3390/jcm10132883
28. Fenemor SP, Gill ND, Sims ST, Beaven CM, Driller MW. Validity of a tympanic thermometer and thermal imaging camera for measuring core and skin temperature during exercise in the heat. *Meas Phys Educ Exerc Sci.* 2020;24(1):49–55. doi:10.1080/1091367X.2019.1667361
29. Chen HY, Chen A, Chen C. Investigation of the Impact of Infrared Sensors on Core Body Temperature Monitoring by Comparing Measurement Sites. *Sensors.* 2020;20(10). doi:10.3390/s20102885
30. Kalasin S, Sangnuang P, Surareungchai W. Intelligent wearable sensors interconnected with advanced wound dressing bandages for contactless chronic skin monitoring: artificial intelligence for predicting tissue regeneration. *Anal Chem.* 2022;94(18):6842–6852. doi:10.1021/acs.analchem.2c00782
31. Farahani M, Shafiee A. Wound healing: from passive to smart dressings. *Adv Healthc Mater.* 2021;10(16):e2100477. doi:10.1002/adhm.202100477
32. Tang N, Zheng Y, Jiang X, et al. Wearable sensors and systems for wound healing-related pH and temperature detection. *Micromachines.* 2021;12(4):430. doi:10.3390/mi12040430
33. Camera F, Marrocco G. Electromagnetic-based correction of bio-integrated RFID sensors for reliable skin temperature monitoring. *IEEE Sens J.* 2021;21(1):421–429.
34. Janus KA, Achtsnicht S, Drinic A, Kopp A, Keusgen M, Schöning MJ. Transient magnesium-based thin-film temperature sensor on a flexible, bioabsorbable substrate for future medical applications. *Appl Res.* 2023. doi:10.1002/appl.202300102
35. Zhang Y, Lin B, Huang R, et al. Flexible integrated sensing platform for monitoring wound temperature and predicting infection. *Microb Biotechnol.* 2021;14(4):1566–1579. doi:10.1111/1751-7915.13821
36. Pang Q, Yang F, Jiang Z, Wu K, Hou R, Zhu Y. Smart wound dressing for advanced wound management: real-time monitoring and on-demand treatment. *Mater Des.* 2023;229:111917. doi:10.1016/j.matdes.2023.111917
37. Lu SH, Samandari M, Li C, et al. Multimodal sensing and therapeutic systems for wound healing and management: a review. *Sensors Actuators Rep.* 2022;4:100075. doi:10.1016/j.snr.2022.100075
38. Ehtesabi H, Kalji SO, Movsesian L. Smartphone-based wound dressings: a mini-review. *Heliyon.* 2022;8(7):e09876. doi:10.1016/j.heliyon.2022.e09876
39. Dabas M, Schwartz D, Beeckman D, Gefen A. Application of artificial intelligence methodologies to chronic wound care and management: a scoping review. *Adv Wound Care.* 2023;12(4):205–240. doi:10.1089/wound.2021.0144
40. Derwin R, Patton D, Strapp H, Moore Z. Wound pH and temperature as predictors of healing: an observational study. *J Wound Care.* 2023;32(5):302–310. doi:10.12968/jowc.2023.32.5.302
41. Siah CJR, Childs C, Chia CK, Cheng KFK. An observational study of temperature and thermal images of surgical wounds for detecting delayed wound healing within four days after surgery. *J Clin Nurs.* 2019;28(11–12):2285–2295. doi:10.1111/jocn.14832

42. Zheng XQ, Huang JF, Lin JL, Chen D, Wu AM. Effects of preoperative warming on the occurrence of surgical site infection: a systematic review and meta-analysis. *Int J Surg.* 2020;77:40–47. doi:10.1016/j.ijsu.2020.03.016
43. Seidelman JL, Mantyh CR, Anderson DJ. Surgical site infection prevention: a review. *JAMA.* 2023;329(3):244–252. doi:10.1001/jama.2022.24075
44. Wyllie FJ, Sutherland AB. Measurement of surface temperature as an aid to the diagnosis of burn depth. *Burns.* 1991;17(2):123–127. doi:10.1016/0305-4179(91)90135-4
45. Liddington MI, Shakespeare PG. Timing of the thermographic assessment of burns. *Burns.* 1996;22(1):26–28. doi:10.1016/0305-4179(95)00076-3
46. Ganon S, Guédon A, Cassier S, Atlan M. Contribution of thermal imaging in determining the depth of pediatric acute burns. *Burns.* 2020;46(5):1091–1099. doi:10.1016/j.burns.2019.11.019
47. Carrière ME, de Haas LEM, Pijpe A, et al. Validity of thermography for measuring burn wound healing potential. *Wound Repair Regen.* 2020;28(3):347–354. doi:10.1111/wrr.12786
48. Fierheller M, Sibbald RG. A clinical investigation into the relationship between increased periwound skin temperature and local wound infection in patients with chronic leg ulcers. *Adv Skin Wound Care.* 2010;23(8):369–379; quiz 380–381. doi:10.1097/01.ASW.0000383197.28192.98
49. Dahllmanns S, Reich-Schupke S, Scholleman F, Stücker M, Leonhardt S, Teichmann D. Classification of chronic venous diseases based on skin temperature patterns. *Physiol Meas.* 2021;42(4):045001. doi:10.1088/1361-6579/abf020
50. Stevens K, Moralejo D, Ersser S, MacLean C. Effectiveness of a foot self-management intervention that utilized commercially available infrared thermometers: mixed methods research incorporating a pilot RCT. *J Tissue Viability.* 2023;32(1):33–38. doi:10.1016/j.jtv.2022.12.005
51. Ghosh A, Ray S, Garg MK, Chowdhury S, Mukhopadhyay S. The role of infrared dermal thermometry in the management of neuropathic diabetic foot ulcers. *Diabet Med.* 2021;38(4):e14368. doi:10.1111/dme.14368
52. Araújo AL, Negreiros FDS, Florêncio RS. Effect of thermometry on the prevention of diabetic foot ulcers. *Latino-Americana de Enfermagem.* 2022;30:e3567.
53. Zhong YF, Wang ZC, Xue YN, et al. The importance of temperature monitoring in predicting wound healing. *J Wound Care.* 2023;32(Sup6a):lxxxvii–xcvi. doi:10.12968/jowc.2023.32.Sup6a.lxxxvii
54. Ena J, Carretero-Gomez J, Arevalo-Lorido JC, Sanchez-Ardila C, Zapatero-Gaviria A, Gómez-Huelgas R. The association between elevated foot skin temperature and the incidence of diabetic foot ulcers: a meta-analysis. *Int J Low Extrem Wounds.* 2021;20(2):111–118. doi:10.1177/1534734619897501
55. Tarigan S, Yusuf S, Syam Y. Effect of interface pressure and skin surface temperature on pressure injury incidence: a turning schedule pilot study. *J Wound Care.* 2021;30(8):632–641. doi:10.12968/jowc.2021.30.8.632
56. Cai F, Jiang X, Hou X, et al. Application of infrared thermography in the early warning of pressure injury: a prospective observational study. *J Clin Nurs.* 2021;30(3–4):559–571. doi:10.1111/jocn.15576
57. Nakagami G, Sanada H, Iizaka S, et al. Predicting delayed pressure ulcer healing using thermography: a prospective cohort study. *J Wound Care.* 2010;19(11):465–466, 468, 470. passim. doi:10.12968/jowc.2010.19.11.79695
58. Schultz GS, Chin GA, Moldawer L, Diegelmann RF. *Principles of Wound Healing.* University of Adelaide Press; 2011.
59. Power G. PH, exudate composition and temperature measurement in wounds - A systematic review; 2016. Available from: [https://repository.rcsi.com/articles/thesis/pH\\_Exudate\\_Composition\\_and\\_Temperature\\_Measurement\\_in\\_Wounds\\_-\\_A\\_Systematic\\_Review/10820606/files/19330319.pdf](https://repository.rcsi.com/articles/thesis/pH_Exudate_Composition_and_Temperature_Measurement_in_Wounds_-_A_Systematic_Review/10820606/files/19330319.pdf). Accessed May 22, 2024.
60. Derwin R. The role of pH and temperature as biomarkers of wound healing; 2022. Available from: [https://repository.rcsi.com/articles/thesis/The\\_Role\\_of\\_pH\\_and\\_Temperature\\_as\\_Biomarkers\\_of\\_Wound\\_Healing/13049678](https://repository.rcsi.com/articles/thesis/The_Role_of_pH_and_Temperature_as_Biomarkers_of_Wound_Healing/13049678). Accessed May 22, 2024.
61. McGuinness W, Vella E, Harrison D. Influence of dressing changes on wound temperature. *J Wound Care.* 2004;13(9):383–385. doi:10.12968/jowc.2004.13.9.26702
62. Whitney JD, Salvadalena G, Higa L, Mich M. Treatment of pressure ulcers with noncontact normothermic wound therapy: healing and warming effects. *J Wound Ostomy Cont Nurs.* 2001;28(5):244–252. doi:10.1067/mjw.2001.117564
63. Kottner J, Black J, Call E, Gefen A, Santamaria N. Microclimate: a critical review in the context of pressure ulcer prevention. *Clin Biomech.* 2018;59:62–70. doi:10.1016/j.clinbiomech.2018.09.010
64. Shi C, Wang C, Liu H, et al. Selection of appropriate wound dressing for various wounds. *Front Bioeng Biotechnol.* 2020;8:182. doi:10.3389/fbioe.2020.00182
65. Zasadziński K, Spalek MJ, Rutkowski P. Modern dressings in prevention and therapy of acute and chronic radiation dermatitis—a literature review. *Pharmaceutics.* 2022;14(6):1204. doi:10.3390/pharmaceutics14061204
66. Legeza VI, Galenko-Yaroshevskii VP, Zinov'ev EV, et al. Effects of new wound dressings on healing of thermal burns of the skin in acute radiation disease. *Bull Exp Biol Med.* 2004;138(3):311–315. doi:10.1007/s10517-005-0029-4
67. Khan AA, Banwell PE, Bakker MC, Gillespie PG, McGrouther DA, Roberts AHN. Topical radiant heating in wound healing: an experimental study in a donor site wound model\*. *Int Wound J.* 2004;1(4):233–240. doi:10.1111/j.1742-4801.2004.00065.x

## Clinical, Cosmetic and Investigational Dermatology

Dovepress

### Publish your work in this journal

Clinical, Cosmetic and Investigational Dermatology is an international, peer-reviewed, open access, online journal that focuses on the latest clinical and experimental research in all aspects of skin disease and cosmetic interventions. This journal is indexed on CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-cosmetic-and-investigational-dermatology-journal>