

Effectiveness of Spironolactone in Reducing Osteoporosis and Future Fracture Risk in Middle-Aged and Elderly Hypertensive Patients

Shuaiwei Song^{1-5,*}, Xintian Cai^{1-5,*}, Junli Hu¹⁻⁵, Qing Zhu¹⁻⁵, Di Shen¹⁻⁵, Huimin Ma¹⁻⁵, Yingying Zhang¹⁻⁵, Rui Ma¹⁻⁵, Wenbo Yang¹⁻⁵, Jing Hong¹⁻⁵, Delian Zhang¹⁻⁵, Nanfang Li¹⁻⁵

¹Hypertension Center of People's Hospital of Xinjiang Uygur Autonomous Region, Urumqi, Xinjiang, 830001, People's Republic of China; ²Xinjiang Hypertension Institute, Urumqi, Xinjiang, 830001, People's Republic of China; ³NHC Key Laboratory of Hypertension Clinical Research, Urumqi, Xinjiang, 830001, People's Republic of China; ⁴Key Laboratory of Xinjiang Uygur Autonomous Region "Hypertension Research Laboratory", Urumqi, Xinjiang, 830001, People's Republic of China; ⁵Xinjiang Clinical Medical Research Center for Hypertension (Cardio-Cerebrovascular) Diseases, Urumqi, Xinjiang, 830001, People's Republic of China

*These authors contributed equally to this work

Correspondence: Nanfang Li, Hypertension Center of People's Hospital of Xinjiang Uygur Autonomous Region, No. 91 Tianchi Road, Urumqi, Xinjiang, 830001, People's Republic of China, Tel +86 8,564,818, Email lnanfang2016@sina.com

Objective: While the role of aldosterone in bone metabolism is well established, the specific effects of the widely used aldosterone antagonist, spironolactone, on bone health are not fully understood. This study aimed to investigate the effects of spironolactone on osteoporosis and future fracture risk in middle-aged and elderly hypertensive patients, revealing its potential benefits for bone health.

Methods: Propensity score matching was employed in this study to create matched groups of spironolactone users and non-users at a 1:4 ratio. We investigated the association between spironolactone use and the risk of osteoporosis using multivariate logistic regression analysis. Furthermore, we conducted multivariate linear regression analysis to explore the relationship between cumulative dosage and the FRAX score. Subgroup analysis was also performed to assess the effects under different stratification conditions.

Results: In both pre-match and post-match analyses, multivariable logistic regression revealed a significant reduction in the risk of osteoporosis in the spironolactone usage group (pre-match: odds ratios [OR] 0.406, 95% confidence interval [CI], 0.280–0.588; post-match: OR 0.385, 95% CI, 0.259–0.571). Furthermore, post-match multivariable linear regression demonstrated a clear negative correlation between cumulative spironolactone dosage and the FRAX score. Subgroup analyses consistently supported these findings.

Conclusion: This study offers evidence supporting the significant positive impact of the antihypertensive drug spironolactone on bone health, resulting in a substantial reduction in the risk of osteoporosis and future fractures in hypertensive patients. Future research should consider conducting large-scale, multicenter, randomized controlled trials to further investigate the long-term effects of spironolactone on bone health in hypertensive patients.

Keywords: hypertensive, spironolactone, aldosterone, osteoporosis, FRAX score

Introduction

Osteoporosis is a common systemic bone disease that results in weakened bones and an increased risk of fractures.¹ Hypertension and osteoporosis are two prevalent chronic diseases that often co-occur in middle-aged and older adults.² Globally, the reported incidence rate of hypertension is approximately 31%, while the incidence rate of osteoporosis can be as high as 21%.^{3,4} As life expectancy increases, the incidence rates of both hypertension and osteoporosis are expected to rise, underscoring the growing healthcare challenge posed by these conditions in an aging global population.^{5,6}

Hypertension is a risk factor for osteoporosis and osteoporotic fractures, significantly impacting morbidity and mortality in both men and women.^{7,8} While lifestyle interventions are crucial for managing blood pressure (BP), pharmacological treatments play a more prominent role in reducing BP and the risk of atherosclerotic cardiovascular disease. Various types

of antihypertensive medications are available, including angiotensin-converting enzyme inhibitors (ACEIs), angiotensin II receptor blockers (ARBs), and spironolactone, all of which are well-known inhibitors of the renin-angiotensin-aldosterone system (RAAS) and have demonstrated effective blood pressure control. Interestingly, an increasing body of cellular and animal research evidence suggests that localized activation of RAAS in bone tissue may contribute to osteoporosis.^{9–11}

Effective prevention and treatment of osteoporosis have traditionally relied on increased vitamin D and calcium intake, bisphosphonates, and nasal spray calcitonin, with limited mention of spironolactone.^{12–14} In recent years, a growing body of research has provided additional evidence supporting the notion that excessive aldosterone secretion within the RAAS system results in increased urinary calcium excretion. This disruption in calcium-phosphate metabolism ultimately elevates the risk of bone mineral loss.^{15–18} Spironolactone, a crucial aldosterone receptor antagonist, effectively suppresses aldosterone secretion.¹⁹ Animal experiments have shown that spironolactone can protect the skeletal health of rats with excessive aldosterone secretion.²⁰ Moreover, a study involving male congestive heart failure (CHF) patients suggested that spironolactone might have the potential to reduce the risk of fractures in this population.²¹ However, despite being a vital antihypertensive medication, there has been no investigation to date into its long-term use in hypertensive patients or its potential impact on bone mineral density (BMD). Additionally, it remains unclear whether spironolactone can reduce the risk of osteoporosis and future fractures, leaving specific outcomes in this regard uncertain.

Therefore, the main aim of this study was to examine the potential impact of extended spironolactone use on the risk of osteoporosis and future fractures in middle-aged and elderly hypertensive patients. To confirm this finding, we will use propensity score matching to rigorously match spironolactone users and non-users in this study.

Materials and Methods

Study Population

Inclusion Criteria

This study selected its participants from a pool of hypertensive patients who underwent BMD screening during hospitalization between January 2021 and December 2023. Initially, 2947 patients met the eligibility criteria.

Exclusion Criteria

The selection process involved several exclusion criteria. Firstly, individuals younger than 40 years old were excluded. Subsequently, patients with a history of previous fractures were also excluded. Additionally, individuals diagnosed with various bone metabolism-related disorders, including hypogonadism, Cushing's syndrome, hyperthyroidism, hyperparathyroidism, as well as those with severe hepatic and renal disorders, were not considered for the study. Finally, individuals who had previously taken medications known to affect BMD were also excluded. After applying these criteria, a total of 2344 participants remained and were enrolled in the final study.

This study was approved by the Research Ethics Committee of the Xinjiang Uygur Autonomous Region People's Hospital (KY2022080905). All the procedures complied with the requirements of the Declaration of Helsinki. All participants provided informed written consent.

Data Collection and Definitions

We collected demographic characteristics, clinical history, lifestyle details, physical examination findings, medication history, and laboratory data of the participants through electronic medical records. Detailed measurements, such as height, weight, body mass index (BMI), smoking status, and blood pressure, can be found in the [Supplementary Materials. Table S1](#) shows the names of the various drugs. Laboratory parameters included measurements of Alanine Aminotransferase (ALT), Aspartate Aminotransferase (AST), Creatinine (Cr), Alkaline Phosphatase (ALP), Thyroid Stimulating Hormone (TSH), as well as serum levels of potassium, calcium, phosphorus, Parathyroid Hormone (PTH), and 25-Hydroxyvitamin D. These measurements were conducted using a fully automated biochemical analyzer, and all hormone assessments followed current guidelines and were based on previous research conducted at our center.^{22–24} Definitions for various diseases can be found in the [Supplementary Materials](#).

Medication Use and Cumulative Drug Dose

Following previous related studies, we categorized individuals as spironolactone users if they had taken spironolactone continuously for a minimum of 6 months in the past.^{21,25} Information regarding participants' medication usage and duration was primarily extracted from their hospital records' medication history. The cumulative drug dose (in mg. months) was calculated as the daily dose (in mg) multiplied by the number of months.

Outcomes

Based on dual-energy X-ray absorptiometry (DXA) scanning, BMD measurements were obtained. Detailed explanations about these measurements are provided in the Supplementary Materials. The unique Chinese FRAX assessment tool algorithm is used to determine the probability of experiencing a major osteoporotic fracture (MOF) or hip fracture (HF) within a 10-year period (www.shef.ac.uk/FRAX).^{26,27} According to the latest guidelines, a BMD T-score of less than -1.0 at any site is considered as decreased bone mass, and a T-score of -2.5 or lower at any site indicates osteoporosis.^{28–30}

Propensity Score Matching

This study aimed to investigate the relationship between spironolactone use and osteoporosis. We employed propensity score matching, following the approach recommended by Lonjon et al.³¹ The propensity score, representing the likelihood of each patient using spironolactone, was generated using a logistic regression model. Our matching variables in the propensity score model included sex, age, BMI, menopausal status, and all antihypertensive and antidiabetic medications. Spironolactone users and non-users were matched using the nearest neighbor method at a 1:4 ratio based on the logit scale. The caliper width was limited to within 0.2 standard deviations, and the matching process was conducted without replacement. To assess the balance between variables in each group before and after matching, we utilized the Standardized Mean Difference (SMD), with an SMD value below 0.10 indicating a balanced distribution.^{32,33}

Statistical Analysis

We categorized study participants into two groups: the spironolactone users group and the non-users group, based on their spironolactone usage. We checked for multicollinearity using variance inflation factors (VIFs), and a VIF value of less than 10 for each variable indicated the absence of multicollinearity (Table S2 and Figure S1). To analyze the relationship between cumulative spironolactone dose and BMD and FRAX scores, we conducted multiple linear regression. Furthermore, we performed multiple logistic regression analyses to compare the risk of decreased bone mass and osteoporosis between the two groups. To assess dose-response relationships, we utilized restricted cubic spline (RCS) analysis, with additional two-stage comparisons based on the turning points of the RCS curves. Finally, to demonstrate the robustness of our results, we conducted both subgroup analysis and sensitivity analysis. For detailed statistical analyses, please refer to the Supplementary Materials.

All data were analyzed using R 4.2.2. Statistical significance was accepted for two-sided $P < 0.05$.

Results

Patient Selection

Figure 1 illustrates the participant selection process, with a total of 1300 participants being finalized for inclusion in our analysis after 1:4 matching (260 participants in the use group and 1040 participants in the non-use group).

Baseline Characteristics Before and After Propensity Score Matching

Table 1 displays the baseline characteristics before and after matching. Among all participants, individuals using spironolactone exhibited relatively higher levels of BMI, serum calcium, and TSH compared to non-users. Conversely, they had lower levels of serum potassium and ALP. In this group, rates of menopause and diabetes mellitus (DM) were comparatively lower, but there was a significantly higher prevalence of primary aldosteronism (PA). Additionally, these individuals were more likely to be taking various medications, including antihypertensives, lipid-lowering drugs, and




Figure 1 Flow chart of patient selection.

antiplatelet agents. The matching improved variable balance, with an absolute SMD < 0.10. The balance before and after propensity score matching is shown in [Figure S2](#).

Relationship Between the Use of Spironolactone and Reduced Bone Mass and Osteoporosis (User Vs Non-User)

Before matching, participants using spironolactone demonstrated a significantly lower risk of reduced bone mass compared to non-users, as evidenced in both Model 1 and the fully adjusted Model 4 ([Table S3](#)). This protective effect of spironolactone also extended to osteoporosis, where users exhibited a 60% lower risk of developing the condition compared to non-users (odds ratio [OR], 0.406; 95% confidence interval [CI], 0.280–0.588) ([Table 2](#)). After the matching process, the analysis consistently indicated a reduced risk of reduced bone mass among spironolactone users (OR, 0.371; 95% CI, 0.273–0.503) ([Table S4](#)). Furthermore, the negative correlation with osteoporosis was further supported by the results from Model 1 and the fully adjusted Model 4, which yielded ORs of 0.434 (95% CI, 0.294–0.624) and 0.385 (95% CI, 0.259–0.571), respectively ([Table 3](#)).

Relationship Between Cumulative Drug Doses and BMD and FRAX Scores

Multivariate linear regression analysis revealed a significant positive correlation between the cumulative dose of spironolactone and BMD at various sites, a correlation that remained significant even after adjustments in multivariable Model 4 ([Table S5](#)). Furthermore, concerning the risk of future fractures, we observed a consistent trend where an increase in the cumulative dose of spironolactone was associated with a reduced risk of future fractures (MOF and HF) across all models ([Table 4](#)). This consistent pattern observed in different analyses suggests that higher doses of spironolactone may have a protective effect on bone health.

Table I Baseline Characteristics Before and After Propensity Score Matching

Characteristic	Before Propensity Score Matching			After Propensity Score Matching		
	Non-user (N=2058)	User (N=286)	SMD	Non-user (N=1040)	User (N=260)	SMD
Age	56.43±10.87	56.99±10.87	0.051	57.42±11.17	56.72±10.68	0.064
Female (%)	1044 (50.73%)	158 (55.24%)	0.091	560 (53.85%)	141 (54.23%)	0.018
BMI (kg/m ²)	27.05±3.74	27.42±3.36	0.104	27.18±3.79	27.30±3.26	0.036
SBP (mmHg)	145.15±17.45	143.50±18.27	0.092	145.37±18.43	143.72±17.98	0.091
DBP (mmHg)	86.76±12.57	85.52±13.62	0.095	86.33±13.07	86.06±13.74	0.020
Current smoking (%)	524 (25.46%)	60 (20.98%)	0.106	241 (23.17%)	55 (21.15%)	0.024
Menopausal (%)	812 (39.46%)	98 (34.27%)	0.267	360 (34.62%)	92 (35.38%)	0.070
Laboratory tests						
Serum potassium (mmol/L)	3.90±0.34	3.81±0.36	0.248	3.89±0.33	3.81±0.36	0.215
PTH (pg/mL)	49.90 (35.80–68.68)	51.90 (40.45–66.20)	0.085	52.33 (38.20–70.28)	50.90 (39.60–65.90)	0.071
Serum calcium (mmol/L)	2.28±0.30	2.32±0.37	0.119	2.29±0.30	2.33±0.38	0.111
25-hydroxyvitamin D (nmol/L)	19.61 (12.39–29.50)	17.60 (11.35–27.45)	0.075	18.91 (12.01–28.58)	17.75 (11.30–27.80)	0.012
Serum phosphorus (mmol/L)	1.16±0.17	1.15±0.16	0.067	1.16±0.17	1.15±0.16	0.033
ALT (U/L)	23.43 (16.20–36.00)	22.00 (16.00–36.30)	0.019	23.00 (16.00–33.00)	22.05 (16.60–37.00)	0.092
AST (U/L)	20.30 (16.71–26.00)	20.00 (16.45–25.00)	0.043	20.20 (16.90–25.56)	20.00 (16.64–25.28)	0.005
Cr (umol/L)	64.65±15.98	66.03±15.45	0.088	65.70±16.60	65.46±15.31	0.015
ALP (U/L)	83.49±28.28	78.88±30.15	0.158	77.99 (61.70–97.00)	79.12 (57.00–103.00)	0.008
TSH (uIU/mL)	2.15 (1.42–3.34)	2.44 (1.74–3.33)	0.106	2.23 (1.45–3.33)	2.38 (1.71–3.33)	0.060
Urea (umol/L)	330.93±94.60	331.43±94.62	0.005	331.02±96.28	328.88±92.98	0.023
Medical history						
PA (%)	171 (8.31%)	138 (48.25%)	1.112	102 (9.81%)	130 (50.00%)	1.097
DM (%)	669 (32.51%)	54 (18.88%)	0.316	218 (20.96%)	54 (20.77%)	0.005
CHD (%)	153 (7.43%)	22 (7.69%)	0.010	83 (7.98%)	19 (7.31%)	0.019
Medications						
Statins (%)	381 (18.51%)	104 (36.36%)	0.408	323 (31.06%)	75 (28.85%)	0.049
Diuretics (%)	200 (9.72%)	62 (21.68%)	0.333	178 (17.12%)	51 (19.62%)	0.063
Beta-blockers (%)	345 (16.76%)	80 (27.97%)	0.271	255 (24.52%)	58 (22.31%)	0.063
Calcium channel blockers (%)	1069 (51.94%)	198 (69.23%)	0.359	700 (67.31%)	172 (66.15%)	0.013
ACEIs/ARBs (%)	824 (40.04%)	168 (58.74%)	0.381	586 (56.35%)	140 (53.85%)	0.053
Antihyperglycemic drug (%)	566 (27.50%)	48 (16.78%)	0.260	194 (18.65%)	47 (18.08%)	0.003
DXA BMD T-scores						
Lumbar 1	−0.93±1.56	−0.08±1.77	0.507	−0.90±1.57	−0.05±1.77	0.505
Lumbar 2	−0.87±1.58	−0.02±1.66	0.527	−0.85±1.61	0.01±1.65	0.531
Lumbar 3	−0.84±1.64	0.06±1.69	0.541	−0.79±1.65	0.09±1.68	0.534
Lumbar 4	−0.73±1.66	0.11±1.78	0.487	−0.68±1.67	0.12±1.76	0.467
Neck	−0.86±1.07	−0.58±0.92	0.280	−0.87±1.07	−0.57±0.92	0.301
Wards	−1.12±1.20	−0.95±1.06	0.148	−1.13±1.18	−0.94±1.06	0.171
Total	−0.20±1.08	0.04±0.95	0.235	−0.19±1.10	0.05±0.95	0.234
FRAX scores (%)						
MOF	3.69±2.83	3.08±1.80	0.257	3.66±2.80	3.03±1.78	0.267
HF	1.47±2.24	0.99±1.37	0.256	1.46±2.24	0.97±1.37	0.264
Outcomes						
Decreased bone mass	1517 (73.75%)	162 (56.64%)	0.350	765 (73.56%)	144 (55.38%)	0.380
Osteoporosis (%)	531 (25.81%)	38 (13.29%)	0.320	264 (25.38%)	33 (12.69%)	0.322

Notes: Data are presented as mean ± standard deviation, median (interquartile range), or as numbers, and percentages.

Abbreviations: BMI, body mass index; SBP, systolic blood pressure; DBP, diastolic blood pressure; PTH, parathyroid hormone; ALT, alanine transaminase; AST, aspartate transaminase; Cr, creatinine; ALP, alkaline phosphatase; TSH, thyroid stimulating hormone; PA, primary aldosteronism; DM, diabetes mellitus; CHD, coronary heart disease; ACEIs, angiotensin-converting enzyme inhibitors; ARBs, angiotensin receptor blockers; Neck, neck of the femur; Wards, Ward's triangle; Total, total femur; MOF, major osteoporotic fracture; HF, hip fracture, SMD Standardized Mean Difference.

Table 2 Effect of User versus Non-User on Osteoporosis Before Matching

Exposure	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)	Model 4 OR (95% CI)
Non-user	Reference	Reference	Reference	Reference
User	0.440 (0.261, 0.703)	0.424 (0.249, 0.683)	0.395 (0.231, 0.642)	0.406 (0.280, 0.588)

Notes: Model 1: no covariates were adjusted. Model 2: age, sex, BMI, and smoking status were adjusted. Model 3: Model 2 plus adjustment for serum potassium, PTH, serum calcium, serum phosphorus, and ALP. Model 4: Model 3 plus adjustment for the the use probability of spironolactone.
Abbreviations: OR, odds ratio; CI, confidence interval; BMI, body mass index; PTH, parathyroid hormone; ALP, alkaline phosphatase.

Table 3 Effect of User versus Non-User on Osteoporosis After Matching

Exposure	Model 1 OR (95% CI)	Model 2 OR (95% CI)	Model 3 OR (95% CI)	Model 4 OR (95% CI)
Non-user	Reference	Reference	Reference	Reference
User	0.434 (0.294, 0.624)	0.430 (0.290, 0.622)	0.386 (0.257, 0.566)	0.385 (0.259, 0.571)

Notes: Model 1: no covariates were adjusted. Model 2: age, sex, BMI, and smoking status were adjusted. Model 3: Model 2 plus adjustment for serum potassium, PTH, serum calcium, serum phosphorus, and ALP. Model 4: Model 3 plus adjustment for the the use probability of spironolactone.
Abbreviations: OR, odds ratio; CI, confidence interval; BMI, body mass index; PTH, parathyroid hormone; ALP, alkaline phosphatase.

Table 4 The Relationship Between Cumulative Drug Dose and FRAX Score

Exposure	Model 1 β (95% CI)	Model 2 β (95% CI)	Model 3 β (95% CI)	Model 4 β (95% CI)
MOF Cumulative dose (per 100 mg.months increase)	-0.076 (-0.099, -0.052)	-0.068 (-0.090, -0.046)	-0.037 (-0.054, -0.012)	-0.035 (-0.056, -0.014)
HF Cumulative dose (per 100 mg.months increase)	-0.041 (-0.059, -0.022)	-0.034 (-0.052, -0.016)	-0.010 (-0.024, -0.010)	-0.007 (-0.027, -0.007)

Notes: Model 1: no covariates were adjusted. Model 2: age, sex, BMI, and smoking status were adjusted. Model 3: Model 2 plus adjustment for serum potassium, PTH, serum calcium, serum phosphorus, and ALP. Model 4: Model 3 plus adjustment for the the use probability of spironolactone.
Abbreviations: MOF, major osteoporotic fracture; HF, hip fracture; β , regression coefficient; CI, confidence interval; BMI, body mass index; PTH, parathyroid hormone; ALP, alkaline phosphatase.

Relationship Between Cumulative Drug Dose, Reduced Bone Mass, and Osteoporosis

We investigated the dose-response relationship between cumulative drug dose and the risks of decreased bone mass and osteoporosis using RCS. [Figure S3](#) illustrates that when cumulative dosages exceed 370 mg.months, participants experience increased BMD and a reduced risk of decreased bone mass ([Tables S6](#) and [S7](#)). A similar pattern was observed with osteoporosis; as illustrated in [Figure 2](#), osteoporosis risk diminishes when the cumulative drug dose surpasses 400 mg.months. The analysis of these turning points reveals that cumulative drug doses above 400 mg.months are associated with a significant increase in BMD and a lower risk of osteoporosis compared to lower dosages ([Tables 5](#) and [S8](#)). These results further highlight the beneficial effects of spironolactone on bone health and may reduce the incidence of osteoporosis.

Subgroup Analysis and Sensitivity Analysis

In our subgroup analyses, we stratified participants based on factors including sex, age, BMI, smoking status, DM, and menopausal status. Across these subgroups, we observed consistent outcomes with respect to both reduced bone mass and osteoporosis ([Figures 3](#) and [S4](#)). Furthermore, when we conducted additional stratification by the type of medication used,




Figure 2 Dose-response relationship between cumulative drug dose and osteoporosis.

this consistency remained (Figures 4 and S5). Finally, in the sensitivity analysis, participants with PA were excluded, and the results remained essentially unchanged, whether conducted before or after matching (Tables S9–S12). These findings collectively suggest that spironolactone has a positive impact on bone mass across various circumstances.

Discussion

The results of the present study reveal a significant association between spironolactone use and both reduced bone mass and a lower risk of osteoporosis in middle-aged and elderly hypertensive patients. This association remained consistent across various subgroup analyses, underscoring the robustness of our findings. Furthermore, we observed a significant and positive correlation between the cumulative dose of spironolactone and BMD at all assessed sites, coupled with a reduced risk of future fractures. These combined findings emphasize that spironolactone, as an antihypertensive medication, offers substantial benefits to skeletal health and effectively mitigates the risk of osteoporosis.

Numerous previous studies have highlighted aldosterone's crucial role in regulating calcium and phosphorus metabolism.^{34–36} These studies reveal a bidirectional interaction between aldosterone and PTH, where aldosterone significantly increases urinary calcium excretion, subsequently stimulating PTH secretion. Furthermore, PTH enhances aldosterone secretion both by elevating calcium concentrations in adrenal glomerular zone cells and through angiotensin II induction. This interaction leads to continuous calcium loss from the body, ultimately resulting in persistent bone mass reduction.³⁷ In an experimental animal study on primary aldosteronism, researchers not only observed mineral loss but also noted a progressive decrease in cortical bone strength.¹¹ These adverse effects, however, were effectively mitigated by spironolactone treatment.^{38,39} A similar US study involving male patients with CHF indicated that spironolactone administration was negatively associated with fracture occurrence, suggesting its potential in reducing fracture risk.²¹

Table 5 The Impact of Cumulative Dose Before and After the RCS Turning Point on Osteoporosis

Osteoporosis	OR (95% CI)	P value
Turning point (mg.months)	400	
Cumulative dose < 400 mg.months	Reference	
Cumulative dose ≥ 400 mg.months	0.337 (0.204, 0.543)	<0.001

Notes: Age, sex, BMI, smoking status, serum potassium, PTH, serum calcium, serum phosphorus, ALP, and the use probability of spironolactone were adjusted.

Abbreviations: RCS, restricted cubic splines; OR, odds ratio; CI, confidence interval; BMI, body mass index; PTH, parathyroid hormone; ALP, alkaline phosphatase.




Figure 3 Association between cumulative drug dose and osteoporosis in various subgroups.




Figure 4 Association between cumulative drug dose and osteoporosis in medication subgroups.

Spironolactone has historically been regarded as an essential antihypertensive drug that plays a crucial role in managing blood pressure and treating HF.^{19,40,41} It is also linked to potential therapeutic benefits in stroke and cardiovascular disease, with a cohort study demonstrating that its administration and cumulative doses significantly lower stroke risks in hypertensive patients.⁴² Recent studies have suggested that spironolactone might positively affect bone health, although these investigations have largely been confined to animal models.^{38,39,43} This leaves the drug’s effects on bone health in humans, particularly its long-term impact on osteoporosis and fracture prevention in hypertensive patients, largely unexplored. Our study fills this gap by confirming spironolactone’s protective effect on BMD, highlighting its significant role in mitigating the risk of osteoporosis and future fractures in middle-aged and elderly hypertensive patients.

In our stratified analyses, we uncovered a notable finding: spironolactone’s effectiveness was significantly higher in menopausal women than in non-menopausal women. This difference in response may stem from the reduced levels of estrogen post-menopause, which tends to obscure spironolactone’s effects in non-menopausal women.⁴⁴ It’s essential to

recognize the pivotal role of estrogen in managing calcium and phosphorus metabolism.^{45–47} A decline in estrogen levels after menopause can cause an increase in bone resorption alongside a decrease in bone formation.^{47,48} This reduction in estrogen consequently leads to a lower bone mineral density, potentially progressing to osteoporosis. Nevertheless, further basic research is required to solidify our understanding of the precise causes and mechanisms behind these observations.

Our findings suggest that spironolactone's beneficial effects on bone health can be attributed to multiple mechanisms. Initially, spironolactone modulates aldosterone action, a hormone crucial for calcium and phosphorus metabolism, thereby potentially improving bone health.^{15,16,19} Additionally, it diminishes urinary calcium loss, aiding in the preservation of calcium balance within the body, a fundamental aspect of maintaining bone strength.^{49,50} Furthermore, the antiandrogenic properties of spironolactone may indirectly enhance estrogen metabolism, thereby elevating estrogen levels and mitigating bone loss in postmenopausal women.^{44,51,52} Moreover, spironolactone contributes to an increase in blood potassium levels. Research indicates that adequate potassium intake can prevent fractures, thus offering protective benefits for bones.^{53–55}

This study is the first to explore the link between spironolactone use and the risk of osteoporosis and future fractures in middle-aged and elderly hypertensive patients, marking a significant transition from laboratory research to clinical application. Its strengths lie in the comprehensive clinical data utilized, ensuring the reliability of results through propensity score matching to minimize group discrepancies, and the employment of two-stage comparisons and subgroup analyses for in-depth evaluation. In interpreting the results of this study, it is important to acknowledge its limitations. The cross-sectional research design precludes establishing causality. Although propensity score matching and multivariate analyses were used, there remains the possibility of residual bias and unmeasured confounding factors. Additionally, the reliance on medical record systems for medication information may introduce information bias. Crucially, the study did not include data on sex hormone levels, which are influential in bone metabolism. Lastly, the study's findings, based solely on Chinese hypertensive patients, may not be generalizable to other populations without careful consideration.

Conclusion

In conclusion, this study demonstrates that the antihypertensive drug spironolactone has a significant impact on the bone health of middle-aged and elderly hypertensive patients. It notably reduces the risk of osteoporosis and future fractures. This finding has substantial implications for clinical practice, suggesting that spironolactone could be a valuable therapeutic option not only for managing hypertension but also for mitigating the associated risks of bone health deterioration. Given the limitations of cross-sectional studies and the specific study population, larger prospective randomized controlled trials are warranted to further confirm these observations.

Funding

This work was sponsored by the Major Science and Technology Projects of the Xinjiang Uygur Autonomous Region (2022A03012-4).

Disclosure

The authors report no conflicts of interest in this work.

References

1. Ensrud KE, Crandall CJ. Osteoporosis. *Ann Intern Med.* 2017;167(3):ITC17–ITC32. doi:10.7326/AITC201708010
2. Cho HW, Jin HS, Eom YB. FGFR1 and FGF genes are associated with height, hypertension, and osteoporosis. *PLoS One.* 2022;17(8):e0273237. doi:10.1371/journal.pone.0273237
3. Mills KT, Bundy JD, Kelly TN, et al. Global disparities of hypertension prevalence and control: a systematic analysis of population-based studies from 90 countries. *Circulation.* 2016;134(6):441–450. doi:10.1161/CIRCULATIONAHA.115.018912
4. Xiao PL, Cui AY, Hsu CJ, et al. Global, regional prevalence, and risk factors of osteoporosis according to the World Health Organization diagnostic criteria: a systematic review and meta-analysis. *Osteoporos Int.* 2022;33(10):2137–2153. doi:10.1007/s00198-022-06454-3
5. Hurley DL, Khosla S. Update on primary osteoporosis. *Mayo Clin Proc.* 1997;72(10):943–949. doi:10.1016/S0025-6196(11)63367-3

6. Lu J, Lu Y, Wang X, et al. Prevalence, awareness, treatment, and control of hypertension in China: data from 1·7 million adults in a population-based screening study (China PEACE Million Persons Project). *Lancet*. 2017;390(10112):2549–2558. doi:10.1016/S0140-6736(17)32478-9
7. Gutzwiller JP, Richterich JP, Stanga Z, et al. Osteoporosis, diabetes, and hypertension are major risk factors for mortality in older adults: an intermediate report on a prospective survey of 1467 community-dwelling elderly healthy pensioners in Switzerland. *BMC Geriatr*. 2018;18(1):115. doi:10.1186/s12877-018-0809-0
8. Johnell O, Kanis JA. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int*. 2006;17(12):1726–1733. doi:10.1007/s00198-006-0172-4
9. Mo C, Ke J, Zhao D, et al. Role of the renin-angiotensin-aldosterone system in bone metabolism. *J Bone Miner Metab*. 2020;38(6):772–779. doi:10.1007/s00774-020-01132-y
10. Kao YT, Huang CY, Fang YA, et al. The association between renin angiotensin aldosterone system blockers and future osteoporotic fractures in a hypertensive population - A population-based cohort study in Taiwan. *Int J Cardiol*. 2020;305:147–153. doi:10.1016/j.ijcard.2019.12.069
11. Chhokar VS, Sun Y, Bhattacharya SK, et al. Loss of bone minerals and strength in rats with aldosteronism. *Am J Physiol Heart Circ Physiol*. 2004;287(5):H2023–6. doi:10.1152/ajpheart.00477.2004
12. Jackson RD, LaCroix AZ, Gass M, et al. Calcium plus vitamin D supplementation and the risk of fractures. *N Engl J Med*. 2006;354(7):669–683. doi:10.1056/NEJMoa055218
13. Black DM, Cummings SR, Karpf DB, et al. Randomised trial of effect of alendronate on risk of fracture in women with existing vertebral fractures. *Fracture Intervent Trial Res*. 1996;348(9041):1535–1541. doi:10.1016/s0140-6736(96)07088-2
14. Overgaard K, Riis BJ, Christiansen C, et al. Nasal calcitonin for treatment of established osteoporosis. *Clin Endocrinol*. 1989;30(4):435–442. doi:10.1111/j.1365-2265.1989.tb00443.x
15. Gao X, Yamazaki Y, Tezuka Y, et al. The crosstalk between aldosterone and calcium metabolism in primary aldosteronism: a possible calcium metabolism-associated aberrant “neoplastic” steroidogenesis in adrenals. *J Steroid Biochem Mol Biol*. 2019;193:105434. doi:10.1016/j.jsbmb.2019.105434
16. Wang A, Wang Y, Liu H, et al. Bone and mineral metabolism in patients with primary aldosteronism: a systematic review and meta-analysis. *Front Endocrinol*. 2022;13:1027841. doi:10.3389/fendo.2022.1027841
17. Salcuni AS, Palmieri S, Carnevale V, et al. Bone involvement in aldosteronism. *J Bone Miner Res*. 2012;27(10):2217–2222. doi:10.1002/jbmr.1660
18. Song S, Cai X, Hu J, et al. Correlation between plasma aldosterone concentration and bone mineral density in middle-aged and elderly hypertensive patients: potential impact on osteoporosis and future fracture risk. *Front Endocrinol*. 2024;15(1373862). doi:10.3389/fendo.2024.1373862
19. Batterink J, Stabler SN, Tejani AM, et al. Spironolactone for hypertension. *Cochrane Database Syst Rev*. 2010;8:CD008169. doi:10.1002/14651858.CD008169.pub2
20. Chhokar VS, Sun Y, Bhattacharya SK, et al. Hyperparathyroidism and the calcium paradox of aldosteronism. *Circulation*. 2005;111(7):871–878. doi:10.1161/01.CIR.0000155621.10213.06
21. Carbone LD, Cross JD, Raza SH, et al. Fracture risk in men with congestive heart failure risk reduction with spironolactone. *J Am Coll Cardiol*. 2008;52(2):135–138. doi:10.1016/j.jacc.2008.03.039
22. Cai X, Song S, Hu J, et al. Body roundness index improves the predictive value of cardiovascular disease risk in hypertensive patients with obstructive sleep apnea: a cohort study. *Clin Exp Hypertens*. 2023;45(1):2259132. doi:10.1080/10641963.2023.2259132
23. Hu J, Cai X, Zhu Q, et al. Relationship between plasma aldosterone concentrations and non-alcoholic fatty liver disease diagnosis in patients with hypertension: a retrospective cohort study. *Diabetes Metab Syndr Obes*. 2023;16:1625–1636. doi:10.2147/DMSO.S408722
24. Cai X, Song S, Hu J, et al. Author Correction: association of the trajectory of plasma aldosterone concentration with the risk of cardiovascular disease in patients with hypertension: a cohort study. *Sci Rep*. 2024;14(1):9827. doi:10.1038/s41598-024-60563-z
25. Chen HY, Ma KY, Hsieh PL, et al. Long-term effects of antihypertensive drug use and new-onset osteoporotic fracture in elderly patients: a population-based longitudinal cohort study. *Chin Med J*. 2016;129(24):2907–2912. doi:10.4103/0366-6999.195472
26. Wen Z, Li Y, Xu L, et al. Triglyceride glucose-body mass index is a reliable indicator of bone mineral density and risk of osteoporotic fracture in middle-aged and elderly nondiabetic Chinese individuals. *J Clin Med*. 2022;11(19):5694. doi:10.3390/jcm11195694
27. Ma H, Cai X, Hu J, et al. Association of systemic inflammatory response index with bone mineral density, osteoporosis, and future fracture risk in elderly hypertensive patients. *Postgrad Med*. 2024;16:1–11. doi:10.1080/00325481.2024.2354158
28. Wang L, Jiang J, Li Y, et al. Global trends and hotspots in research on osteoporosis rehabilitation: A bibliometric study and visualization analysis. *Front Public Health*. 2022;10:1022035. doi:10.3389/fpubh.2022.1022035
29. Watts NB, Camacho PM, Lewiecki EM, et al. American association of clinical endocrinologists/American College of Endocrinology Clinical Practice guidelines for the diagnosis and treatment of postmenopausal Osteoporosis—2020 Update. *Endocr Pract*. 2021;27(4):379–380. doi:10.1016/j.eprac.2021.02.001
30. Tang Y, Wang S, Yi Q, et al. High-density lipoprotein cholesterol is negatively correlated with bone mineral density and has potential predictive value for bone loss. *Lipids Health Dis*. 2021;20(1):75. doi:10.1186/s12944-021-01497-7
31. Lonjon G, Porcher R, Ergina P, et al. Potential pitfalls of reporting and bias in observational studies with propensity score analysis assessing a surgical procedure: a methodological systematic review. *Ann Surg*. 2017;265(5):901–909. doi:10.1097/SLA.0000000000001797
32. Austin PC. A tutorial and case study in propensity score analysis: an application to estimating the effect of in-hospital smoking cessation counseling on mortality. *Multivariate Behav Res*. 2011;46(1):119–151. doi:10.1080/00273171.2011.540480
33. Gu WJ, Duan XJ, Liu XZ, et al. Association of magnesium sulfate use with mortality in critically ill patients with sepsis: a retrospective propensity score-matched cohort study. *Br J Anaesth*. 2023;131(5):861–870. doi:10.1016/j.bja.2023.08.005
34. Rossi E, Sani C, Perazzoli F, et al. Alterations of calcium metabolism and of parathyroid function in primary aldosteronism, and their reversal by spironolactone or by surgical removal of aldosterone-producing adenomas. *Am J Hypertens*. 1995;8(9):884–893. doi:10.1016/0895-7061(95)00182-O
35. Pilz S, Kienreich K, Drechsler C, et al. Hyperparathyroidism in patients with primary aldosteronism: cross-sectional and interventional data from the GEOH study. *J Clin Endocrinol Metab*. 2012;97(1):E75–9. doi:10.1210/jc.2011-2183
36. Jiang Y, Zhang C, Ye L, et al. Factors affecting parathyroid hormone levels in different types of primary aldosteronism. *Clin Endocrinol*. 2016;85(2):267–274. doi:10.1111/cen.12981
37. Ceccoli L, Ronconi V, Giovannini L, et al. Bone health and aldosterone excess. *Osteoporos Int*. 2013;24(11):2801–2807. doi:10.1007/s00198-013-2399-1

38. Law PH, Sun Y, Bhattacharya SK, et al. Diuretics and bone loss in rats with aldosteronism. *J Am Coll Cardiol*. 2005;46(1):142–146. doi:10.1016/j.jacc.2005.03.055
39. Runyan AL, Chhokar VS, Sun Y, et al. Bone loss in rats with aldosteronism. *Am J Med Sci*. 2005;330(1):1–7. doi:10.1097/00000441-200507000-00001
40. Beldhuis IE, Myhre PL, Bristow M, et al. Spironolactone in patients with heart failure, preserved ejection fraction, and worsening renal function. *J Am Coll Cardiol*. 2021;77(9):1211–1221. doi:10.1016/j.jacc.2020.12.057
41. Tsujimoto T, Kajio H. Spironolactone use and improved outcomes in patients with heart failure with preserved ejection fraction with resistant hypertension. *J Am Heart Assoc*. 2020;9(23):e018827. doi:10.1161/JAHA.120.018827
42. Cai X, Li N. Association between use of spironolactone and risk of stroke in hypertensive patients: a cohort study. *Pharmaceuticals*. 2022;16(1):57. doi:10.3390/ph16010057
43. Altieri B, Muscogiuri G, Paschou SA, et al. Adrenocortical incidentalomas and bone: from molecular insights to clinical perspectives. *Endocrine*. 2018;62(3):506–516. doi:10.1007/s12020-018-1696-z
44. Seki K, Nagasaki M, Yoshino T, et al. Radiographical Diagnostic Evaluation of Mandibular Cortical Index Classification and Mandibular Cortical Width in Female Patients Prescribed Antiosteoporosis Medication: A Retrospective Cohort Study. *Diagnostics (Basel)*. 2024;14(10):1009. doi:10.3390/diagnostics14101009
45. Alemany M. The roles of androgens in humans: biology, metabolic regulation and health. *Int J Mol Sci*. 2022;23(19):11952. doi:10.3390/ijms231911952
46. Josse RG. Prevention and management of osteoporosis: consensus statements from the scientific advisory board of the osteoporosis society of Canada. 3. Effects of ovarian hormone therapy on skeletal and extraskelatal tissues in women. *CMAJ*. 1996;155(7):929–934.
47. Bansal N, Katz R, de Boer IH, et al. Influence of estrogen therapy on calcium, phosphorus, and other regulatory hormones in postmenopausal women: the mesa study. *J Clin Endocrinol Metab*. 2013;98(12):4890–4898. doi:10.1210/jc.2013-2286
48. Raisz LG. Pathogenesis of postmenopausal osteoporosis. *Rev Endocr Metab Disord*. 2001;2(1):5–12. doi:10.1023/a:1010074422268
49. Liu Y, Zhou L, Liu Z, et al. Higher blood urea nitrogen and urinary calcium: new risk factors for diabetes mellitus in primary aldosteronism patients. *Front Endocrinol*. 2020;11:23. doi:10.3389/fendo.2020.00023
50. Bayomy O, Zaheer S, Williams JS, et al. Disentangling the relationships between the renin-angiotensin-aldosterone system, calcium physiology, and risk for kidney stones. *J Clin Endocrinol Metab*. 2020;105(6):1937–1946. doi:10.1210/clinem/dgaa123
51. Xue B, Johnson AK, Hay M. Sex differences in angiotensin II- and aldosterone-induced hypertension: the central protective effects of estrogen. *Am J Physiol Regul Integr Comp Physiol*. 2013;305(5):R459–63. doi:10.1152/ajpregu.00222.2013
52. Olatunji LA, Adeyanju OA, Michael OS, et al. Ameliorative effect of low-dose spironolactone on obesity and insulin resistance is through replenishment of estrogen in ovariectomized rats. *Can J Physiol Pharmacol*. 2019;97(1):65–74. doi:10.1139/cjpp-2018-0416
53. Büssemaier E, Hillebrand U, Hausberg M, et al. Pathogenesis of hypertension: interactions among sodium, potassium, and aldosterone. *Am J Kidney Dis*. 2010;55(6):1111–1120. doi:10.1053/j.ajkd.2009.12.022
54. Lemann J, Pleuss JA, Gray RW, et al. Potassium administration reduces and potassium deprivation increases urinary calcium excretion in healthy adults [corrected]. *Kidney Int*. 1991;39(5):973–983. doi:10.1038/ki.1991.123
55. Bushinsky DA, Gavrillov K, Chabala JM, et al. Effect of metabolic acidosis on the potassium content of bone. *J Bone Miner Res*. 1997;12(10):1664–1671. doi:10.1359/jbmr.1997.12.10.1664

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>