

Radiofrequency vs Steroid Injections for Spinal Facet and Sacroiliac Joint Pain: A Systematic Review and Meta-Analysis

Bo Xu¹, Xudong Zhao², Lei Zhang¹, Shouhan Feng³, Jinxia Li¹, Ye Xu⁴

¹Acupuncture and Rehabilitation Department, Huzhou Traditional Chinese Medicine Hospital Affiliated to Zhejiang Chinese Medical University, Huzhou, Zhejiang, People's Republic of China; ²Physical Therapy Center, Huzhou Third Municipal Hospital, the Affiliated Hospital of Huzhou University, Huzhou, Zhejiang, People's Republic of China; ³Oncology Department, Huzhou Traditional Chinese Medicine Hospital Affiliated to Zhejiang Chinese Medical University, Huzhou, Zhejiang, People's Republic of China; ⁴Rehabilitation Department, Huzhou Third Municipal Hospital, the Affiliated Hospital of Huzhou University, Huzhou, Zhejiang, People's Republic of China

Correspondence: Ye Xu, Rehabilitation Department, Huzhou Third Municipal Hospital, the Affiliated Hospital of Huzhou University, No. 2088 Tiaoxi East Road, Huzhou City, Zhejiang Province, People's Republic of China, Email 1526827775@163.com

Purpose: Pain management for spinal facet joint (SFJ) and sacroiliac joint (SIJ) pain is challenging, often requiring interventions like radiofrequency ablation (RFA) or corticosteroid injections (CI). This study aims to assess and compare the effectiveness of CI and RFA in treating SFJ and SIJ pain. We combine these treatments due to their shared pathophysiology, similar therapeutic interventions, and the necessity for an integrated approach to spinal pain management.

Patients and methods: Literature search from PubMed, Scopus, CENTRAL and Google Scholar for published studies upto 31st December 2023, and reporting data of patients who were treated using CI or RFA for SFJ and SIJ pain. Pooled standardized mean difference (SMD) with a 95% Confidence Interval (CI) was calculated.

Results: Our meta-analysis incorporated thirteen studies. Overall, patients, treated with CI had a higher pain intensity score compared to patients treated with RFA (SMD=0.92; 95% CI: 0.19 to 1.65) at 3 months, and at 6 months (SMD=1.53; 95% CI: 0.66 to 2.40) after the treatment. No significant association was reported at 12 months (SMD=1.47; 95% CI: -0.03 to 2.97). Subgroup analysis based on joint types revealed increased pain intensity scores in patients who were treated with CI for SIJ (SMD=1.25; 95% CI: 0.39 to 2.11) and SFJ (SMD=1.33; 95% CI: 0.09 to 2.57) pain. A negative but not significant effect was detected in patients, treated with CI for cervical joint pain (SMD=-0.40; 95% CI: -0.90 to 0.10). Patients treated with CI exhibited higher functional disability score compared to patients treated with RFA at 3 months (SMD=1.28; 95% CI: 0.20 to 2.35) post-treatment.

Conclusion: This study suggests that RFA may offer superior pain relief with longer duration compared to steroid injections for spinal facet and sacroiliac joint pain. Decision regarding specific interventions should be individualized and consider patient preferences, clinical context, and potential risks.

Keywords: radiofrequency ablation, steroid injections, spinal facet pain, sacroiliac joint pain, systematic review, meta-analysis, pain management

Introduction

Pain management for spinal facet joint (SFJ) and sacroiliac joint (SIJ) pain is challenging, and interventions, aimed to alleviate discomfort and improve patients' quality of life are crucial.^{1,2}

Over 25% of individuals with chronic lower back pain are reported to have SFJ, or more specifically, lumbar facet joint pain that is predominantly linked to degenerative osteoarthritis.^{3,4} SIJ pain accounts for up to 20% of low back pain cases, and poses its own set of challenges.⁵ Valid clinical and diagnostic tests for SIJ pain remain elusive, leading to diverse diagnostic approaches.⁶

Both SFJ and SIJ pain result from similar underlying mechanisms, including joint degeneration, inflammation, and nerve irritation, suggesting that therapies targeting one condition may also be effective for the other. An integrated

approach is essential for effectively managing spinal pain, given the interconnected nature of spinal structures and the common occurrence of multi-faceted pain syndromes. Although the SIJ and facet joints differ in size and function—the SIJ being larger and primarily responsible for load transfer between the spine and lower limbs, while facet joints facilitate spinal movement and stability—there may be overlapping pain mechanisms and treatment responses. Investigating these conditions concurrently allows for the evaluation of shared diagnostic tools and therapeutic approaches, potentially leading to more efficient and effective management strategies. Additionally, comparing these distinct but related pain sources can highlight similarities and differences in their pathophysiology, enhancing the precision of interventions tailored to each condition.

Recently, radiofrequency ablation (RFA) and corticosteroid injections (CI) have emerged as prominent therapeutic modalities in pain management of such patients.^{7–9} The primary interventions for both conditions, RFA and CI, work by reducing inflammation and disrupting pain signals, supporting a combined therapeutic strategy. Injection with corticosteroids have shown promise in managing SFJ-related lower back pain.¹⁰ Additionally, the application of RFA in patients with SFJ pain has demonstrated effectiveness in reducing pain scores and enhancing overall quality of life.^{11,12} Current treatment options for patients with SIJ pain include pharmacotherapy, chiropractic manipulation, CI injections, and surgical fixation.¹³ Recently, RFA has gained popularity for SIJ pain management, with studies reporting promising results.^{14,15}

However, due to variability in methodologies, including differences in techniques and selection criteria, the comparative effectiveness of RFA and CI in pain management of patients with SFJ and SIJ problem is still unclear. Given the significant impact of chronic spinal pain on quality of life, physical function, and healthcare resources, identifying the optimal treatment modality is essential. Insurance coverage for these procedures often varies, with some insurers being more inclined to cover steroid injections over RFA or vice versa. This variability can affect patients' access to the most effective treatments. By providing a systematic comparison of the efficacy of these interventions, our study offers valuable insights that can inform insurance policy decisions. This meta-analysis aims to comparing the efficiency of CI and RFA in managing pain associated with SFJ and SIJ. Our results may provide valuable insights into the optimal approach for pain management of these conditions, guiding clinicians in making informed decisions for the benefit of their patients, reducing reliance on opioids, and enhancing overall management strategies for spinal pain.

Methods

Protocol Registration

We adhered to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)¹⁶ guidelines. The study is registered with PROSPERO with the identifier no. **CRD42024496067**.

Search Strategy

An electronic search was conducted on PubMed, Scopus, CENTRAL (Cochrane Central Register of Controlled Trials), and Google Scholar databases for English language papers published up to 31st December 2023. The search key terms included variations of “radiofrequency ablation”, such as “RF ablation” and “Radiofrequency”, and terms related to steroid injections, such as “Steroid injections” and “Corticosteroid injections”. The focus extended to spinal facet joint pain, represented by terms like “Spinal facet joint pain” and “Facet joint syndrome”, as well as sacroiliac joint pain, identified by terms like “Sacroiliac joint pain” and “Sacroiliitis”. Additionally, we included terms associated with osteoarthritis, such as “osteoarthritis”, “degenerative joint disease”, and “OA”, and terms related to axial pain, like “axial pain”, “axial back pain”, and “axial spine pain”. Our focus also extended to terms representing spinal facet joint pain, including “spinal facet joint pain” and “facet joint syndrome”, and sacroiliac joint pain, identified by terms like “sacroiliac joint pain” and “sacroiliitis”. Bibliography of retrieved studies and relevant reviews were further screened for the potential inclusion of additional studies.

Inclusion Criteria (PICOS)

Population (P): Patients with spinal facet joint (SFJ) or sacroiliac joint (SIJ)

Intervention (I): Radiofrequency ablation

Comparison (C): Corticosteroid injections
Outcome (O): Pain intensity
Study Design (S): Randomized controlled trials (RCTs).

Exclusion Criteria

1. Non-English or non-human studies
2. Studies lacking adequate data or with short follow-up periods
3. Mixed interventions, inappropriate comparators, or high risk of bias
4. Irrelevant conditions or duplicate data, published beyond a predefined date.

Data Extraction

During the initial phase of our systematic review, databases were searched, and titles and abstracts of identified studies were independently assessed by the two authors. Subsequently, full texts of papers meeting eligibility criteria were obtained for further analysis. The comprehensive data extraction process included key details such as author information, publication year, age distribution within the studied population, sample size, intervention specifics, pain intensity scores, functional disability scores, and any reported adverse events.

Outcomes

The primary outcome, pain intensity, was evaluated by visual analogue scale (VAS) or numerical rating scale (NRS). Pain intensity measurements were categorized into three distinct follow-up periods (3, 6, and 12 months after the procedure). Functional disability was measured using the Oswestry Disability Index (ODI) at 3 months after the treatment.

Quality Assessment

We used Grading of Recommendations, Assessment, Development, and Evaluations (GRADE), and specifically, GRADEpro tool, to assess evidence certainty for the outcomes. This tool considers factors such as study design, risk of bias, inconsistency, indirectness, imprecision, and publication bias.

Publication Bias

Publication bias was assessed by funnel plot analysis¹⁷ and Egger's regression test.¹⁸

Statistical Analysis

All statistical analyses were done by STATA, version 12.0 (Stata Statistical Software, Release 12; StataCorp LP, College Station, TX) and Review Manager (RevMan v5.3 2014; Cochrane, London, England).

Results were presented as standardized mean difference (SMD) with 95% confidence interval (CI) for pain intensity and functional disability. Heterogeneity was assessed by the I^2 statistic. In cases of significant ($I^2 > 50\%$) heterogeneity, random effects model was used. For $I^2 < 50\%$, fixed-effect model was used. Sensitivity analyses were done to validate the pooled observed effect. $P < 0.05$ was statistically significant.

Results

The PRISMA flow diagram of our study (Figure 1) illustrates the systematic process of identifying and selecting relevant studies. Our initial systematic search across various databases yielded a total of 117 studies. After removing duplicate entries, we were left with 46 unique articles. These full texts were then meticulously screened for eligibility based on our inclusion and exclusion criteria. During this screening process, 31 studies were deemed irrelevant as they did not meet our criteria, and 2 studies were excluded because they did not report outcomes of interest. Finally, 13 studies^{19–31} were incorporated in the analysis.




Figure 1 PRISMA 2020 Flow diagram for the selection of studies and specific reasons for exclusion from the present meta-analysis.

Note: *Databases: PubMed, Scopus, CENTRAL and Google Scholar; **31 studies were deemed irrelevant as they did not meet our criteria; irrelevant populations (n=14), different interventions (n=5), lacking a comparator group (n=7), and non-randomized design (n=5).

Characteristics of Studies

As shown in Table 1, eight out of 13 studies focused on patients with lumbar pain, two on cervical pain, and three on sacroiliac joint pain. The publication years of the studies ranged from 2012 to 2023, with sample sizes varying between 11 and 50 participants, and follow-up durations spanning from 6 to 12 months. Six studies were conducted in Asian populations, while seven were carried out in Caucasian populations.

All studies incorporated combined regimen of corticosteroid and local anesthetics for the corticosteroid groups. Methylprednisolone was used in six trials, typically in dosages ranging from 20 mg to 40 mg and often combined with varying amounts of bupivacaine. Betamethasone, used in three trials, had dosages ranging from 3 mg to 12 mg, while dexamethasone, used in two trials, was consistently dosed at 10 mg. Additionally, triamcinolone was used at 10 mg. These steroids were frequently combined with local anesthetics such as bupivacaine or lidocaine to enhance their effectiveness. Eight trials utilized pulsed radiofrequency (PRF), and five trials applied continuous RF (CRF). Adverse events were not reported in seven trials. A study by Lim et al²⁵ reported an increase in lower back pain (LBP) for two patients during the follow-up period, while another study by Do et al¹⁹ reported hyperglycemia in one patient.

Main Findings

Pain Intensity at 3 Months

Nine studies reported data for pain intensity score at 3 months (Figure 2a). CI correlated with higher pain intensity score compared to RFA (SMD=0.92; 95% CI: 0.19 to 1.65), with substantial heterogeneity ($I^2 = 93\%$). Subgroup analysis based on type of joints showed that CI correlated with higher pain intensity score compared to RFA in patients with sacroiliac region (SMD=1.25; 95% CI: 0.39 to 2.11) and lumbar region (SMD=1.33; 95% CI: 0.09 to 2.57) pain.

Table 1 Characteristics of Included Studies Investigating for Radiofrequency vs Steroid Injections for Spinal Facet and Sacroiliac Joint Pain

| S. No | Study | Country | Joint Type | Inclusion Criteria | Sample Size | | Male/Female | | Procedures | | Adverse Events | | Mean Age ± SD | | Measured Scale | Imaging Modality | Follow up Duration (months) |
|-------|-------------------------------------|---------|------------|---|-------------|-----|-------------|-------|---|---|----------------------------|--|---------------|--------------|---------------------|------------------|-----------------------------|
| | | | | | CI | RFA | CI | RFA | CI | RFA | CI | RFA | CI | RFA | | | |
| 1. | Civelek et al, 2012 ²⁰ | Turkey | Lumbar | Failure to respond 6 wks after therapy | 50 | 50 | 11/39 | 17/33 | MBB mixture of MPS40 mg + 2 mL of bupivacaine 0.25–0.50% | MBB CRF 80°C in 120 sec | None | Neuropathy in 2 subjects, Minor burns in few subjects; | 56.5 ± 17.7 | 51.8 ± 17 | VNS, NASS and EQ-5D | Fluoroscopy | 12 |
| 2. | Duger et al, 2012 ²¹ | Turkey | Lumbar | ≥ 6 months history of LBP | 40 | 40 | - | - | IA 5 mL mixture of MPS20 mg + 5 mg bupivacaine | PRF at 40°C in 360 sec | - | - | - | - | VAS | - | - |
| 3. | Lakemeier et al, 2013 ²⁴ | Germany | Lumbar | ≥ 24 months history of non-specific LBP | 26 | 26 | 16/10 | 17/9 | IA mixture of 1 mL of betamethasone 3mg + 0.5 mL of bupivacaine 0.5% | MBB CRF at 80°C in 90 sec | None | None | 56.3 ± 10.8 | 57.6 ± 12.8 | RMQ,VAS, ODI | Fluoroscopy | 6 |
| 4. | Hashemi et al, 2014 ²³ | Iran | Lumbar | ≥ 6 months history of LBP | 40 | 40 | 28/12 | 29/11 | MBB using MPS 40 mg+ 0.5 mL bupivacaine 0.5% | MBB PRF at 42°C in 120 sec (2 shots of 45 V/s) | NR | NR | 63.85 ± 11.46 | 64.32 ± 13.2 | NRS,ODI | Fluoroscopy | 12 |
| 5. | Martinez et al, 2016 ²² | Spain | Sacroiliac | Intense SIJ > 3 months | 20 | 20 | 6/14 | 5/15 | 2 USG guided IA SIJ blocks with 3 mL 5% levobupivacaine and 12 mg betamethasone sodium, with a 7-day interval | Six 10 cm,10 mm active tip, 22G cannula and 4 electrodes were used to create 2 overlapping bipolar lesions with 4 cannula (90-C,3 min). | None | Moderate Pain | 52.6 ± 14.1 | 53.2 ± 13.9 | VAS | USG | 12 |
| 6. | Zhou et al, 2016 ³¹ | China | Lumbar | ≥ 6 months history of LBP | 40 | 40 | 23/17 | 21/19 | MBB and IA injection using 5 mL mixture of betamethasone 6 mg + lidocaine 20 mg | MBB CRF at 80°C in 90 sec | None | None | 54.6 ± 7.5 | 56.5 ± 8.7 | VAS, Schober index | X-ray | 6 |
| 7. | Shin et al, 2017 ²⁹ | Korea | Cervical | Complaint of suboccipital neck pain predominantly for at least 6 months | 11 | 12 | 3/8 | 4/8 | 2% lidocaine, 0.75 mL + Triamcinolone 10 mg, 0.25 mL | PRF 5 Hz and 5 ms pulse width for 360 sec at 55 V | None | None | 47.6 ± 6.6 | 47.6 ± 6.8 | NRS | Fluoroscopy | 6 |
| 8. | Do et al, 2017 ¹⁹ | Korea | Lumbar | 6 months history of LBP | 30 | 30 | 12/18 | 12/18 | IA 0.5 mL mixture of dexamethasone 10 mg + 0.25 mL of bupivacaine 0.125% | IA PRF at 42°C in 360 sec (5 Hz, a5-ms pulsed width at 55 V) | Hyperglycemia in 1 subject | None | 63 ± 10.9 | 66.9 ± 9.6 | NRS | Fluoroscopy | 6 |

(Continued)

Table I (Continued).

| S. No | Study | Country | Joint Type | Inclusion Criteria | Sample Size | | Male/Female | | Procedures | | Adverse Events | | Mean Age \pm SD | | Measured Scale | Imaging Modality | Follow up Duration (months) |
|-------|-------------------------------------|---------|------------|---|-------------|-----|-------------|-------|--|---|---------------------|------|-------------------|-----------------|----------------|------------------|-----------------------------|
| | | | | | CI | RFA | CI | RFA | CI | RFA | CI | RFA | CI | RFA | | | |
| 9. | Yasar et al, 2018 ³⁰ | Turkey | Lumbar | ≥ 3 months history of LBP | 50 | 50 | 34/16 | 40/10 | 2.5 mL mixture of MPS40 mg + bupivacaine 0.25% | PRF at 42°C in 120sec (2 shots of 45 V/s) | NR | NR | 43.1 \pm 8.3 | 47.4 \pm 11.1 | VAS, ODI | x-ray | 12 |
| 10. | Dutta et al, 2018 ²² | India | Sacroiliac | LBP ≥ 3 months | 15 | 15 | 10/5 | 4/11 | 3 mL solution containing 2 mL of bupivacaine 0.5% and 1 mL of 40 mg/mL DEPO-MEDROL | PRF parameters were 45 V for 180 s at all levels using the RF generator. Temperatures varied from 38°C to 42°C. | None | None | 41.6 (27–60) | 43.1 (19–76) | NRS,ODI | Fluoroscopy | 6 |
| 11. | Salman et al, 2016 ²⁸ | Egypt | Sacroiliac | LBP for 6 months | 15 | 15 | 7/8 | 5/10 | 3 mL solution containing 2 mL of Lidocaine 2%- and 1-mL bupivacaine 0.5% | RF probes were sequentially inserted into the cannula and 90s 80 °C lesions were performed | NR | NR | 51.8 \pm 13.1 | 51.9 \pm 13.6 | VAS | Fluoroscopy | 6 |
| 12. | Lim et al, 2017 ²⁵ | Korea | Cervical | 3-month history of axial cervical pain without radicular symptoms | 20 | 20 | 10/10 | 7/13 | 0.3 mL of contrast into CFJ space, injected 10 mg (0.25 mL) of dexamethasone, mixed with 0.25 mL of 0.125% bupivacaine | 5 Hz and a 5-ms pulsed width, for 360 seconds, at 55 V, under the condition that the electrode tip temperature did not exceed 42°C. | Minor in 2 patients | None | 52.7 \pm 12.1 | 52.8 \pm 12.7 | NRS | Fluoroscopy | 6 |
| 13. | McCormick et al, 2023 ²⁷ | USA | Lumbar | Unilateral or bilateral axial (non-radicular) LBP for 3 months | 12 | 20 | 4/8 | 10/10 | 0.5 mL of 40 mg/mL Kenalog and 0.5mL of 2% preservative-free lidocaine | C-RFA lesions performed for 165 seconds, with the RFA generator temperature set to 60 °C (intralesional temperature >80 °C) | None | None | 62.4 \pm 14.6 | 62.2 \pm 12.4 | NPRS | Fluoroscopy | 12 |

Abbreviations: CI, corticosteroid injection; CRF, continuous radiofrequency; IA, intraarticular; LBP, Low back pain; LFJ, lumbar facet joint; MBB, medial branch block; NRS, numeric rating scale; VAS, visual analog scale, ODI, Oswestry Disability Index; RMQ, Roland, Morris Low Back Pain and Disability Questionnaire; EQ-5D, EuroQol, 5 Dimension; PRF, pulsed radiofrequency; SFJ, spinal facet joint; SIJ, sacroiliac joint; IA, intraarticular; MBB, medial branch block; NR, Not Reported; Sec, seconds; MPS, Methylprednisolone; USG, Ultrasonography; ms, millisecond; v, Volts; wks, weeks; NPRS, numerical pain rating scale.




Figure 2 (a-d) Forest Plot for comparing the pain intensity in Radiofrequency vs steroid injections for spinal facet and sacroiliac joint pain across various time points (a) Pain intensity at 3 months, (b) Pain intensity at 6 months, (c) Pain intensity at 12 months and (d) Functional disability at 3 months.

Negative, although not statistically significant effect of CI was reported in patients with cervical region pain (SMD=-0.40; 95% CI: -0.90 to 0.10).

Pain Intensity at 6 Months

Eleven studies reported data of pain intensity score at 6 months (Figure 2b). CI led to higher pain intensity score than RFA (SMD=1.53; 95% CI: 0.66 to 2.40), with substantial heterogeneity ($I^2 = 95.4\%$). Subgroup analysis based on the type of joints showed higher pain intensity score in CI group compared to the RFA group in patients with the sacroiliac region (SMD=1.62; 95% CI: 0.79 to 2.45) and lumbar region (SMD=1.94; 95% CI: 0.92 to 2.97) pain. CI treatment was associated with negative effect in patients with the cervical region pain (SMD=-0.24; 95% CI: -0.74 to 0.26), but the difference was not statistically significant.

Pain Intensity at 12 Months

Five studies reported data of pain intensity at 12 months after the treatment (Figure 2c). Pain scores on the CI and the RFA groups were comparable (SMD=1.47; 95% CI: -0.03 to 2.97), with substantial heterogeneity ($I^2 = 97\%$). Subgroup analysis based on type of joints also showed comparable pain intensity scores in patients with sacroiliac region (SMD=0.10; 95% CI: -0.52 to 0.72) and lumbar region (SMD=1.82; 95% CI: -0.03 to 3.66) pain. No data were available for pain scores of patients with cervical region pain.

Functional Disability at 3 Months

Five studies reported data of functional disability scores at 3 months after the treatment. As shown in Figure 2d, patients in the CI group had a higher functional disability score compared to RFA group (SMD=1.28; 95% CI: 0.20 to 2.35), with substantial heterogeneity ($I^2 = 93.6\%$). Subgroup analysis based on type of joints also showed the CI correlated with significantly higher functional disability than RF ablation in patients with the sacroiliac region (SMD=0.75; 95% CI: 0.01 to 1.50) and lumbar region (SMD=1.41; 95% CI: 0.08 to 2.73) pain. No data were available for patients with the cervical region pain.

Sensitivity Analysis

Sensitivity analysis demonstrated that the pooled SMD estimates were not significantly influenced by the removal of any individual study (Figure 3). These findings indicate the robustness of our meta-analysis results, reassuring that they are not driven by any single study, enhancing confidence in the validity of our findings.

Publication Bias

Funnel plot analysis revealed no apparent signs of asymmetry, with studies evenly distributed around the estimated effect. This suggests the absence of significant publication bias that could impact the results. Funnel plots for pain intensity at 3 months ($p = 0.55$), 6 months ($p = 0.35$), and 12 months ($p = 0.30$) and functional disability at 3 months ($p = 0.69$) are shown in Figure 4a–d.

Quality Assessment

Figure 5 illustrates the assessment of bias across various domains. The evaluation considered the method of randomization, adherence to interventions, management of missing data, precision in measurement, and reporting acceptability in a total of nine studies. Table 2 provides a summary of findings with each outcome based on GRADEpro, enhancing the understanding of the reliability of findings from the studies.

Discussion

Our findings emphasize the potential superiority of RFA in providing sustained pain relief for patients with chronic LBP. Chronic conditions such as LBP often necessitate long-term investment in conservative treatments, which are frequently




Figure 3 Sensitivity plot for comparing the pain intensity in radiofrequency vs steroid injections for spinal facet and sacroiliac joint pain.




Figure 4 (a-d) Funnel Plot of for comparing the pain intensity in radiofrequency vs steroid injections for spinal facet and sacroiliac joint pain at various time point.

limited in their efficacy. In this context, radiofrequency thermocoagulation presents itself as both an effective and relatively cost-effective alternative to traditional surgery.¹² Conventional radiofrequency thermocoagulation induces localized injury with well-defined boundaries by applying radiofrequency current to heat adjacent tissue.³³ Studies have shown that temperatures up to 80°C can lead to the dissolution and necrosis of unmyelinated nerve C fibers, blocking neuralgia-conducting nerves for extended periods. In contrast, pulsed radiofrequency does not induce neuro-destructive processes like conventional radiofrequency,³⁴ a distinction particularly relevant for sensitive nerves in regions such as the head, neck, dorsal root ganglia, and trigeminal ganglia.

Intraarticular (IA) injections, while potentially offering greater accuracy due to varied facet joint innervation, pose challenges due to their difficulty and increased pain compared to medial branch block (MBB). The choice between IA and MBB approaches may necessitate a trade-off between accuracy and patient comfort.³⁵ The included studies employed diverse techniques for RFA, lacked standardized approaches, and reported varied adverse events. Different pain assessment scales and follow-up intervals were utilized, with no consensus on a superior technique or standardization. Stricter criteria for MBBs have been associated with improved outcomes in lumbar RFA, while more relaxed standards have shown reduced efficacy. The absence of a standardized approach or superior technique indicates a lack of conclusive evidence supporting one method over another, highlighting the need for further research and standardization in RFA procedures to enhance clarity for clinicians and improve patient outcomes.

The variations in steroid dosages and methodologies across the included studies present significant challenges in drawing firm conclusions about their efficacy and safety. Different types and dosages of steroids were combined with various local anesthetics in differing concentrations, making direct comparisons difficult. The methodologies employed

| | Random sequence generation (selection bias) | Allocation concealment (selection bias) | Blinding of participants and personnel (performance bias) | Blinding of outcome assessment (detection bias) | Incomplete outcome data (attrition bias) | Selective reporting (reporting bias) | Other bias |
|-------------------------------------|---|---|---|---|--|--------------------------------------|------------|
| Civelek et al. 2012 ²⁰ | + | - | + | + | - | + | + |
| Do et al. 2017 ¹⁹ | + | + | ? | + | ? | + | + |
| Duger et al. 2012 ²¹ | + | - | ? | ? | - | ? | ? |
| Dutta et al. 2018 ²² | + | + | + | ? | + | + | + |
| Hashemi et al. 2018 ²³ | ? | ? | ? | ? | + | + | + |
| Lakemeier et al. 2013 ²⁴ | + | + | + | + | + | + | + |
| Lim et al. 2017 ²⁵ | ? | ? | + | ? | + | + | ? |
| Martinez et al. 2016 ²⁶ | + | ? | ? | ? | ? | + | ? |
| McCormick et al. 2023 ²⁷ | - | - | - | - | + | + | ? |
| Salman et al. 2016 ²⁸ | + | + | + | + | + | + | + |
| Shin et al. 2017 ²⁹ | + | + | + | + | + | + | + |
| Yasar et al. 2018 ³⁰ | + | + | + | + | + | + | ? |
| Zhou et al. 2016 ³¹ | ? | + | + | + | + | + | ? |

Figure 5 Risk of Bias Summary.

Table 2 Summary of Evidences Using GRADEpro

| Certainty assessment | | | | | | | № of Patients | | Effect | | Certainty | Importance |
|---|-------------------|--------------|---------------|--------------|-------------|----------------------|----------------|--------------|-------------------|---|-----------|------------|
| № of Studies | Study Design | Risk of Bias | Inconsistency | Indirectness | Imprecision | Other Considerations | [intervention] | [comparison] | Relative (95% CI) | Absolute (95% CI) | | |
| Pain Intensity at 3 months (follow-up: median 12 months; Scale from: 0 to 10) | | | | | | | | | | | | |
| 9 | Randomised trials | Not serious | Not serious | Not serious | Not serious | None | 339 | 348 | - | SMD 0.92 SD higher (0.19 higher to 1.65 higher) | ⊕⊕⊕⊕High | |
| Pain Intensity at 6 months (follow-up: mean 12 months; Scale from: 0 to 10) | | | | | | | | | | | | |
| 11 | Randomised trials | Not serious | Not serious | Not serious | Not serious | None | 369 | 378 | - | SMD 1.53 SD higher (0.66 higher to 2.4 higher) | ⊕⊕⊕⊕High | |
| Pain Intensity at 12 Months (follow-up: mean 12 months; Scale from: 0 to 10) | | | | | | | | | | | | |
| 5 | Randomised trials | Not serious | Not serious | Not serious | Not serious | None | 192 | 200 | - | SMD 1.47 SD higher (0.03 higher to 2.97 higher) | ⊕⊕⊕⊕High | |

in these studies also differed, impacting the ability to assess long-term outcomes. Additionally, the inclusion criteria and patient populations were not uniform, which could influence the generalizability of the results. The lack of standardized measures for outcomes and the reporting of adverse events further complicates the analysis, making it challenging to perform a meta-analysis or systematic review that provides definitive conclusions.

Corticosteroids can exert their anti-inflammatory effects to alleviate synovial inflammation and attenuate the excitability of nociceptive nerve fibers, thus providing pain relief. However, potential adverse effects, such as inhibition of the pituitary-adrenal axis, hyperadrenocorticism, osteoporosis, avascular necrosis, and steroid myopathy, pose safety concerns.^{36,37} In contrast, RFA therapy eliminates the need for corticosteroid administration, thereby avoiding these side effects.

It is essential to acknowledge limitations for a comprehensive interpretation of our findings. Small sample sizes observed in most of the included studies may compromise the generalizability and statistical robustness of the results. Larger sample sizes are necessary to improve reliability and accommodate potential variability in patient responses. Additionally, there is substantial heterogeneity among the studies due to variations in patient populations, procedural techniques, and outcome measures. The diversity in procedures and agents used across trials contributes to this heterogeneity, particularly evident within the comparison between intervention groups involving IA and MBB approaches. Moreover, the choice between pulsed and continuous RFA methods may impact outcomes, potentially influencing the depth of analysis. These limitations should be considered when interpreting the study findings, and further research is needed to address these challenges effectively.

Conclusion

This study suggests that RFA may offer superior pain relief and a longer duration of efficacy compared to steroid injections for spinal facet and sacroiliac joint pain. However, the choice between these interventions should be individualized, considering patient preferences, clinical context, and potential risks. Further research, particularly large-scale randomized controlled trials, is needed to strengthen the evidence and refine clinical recommendations.

Funding

Huzhou City Science and Technology Plan Project (2022GY21).

Disclosure

The author(s) report no conflicts of interest in this work.

References

1. Won HS, Yang M, Kim YD. Facet joint injections for management of low back pain: a clinically focused review. *Anesth Pain Med.* 2020;15(1):8–18. doi:10.17085/apm.2020.15.1.8
2. Rashbaum RF, Ohnmeiss DD, Lindley EM, Kitchel SH, Patel VV. Sacroiliac joint pain and its treatment. *Clin Spine Surg.* 2016;29(2):42–48. doi:10.1097/BSD.0000000000000359
3. Kalichman L, Li L, Kim DH, et al. Facet joint osteoarthritis and low back pain in the community-based population. *Spine.* 2008;33(23):2560–2565. doi:10.1097/BRS.0b013e318184ef95
4. Perolat R, Kastler A, Nicot B, et al. Facet joint syndrome: from diagnosis to interventional management. *Insights Imag.* 2018;9(5):773–789. doi:10.1007/s13244-018-0638-x
5. Cohen SP, Chen Y, Neufeld NJ. Sacroiliac joint pain: a comprehensive review of epidemiology, diagnosis and treatment. *Expert Rev Neurother.* 2013;13(1):99–116. doi:10.1586/ern.12.148
6. Falowski S, Sayed D, Pope J, et al. A review and algorithm in the diagnosis and treatment of sacroiliac joint pain. *J Pain Res.* 2020;13:3337–3348. doi:10.2147/JPR.S279390
7. Li SJ, Zhang SL, Feng D. A comparison of pulsed radiofrequency and radiofrequency denervation for lumbar facet joint pain. *J Orthop Surg Res.* 2023;18(1):331. doi:10.1186/s13018-023-03814-5
8. Kwak DG, Kwak SG, Lee AY, Chang MC. Outcome of intra-articular lumbar facet joint corticosteroid injection according to the severity of facet joint arthritis. *Exp Ther Med.* 2019;18(5):4132–4136. doi:10.3892/etm.2019.8031
9. Çetin A, Yektaş A. Evaluation of the short- and long-term effectiveness of pulsed radiofrequency and conventional radiofrequency performed for medial branch block in patients with lumbar facet joint pain. *Pain Res Manag.* 2018;2018:7492753. doi:10.1155/2018/7492753
10. Benoist M, Boulu P, Hayem G. Epidural steroid injections in the management of low-back pain with radiculopathy: an update of their efficacy and safety. *Eur Spine J.* 2012;21(2):204–213. doi:10.1007/s00586-011-2007-z

11. Das G, Surange P, Agarwal A, et al. Radiofrequency ablation in chronic pain syndromes: an evidence- and consensus-based Indian Society for the Study of Pain Guidelines, 2022. *Indian J Pain*. 2022;36:S2–3.
12. Leggett LE, Soril LJJ, Lorenzetti DL, et al. Radiofrequency ablation for chronic low back pain: a systematic review of randomized controlled trials. *Pain Res Manag*. 2014;19(5):e146–153. doi:10.1155/2014/834369
13. Aranke M, McCrudy G, Rooney K, et al. Minimally invasive and conservative interventions for the treatment of sacroiliac joint pain: a review of recent literature. *Orthop Rev*. 2022;14(4):34098. doi:10.52965/001c.34098
14. Tseng C, Chen KT, Fong YC, et al. Biportal endoscopic radiofrequency ablation of the sacroiliac joint complex in the treatment of chronic low back pain: a technical note with 1-year follow-up. *Diagnostics*. 2023;13(2):229. doi:10.3390/diagnostics13020229
15. Lowe M, Okunlola O, Raza S, et al. Radiofrequency ablation as an effective long-term treatment for chronic sacroiliac joint pain: a systematic review of randomized controlled trials. *Cureus*. 2022;14(6):e26327. doi:10.7759/cureus.26327
16. Moher D, Shamseer L, Clarke M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev*. 2015;4(1):1. doi:10.1186/2046-4053-4-1
17. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics*. 1994;50(4):1088–1101. doi:10.2307/2533446
18. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ*. 1997;315(7109):629–634. doi:10.1136/bmj.315.7109.629
19. Do KH, Ahn SH, Cho YW, Chang MC. Comparison of intra-articular lumbar facet joint pulsed radiofrequency and intra-articular lumbar facet joint corticosteroid injection for management of lumbar facet joint pain: a randomized controlled trial. *Medicine*. 2017;96(13):e6524. doi:10.1097/MD.0000000000006524
20. Civelek E, Cansever T, Kabatas S, et al. Comparison of effectiveness of facet joint injection and radiofrequency denervation in chronic low back pain. *Turk Neurosurg*. 2012;22(2):200–206. doi:10.5137/1019-5149.JTN.5207-11.1
21. Duger C, Kol I, Kaygusuz K, Gursoy S, Mimaroglu C. Effects of facet joint nerve block addition to radiofrequency in the treatment of low back pain. *HealthMED*. 2012;6:2052–2056.
22. Dutta K, Dey S, Bhattacharyya P, Agarwal S, Dev P. Comparison of efficacy of lateral branch pulsed radiofrequency denervation and intraarticular depot methylprednisolone injection for sacroiliac joint pain. *Pain Physician*. 2018;21(5):489–496.
23. Hashemi M, Hashemian M, Mohajerani SA, Sharifi G. Effect of pulsed radiofrequency in treatment of facet-joint origin back pain in patients with degenerative spondylolisthesis. *Eur Spine J*. 2014;23(9):1927–1932. doi:10.1007/s00586-014-3412-x
24. Lakemeier S, Lind M, Schultz W, et al. A comparison of intraarticular lumbar facet joint steroid injections and lumbar facet joint radiofrequency denervation in the treatment of low back pain: a randomized, controlled, double-blind trial. *Anesth Analg*. 2013;117(1):228–235. doi:10.1213/ANE.0b013e3182910c4d
25. Lim JW, Cho YW, Lee DG, Chang MC. Comparison of intraarticular pulsed radiofrequency and intraarticular corticosteroid injection for management of cervical facet joint pain. *Pain Physician*. 2017;20(6):E961–E967.
26. Martinez CH, Richardson CR, Han MK, Cigolle CT. Chronic obstructive pulmonary disease, cognitive impairment, and development of disability: the health and retirement study. *Ann Am Thorac Soc*. 2014;11(9):1362–1370. doi:10.1513/AnnalsATS.201405-187OC
27. McCormick ZL, Conger A, Kendall R, et al. A pragmatic randomized prospective trial of cooled radiofrequency ablation of the medial branch nerves versus facet joint injection of corticosteroid for the treatment of lumbar facet syndrome: 12 month outcomes. *Pain Med*. 2023;24(12):1318–1331. doi:10.1093/pm/pnad107
28. Salman O, Sayed Gad G, Mohamed A, Rafae H, Abdelfatah A. Randomized, controlled blind study comparing sacroiliac intra-articular steroid injection to radiofrequency denervation for sacroiliac joint pain. *Egypt J Anaesth*. 2015;32. doi:10.1016/j.egja.2015.07.005
29. Shin SM, Kwak SG, Lee DG, Chang MC. Clinical effectiveness of intra-articular pulsed radiofrequency compared to intra-articular corticosteroid injection for management of atlanto-occipital joint pain: a prospective randomized controlled pilot study. *Spine*. 2018;43(11):741–746. doi:10.1097/BRS.0000000000002414
30. Yasar D, Korgun O, Emine D. Radiofrequency and methylprednisolone in treatment of lower back pain caused by facet joint syndrome: comparison of the outcomes. *Asian J Neurosurg*. 2018;13(2):283–287. doi:10.4103/1793-5482.228569
31. Zhou Q, Zhou F, Wang L, Liu K. An investigation on the effect of improved X-rays-guided radiofrequency thermocoagulation denervation on lumbar facet joint syndrome. *Clin Neurol Neurosurg*. 2016;148:115–120. doi:10.1016/j.clineuro.2016.07.018
32. Cohen SP, Moon JY, Brummett CM, White RL, Larkin TM. Medial branch blocks or intra-articular injections as a prognostic tool before lumbar facet radiofrequency denervation: a multicenter, case-control study. *Reg Anesth Pain Med*. 2015;40(4):376–383. doi:10.1097/AAP.000000000000229
33. Shealy CN. Percutaneous radiofrequency denervation of spinal facets. Treatment for chronic back pain and sciatica. *J Neurosurg*. 1975;43(4):448–451. doi:10.3171/jns.1975.43.4.0448
34. Byrd D, Mackey S. Pulsed radiofrequency for chronic pain. *Curr Pain Headache Rep*. 2008;12(1):37–41. doi:10.1007/s11916-008-0008-3
35. Cohen SP, Bhaskar A, Bhatia A, et al. Consensus practice guidelines on interventions for lumbar facet joint pain from a multispecialty, international working group. *Reg Anesth Pain Med*. 2020;45(6):424–467. doi:10.1136/rapm-2019-101243
36. Brinks A, Koes BW, Volkers ACW, Verhaar JAN, Bierma-Zeinstra SMA. Adverse effects of extra-articular corticosteroid injections: a systematic review. *BMC Musculoskelet Disord*. 2010;11(1):206. doi:10.1186/1471-2474-11-206
37. Kamel SI, Rosas HG, Gorbachova T. Local and systemic side effects of corticosteroid injections for musculoskeletal indications. *AJR Am J Roentgenol*. 2023;222(3). doi:10.2214/AJR.23.30458

Journal of Pain Research

Dovepress

Publish your work in this journal

The Journal of Pain Research is an international, peer reviewed, open access, online journal that welcomes laboratory and clinical findings in the fields of pain research and the prevention and management of pain. Original research, reviews, symposium reports, hypothesis formation and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-pain-research-journal>