

# A Review of the Risk Factors Associated with Endometrial Hyperplasia During Perimenopause

Lianping Wang<sup>1</sup>, Wengong Wei<sup>2</sup>, Meiling Cai<sup>2</sup>

<sup>1</sup>Nursing Department, Qingpu Branch, Zhongshan Hospital, Fudan University, Shanghai, 201700, People's Republic of China; <sup>2</sup>Department of Obstetrics and Gynecology, Qingpu Branch, Zhongshan Hospital, Fudan University, Shanghai, 201700, People's Republic of China

Correspondence: Meiling Cai, Department of Obstetrics and Gynecology, Qingpu Branch, Zhongshan Hospital, Fudan University, Shanghai, 201700, People's Republic of China, Tel +8618116016012, Email cai.meiling@qphospital.co

**Background:** Endometrial hyperplasia, characterized by excessive growth leading to endometrial thickening, is commonly observed in the premenopausal period. Its prevalence in postmenopausal women is approximately 15%, peaking between ages 50 and 60. This condition often manifests as abnormal uterine bleeding and can progress to malignancy, with varying risks depending on the type of hyperplasia.

**Purpose:** This study aims to investigate the factors influencing endometrial thickness during the perimenopausal period and raise awareness among healthcare professionals about the importance of evaluating and caring for individuals with endometrial hyperplasia.

**Methods:** Studies examining the association between various factors such as diabetes mellitus, hypertension, age, estrogen replacement therapy, anovulatory disorders, smoking, medications, genetic factors, and endocrine-related proteins and the development of endometrial hyperplasia were reviewed. The literature search encompassed relevant databases, including PubMed, Scopus, and Web of Science.

**Results:** Research findings indicate significant associations between changes in gene expression of several factors and the development of endometrial hyperplasia. Notably, the risk of progression to cancer varies between non-atypical and atypical hyperplasia cases. Factors such as diabetes mellitus, hypertension, age, estrogen replacement therapy, anovulatory disorders, smoking, medications, Lynch syndrome, tamoxifen use, and alterations in gene expression of TNF- $\alpha$ , EGF, IGF-1, IGF-1R, and PTEN have been implicated in the pathogenesis of endometrial hyperplasia.

**Conclusion:** This study underscores the importance of understanding the factors influencing endometrial thickness during the perimenopausal period. It emphasizes the pivotal role of healthcare professionals in evaluating and caring for individuals with this condition.

**Keywords:** endometrial hyperplasia, premenopausal period, postmenopausal women, pathogenesis, malignancy risk

## Introduction

The endometrium is an epithelial layer located inside the uterus and is a dynamic tissue undergoing various changes in a woman of reproductive age.<sup>1,2</sup> These changes include proliferation, secretion, and menstruation. There exists a complex interaction between the female sex hormones estradiol and progesterone. While estradiol stimulates the thickening of the uterus, progesterone promotes differentiation and secretion of epithelial cells.<sup>3-5</sup> The delicate balance between endometrial proliferation and apoptosis is influenced by hormonal and molecular factors as well as environmental factors such as age. Therefore, the endometrial tissue is predisposed to various abnormalities.<sup>6</sup>

Perimenopause generally refers to the transition period from premenopause to menopause in women.<sup>7-9</sup> The average duration of this period varies between 4 and 11 years, starting from the last menstrual cycle.<sup>10</sup> While most women show menopausal symptoms in their 40s, some exhibit symptoms in their 50s, and only 10% show symptoms in their 30s.<sup>7,11,12</sup>

Endometrial hyperplasia (EH) is a condition commonly seen in the menopausal period, characterized by thickening of the endometrial tissue.<sup>13</sup> EH can be diagnosed in women with postmenopausal bleeding. An endometrial thickness of more than 5 mm in women with postmenopausal bleeding is an undesirable condition, and asymptomatic postmenopausal

women with increased vascularity or endometrial thickness exceeding 11 mm should be evaluated for malignancy risk.<sup>14,15</sup> EH is observed in 15% of postmenopausal women, with its peak occurrence between the ages of 50 and 60. EH develops in 1.3% of women of reproductive age.<sup>13,16</sup>

Common symptoms of EH include menorrhagia, intermenstrual bleeding, postmenopausal bleeding, and abnormal uterine bleeding during hormone replacement therapy or tamoxifen use.<sup>13</sup> Some types of EH may be associated with abnormal epithelial cell proliferation and a tendency to spread to stroma and surrounding tissues, increasing the risk of malignancy.<sup>17–20</sup> Studies have investigated the risk factors associated with EH, but further research is needed.<sup>13,16,21–26</sup> Evaluation and risk assessment of women in the perimenopausal period are also critical.

Early detection of endometrial hyperplasia can reduce mortality from endometrial carcinoma, decrease oncology treatment costs, and contribute to the effective utilization of resources. This review aims to identify risk factors that may contribute to endometrial hyperplasia and emphasize the importance of early diagnosis. Additionally, increasing awareness among healthcare professionals working in women's health regarding endometrial hyperplasia and its risk factors is crucial.

## Risk Factors for Endometrial Hyperplasia (EH) During the Perimenopausal Period

### Regulation of Endometrial Function by Gonadal Hormones and Implications for EH

Gonadal hormones are secreted in a regulated manner throughout the menstrual cycle, regulating the functions of the endometrium. Endometrial thickness is minimal at the end of menstruation, increases during the follicular phase due to estradiol secretion from the dominant follicle, and reaches maximum thickness in the mid-luteal phase due to increased glandular secretion influenced by progesterone.<sup>27</sup> Estrogens play a role in regulating events such as endometrial cell proliferation, angiogenesis, and inflammation.<sup>28,29</sup> In addition to promoting endometrial cell proliferation, estrogen hormones induce morphometric changes, including alterations in glandular epithelial type, gland numbers and shapes, gland-to-stroma ratio, and epithelial cell morphology.<sup>13,30</sup>

EH results from changes in the shape and size of glandular tissue and an increase in the endometrial gland-to-stroma ratio, representing a pre-cancerous, non-physiological, and non-invasive proliferation of the endometrium.<sup>17–19</sup> According to the World Health Organization's 2014 classification, EH is divided into two groups: non-atypical (endometrial intraepithelial neoplasia/EIN) and atypical endometrial hyperplasia.<sup>31</sup> Non-atypical hyperplasia are benign changes that regress when hormonal changes or endocrine disorders return to normal. However, if endocrine disorders persist long-term, invasive disorders may develop. Atypical endometrial hyperplasia contain many of the mutations typical of invasive endometrioid endometrial cancer (EC).<sup>31</sup> EH is clinically characterized by abnormal uterine bleeding, especially in women.<sup>13</sup> Pathologically, some types of endometrial hyperplasia can be precursors to malignancy.<sup>32</sup> It is noted that the risk of untreated EH progressing to cancer can range from 3% in cases of non-atypical EH to 29% in cases of atypical EH.<sup>33</sup>

AEH is recognized as a precancerous lesion with a significantly higher risk of progression to endometrial carcinoma (EC) compared to non-atypical EH. AEH is associated with a 29% risk of progression to EC, while non-atypical EH carries a much lower risk, approximately 1–3%. This distinction is crucial in clinical decision-making, as AEH often warrants more aggressive treatment, including surgical intervention, whereas non-atypical EH may be managed conservatively.<sup>34</sup>

The International Federation of Gynecology and Obstetrics (FIGO) has recently updated its classification criteria for EC, which now places greater emphasis on the histopathological characteristics of AEH. The new classification recognizes AEH as a distinct entity with a clear progression pathway to EC, necessitating more vigilant monitoring and early therapeutic intervention. This change in classification further reinforces the need to accurately diagnose and appropriately manage AEH to prevent the progression to invasive carcinoma.<sup>35,36</sup>

Chronic stimulation of the endometrial tissue by estrogen due to progesterone deficiency plays a significant role in the development of hyperplasia.<sup>16</sup> EH is considered a precursor to type 1 endometrial carcinoma (EC) and is generally accepted as a precursor to low-grade tumors typically associated with estrogen.<sup>37</sup> Risk factors for EH are similar to those

for type 1 endometrial cancer. Conditions associated with hormonal imbalance, such as polycystic ovary syndrome (PCOS), which causes chronic anovulation, early menarche, late menopause, nulliparity, infertility, and prolonged estrogen use/exposure without progesterone, increase the risk of EH.<sup>16,38</sup> Parity is protective against EC but not against EH.<sup>39</sup> Reproductive factors may differ for EH and EC.<sup>40</sup> EH may occur postmenopausally due to cessation of ovulation and lack of progesterone production in the ovaries, and it may also occur during the perimenopausal period with irregular menstruation.<sup>13</sup>

Furthermore, it has been suggested that the increased development of dominant follicles during the luteal phase, associated with decreased levels of progesterone and inhibin A about advancing age, may contribute to increased endometrial thickness.<sup>21</sup> In the postmenopausal period, estrone (E1), a form of estrogen derived from androgens in adrenal and adipose tissues, replaces estradiol (E2), an ovarian-derived estrogen.<sup>41</sup> Adipocytes in adipose tissue convert adrenal-derived androstenedione to E1 via the action of the aromatase enzyme. Factors such as estrogen replacement therapy, diabetes, history of tamoxifen, and oral contraceptive use are also indicated to be associated with endometrial thickening and endometrial malignancy.<sup>22</sup>

## High-Risk Populations for the Development of EH

Two particularly high-risk patient populations for the development of EH have been identified. The first comprises peri- or postmenopausal women who are obese due to excessive adiposity in the abdominal region. This leads to a significant conversion of androgens in adipose tissue to estrogen by the action of the aromatase enzyme.<sup>42</sup> It is known that insulin-like growth factor 1 (IGF-1) and its binding protein (IGF binding protein-1) support endometrial cell growth. Elevated levels of IGF have been reported in obese women, which may contribute to EH development by predisposing them to endometrial cancer.<sup>43</sup> Additionally, in obese women with diabetes mellitus (DM), high insulin resistance has been observed, leading to a decrease in sex hormone-binding globulin (SHBG) concentration due to increased insulin levels, resulting in elevated estrogen levels.<sup>44–46</sup>

The second high-risk group for EH development consists of premenopausal women with polycystic ovary syndrome (PCOS) characterized by hyperandrogenic activity. While endometrial stimulation by estrogens is considered a primary risk factor for EH development, other factors, such as immunosuppression, may also play a role.<sup>42,47</sup> The chronic exposure of the endometrium to high levels of estrogen without the counterbalancing effect of progesterone leads to increased cell proliferation and a heightened risk of malignant transformation.<sup>48</sup>

Most cases of EH occur in the presence of chronic exposure to estrogen, which is not counteracted by progesterone, as seen in conditions like PCOS and obesity.<sup>49</sup> PCOS is a common endocrinopathy associated with insulin resistance, hyperandrogenism, negative cardiovascular risk factors, and infertility-related issues.<sup>50–52</sup> Hyperandrogenism in PCOS can lead to acne, hirsutism, and alopecia. Moreover, due to the conversion of androgens to estrogens and excess estrogen synthesis, ovarian dysfunction, chronic oligomenorrhea, infertility, endometrial hyperplasia, and endometrial cancer can occur.<sup>53–55</sup>

Studies have shown that the risk of EH increases up to threefold in obesity and is a common condition in PCOS, where EH frequency ranges from 35.7% to 48.8%. Furthermore, obesity, diabetes mellitus, and hypertension are associated with EH.<sup>56,57</sup> Obesity-induced chronic inflammation, resulting in increased estrogen levels, is suggested to contribute to EH and malignancy.<sup>13,56,58–64</sup> Additionally, factors such as high BMI and hypertensive and anti-inflammatory drug use, as well as hormone replacement therapy, are associated with EH development.<sup>65,66</sup> Studies differentiate between the effects of steroidal anti-inflammatory drugs, like corticosteroids, which may exert a broader influence on metabolic processes and potentially alter the risk of endometrial hyperplasia (EH), compared to non-steroidal anti-inflammatory drugs (NSAIDs), which primarily target inflammation pathways.<sup>67,68</sup> Additionally, the literature has discussed the impacts of various classes of antihypertensive drugs—such as ACE inhibitors, beta-blockers, and diuretics—on EH risk, providing relevant insights into how these medications may influence the development or management of EH.<sup>67</sup> Research indicates that hypertensive patients often have a higher frequency of endometrial thickening and hyperplasia. A study showed that hyperplasia was the most common positive histological finding among hypertensive women, further supporting the association between hypertension, EH, and the pharmacological agents used to manage these conditions.<sup>67</sup>

However, no significant differences in EH development were found between smokers and non-smokers, individuals with hypertension and normotensives, or individuals with and without diabetes mellitus. While the results regarding age were insufficient, a significant relationship was established between BMI and endometrial hyperplasia, indicating that a high BMI may increase the risk of EH.<sup>69–71</sup>

In postmenopausal women, a study examining factors affecting endometrial thickness found no relationship between diseases such as DM and hypertension and endometrial thickness, but a relationship was observed between BMI, ovarian volume, uterine volume, tamoxifen use, serum E2 levels, and endometrial thickness.<sup>72</sup> Another study found no significant relationship between DM and high BMI and increased endometrial thickness, but a significantly higher endometrial thickness was observed in patients with hypertension and non-smokers. A weak significant relationship was found between hormone replacement therapy and increased endometrial thickness. This study emphasized the importance of BMI as a significant parameter affecting endometrial thickness in postmenopausal women.<sup>73</sup>

In a study comparing the effects of age, endometrial thickness, and BMI on malignancy development in postmenopausal women with postmenopausal bleeding, a significant relationship was found between endometrial thickness and age and malignancy development, while no significant relationship was found between BMI and malignancy. These findings may vary depending on changing diagnostic criteria and the nature of PCOS. The threefold increased risk of EC in PCOS and the connection between EH and EC suggest a relationship between EH and PCOS.<sup>74–77</sup> Recent evidence suggests that asymptomatic women, particularly those with underlying risk factors such as obesity, may still harbor a significant risk of developing EC. The recent findings underscore the need for updated screening guidelines and risk stratification models to identify high-risk individuals who might benefit from early intervention.<sup>78,79</sup>

In hypertensive patients, endometrial hyperplasia (EH) is frequently encountered, and the use of hypertensive medications has been associated with EH development.<sup>21,80–82</sup> It is speculated that diabetes mellitus (DM) may not only cause EH but also predispose existing hyperplasia to malignant transformation. A meta-analysis has shown the likelihood of malignancy in women with DM who have EH.<sup>83</sup> In another study examining the relationship between gestational diabetes mellitus (GDM) and EH and endometrial carcinoma (EC), it was found that GDM is associated with both EH development and EC. However, the relationship between GDM and EC was reported to be stronger than that between GDM and EH.<sup>84</sup> The use of tamoxifen, a selective estrogen receptor modulator (SERM) used in cancer treatment, poses a risk for EH, endometrial polyps, abnormal vaginal bleeding, and EC development due to its estrogenic effect on the endometrium.<sup>85</sup> The studies indicate that women undergoing tamoxifen therapy have an increased risk of developing EH and EC. The mechanism is believed to be related to the agonistic effects of tamoxifen on the endometrial tissue, which may lead to hyperplasia and subsequent malignancy. This risk highlights the necessity of regular endometrial surveillance in patients receiving tamoxifen or similar agents.<sup>86,87</sup> Women with Lynch syndrome may develop atypical EH at an early age, as there are changes in estrogen levels affecting the expression of DNA repair genes in these women.<sup>25</sup> Additionally, peripheral estrogen synthesis may increase due to adrenal cortical tumors that secrete androgens, leading to EH.<sup>13</sup>

Apart from estrogen stimulation of the endometrium, EH can also occur due to other reasons, such as immunosuppression and infection.<sup>88</sup> Endometrial inflammation leads to an imbalance in the cytokine system, developing EH. Inflammation causes a decrease in factors such as tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) and epidermal growth factor (EGF), while insulin-like growth factor-1 receptor (IGF-1R) levels increase.<sup>89–91</sup> It has been stated that the expression of IGF-1R is much higher in EH and malignancy cases compared to normal conditions.<sup>92,93</sup> Only in adenomatous hyperplasia has a decrease in the expression of the IGF-1 gene been noted.<sup>89</sup> The synthesis of IGF-1 is induced by estradiol and affects uterine growth.<sup>94</sup> Another factor in EH development is phosphatase and tensin homolog (PTEN). PTEN is a tumor suppressor gene located on chromosome 10q23, encoding a phosphatase with protein and lipid effects. As an antagonist in cell signaling pathways induced by growth factors, PTEN regulates cell proliferation and apoptosis.<sup>95,96</sup> While PTEN expression increases in both glandular epithelium and stromal compartments during the proliferative phase, it decreases in the glandular epithelium compartment during the secretory phase.<sup>97</sup> A mutation causing loss of function in the PTEN gene may lead to upregulation of endometrial glandular proliferation and is associated with EH and cancer development.<sup>26,98</sup> Among cases, isolated PTEN-null glands have been shown in 43% of

macroscopically standard premenopausal endometrial samples. Due to genetic mutation and deletion, these glands cannot express the PTEN protein and are mainly found between menstrual cycles.<sup>99</sup>

Endometrial hyperplasia is essential not only due to the damaging physical and psychological symptoms it causes but also because it can be a precursor to cancer. The risk of endometrial hyperplasia progressing to cancer can range from 3% in cases of non-atypical EH to 29% in cases of atypical EH.<sup>33</sup> At this point, evaluating risk factors that may lead to the development of endometrial hyperplasia and intervening early can be lifesaving.<sup>100–103</sup>

## Conclusion

A myriad of risk factors influence endometrial hyperplasia. Research has extensively explored various contributors, including diabetes mellitus (DM), hypertension (HT), age, estrogen replacement therapy, anovulatory disorders such as polycystic ovary syndrome (PCOS), smoking, medical treatments, Lynch syndrome, and tamoxifen use. Among these, obesity has emerged as a particularly significant factor in the development of endometrial hyperplasia. Additionally, studies have examined the relationship between genetic expression changes—specifically in genes such as TNF- $\alpha$ , EGF, IGF-1, IGF-1R, and PTEN—and the onset of endometrial hyperplasia. Notably, IGF-1 and PTEN are thought to play pivotal roles. However, despite extensive research on these risk factors, further studies are necessary to achieve more comprehensive and conclusive results.

Given that certain types of endometrial hyperplasia can cause significant physical and psychological distress and serve as precursors to cancer, early intervention is essential for long-term health. With the rising incidence of endometrial hyperplasia during the postmenopausal period, it is critical to conduct thorough evaluations of women during this time. Additionally, comprehensive assessments and risk evaluations should be carried out during the premenopausal period. Early detection and intervention in cases of malignancies associated with endometrial hyperplasia can alleviate the physical and economic burden on patients, enhance women's health, and reduce mortality rates related to endometrial cancer.

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. Jabbour HN, Kelly RW, Fraser HM, et al. Endocrine regulation of menstruation. *Endocrine Reviews*. 2006;27(1):17–46. doi:10.1210/er.2004-0021
2. Naz MSG, Dovom MR, Tehrani FR. The menstrual disturbances in endocrine disorders: a narrative review. *J Clin Endocrinol Metab*. 2020;18(4):1.
3. Medh RD, Thompson EB. Hormonal regulation of physiological cell turnover and apoptosis. *Cell Tissue Res*. 2000;301:101–124. doi:10.1007/s004419900159
4. Islam MS, Afrin S, Jones SI, et al. Selective progesterone receptor modulators—mechanisms and therapeutic utility. *Endocrin Rev*. 2020;41(5):bnaa012. doi:10.1210/endrev/bnaa012
5. Chodankar RR, Murray A, Nicol M, et al. The endometrial response to modulation of ligand-progesterone receptor pathways is reversible. *Fertil Steril*. 2021;116(3):882–895. doi:10.1016/j.fertnstert.2021.02.008
6. Cuilan G, Xiaomei Y, Shuang X, XiLing Y, Yan Z. Obesity, endogenous hormones, and endometrial cancer risk. *Ethiop J Health Dev*. 2021;35(3):1.
7. Woods NF, Mitchell ES, Coslov N, et al. Transitioning to the menopausal transition: a scoping review of research on the late reproductive stage in reproductive aging. *Menopause*. 2021;28(4):447–466. doi:10.1097/GME.0000000000001707
8. Bhalerao S, Vora P. Stages of reproductive aging and correlation with symptoms. *ICOG Campus*. 2023;1:78.
9. Bhattacharjee NV, Schumacher A, Aali A, et al. Global fertility in 204 countries and territories, 1950–2021, with forecasts to 2100: a comprehensive demographic analysis for the global burden of disease study 2021. *Lancet*. 2024;1:1.
10. Harlow SD, Paramsothy P. Menstruation and the menopausal transition. *Obstet Gynecol Clin*. 2011;38(3):595–607. doi:10.1016/j.ogc.2011.05.010
11. Chon SJ, Umair Z, Yoon M-S. Premature ovarian insufficiency: past, present, and future. *Front Cell Develop Biol*. 2021;9:672890. doi:10.3389/fcell.2021.672890
12. Nikolaou D, Templeton A. Early ovarian ageing: a hypothesis: detection and clinical relevance. *Hum Reprod*. 2003;18(6):1137–1139. doi:10.1093/humrep/deg245
13. Ring KL, Mills AM, Modesitt SC. Endometrial hyperplasia. *Obstetrics Gynecol*. 2022;140(6):1061–1075. doi:10.1097/AOG.0000000000004989
14. Aggarwal A, Hatti A, Tirumuru SS, et al. Management of asymptomatic postmenopausal women referred to outpatient hysteroscopy service with incidental finding of thickened endometrium—a UK district general hospital experience. *J Minim Invasive Gynecol*. 2021;28(10):1725–1729. doi:10.1016/j.jmig.2021.02.012
15. Giri S, Nayak B, Mohapatra J. Thickened endometrium: when to intervene? A clinical conundrum. *J Obstet Gynaecol India*. 2021;71(3):216–225. doi:10.1007/s13224-020-01415-4
16. Mencaglia L, Ciociola F, Magnolfi S. Endometrial hyperplasia. *Hysteroscopy*. 2018;1:261–267.



17. Prip CM, Stentebjerg M, Bennetsen MH, et al. Risk of atypical hyperplasia and endometrial carcinoma after initial diagnosis of non-atypical endometrial hyperplasia: a long-term follow-up study. *PLoS One*. 2022;17(4):e0266339. doi:10.1371/journal.pone.0266339
18. Suresh K. Enigma of endometriosis-diagnosis and management dilemma. *J Clin Med Img Case Rep*. 2023;3(6):1599.
19. Daud S, Jalil SSA, Griffin M, et al. Endometrial hyperplasia—the dilemma of management remains: a retrospective observational study of 280 women. *Eur J Obstetrics Gynecol Reprod Biol*. 2011;159(1):172–175. doi:10.1016/j.ejogrb.2011.06.023
20. Raffone A, Travaglino A, Flacco ME, et al. Clinical predictive factors of response to treatment in patients undergoing conservative management of atypical endometrial hyperplasia and early endometrial cancer. *J Adolesc Young Adult Oncol*. 2021;10(2):193–201. doi:10.1089/jayao.2020.0100
21. Baerwald A, Vanden Brink H, Lee C, et al. Endometrial development during the transition to menopause: preliminary associations with follicular dynamics. *Climacteric*. 2020;23(3):288–297. doi:10.1080/13697137.2020.1721455
22. Ray S, Zohorinia S, Bhattacharyya D, et al. Risk factors for endometrial cancer among post-menopausal women in South Africa. *Asian Pac J Cancer Biol*. 2019;4(2):41–45. doi:10.31557/apjcb.2019.4.2.41-45
23. Dreisler E. Endometrial thickness in postmenopausal women. *J Case Rep Womens Health*. 2022;35:e00431.
24. Ai F, Qin X, Zhou L, et al. Analysis of factors related to endometrial cancer in postmenopausal women with endometrial thickening. *Menopause*. 2023;30(9):920–926. doi:10.1097/GME.0000000000002232
25. Lucas E, Chen H, Molberg K, et al. Mismatch repair protein expression in endometrioid intraepithelial neoplasia/atypical hyperplasia: should we screen for Lynch syndrome in precancerous lesions? *Int J Gynecol Pathol*. 2019;38(6):533–542. doi:10.1097/PGP.0000000000000557
26. Wang X, Wendel JRH, Emerson RE, et al. Pten and Dicer1 loss in the mouse uterus causes poorly differentiated endometrial adenocarcinoma. *Oncogene*. 2020;39(40):6286–6299. doi:10.1038/s41388-020-01434-5
27. Baerwald A, Pierson R. Ovarian follicular waves during the menstrual cycle: physiologic insights into novel approaches for ovarian stimulation. *Fertil Steril*. 2020;114(3):443–457. doi:10.1016/j.fertnstert.2020.07.008
28. Gibson DA, Saunders PT. Estrogen dependent signaling in reproductive tissues—a role for estrogen receptors and estrogen related receptors. *Mol Cell Endocrinol*. 2012;348(2):361–372. doi:10.1016/j.mce.2011.09.026
29. Yu K, Huang Z-Y, Xu X-L, et al. Estrogen receptor function: impact on the human endometrium. *Front Endocrinol*. 2022;13:827724. doi:10.3389/fendo.2022.827724
30. Turashvili G, Hanley K. Practical updates and diagnostic challenges in endometrial carcinoma. *Arch Pathol Lab Med*. 2024;148(1):78–98. doi:10.5858/arpa.2022-0280-RA
31. Emons G, Beckmann M, Schmidt D, et al. New WHO classification of endometrial hyperplasias. *Geburtshilfe Frauenheilkd*. 2015;75(02):135–136. doi:10.1055/s-0034-1396256
32. Dugo E, Piva F, Giulietti M, Giannella L, Ciavattini A, et al. Copy number variations in endometrial cancer: from biological significance to clinical utility. *Int J Gynecologic Cancer*. 2024;1:ijgc-2024-005295.
33. Gallos ID, Ganesan R, Gupta JK. Prediction of regression and relapse of endometrial hyperplasia with conservative therapy. *Obstetrics Gynecol*. 2013;121(6):1165–1171. doi:10.1097/AOG.0b013e31828cb563
34. Nees LK, Heublein S, Steinmacher S, et al. Endometrial hyperplasia as a risk factor of endometrial cancer. *Arch Gynecol Obstet*. 2022;306(2):407–421. doi:10.1007/s00404-021-06380-5
35. Berek JS, Matias-Guiu X, Creutzberg C, et al. FIGO staging of endometrial cancer: 2023. *Int J Gynaecol Obstet*. 2023;162(2):383–394. doi:10.1002/ijgo.14923
36. Yu C, Yuan X, Yao Q, et al. Clinical application of FIGO 2023 staging system of endometrial cancer in a Chinese cohort. *BMC Cancer*. 2024;24(1):862. doi:10.1186/s12885-024-12633-8
37. Murali R, Soslow RA, Weigelt B. Classification of endometrial carcinoma: more than two types. *Lancet Oncol*. 2014;15(7):e268–e278. doi:10.1016/S1470-2045(13)70591-6
38. Epplen M, Reed SD, Voigt LF, et al. Risk of complex and atypical endometrial hyperplasia in relation to anthropometric measures and reproductive history. *Am J Epidemiol*. 2008;168(6):563–570. doi:10.1093/aje/kwn168
39. Wu Q-J, Li -Y-Y, Tu C, et al. Parity and endometrial cancer risk: a meta-analysis of epidemiological studies. *Sci Rep*. 2015;5(1):14243. doi:10.1038/srep14243
40. Kahn JL, Buckingham L, Koelper NC, et al. Risk factors for atypical hyperplasia and endometrial cancer in the infertility population: a case-control study. *F&S Reports*. 2021;2(1):104–108. doi:10.1016/j.xfre.2020.10.005
41. Vanden Brink H, Robertson DM, Lim H, et al. Associations between antral ovarian follicle dynamics and hormone production throughout the menstrual cycle as women age. *J Clin Endocrinol Metab*. 2015;100(12):4553–4562. doi:10.1210/jc.2015-2643
42. Bobrowska K, Kamiński P, Cyganek A, et al. High rate of endometrial hyperplasia in renal transplanted women. *Transplant Proc*. 2006;38(1):177–179. doi:10.1016/j.transproceed.2005.12.007
43. Merritt MA, Strickler HD, Hutson AD, et al. Sex hormones, insulin, and insulin-like growth factors in recurrence of high-stage endometrial cancer. *Cancer Epidemiol Biomarkers Prev*. 2021;30(4):719–726. doi:10.1158/1055-9965.EPI-20-1613
44. Aleksioska Papestiev I, Antovska V, Dabeski D, et al. The impact of obesity and fat distribution on endometrial cancer risk in postmenopausal patients. *J Morphol Sci*. 2023;6(2):157–164. doi:10.55302/JMS2362157ap
45. Sağnıç S. Obesity and Endometrial Cancer. *Role Obes Human Health Dis*. 2021;1:53.
46. Shenbagadevi S. *Effect of Body Mass Index on Endometrial Thickness and Serum Estradiol in Asymptomatic Post Menopausal Women: A Cross Sectional Study*. Chennai: Madras Medical College; 2020.
47. Al-Kaif LA, Al-Ameri H, Alfatlawi WRO, et al. Detection of CTLA-4 level and humeral immune response after the second dose of COVID-19 vaccine in certain Iraqi provinces participants. *PLoS One*. 2024;19(1):e0296521. doi:10.1371/journal.pone.0296521
48. Zhao H, Yang H, Li J, et al. Association of urinary levels of estrogens and estrogen metabolites with the occurrence and development of endometrial hyperplasia among premenopausal women. *Reprod Sci*. 2023;30(10):3027–3036. doi:10.1007/s43032-023-01229-5
49. Charalampakis V, Tahrani AA, Helmy A, et al. Polycystic ovary syndrome and endometrial hyperplasia: an overview of the role of bariatric surgery in female fertility. *Eur J Obstet Gynecol Reprod Biol*. 2016;207:220–226. doi:10.1016/j.ejogrb.2016.10.001
50. Conway G, Dewailly D, Diamanti-Kandarakis E, et al. The polycystic ovary syndrome: a position statement from the European society of endocrinology. *Eur J Endocrinol*. 2014;171(4):P1–P29. doi:10.1530/EJE-14-0253

51. Akkaif MA, Sha'aban A, Cesaro A, et al. The impact of SARS-CoV-2 treatment on the cardiovascular system: an updated review. *Inflammopharmacology*. 2022;30(4):1143–1151. doi:10.1007/s10787-022-01009-8
52. Akkaif MA, Bitar AN, Al-Kaif LAIK, et al. The management of myocardial injury related to SARS-CoV-2 pneumonia. *J Cardiovasc Dev Dis*. 2022;9(9):307. doi:10.3390/jcdd9090307
53. Chang S, Dunaif A. Diagnosis of polycystic ovary syndrome: which criteria to use and when? *Endocrinol Metab Clin*. 2021;50(1):11–23. doi:10.1016/j.ecl.2020.10.002
54. Carmina E, Dreno B, Lucky WA, et al. Female adult acne and androgen excess: a report from the multidisciplinary androgen excess and PCOS committee. *J Endocrin Soc*. 2022;6(3):bvac003. doi:10.1210/jendso/bvac003
55. Joham AE, Norman RJ, Stener-Victorin E, et al. Polycystic ovary syndrome. *Lancet Diabetes Endocrinol*. 2022;10(9):668–680. doi:10.1016/S2213-8587(22)00163-2
56. Holm NSL, Glinborg D, Andersen MS, Schledermann D, Ravn P. The prevalence of endometrial hyperplasia and endometrial cancer in women with polycystic ovary syndrome or hyperandrogenism. *Acta obstetrica et gynecologica Scandinavica*. 2012;91(10):1173–1176. doi:10.1111/j.1600-0412.2012.01458.x
57. Park B, Lee H, Park S, et al. Trend changes and factor analysis of endometrial hyperplasia in patients with polycystic ovarian syndrome based on the Korean national health insurance database. *BMC Women's Health*. 2022;22(1):439. doi:10.1186/s12905-022-02015-2
58. Shafiee MN, Khan G, Ariffin R, et al. Preventing endometrial cancer risk in polycystic ovarian syndrome (PCOS) women: could metformin help? *Gynecol Oncol*. 2014;132(1):248–253. doi:10.1016/j.ygyno.2013.10.028
59. Modesitt SC, Hallowell PT, Slack-Davis JK, et al. Women at extreme risk for obesity-related carcinogenesis: baseline endometrial pathology and impact of bariatric surgery on weight, metabolic profiles and quality of life. *Gynecol Oncol*. 2015;138(2):238–245. doi:10.1016/j.ygyno.2015.05.015
60. Armstrong AJ, Hurd WW, Elguero S, Barker NM, Zanotti KM. Diagnosis and management of endometrial hyperplasia. *J Minim Invasive Gynecol*. 2012;19(5):562–571. doi:10.1016/j.jmig.2012.05.009
61. Gupta JK, Rimmer M. Diagnosis and management of endometrial hyperplasia: a UK national audit of adherence to national guidance 2012–20. *PLoS Med*. 2024;21(2):1–18.
62. Tingthanatikul Y, Choktanasiri W, Rochanawutanon M, et al. Prevalence and clinical predictors of endometrial hyperplasia in anovulatory women presenting with amenorrhea. *Gynecol Endocrinol*. 2006;22(2):101–105. doi:10.1080/09513590600585997
63. Akkaif MA, Daud NA, Noor DA, Sha'aban A, Kader MA, Ibrahim B. The impact of CYP2C19 genotype on the platelet reactivity index (PRI) among chronic coronary syndromes (CCS) patients undergoing percutaneous coronary intervention (PCI): affectability of rapid genetic testing. *Cardiovasc Drugs Ther*. 2024;38:1–10. doi:10.1007/s10557-024-07543-7
64. Cai J, Sun X, Li M, et al. Dietary factors in relation to hypertension: a mendelian randomization study. *J Health Popul Nutr*. 2024;43(1):1–10. doi:10.1186/s41043-024-00575-7
65. Wise MR, Jordan V, Lagas A, et al. Obesity and endometrial hyperplasia and cancer in premenopausal women: a systematic review. *Am J Clin Exp Obstet Gynecol*. 2016;214(6):689. e1–689. e17. doi:10.1016/j.ajog.2016.01.175
66. Ibrahim NA, Abdullah SA, Rashid KN. Relation assessment between some factors and diseases with endometrial hyperplasia in Salahuddin province. *Med J Tikrit*. 2018;24(2): 34–43.
67. Vorgias G, Strigou S, Varhalama E, et al. The effect of hypertension and anti-hypertensive drugs on endometrial thickness and pathology. *Eur J Obstet Gynecol Reprod Biol*. 2006;125(2):239–242. doi:10.1016/j.ejogrb.2005.07.031
68. Indraccolo U, Matteo M, Greco P, et al. ACE inhibitors and angiotensin receptor blockers could promote the onset of endometrial polyps in hypertensive women. *Int J Clin Pharmacol Ther*. 2013;51(12):963–968. doi:10.5414/CP201962
69. Sanni OB, Kunzmann AT, Murray LJ, McCluggage WG, Coleman HG. Risk factors (excluding hormone replacement therapy) for endometrial hyperplasia: a systematic review. *Epidemiology*. 2016;06. doi:10.4172/2161-1165.1000229
70. Nees LK, Heublein S, Steinmacher S, et al. Endometrial hyperplasia as a risk factor of endometrial cancer. *Arch Gynecol Obstet*. 2022;1:1–15.
71. Petersdorf K, Groettrup-Wolfers E, Overton PM, et al. Endometrial hyperplasia in pre-menopausal women: a systematic review of incidence, prevalence, and risk factors. *Eur J Obstet Gynecol Reprod Biol*. 2022;271:158–171. doi:10.1016/j.ejogrb.2022.02.015
72. Hebbat S, Chaya V, Rai L, et al. Factors influencing endometrial thickness in postmenopausal women. *Ann Med Health Sci Res*. 2014;4(4):608–614. doi:10.4103/2141-9248.139340
73. Yurtsever ÇY, Aydın DS, Gültaş A, Ateşer G, Bacanakgil BH, Boran AB. Asemptomatik postmenopozal hastalarda vücut kitle endeksinin endometrial kalınlığa etkisi. *Istn Med J*. 2015;16(1):14–17.
74. Saccardi C, Vitagliano A, Marchetti M, et al. Endometrial cancer risk prediction according to indication of diagnostic hysteroscopy in post-menopausal women. *Diagnostics*. 2020;10(5). doi:10.3390/diagnostics10050257
75. Jha S, Singh A, Sinha HH, et al. Rate of premalignant and malignant endometrial lesion in “low-risk” premenopausal women with abnormal uterine bleeding undergoing endometrial biopsy. *Obstet Gynecol Sci*. 2021;64(6):517. doi:10.5468/ogs.21150
76. Akalyaa K. *Correlation of Body Mass Index and Abnormal Uterine Bleeding in Premenopausal Women*. Rajiv Gandhi University of Health Sciences (India); 2020.
77. Deurenberg-Yap M, Schmidt G, van Staveren WA, et al. The paradox of low body mass index and high body fat percentage among Chinese, Malays and Indians in Singapore. *Int J Obesity*. 2000;24(8):1011–1017. doi:10.1038/sj.ijo.0801353
78. Vitale SG, Riemma G, Haimovich S, et al. Risk of endometrial cancer in asymptomatic postmenopausal women in relation to ultrasonographic endometrial thickness: systematic review and diagnostic test accuracy meta-analysis. *Am J Obstet Gynecol*. 2023;228(1):22–35.e2. doi:10.1016/j.ajog.2022.07.043
79. Wolfman W, Bougie O, Chen I, et al. Guideline no. 451: asymptomatic endometrial thickening in postmenopausal women. *J Obstet Gynaecol Can*. 2024;46(7):102591. doi:10.1016/j.jogc.2024.102591
80. Takreem A, Danish N, Razaq S. Incidence of endometrial hyperplasia in 100 cases presenting with polymenorrhagia/menorrhagia in perimenopausal women. *J Ayub Med Coll Abbottabad*. 2009;21(2):60–63.
81. Abd Al-Kadir IT, Amen NS. Evaluation of endometrial thickness in asymptomatic postmenopausal women with hypertension. *Iraqi Med J*. 2020;66(1):30–35.

82. Yao L, Li C, Cheng J. The relationship between endometrial thickening and endometrial lesions in postmenopausal women. *Arch Gynecol Obstet*. 2022;306(6):2047–2054. doi:10.1007/s00404-022-06734-7
83. Raffone A, Travaglini A, Saccone G, et al. Diabetes mellitus is associated with occult cancer in endometrial hyperplasia. *Pathol Oncol Res*. 2020;26:1377–1384. doi:10.1007/s12253-019-00684-3
84. Wartko PD, Beck TL, Reed SD, et al. Association of endometrial hyperplasia and cancer with a history of gestational diabetes. *Cancer Causes Control*. 2017;28:819–828. doi:10.1007/s10552-017-0908-9
85. Kedar RP, Bourne TH, Powles TJ, et al. Effects of tamoxifen on uterus and ovaries of postmenopausal women in a randomised breast cancer prevention trial. *Obstetrical Gynecol Surv*. 1995;50(2):110–112. doi:10.1097/00006254-199502000-00016
86. Vitale SG, Angioni S, D'Alterio MN, et al. Risk of endometrial malignancy in women treated for breast cancer: the BLUSH prediction model - evidence from a comprehensive multicentric retrospective cohort study. *Climacteric*;2024. 1–7. doi:10.1080/13697137.2024.2376189
87. Ryu KJ, Kim MS, Lee JY, et al. Risk of endometrial polyps, hyperplasia, carcinoma, and uterine cancer after tamoxifen treatment in premenopausal women with breast cancer. *JAMA Network Open*. 2022;5(11):e2243951. doi:10.1001/jamanetworkopen.2022.43951
88. Hyperplasia E. Management of (green-top guideline no. 67). *RCOG/BSGE Joint Guideline*. 2016.
89. Sukhikh G, Zhdanov AV, Davydova MP, et al. Disorders in cytokine gene expression in endometrial hyperplasia and effect of hormone therapy. *Bull Exp Biol Med*. 2005;139:235–237. doi:10.1007/s10517-005-0257-7
90. Khan D, Patel R. Role of immune dysregulation in the pathogenesis of endo-metrial hyperplasia. *World J Pharm Res*. 2022;11(7):143–158.
91. Al-Kaif LA, Al-Khafaji YK, Shandaway S, et al. Interleukin-8 and-17 levels in the sera of vaccinated subjects receiving a booster dose of measles virus: a follow-up study in Iraq. *Med J Babylon*. 2023;20(2):422–425. doi:10.4103/MJBL.MJBL\_566\_23
92. Stavropoulos A, Varras M, Philippou A, et al. Immunohistochemical expression of insulin-like growth factor-1Ec in primary endometrial carcinoma: association with PTEN, p53 and survivin expression. *Oncol Lett*. 2020;20(6):1. doi:10.3892/ol.2020.12258
93. McCampbell AS, Broaddus RR, Loose DS, et al. Overexpression of the insulin-like growth factor I receptor and activation of the AKT pathway in hyperplastic endometrium. *Clin Cancer Res*. 2006;12(21):6373–6378. doi:10.1158/1078-0432.CCR-06-0912
94. Bruchim I, Sarfstein R, Werner H. The IGF hormonal network in endometrial cancer: functions, regulation, and targeting approaches. *Front Endocrinol*. 2014;5:91120. doi:10.3389/fendo.2014.00076
95. Latta E, Chapman WB. PTEN mutations and evolving concepts in endometrial neoplasia. *Curr Opin Obstet Gynecol*. 2002;14(1):59–65. doi:10.1097/00001703-200202000-00010
96. Sanderson PA, Esnal-Zufiaurre A, Arends MJ, et al. Improving the diagnosis of endometrial hyperplasia using computerized analysis and immunohistochemical biomarkers. *Front Reprod Health*. 2022;4:896170. doi:10.3389/frph.2022.896170
97. Russo A, Czarnecki AA, Dean M, et al. PTEN loss in the fallopian tube induces hyperplasia and ovarian tumor formation. *Oncogene*. 2018;37(15):1976–1990. doi:10.1038/s41388-017-0097-8
98. Maru Y, Hippo Y. Two-way development of the genetic model for endometrial tumorigenesis in mice: current and future perspectives. *Front Genetics*. 2021;12:798628. doi:10.3389/fgene.2021.798628
99. Mutter GL, Ince TA, Baak JP, et al. Molecular identification of latent precancers in histologically normal endometrium. *Cancer Res*. 2001;61(11):4311–4314.
100. Piergentili R, Gullo G, Basile G, et al. Circulating miRNAs as a tool for early diagnosis of endometrial cancer—implications for the fertility-sparing process: clinical, biological, and legal aspects. *Int J Mol Sci*. 2023;24(14):11356. doi:10.3390/ijms241411356
101. Chelmow D, Brooks R, Cavens A, et al. Executive summary of the uterine cancer evidence review conference. *Obstetrics Gynecol*. 2022;139(4):626–643. doi:10.1097/AOG.0000000000004711
102. Özcan H, Demir Doğan M. Gynecological cancer awareness among women. *Indian J Gynecol Oncol*. 2021;19:1–9. doi:10.1007/s40944-020-00481-w
103. Mitra S, Lami MS, Ghosh A, et al. Hormonal therapy for gynecological cancers: how far has science progressed toward clinical applications? *Cancers*. 2022;14(3):759. doi:10.3390/cancers14030759

## International Journal of Women's Health

Dovepress

## Publish your work in this journal

The International Journal of Women's Health is an international, peer-reviewed open-access journal publishing original research, reports, editorials, reviews and commentaries on all aspects of women's healthcare including gynecology, obstetrics, and breast cancer. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-journal-of-womens-health-journal>