


Impact of Indoor Air Pollutants on the Cardiovascular Health Outcomes of Older Adults: Systematic Review

Nomagugu Ndlovu ^{*}, Benedicta Ngwenchi Nkeh-Chungag^{*}

Department of Biological and Environmental Sciences, Walter Sisulu University, Mthatha, South Africa

^{*}These authors contributed equally to this work

Correspondence: Nomagugu Ndlovu, Email nondlovu@wsu.ac.za

Abstract: Indoor air pollution accounts for approximately 3.8 million inopportune deaths annually at global level. Due to spending more time indoors, children and older adults are especially susceptible to the health risks of indoor air pollution. This review seeks to summarise existing knowledge on the cardiovascular health effects of three common indoor air pollutants, namely carbon monoxide (CO), particulate matter (PM_{2.5} and PM₁₀), and Nitrogen dioxide (NO₂), focusing on older adults. We systematically reviewed the literature (PROSPERO CRD42024479220) on PubMed, Google Scholar, Scopus, Web of Science and Embase. The search yielded 20,914 records. Two independent reviewers screened the articles using titles, abstracts, and full-length articles written in English. Upon a detailed assessment of all the records, the review considered 38 full-length articles. Several studies reported mortality, myocardial infarction, stroke, increased hospitalisation and increased emergency room visits due to exposure to indoor air pollution. A few studies reported arrhythmias, hypertension and Ischaemic heart disease due to exposure to indoor air pollutants. The increased mortality, morbidity, hospitalization, and emergency rooms visits resulting from indoor air pollution associated CVDs makes indoor air pollution a health risk for older adults. There is, therefore, a need to synthesize information on studies related to how the selected indoor air pollutants affected the cardiovascular health of older adults.

Keywords: particulate matter, PM_{2.5}, PM₁₀, nitrogen dioxide, myocardial infarction, stroke, hospitalisation, mortality, older adults

Introduction

Indoor air pollution is an emerging problem which negatively affects health of people worldwide. While air pollutants are a natural occurrence, their types and concentrations have skyrocketed due to industrialization and urbanization, driven by human activities.¹ Air pollution is a major global health threat, contributing to millions of deaths annually.² Outdoor pollutants alone cause over 4 million deaths, while indoor air pollution is responsible for an additional 2.3 million.³ This exposure raises the risk of various cardiovascular, respiratory, and neurological diseases.³ While the specific sources of indoor air pollution can differ depending on location, some common culprits are inefficient fuel burning for cooking and heating. The most used biomass fuels include wood, crop residues, and animal dung, while the solid mass fuels include charcoal and coal.⁴ Cooking and heating with these usually take place indoors using inefficient stoves, leading to incomplete combustion due to the limited oxygen supply. The result of incomplete combustion of these fuels is the production of carbon monoxide, black carbon, and complex organic carbon compounds.⁵ The situation is further aggravated by the inadequate ventilation prevalent in many kitchens in low income settings of developing countries. Indoor cooking with solid fuels often generates smoke containing particulate matter (PM) levels as high as 1000 mg/m³, with even higher concentrations having been documented.⁴ These levels far exceed the national ambient air quality standards for PM set by the US Environmental Protection Agency (EPA) and the guidelines established by the World Health Organization (WHO). In addition, use of household products such as cleaning detergents, pesticides, and solvents

often emit volatile organic compounds (VOCs) that can pollute the air.⁵ In 2015, nearly a quarter (24%) of deaths from ischemic heart disease, 23% of lung cancer deaths worldwide and over one-fifth (21%) of deaths from stroke were attributed to air pollution.⁶

Cardiovascular diseases (CVDs) are among the top causes of mortality and disability globally with over 17.9 million deaths recorded in 2019^{7,8} and this is predicted to escalate to 32.3 million in 2050. The risk of death from cardiovascular disease (CVD) increases dramatically with age.⁹ After 50, the number of fatalities from CVD roughly doubles with each passing decade.⁹ A disproportionate burden, three-quarters (75%) of these CVD deaths, falls on low- and middle-income countries (LMICs).⁷ Limited access to primary care in these regions hinders early detection and treatment, leading to delayed diagnoses and tragically, earlier deaths.⁷ The growing burden of cardiovascular diseases (CVDs) and their associated risk factors stems from a complex interplay of factors.¹⁰ This includes the ongoing shift in population demographics known as the epidemiological transition.¹⁰ Factors like increasing urbanization, changing lifestyles and diets, socioeconomic development, and modernization all play a role.¹⁰ Furthermore, growing evidence suggests that environmental factors significantly contribute to CVD risk. As much as a quarter of all ischemic heart disease (IHD) cases may be linked to an unhealthy environment – factors like air pollution and even traffic noise can have a significant impact.¹

People from middle- and low-income countries particularly experience the highest rates of health challenges related to poor air quality¹¹ because they lack the resources to obtain cleaner fuels and devices.¹² The key pollutants identified are particulate matter, sulphur dioxide, nitrogen dioxide, volatile organic compounds carbon monoxide, and hydrocarbons.¹³ The most common forms of indoor air pollution in low-income settings come from the biomass fuels which produce particulate matter (PM_{2.5} and PM₁₀) and carbon monoxide (CO).¹⁴ African cities are believed to have much higher levels of fine particulate matter (PM_{2.5}) compared to most European and North American cities.¹¹ PM_{2.5} concentrations in African cities are estimated to be around five times higher, reaching 100 micrograms per cubic meter (mg/m³), whereas most European and North American cities have levels below 20mg/m³.¹¹ In low-income countries, a significant portion of the population, an estimated 57%, is directly exposed to unhealthy concentrations of PM_{2.5}, a dangerous air pollutant.² Long-term exposure to particulate matter has been linked to oxidative stress and inflammation in the respiratory system, contributing to respiratory and cardiovascular diseases (CVDs).^{15,16} This oxidative stress occurs either through direct inhalation of reactive oxygen species (ROS) or indirectly via toxic aerosol compounds from sources like combustion. Inhaled particles can reach the alveoli, triggering inflammatory responses, while some ultrafine particles may enter the bloodstream, affecting cardiovascular function and the autonomic nervous system.¹⁵ Numerous studies have found that air pollution is associated with increased inflammatory biomarkers and subclinical effects such as oxidative stress and autonomic imbalance, emphasizing its impact on public health.¹⁷

Age is a well-established risk factor for a range of cardiovascular diseases (CVDs). This includes conditions like coronary artery disease, high blood pressure (hypertension), heart failure, and stroke.¹⁸ The global burden of cardiovascular disease (CVD) is increasing with age, and this trend is particularly concerning in low-income countries. Adults in these settings experience disproportionately higher rates of CVD compared to those in wealthier nations.¹⁹ Overtime, the body undergoes aging, a process marked by gradual buildup of cellular damage, a decline in how well our organs and systems function, and a greater risk of developing various diseases.¹⁸ With age comes a heightened risk of chronic illnesses like heart disease, diabetes, and cancer.²⁰ These pre-existing conditions in older adults can significantly worsen the impact of air pollutants on their health.²¹

Methods

The protocol for this review was registered with the International prospective register of systematic reviews (PROSPERO) in accordance with PRISMA guidelines (PROSPERO CRD42024479220).

Search Strategy

Online databases, particularly PubMed, Google Scholar, Scopus, Web of Science and Embase were used to conduct the systematic review between November 2023 and March 2024. Full Length prospective cohort studies, retrospective cohort studies, case-control studies, cross-sectional studies, randomized controlled trials (RCTs) and longitudinal studies with follow-up periods sufficient to assess cardiovascular health outcomes were used in this systematic review. Only those

Table 1 Search Strategy

Category	Keywords used
Population	"Elderly" OR "Seniors" OR "Geriatric population" OR "Aging population" OR "Older individuals".
Exposure	"Particulate matter (PM _{2.5} & PM ₁₀)" OR "Nitrogen dioxide" OR "Carbon monoxide".
Comparator	"Low pollution areas" OR "Clean air environments" OR "Reduced exposure to pollutants" OR "Control groups".
Outcome	"Cardiovascular disease" OR "Stroke" OR "Myocardial infarction" OR "Hypertension" OR "Arrhythmia" OR "Heart failure" OR "Mortality"

articles reported in English between 2013 and 2024 were considered for the review. The Population Exposure Comparison Outcome (PECO) tool was used to develop the search criteria as shown in Table 1. Briefly, the population of interest was older adults aged 65 years and above, the exposure was indoor air pollutants (NO₂; SO₂; PM_{2.5}; PM₁₀), comparator population was adults above 65 years living in less polluted/cleaner environments and the outcome was cardiovascular health. The retrieved documents were manually screened by two reviewers independently. Instances of discordance were addressed through deliberation to facilitate merging of extracted data. Ancillary sources, such as editorials, reviews, and case reports, were excluded from consideration, and materials classified as grey literature were also omitted from the analysis.

Data Extraction

The study titles were screened and records meeting the predetermined criteria were subsequently transferred to Microsoft Excel spreadsheet. Data such as the study title, abstract, and year of publication was extracted from these chosen studies. We removed duplicate entries and irrelevant studies. This process ensured only full-text articles relevant to our systematic review were included for further analysis.

Results

The details of the study selection process are shown in Figure 1. Results for the systematic review of literature are shown in Table 2.

All thirty-four studies included in this review were from only 3 continents (Asia, Europe and North America). More than half of the studies (22) included in this systematic review were from Asia, 8 from North America and the remaining 8 were from Europe. The most reported cardiovascular health outcomes from all the studies in descending order included mortality,^{26,27,29,34,35,40,51,55} myocardial infarction,^{23–25,34,38,39,46,55} stroke,^{24,26,34,41,44,49,54} increased CVD hospital admissions,^{22,28,41,42,47,49} heart failure,^{24,32,34,41,47} Ischaemic heart disease,^{26,32,41,53} increased emergency room visits (4 studies),^{37,45,48,50} hypertension^{15–17,30,56} and arrhythmia.^{24,39,48} The study conducted in the United States by Hayes et al revealed that a 10µg/m³ increase in PM_{2.5} was associated with a 16% increase in Ischaemic Heart disease-related mortality and a 14% increase in stroke-related mortality.²⁶ A study done in the Hubei province of China by Liu et al reported that short-term exposure to PM_{2.5}, PM₁₀, and NO₂ was associated with an increased risk of myocardial infarction mortality, and the association was stronger with NO₂.⁵⁵ Interestingly, 5 out of 8 studies reporting on mortality attributed it to NO₂.^{27,29,51,53,55} All studies reporting myocardial infarction as a resultant cardiovascular health outcome of indoor air pollution, except for one study by Atkinson et al, attributed the increased risk of myocardial infarction to exposure to PM_{2.5}.^{23,25,34,38,39,46,55} More specifically, the study done in Belgium on adults ≥ 75 years by Argacha et al reported that exposure to PM_{2.5} and NO₂ increased the risk of ST-elevation myocardial infarction (STEMI). Furthermore, they reported that the risk related to PM_{2.5} appears to be greater in the elderly compared to younger individuals.²³ A study conducted in Ontario by Bai et al reported a linear relationship between exposure to PM_{2.5} and acute myocardial infarction (AMI).²⁵ Contrary to these reports, Atkinson et al's study on older adults in the UK reported a weak association between indoor air pollution exposure and myocardial infarction.¹⁷




Figure 1 Study selection process.

Discussion

This systematic review investigated the impact of indoor air pollution on the cardiovascular health outcomes of older adults. Most studies came from Asia, possibly because of the high population density² and rapid industrialisation in countries like China, India, and South Korea⁵⁷ which has prompted extensive studies to understand its impacts. Majority (nearly 99%) of people are exposed to air pollution exceeding the safety limits set by the World Health Organization (WHO),³ with low- and middle-income countries being the most exposed.¹² Low-income countries face a double burden of air pollution due to weaker regulations, older machinery, subsidized fossil fuels, congested transportation, rapid industrial growth, and agricultural practices that all contribute to significantly higher pollution levels.² Grey literature has shown that airborne pollutants in Sub-Saharan African cities are primarily composed of gases and vapours (over 90%), namely particulate matter, NO₂, volatile organic compounds (eg, benzene), SO₂ and CO.¹³ Our systematic review did not find any studies reporting on the impact of indoor air pollution on the cardiovascular health outcomes of older adults in Africa despite the continent being flagged as potentially exposed to high pollution, as highlighted in grey literature.^{12,13} However, some studies have been done on indoor air quality and CVD risk in children in South Africa.^{58,59} The present findings indicate a notable deficiency in comprehension and awareness on the adverse impacts on cardiovascular well-being attributable to

Table 2 Cardiovascular Health Outcomes of Older Adults Exposed to Indoor Air Pollution

Country	Study Design	Sample Size	Population	Exposure	Outcome	Reference
China	Longitudinal	460 938	≥65 years	Short term PM _{2.5}	• Increased admission due to CHDs	[22]
Belgium	Case crossover	2598	≥75 years	10µg/m ³ of PM _{2.5} , NO ₂	• Increased the risk of MI	[23]
United Kingdom	Retrospective cohort	8336 557	40–89 years	Long term, PM, SO ₂ & NO ₂	• Weak association between exposure MI, stroke, or arrhythmia • Increased development of HF	[24]
Ontario, Canada	Retrospective Cohort	5100 000	35–85 years	PM _{2.5}	• Increased risk for acute MI	[25]
US	Longitudinal	565 477 men and women	50–71 years	10 µg/m ³ PM _{2.5}	• Increased mortality due to IHD and stroke	[26]
China	Longitudinal	21,816	≥75 years	PM ₁₀ , SO ₂ , and NO ₂	• Increased CVD mortality	[27]
Iran	Longitudinal	84,114	65–74 years	PM ₁₀ , NO ₂ , SO ₂ , CO	• Increased hospital admissions for CVD	[28]
Britain	Longitudinal	7569	40–69 years	Long term PM _{2.5} , PM ₁₀ , PMcoarse, SO ₂ , NO ₂	• Increased CVD Mortality	[29]
China	Prospective cohort	3754	≥65years	Biomass fuel	• Increased risk of hypertension	[30]
Shiraz, Iran	Case crossover		≥65 years	Short term exposure, PM ₁₀ , SO ₂ , and NO ₂	• Increased CVD morbidity	[31]
New York City	Case control	837523	67±15 years	NO ₂ , PM _{2.5} , SO ₂	• Increased risk of CVD, IHD, and HF	[32]
US	Prospective cohort study	6795	45–84 years	PM _{2.5}	• Increased coronary calcification, increased progression of atherosclerosis	[33]
Korea	Retrospective cohort study	900845	≥65 years	PM _{2.5}	• Increased cardiovascular mortality, • Acute MI • Congestive heart failure • Stroke.	[34]
Northern California	Retrospective cohort	169714	60.8±15.6	PM _{2.5}	• Increased risk of cardiovascular mortality	[35]
US		502	29–89 years	PM _{2.5}	• Increased risk of AF	[36]
Beijing	Longitudinal	82,430	≥65 years	NO ₂ , SO ₂ , PM ₁₀	• Increased cardiovascular emergency room admissions	[37]
North Carolina	Retrospective cohort	5679	60.8±12.1	Daily, long term PM _{2.5}	• Increased CAD and incidence of MI	[38]
Wales	Case cross over		60–81 years	PM _{2.5} NO ₂	• Arrhythmias and AF • Increased risk for MI	[39]
North-Eastern Iran	Retrospective Cohort	50045	40–75 years	Biomass fuels	• Increased risk for CVD mortality	[40]
Vietnam	Case crossover	135 101	≥65 years	PM _{2.5} ; PM ₁ ; SO ₂	• Increased daily hospital admissions due to CVD conditions (IHD, CF, stroke)	[41]
Thailand	Retrospective cohort	503 105	≥65 years	NO ₂ , SO ₂ , PM ₁₀	• Increase in CVD hospital admission	[42]
Vietnam	Retrospective cohort	N/A	≥65 years	PM ₁₀ ; NO ₂ & SO ₂	• Increased cardiovascular admissions	[43]
Western Germany	Prospective Cohort	4105	45–76 years	PM _{2.5} and PM ₁₀	• Increased risk of stroke	[44]
New Mexico	Case crossover	4739	≥65 years	PM _{2.5} and PM ₁₀	• Increased emergency room visits for CVDs	[45]

(Continued)

Table 2 (Continued).

Country	Study Design	Sample Size	Population	Exposure	Outcome	Reference
Finland	Case crossover	N/A	65.5±12.5 years	PM _{2.5}	<ul style="list-style-type: none"> Increased risk of acute MI 	[46]
China	Prospective cohort	4866	65.2±13.5 years	PM _{2.5} , PM ₁₀ , CO, SO ₂	<ul style="list-style-type: none"> Increased risk of HF readmission 	[47]
Beijing China	Time series	N/A	45–75 years	SO ₂ , NO ₂ , PM ₁₀ PM _{2.5}	<ul style="list-style-type: none"> Increased CVD emergency room visits Increased arrhythmia 	[48]
Beijing, China	Retrospective cohort	63,956	≥65 years	Short term PM _{2.5}	<ul style="list-style-type: none"> Increased hospitalisation for Ischaemic stroke 	[49]
Shanghai, China	Cross sectional	N/A	≥65 years	PM _{2.5} , PM ₁₀	<ul style="list-style-type: none"> Increased risk of CHD outpatients and emergency department visit 	[50]
Hong Kong, China	Prospective cohort	66820	≥65 years	Long term PM _{2.5} , NO ₂	<ul style="list-style-type: none"> Elevated risk of cardiovascular mortality. 	[51]
China	Cross sectional	24,845	18–74 years	PM ₁ , PM _{2.5}	<ul style="list-style-type: none"> Increased prevalence of CVDs 	[52]
Wuhan, China	Case cross over	8955	≥65 years	PM ₁₀ , NO ₂ , SO ₂ NO ₂	<ul style="list-style-type: none"> Increased risk of IHD Cardiovascular mortality 	[53]
Okayama, Japan	Case crossover	10,949	≥65 years	Particulate matter	<ul style="list-style-type: none"> Increased risk of haemorrhagic and ischemic stroke 	[54]
Hubei Province, China	Case cross over	15 1608	70–85 years	PM _{2.5} , PM ₁₀ , and NO ₂	<ul style="list-style-type: none"> Increased MI mortality. The association was stronger with NO₂ 	[55]
Germany	Randomised controlled exposure	54	65–79 years	PM ₁ , PM _{2.5} and PM ₁₀	<ul style="list-style-type: none"> PM_{2.5} and PM₁₀ increased systolic blood pressure 	[15]
Beijing, China	Randomised crossover intervention	35	Average 66.26 years	PM _{2.5} and black carbon	<ul style="list-style-type: none"> PM_{2.5} increased BP 	[56]
Chongqing, China	Randomised double-blind crossover trial	24	61–97 years	PM ₁ , PM _{2.5} and PM ₁₀	<ul style="list-style-type: none"> Increased markers of blood inflammation and heart rate 	[17]
Germany	Randomised controlled exposure	55	65–79 years	PM ₁ , PM _{2.5} and PM ₁₀	<ul style="list-style-type: none"> PM₁ and PM_{2.5} increased arterial stiffness indices 	[17]

Abbreviations: NO₂, Nitrogen dioxide; SO₂, Sulphur dioxide; CO, Carbon monoxide; PM, Particulate matter; PM_{2.5}, Particulate matter 2.5; PM₁₀, Particulate matter 10; MI, Myocardial infarction; CF, Cardiac failure; CVD, cardiovascular diseases; CHD, coronary heart diseases; CAD, coronary artery diseases; AF, Atrial fibrillation; IHD, Ischaemic heart diseases; HF, Heart failure; BP, Blood pressure.

air pollution across all life stages in the African continent. It is known that cardiovascular aging is a major trigger for cardiovascular diseases in elderly people. As cardiovascular function declines with age, the risk of atherosclerosis, stroke, heart attack (myocardial infarction), and other diseases with poor outcomes all rise significantly.⁹ Stroke has previously been reported as the major contributor to the total CVD deaths in people older than 85 years.⁶⁰ While ischemic stroke is more common overall, intracerebral hemorrhage causes the most disability-adjusted life years (DALYs) and deaths in people under 70.⁶¹ Our study confirms the reports by previous studies; mortality, myocardial infarction, and stroke were the top 3 reported cardiovascular health outcomes of older adults in this systematic review, and in some studies, mortality was attributed to myocardial infarction or stroke. This illustrates the extent of deleterious effects of exposure to atmospheric pollutants on cardiovascular health outcomes among the elderly demography.

Particulate matter (PM), particularly PM_{2.5} was the most reported air pollutant with most studies reporting myocardial infarction. This could be because PM_{2.5}, the major component of air pollution, easily enters the bloodstream due to its tiny size, potentially affecting organs more readily than larger PM₁₀ particles. Although there are finer particles than PM_{2.5}, known as the ultra-fine particles with an aerodynamic diameter of less than 0.1µm, also known as UFPs, the latter are less abundant in the atmosphere than PM_{2.5}.⁶² However, both PM_{2.5} and even finer particles called UFPs can travel deep into the lungs and enter the bloodstream, potentially causing harm throughout the body.⁶³ Inhaling air pollution, particularly PM_{2.5}, can trigger harmful consequences within the lungs. These tiny particles reach the alveoli, the air sacs where oxygen exchange occurs. PM can directly generate reactive oxygen species (ROS), molecules that damage cells or indirectly cause their production.⁶⁴ This process, known as oxidative stress, can also be initiated by inhaling other toxic compounds in air pollution.⁵ The lungs respond to PM exposure by activating immune cells called alveolar macrophages. These macrophages release inflammatory molecules (cytokines) that can travel throughout the body, potentially affecting heart function and blood vessel control. PM enters the body through the respiratory tract, with larger particles (>50µm) often trapped by the nose. PM₁₀ may lodge in the upper airways, while the tiny PM_{2.5} particles can penetrate deep into the lungs and even enter the bloodstream. It's important to note that carbon monoxide (CO), another air pollutant, has a different mechanism of harm. Unlike PM, CO toxicity does not involve oxidative stress but rather its ability to disrupt oxygen transport in the blood at very high concentrations (not typically found in ambient air).⁶⁴ When inhaled, CO travels to the lungs and reaches the alveoli.⁶⁵ Once in the alveoli, CO diffuses across the thin alveolar-capillary membrane into the bloodstream, displacing O₂ in haemoglobin and binds irreversibly to haemoglobin thus depriving organs of oxygen leading to hypoxia.⁶⁴ This systematic review reported CO in only 2 studies,^{28,47} possibly because it usually does not occur in ambient air. A study by Shah et al found a link between rising carbon monoxide (CO) levels and an increased risk of hospitalization or death from heart failure (HF).⁶⁴ This risk increased by 3.52% for every one part per million increase in CO levels.⁶⁴ The confidence interval (CI) indicates a 2.52% to 4.54% increase in risk.⁶⁴ One of the studies included in this review also reported that CO increased CVD hospitalisations,²⁸ while that by Shi et al reported an increased risk in HF readmissions.⁴⁷

Nitrogen Dioxide (NO₂) is a gas formed from both natural processes and human activities like burning fossil fuels in power plants and cars.^{66,67} It can also contribute to the formation of other harmful air pollutants.⁶⁶ Studies have shown a link between NO₂ exposure and negative health effects, particularly on the heart. In people with weakened heart muscles (dilated cardiomyopathy), NO₂ has been associated with structural changes in the heart itself.⁶⁸ Long-term exposure to NO₂ may also increase the risk of death from heart disease and other causes, with one study finding a 13% rise in cardiovascular deaths for every 10 micrograms per cubic meter increase in annual NO₂ levels.⁶⁹ Research by Huang et al further suggests that long-term NO₂ exposure can increase the risk of death from various causes, including respiratory and cardiovascular problems⁶⁷.

In the current study, NO₂ was the most dominant air pollutant in mortality reports,^{27,29,51,53,55} confirming the previously reported association of this pollutant with increased mortality. A study by Zhang et al in China investigating the link between air pollution and cardiovascular deaths found no statistically significant effect of NO₂ on CVD mortality.²⁷ This suggests that the impact of NO₂ on heart health may vary depending on other factors not explored in any of the studies included in this systematic review. Later in 2021, Liu et al carried out a study in China on short term exposure to air pollution and mortality from MI. Their study found that the risk from PM_{2.5} and PM₁₀ exposures rose sharply until certain levels (PM_{2.5} at 33.3 mg/m³ and PM₁₀ at 57.3 mg/m³), which levelled off. In contrast, the risk from NO₂ exposure increased steadily nearly linearly.⁵⁵ This implies minimal risk associated with PM_{2.5} and PM₁₀ exposure levels beyond the threshold points. Meanwhile, with the increase in NO₂ levels, the risk of adverse health outcomes

increases consistently, without a noticeable breakpoint where the risk changes dramatically. Interestingly, the study also found that the link between NO_2 exposure and death from heart attack (MI) was stronger in older adults compared to younger people.⁵⁵ This contradicts previous findings suggesting that NO_2 has a greater impact on younger patients.²³ While NO_2 is often used to indicate overall air pollution, research is ongoing to determine its exact independent contributions to health problems. However, increasing evidence suggests that NO_2 can have harmful effects on its own beyond simply reflecting the presence of other pollutants.^{55,67,69} The connection between exposure to these pollutants and adverse cardiovascular outcomes in older adults is shown in Figure 2 and the two possible mechanisms linking air pollutants to CVDs are summarized in Figure 3.




Figure 2 Connection between exposure to indoor air pollutants and adverse cardiovascular outcomes in older adults. An illustration of the cardiovascular health impacts of indoor air pollution from the use of biomass and solid fuels for cooking. Pollutants such as particulate matter, sulphur dioxide, nitrogen dioxide, and carbon monoxide are released during combustion and inhaled into the body. These pollutants contribute to cardiovascular conditions, including stroke, ischemic heart disease (IHD), arrhythmia, heart failure, myocardial infarction, and ultimately increased mortality.




Figure 3 Two possible mechanisms by which air pollutants affect the cardiovascular system. 1: PM_{10} and $\text{PM}_{2.5}$ can penetrate deep into the respiratory system and even enter the bloodstream. Once in the body, they can trigger inflammation. PM-induced inflammation can lead to endothelial dysfunction, oxidative stress, and activation of immune cells. Inflammation plays a crucial role in the development and progression of cardiovascular diseases such as atherosclerosis, hypertension, and heart failure. 2: NO_2 can impact the autonomic nervous system, which regulates involuntary bodily functions such as heart rate, blood pressure, and respiratory rate. This imbalance in autonomic tone can contribute to the development of cardiovascular diseases by promoting hypertension, arrhythmias, and myocardial ischemia.

Conclusion

Studies have linked exposure to indoor air pollutants with a higher risk of cardiovascular problems in older adults. This includes an increased chance of death from cardiovascular disease (CVD), heart attack (MI), and stroke. Additionally, older adults exposed to indoor air pollution may experience more hospital admissions and emergency room visits for cardiovascular issues. PM_{2.5} was associated with most reports of myocardial infarction, while NO₂ was associated with most mortalities reported by studies in this review. These findings imply that exposure to PM₁₀, PM_{2.5} and NO₂ can potentially reduce the quality of life of older adults. The paucity of data on the impact of indoor air pollutants on cardiovascular health in some continents poses a public health challenge as individuals may be exposed unknowingly leading to potential cardiovascular problems such as MI, stroke and even death. Additionally, in the absence of comprehensive data, public health policies and regulations may not adequately address indoor air quality issues, resulting in failure to mitigate exposure to harmful pollutants, leaving populations vulnerable to associated health risks. There is need for epidemiological studies in the understudied continents to identify the health risks associated with exposure to indoor air pollutants.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Münzel T, Hahad O, Sørensen M, et al. *Environmental Risk Factors and Cardiovascular Diseases: A Comprehensive Expert Review. Vol. 118.* Cardiovascular Research. Oxford University Press; 2022:2880–2902.
2. Rentschler J, Leonova N. Global air pollution exposure and poverty. *Nat Commun.* 2023;14(1). doi:10.1038/s41467-023-39797-4
3. Taghizadeh-Hesary F, Taghizadeh-Hesary F. The impacts of air pollution on health and economy in Southeast Asia. *Energies.* 2020;13(7):1812.
4. Balmes JR. Household air pollution from domestic combustion of solid fuels and health. *J Allergy Clin Immunol Mosby Inc.* 2019;143(6):1979–1987. doi:10.1016/j.jaci.2019.04.016
5. Caldeira D, Franco F, Bravo Baptista S, et al. Air pollution and cardiovascular diseases: a position paper. *Revista Portuguesa de Cardiologia.* 2022;41(8):709–717. doi:10.1016/j.repc.2022.05.006
6. Mannucci PM, Harari S, Franchini M. *Novel Evidence for a Greater Burden of Ambient Air Pollution on Cardiovascular Disease.* Vol. 104. Haematologica. Ferrata Storti Foundation; 2019:2349–2357.
7. Okop K, Delobelle P, Lambert EV, et al. Implementing and Evaluating Community Health Worker-Led Cardiovascular Disease Risk Screening Intervention in Sub-Saharan Africa Communities: a Participatory Implementation Research Protocol. *Int J Environ Res Public Health.* 2023;20(1):298.
8. An Z, Jin Y, Li J, Li W, Wu W. Impact of Particulate Air Pollution on Cardiovascular Health. *Current Allergy and Asthma Reports Current Medicine Group LLC I.* 2018;18:1–7.
9. Kuntic M, Kuntic I, Hahad O, Lelieveld J, Münzel T, Daiber A. Impact of air pollution on cardiovascular aging. *Mech Ageing Dev.* 2023;214.
10. Odukoya JO, Odukoya JO, Mmutlane EM, Ndinteh DT. Ethnopharmacological Study of Medicinal Plants Used for the Treatment of Cardiovascular Diseases and Their Associated Risk Factors in sub-Saharan Africa. *Plants.* 2022;11(10):1387. doi:10.3390/plants11101387
11. Amegah AK, Agyei-Mensah S. *Urban Air Pollution in Sub-Saharan Africa: Time for Action.* Vol. 220. Environmental Pollution. Elsevier Ltd; 2017:738–743.
12. Burning Opportunity: clean Household Energy for Health, Sustainable Development, and Wellbeing of Women and Children [Internet]. 2016. Available from: www.who.int. Accessed September 29, 2024.
13. Schwela D Review of Urban Air Quality in Sub-Saharan Africa Region-Air Quality profile of SSA countries. World Bank; 2012.
14. Tumwesige V, Okello G, Semple S, Smith J. Impact of partial fuel switch on household air pollutants in sub-Saharan Africa. *Environ. Pollut.* 2017;231:1021–1029. doi:10.1016/j.envpol.2017.08.118
15. Soppa VJ, Schins RPF, Hennig F, et al. Arterial blood pressure responses to short-term exposure to fine and ultrafine particles from indoor sources – a randomized sham-controlled exposure study of healthy volunteers. *Environ Res.* 2017;158:225–232. doi:10.1016/j.envres.2017.06.006
16. Soppa VJ, Shinnawi S, Hennig F, et al. Effects of short-term exposure to fine and ultrafine particles from indoor sources on arterial stiffness – a randomized sham-controlled exposure study. *Int J Hyg Environ Health.* 2019;222(8):1115–1132. doi:10.1016/j.ijheh.2019.08.002
17. Guo M, Du C, Li B, et al. Reducing particulates in indoor air can improve the circulation and cardiorespiratory health of old people: a randomized, double-blind crossover trial of air filtration. *Science of the Total Environment.* 2021;798:149248.
18. Jaul E, Barron J. Age-Related Diseases and Clinical and Public Health Implications for the 85 Years Old and Over Population. In: *Frontiers in Public Health.* Vol. 5. Frontiers Media S.A.; 2017.
19. Gaziano TA, Bitton A, Anand S, Abrahams-Gessel S, Murphy A. Growing Epidemic of Coronary Heart Disease in Low- and Middle-Income Countries. *Curr Probl Cardiol.* 2010;35(2):72–115. doi:10.1016/j.cpcardiol.2009.10.002
20. Prasad S, Sung B, Aggarwal BB. *Age-Associated Chronic Diseases Require Age-Old Medicine: Role of Chronic Inflammation.* Vol. 54. Preventive Medicine; 2012.
21. Tilstra MH, Tiwari I, Niwa L, et al. Risk and resilience: how is the health of older adults and immigrant people living in Canada impacted by climate-and air pollution-related exposures? *Int J Environ Res Pub Health MDPI.* 2021;18:10575.

22. Amsalu E, Wang T, Li H, et al. Acute effects of fine particulate matter (PM_{2.5}) on hospital admissions for cardiovascular disease in Beijing, China: a time-series study. *Environ Health*. 2019;18(1). doi:10.1186/s12940-019-0506-2.
23. Argacha JF, Collart P, Wauters A, et al. Air pollution and ST-elevation myocardial infarction: a case-crossover study of the Belgian STEMI registry 2009–2013. *Int J Cardiol*. 2016;223:300–305. doi:10.1016/j.ijcard.2016.07.191
24. Atkinson RW, Carey IM, Kent AJ, Van Staa TP, Ross Anderson H, Cook DG. Long-term exposure to outdoor air pollution and incidence of cardiovascular diseases. *Epidemiology*. 2013;24(1):44–53. doi:10.1097/EDE.0b013e318276ccb8
25. Bai L, Shin S, Burnett RT, et al. Exposure to ambient air pollution and the incidence of congestive heart failure and acute myocardial infarction: a population-based study of 5.1 million Canadian adults living in Ontario. *Environ Int*. 2019;123:132. doi:10.1016/j.envint.2018.11.066
26. Hayes RB, Lim C, Zhang Y, et al. PM_{2.5} air pollution and cause-specific cardiovascular disease mortality. *Int J Epidemiol*. 2020;49(1):25–35. doi:10.1093/ije/dyz114
27. Zhang C, Ding R, Xiao C, et al. Association between air pollution and cardiovascular mortality in Hefei, China: a time-series analysis. *Environ. Pollut*. 2017;229:790–797. doi:10.1016/j.envpol.2017.06.022
28. Dastoorpoor M, Sekhavatpour Z, Masoumi K, et al. Air pollution and hospital admissions for cardiovascular diseases in Ahvaz, Iran. *Science of the Total Environment*. 2019;652:1318–1330. doi:10.1016/j.scitotenv.2018.10.285
29. Dehbi HM, Blangiardo M, Gulliver J, et al. Air pollution and cardiovascular mortality with over 25 years follow-up: a combined analysis of two British cohorts. *Environ Int*. 2017;99:275–281. doi:10.1016/j.envint.2016.12.004
30. Deng Y, Gao Q, Yang D, et al. Association between biomass fuel use and risk of hypertension among Chinese older people: a cohort study. *Environ Int*. 2020;138:105620.
31. Gharehchahi E, Mahvi AH, Amini H, et al. Health impact assessment of air pollution in Shiraz, Iran: a two-part study. *J Environ Health Sci Eng*. 2013;11(1). doi:10.1186/2052-336X-11-11.
32. Humphrey JL, Kinnee EJ, Robinson LF, Clougherty JE. Disentangling impacts of multiple pollutants on acute cardiovascular events in New York city: a case-crossover analysis. *Environ Res*. 2024;242:117758.
33. Kaufman JD, Adar SD, Barr RG, et al. Association between air pollution and coronary artery calcification within six metropolitan areas in the USA (the Multi-Ethnic Study of Atherosclerosis and Air Pollution): a longitudinal cohort study. *Lancet*. 2016;388(10045):696–704. doi:10.1016/S0140-6736(16)00378-0
34. Kim H, Kim J, Kim S, et al. Cardiovascular effects of long-term exposure to air pollution: a population-based study with 900 845 person-years of follow-up. *J Am Heart Assoc*. 2017;6(11). doi:10.1161/JAHA.117.007170.
35. Alexeeff SE, Deosaransingh K, Liao NS, van Den Eeden SK, Schwartz J, Sidney S. Particulate matter and cardiovascular risk in adults with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2021;204(2):159–167. doi:10.1164/rccm.202007-2901OC
36. Link MS, Luttmann-Gibson H, Schwartz J, et al. Acute exposure to air pollution triggers atrial fibrillation. *J Am Coll Cardiol*. 2013;62(9):816–825. doi:10.1016/j.jacc.2013.05.043
37. Ma Y, Zhao Y, Yang S, et al. Short-term effects of ambient air pollution on emergency room admissions due to cardiovascular causes in Beijing, China. *Environ. Pollut*. 2017;230:974–980. doi:10.1016/j.envpol.2017.06.104
38. McGuinn LA, Ward-Caviness CK, Neas LM, et al. Association between satellite-based estimates of long-term PM_{2.5} exposure and coronary artery disease. *Environ Res*. 2016;145:9–17. doi:10.1016/j.envres.2015.10.026
39. Milojevic A, Wilkinson P, Armstrong B, Bhaskaran K, Smeeth L, Hajat S. Short-term effects of air pollution on a range of cardiovascular events in England and Wales: case-crossover analysis of the MINAP database, hospital admissions and mortality. *Heart*. 2014;100(14):1093–1098. doi:10.1136/heartjnl-2013-304963
40. Mitter SS, Vedanthan R, Islami F, et al. Household fuel use and cardiovascular disease mortality. *Circulation*. 2016;133(24):2360–2369. doi:10.1161/CIRCULATIONAHA.115.020288
41. Nhung NTT, Schindler C, Chau NQ, et al. Exposure to air pollution and risk of hospitalization for cardiovascular diseases amongst Vietnamese adults: case-crossover study. *Science of the Total Environment*. 2020;703:136637.
42. Phosri A, Ueda K, Phung VLH, Tawatsupa B, Honda A, Takano H. Effects of ambient air pollution on daily hospital admissions for respiratory and cardiovascular diseases in Bangkok, Thailand. *Science of the Total Environment*. 2019;651:1144–1153. doi:10.1016/j.scitotenv.2018.09.183
43. Phung D, Hien TT, Linh HN, et al. Air pollution and risk of respiratory and cardiovascular hospitalizations in the most populous city in Vietnam. *Science of the Total Environment*. 2016;557–558:322–330. doi:10.1016/j.scitotenv.2016.03.070
44. Rodins V, Lucht S, Ohlwein S, et al. Long-term exposure to ambient source-specific particulate matter and its components and incidence of cardiovascular events – the Heinz Nixdorf Recall study. *Environ Int*. 2020;142:105854.
45. Rodopoulou S, Chalbot MC, Samoli E, DuBois DW, San Filippo BD, Kavouras IG. Air pollution and hospital emergency room and admissions for cardiovascular and respiratory diseases in Doña Ana County, New Mexico. *Environ Res*. 2014;129:39–46. doi:10.1016/j.envres.2013.12.006
46. Rosenthal FS, Kuisma M, Lanki T, et al. Association of ozone and particulate air pollution with out-of-hospital cardiac arrest in Helsinki, Finland: evidence for two different etiologies. *J Expo Sci Environ Epidemiol*. 2013;23(3):281–288. doi:10.1038/jes.2012.121
47. Shi Y, Zhang L, Li W, et al. Association between long-term exposure to ambient air pollution and clinical outcomes among patients with heart failure: findings from the China PEACE Prospective Heart Failure Study. *Ecotoxicol Environ Saf*. 2021;222:112517.
48. Su C, Bretnier S, Schneider A, et al. Short-term effects of fine particulate air pollution on cardiovascular hospital emergency room visits: a time-series study in Beijing, China. *Int Arch Occup Environ Health*. 2016;89(4):641–657. doi:10.1007/s00420-015-1102-6
49. Tian Y, Xiang X, Wu Y, et al. Fine Particulate Air Pollution and First Hospital Admissions for Ischemic Stroke in Beijing, China. *Sci Rep*. 2017;7(1):3897.
50. Ye Z, Li X, Han Y, Wu Y, Fang Y. Association of long-term exposure to PM_{2.5} with hypertension and diabetes among the middle-aged and elderly people in Chinese mainland: a spatial study. *BMC Public Health*. 2022;22(1). doi:10.1186/s12889-022-12984-6
51. Yang Y, Tang R, Qiu H, et al. Long term exposure to air pollution and mortality in an elderly cohort in Hong Kong. *Environ Int*. 2018;117:99–106. doi:10.1016/j.envint.2018.04.034
52. Yang BY, Guo Y, Markevych I, et al. Association of Long-term Exposure to Ambient Air Pollutants with Risk Factors for Cardiovascular Disease in China. *JAMA Network Open*. 2019;2(3):e190318. doi:10.1001/jamanetworkopen.2019.0318
53. Liu Y, Chen X, Huang S, et al. Association between air pollutants and cardiovascular disease mortality in Wuhan, China. *Int J Environ Res Public Health*. 2015;12(4):3506–3516. doi:10.3390/ijerph120403506

54. Rowley HA. *The Alphabet of Imaging in Acute Stroke Does It Spell Improved Selection and Outcome?* In: Stroke. Lippincott Williams and Wilkins; 2013.
55. Liu Y, Pan J, Fan C, et al. Short-Term Exposure to Ambient Air Pollution and Mortality From Myocardial Infarction. *J Am Coll Cardiol.* 2021;77(3):271–281. doi:10.1016/j.jacc.2020.11.033
56. Liu S, Chen J, Zhao Q, et al. Cardiovascular benefits of short-term indoor air filtration intervention in elderly living in Beijing: an extended analysis of BIAPSY study. *Environ Res.* 2018;167:632–638. doi:10.1016/j.envres.2018.08.026
57. Iqbal M, Kalim R, Arshed N. Evaluating industrial competitiveness strategy in achieving environmental sustainability. *Competitiveness Rev.* 2024;34(2):353–369. doi:10.1108/CR-12-2022-0191
58. Chungag A, Engwa GA, Sewani-Rusike CR, Nkeh-Chungag BN. Effect of seasonal variation on the relationship of indoor air particulate matter with measures of obesity and blood pressure in children. *J Health Pollut.* 2021;11(30):210610. doi:10.5696/2156-9614-11.30.210610
59. Lung. function indices in children. doi:10.1016/S0140-
60. Sidney S, Go AS, Jaffe MG, Solomon MD, Ambrosy AP, Rana JS. Association between Aging of the US Population and Heart Disease Mortality from 2011 to 2017. *JAMA Cardiol.* 2019;4(12):1280–1286. doi:10.1001/jamacardio.2019.4187
61. Pacheco-Barrios K, Giannoni-Luza S, Navarro-Flores A, et al. Burden of Stroke and Population-Attributable Fractions of Risk Factors in Latin America and the Caribbean. *J Am Heart Assoc.* 2022;11(21). doi:10.1161/JAHA.122.027044.
62. Anderson JO, Thundiyil JG, Stolbach A. Clearing the Air: a Review of the Effects of Particulate Matter Air Pollution on Human Health. *J Med Toxicol.* 2012;8(2):166–175. doi:10.1007/s13181-011-0203-1
63. Lv S, Shi Y, Xue Y, et al. Long-term effects of particulate matter on incident cardiovascular diseases in middle-aged and elder adults: the CHARLS cohort study. *Ecotoxicol Environ Saf.* 2023;262:115181.
64. Shah ASV, Langrish JP, Nair H, et al. Global association of air pollution and heart failure: a systematic review and meta-analysis. *Lancet.* 2013;382(9897):1039–1048. doi:10.1016/S0140-6736(13)60898-3
65. Chu LM, Shaefi S, Byrne JD, Alves de Souza RW, Otterbein LE. Carbon monoxide and a change of heart. *Redox Biol.* 2021;48:102183.
66. Pérez Velasco R, Jarosińska D. Update of the WHO global air quality guidelines: systematic reviews – an introduction. *Environ Int.* 2022;170:107556.
67. Huang S, Li H, Wang M, et al. *Long-Term Exposure to Nitrogen Dioxide and mortality: A Systematic Review and Meta-Analysis.* Vol. 776. Science of the Total Environment. Elsevier B.V.; 2021.
68. Fecht D, Chadeau-hyam MARC, Owen RUTH, et al. Exposure to Elevated Nitrogen Dioxide Concentrations and Cardiac Remodeling in Patients With Dilated Cardiomyopathy. *J Card Fail.* 2022;28(6):924–934. doi:10.1016/j.cardfail.2021.11.023
69. Faustini A, Rapp R, Forastiere F. Nitrogen dioxide and mortality: review and meta-analysis of long-term studies. *Eur Resp J Eur Resp Soc.* 2014;44(3):744–753. doi:10.1183/09031936.00114713

Clinical Interventions in Aging

Dovepress

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>