

# Profile of Trofinetide in the Treatment of Rett Syndrome: Design, Development and Potential Place in Therapy

Laura Camillo<sup>1,\*</sup>, Marco Pozzi<sup>1,\*</sup>, Pia Bernardo<sup>2,\*</sup>, Simone Pisano<sup>3</sup>, Maria Nobile<sup>1</sup>

<sup>1</sup>Scientific Institute IRCCS Eugenio Medea, Bosisio Parini, LC, Italy; <sup>2</sup>Department of Neurosciences, Santobono-Pausilipon Children's Hospital, Naples, NA, Italy; <sup>3</sup>Department of Translational Medical Sciences, University Federico II, Naples, NA, Italy

\*These authors contributed equally to this work

Correspondence: Marco Pozzi, Scientific Institute IRCCS Eugenio Medea, Via Don Luigi Monza, 20, Bosisio Parini, LC, 23842, Italy, Tel +39 031 877 919, Email marco.pozzi@lanostrafamiglia.it

**Abstract:** Trofinetide is a first-in-class pharmacological treatment proposed for patients with Rett Syndrome. It is a long half-life derivative of glycine-proline-glutamate, the tripeptide normally excised from Insulin-like Growth Factor 1 upon degradation. Due to containing glutamate and glycine in its structure, trofinetide is thought to act through NMDA receptor modulation, thus providing a normalization of neuronal activity and survival. Trofinetide was tested in a series of short and long-term trials, showing good efficacy at improving scores on the Clinical Global Impression-Improvement scale and Rett Syndrome Behavior Questionnaire, with specific effect only on some subscales, ie General Mood subscale and Repetitive Face Movement subscale. No effects were documented on other subscales or on epilepsy, heart and bone-related symptoms. The main adverse effects of trofinetide, severe enough to determine discontinuation, include diarrhea, vomiting, and consequent weight loss. These may be scarcely avoidable, given the need to assume a very large amount of trofinetide per day. Other inherent limitations of use possibly regard the limited duration of drug supplies, as one bottle may last three days only, depending on weight, and the relatively high cost per bottle. Trofinetide has no direct competitors: single symptoms of the Rett Syndrome, for instance, seizures or aggressive behaviors, are currently treated with drugs that have been developed for patients without the Rett Syndrome. This leads to suboptimal efficacy and increased risk of adverse effects. The place in therapy of trofinetide is yet to be determined, based on the results of clinical trials, on its practical usability, and on the windows of opportunity for intervention. Moreover, trofinetide may be curative if given early enough during brain development, or merely symptomatic if given to young adults, and no data exist on this aspect. The place in therapy of trofinetide will require reassessment after competing treatments enter the market.

**Keywords:** IGF-1, GPE, GH, NMDA, efficacy, safety

## Introduction

Trofinetide is the first drug approved specifically for the treatment of patients with Rett Syndrome. It is a long-acting derivative of glycine-proline-glutamate (GPE), the tripeptide generated by enzymatic cleavage of Insulin-like Growth Factor 1 (IGF-1). To date, several reviews are available that deal with the chemical and preclinical development of this compound.<sup>1,2</sup> However, the literature is lacking with respect to defining the clinical needs of patients affected by the Rett Syndrome and there is no review available on the current therapeutic options. Therefore, in this narrative review, we will provide an overview of the clinical picture and the therapeutic alternatives already available for clinicians. This will allow us to focus on the place in therapy of trofinetide, with respect to the complex symptoms of the Rett Syndrome and the heterogeneous picture of concomitant drug treatments in routine clinical use.

## Materials and Methods

The initial literature analysis was performed using the Epistemic AI platform<sup>3</sup> using the results obtained by searching what drugs are used and being developed for the Rett Syndrome.

The narrative literature review was conducted in PubMed database using the search terms (Rett syndrome) (treatment) (drug); this search resulted in 489 articles. Inclusion criteria were being published in the last 20 years (444 articles retained), being conducted on human populations (297 articles retained) and written in English (280 articles retained). We screened articles and excluded all those who were not clinical studies, clinical trials, randomized controlled trials or meta-analysis. Finally, we analyzed the subset of 38 selected articles. After reading the full texts, 12 articles were selected based on the relevance of the studied drug treatments.

## Rett Syndrome

Rett Syndrome (RTT; OMIM entry #312750) is a genetic neurodevelopmental disorder predominantly affecting females, with a prevalence ranging from 1:10,000 to 1:23,000 female live births. Though RTT diagnosis is mainly clinical, more than 95% of cases are caused by mutations in the gene encoding the methyl-CpG binding protein 2 (*MECP2*), a transcriptional regulator involved in chromatin remodeling and the modulation of RNA splicing.<sup>16,17</sup> The variation spectrum of *MECP2* mutations appears to be independent of ethnicity.<sup>18</sup>

## RTT Presentation

RTT is characterized by a unique developmental regression period, occurring after apparently normal early development, with the onset of autistic-like features including intense stereotypic midline hand movements, impairment of communicative abilities and of fine motor skills.<sup>19</sup> Other clinical features include postnatal deceleration of head growth, epilepsy, cognitive impairment, scoliosis, feeding difficulties, growth restriction, sleep disturbance, bruxism, autonomic and motor dysfunction. RTT diagnosis is clinically based on distinct criteria revised in 2010, independent of molecular findings.<sup>19</sup> With respect to previous criteria of 2002, the revision recommended the presence of regression plus four main criteria to be required for the diagnosis of typical RTT. The four main criteria included:

1. Partial or complete loss of acquired purposeful hand skills;
2. Partial or complete loss of acquired spoken language;
3. Gait abnormalities: impairment (dyspraxia) or absence of ability;
4. Stereotypic hand movements such as hand wringing/squeezing, clapping/tapping, mouthing and washing/rubbing automatisms.

The post-natal deceleration in head growth was eliminated from the necessary criteria because this feature is not found in all individuals with typical RTT.<sup>19</sup>

## RTT Staging

The symptoms evolve with development and typically progress through four stages.<sup>17,19,20</sup>

Stage 1 is known as the “early stagnation stage” (6–18 months). During this phase, the symptoms may be very subtle at first and may go unnoticed. The main symptoms are issues with sitting, crawling, and walking, lack of interest in toys, problems in speech development, low muscle tone, or hypotonia.

Stage 2 is known as the “rapid destructive phase” (1–4 years) characterized by loss of previously acquired motor, communication, and social skills, and loss of the ability to use hands purposefully. Instead, patients show abnormal hand movements such as clasping or squeezing, clapping or tapping, and hand-washing movements, irritability characterized by children screaming or crying without reason, mobility and coordination problems including unsteady gait, reduced head circumference (acquired microcephaly), and emergence of breathing issues such as rapid breathing and breath holding.

Stage 3 is known as the “pseudostationary stage” or “plateau phase” that usually begins between ages 2 and 10 and can last for many years. During this phase, there may be improvements in hand usage, alertness, communication skills, walking ability, and irritability (less crying). The main symptoms of stage 3 are seizures, worsening of breathing problems, development of cardiac arrhythmias (abnormal heart rhythm) in some children, teeth grinding, difficulty to gain and maintain weight.

Stage 4 is known as the “late motor deterioration stage”. This usually begins at about age 10 and can last for years or decades. The main symptoms during this phase are reduced mobility, muscle weakness and stiffness, stiffness in joints, scoliosis (abnormal curvature of the spine), and loss of walking ability.<sup>21</sup>

## Rett Syndrome Behavior Questionnaire (RSBQ)

The RSBQ is a validated assessment tool that has been accepted by the FDA as a primary outcome measure for clinical studies on RTT.<sup>22</sup> It is a caregiver-rated scale that assesses neurological and behavioral symptoms in patients between 2 and 47 years of age. The RSBQ consists of 45 items (Table 1), 38 of which are divided into 8 subscales corresponding to the symptomatological domains of RTT such as general mood, breathing problems, hand behaviors, repetitive movements, night-

**Table 1** RSBQ Questionnaire

Domains/ Subscales	Item	Max Score
General mood	<ul style="list-style-type: none"> <li>• Spells of screaming for no apparent reason during the day</li> <li>• Abrupt changes in mood</li> <li>• Certain days/periods where she performs much worse than others</li> <li>• There are times when she appears miserable for no apparent reason</li> <li>• Screams hysterically for long periods of time and cannot be consoled</li> <li>• There are times when she is irritable for no apparent reason</li> <li>• Spells of inconsolable crying for no apparent reason during the day</li> <li>• Vocalizes for no apparent reason</li> </ul>	16
Breathing problems	<ul style="list-style-type: none"> <li>• There are times when breathing is deep and fast (hyperventilation)</li> <li>• There are times when breath is held</li> <li>• Air or saliva is expelled from mouth with force</li> <li>• Swallows air</li> <li>• Abdomen fills with air and sometimes feels hard</li> </ul>	10
Hand behaviors	<ul style="list-style-type: none"> <li>• Does not use hands for purposeful grasping</li> <li>• Hand movements are uniform and monotonous</li> <li>• Has frequent naps during the day</li> <li>• Restricted repertoire of hand movement</li> <li>• Has difficulty in breaking/stopping hand stereotypies</li> <li>• The amount of time spent looking at objects is longer than the time spent holding or manipulating them</li> </ul>	12
Repetitive face movements	<ul style="list-style-type: none"> <li>• Makes repetitive movements involving fingers around tongue</li> <li>• Makes mouth grimaces</li> <li>• Makes repetitive tongue movements</li> <li>• Makes grimacing expressions with face</li> </ul>	8
Body rocking and expressionless face	<ul style="list-style-type: none"> <li>• Expressionless face</li> <li>• Seems to look through people into the distance</li> <li>• Uses eye gaze to convey feelings, needs, and wishes</li> <li>• Rocks self when hands are prevented from moving</li> <li>• Tendency to bring hands together in front of chin or chest</li> <li>• Rocks body repeatedly</li> </ul>	12

(Continued)

**Table 1** (Continued).

Domains/ Subscales	Item	Max Score
Nighttime behaviors	<ul style="list-style-type: none"> <li>• Spells of screaming for no apparent reason during the night</li> <li>• Spells of laughter for no apparent reason during the night</li> <li>• Spells of inconsolable crying for no apparent reason during the night</li> </ul>	6
Fear/Anxiety	<ul style="list-style-type: none"> <li>• Spells of apparent anxiety/fear in unfamiliar situations</li> <li>• Seems frightened when there are sudden changes in own body position</li> <li>• There are times when parts of body are held rigid</li> <li>• Spells of apparent panic</li> </ul>	8
Walking/ Standing	<ul style="list-style-type: none"> <li>• Although can stand independently tends to lean on objects or people</li> <li>• Walks with stiff legs</li> </ul>	4
Uncategorized	<ul style="list-style-type: none"> <li>• Spells of laughter for no apparent reason during the day</li> <li>• Has wounds on hands as result of repetitive hand movements</li> <li>• Shifts gaze with slow horizontal turn of head</li> <li>• Makes repetitive hand movements apart</li> <li>• Appears isolated</li> <li>• Grinds teeth</li> <li>• Vacant “staring” spells</li> </ul>	7

**Note:** Reproduced with permission from Mount RH, Charman T, Hastings RP, Reilly S, Cass H. The Rett Syndrome Behaviour Questionnaire (RSBQ): refining the behavioural phenotype of Rett syndrome. *J Child Psychol Psychiatry*. 2002 Nov;43(8):1099-110. Copyright 2002, John Wiley and Sons.<sup>24</sup>

time behaviors, anxiety, and eye movements. Each item is rated on a Likert scale as 0 (behavior “not true”), 1 (behavior “somewhat or sometimes true”) or 2 (behavior “often true”), with the total score ranging from 0 to 90, indicating the severity and frequency of symptoms. The RSBQ has been correlated with functioning and quality of life.<sup>22,23</sup>


## RTT and Putative Mechanisms of Action

The typical RTT is determined by loss-of-function mutations of the methyl-CpG binding protein 2 (*MECP2*) gene. Other mutations of *MECP2* and/or dysregulations of its expression lead to atypical forms of RTT; moreover, other syndromes similar to RTT can be determined, for instance, by mutations of *CDKL5* or *FOXG1* genes, which are involved with the same biological pathways targeted by *MECP2* mutations.<sup>12,25</sup>

These various disorders have common and distinctive features regarding both the biological etiopathogenesis and clinical manifestations. Key etiopathogenic mechanisms include altered expression of the chloride potassium symporter 5 (*KCC2*), the vesicular glutamate transporter 1 (vGlut1), an orphan glutamate receptor subunit-1 (GluD1), the postsynaptic density protein 95 (PSD-95), and putatively the protein kinase B (Akt).<sup>26</sup> It is not yet clear whether Akt dysregulation is etiopathogenic for RTT or an associated feature, but reduced Akt signaling is reported consistently in RTT models and patients.<sup>15</sup> Moreover, several studies demonstrated beneficial effects of Akt activation obtained through increased signaling at various steps in the pathway CDKL5 – brain derived neurotrophic factor (BDNF) – FOXG1 that eventually regulates synaptic structure.<sup>15</sup> The pathway of GH – IGF-1 is also likely involved, due to its pleiotropic activity on the growth and trophism of all body tissues<sup>27</sup> and it has been extensively targeted by pharmacological trials, together with the NMDA glutamate receptors that mediate neuronal survival and health, especially in the context of energetic and functional stress.<sup>28</sup>

## Therapeutic Approaches to RTT

The syndromic nature, the multi-systemic symptoms and the peculiar natural history make RTT difficult to treat, with limited therapeutic options available. However, despite its rarity, RTT represents one of the rare diseases that has the most advanced development of pharmacological treatments. Pharmacological studies on RTT dramatically increased over the past few decades, with more than 60 clinical trials finished or in progress. Yet, no treatment is available to cure the



**Figure 1** Chemical structures of GPE and Trofinetide.

**Note:** The orange arrow indicates the differences between trofinetide and GPE.

disease. Considering the mechanisms of action of the *MECP2* gene and the broad phenotypic spectrum, pharmacological strategies focused on targets downstream *MECP2* will likely require multiple drugs to effectively treat the full spectrum of RTT symptoms.<sup>29</sup> However, even the improvement of one of the core symptoms of the disease can be considered a valid objective and a useful pharmacological target. In this regard, we can include as main objectives of drugs the dysregulation of breathing, cardiac dysfunction, bone mass loss and fractures, sleep disorders, epilepsy and behavioral alterations. The novel and potentially advantageous approach chosen by the developers of trofinetide and other competing treatments is to try and compensate or revert the degenerative course of RTT, choosing to address directly the pathway of IGF-1 and on NMDA receptors, rather than focusing on single organ defects (Figure 1).<sup>30</sup>

## Focus on IGF-1 and GPE

Considering the dysregulated genes and pathways mentioned above, growth hormone (GH), insulin-like growth factor 1 (IGF-1), and NMDA glutamate receptor targeting treatments were shown to produce benefits on survival and functional parameters, both in vitro and in vivo,<sup>31,32</sup> and ultimately in patients affected by RTT.<sup>33</sup>

## IGF-1 and Neuronal Health

IGF-1 is a growth factor belonging to the GH/somatostatin axis. IGF-1 is produced mainly by energy-managing cells, including hepatocytes and pancreatic cells. The main actions of IGF-1 are exerted through activation of tyrosine kinase receptors in most peripheral tissues, and effects are related to the stimulation of growth and the inter-regulation of growth with energy metabolism. IGF-1 can cross the blood–brain barrier and have profound effects on neuronal growth and trophism, as well. IGF-1 is a driver of, and a factor associated with, neuronal regeneration,<sup>34</sup> survival and function in the adult brain.<sup>35–38</sup> It is recognized that IGF-1 concentrations decline with both ageing and degenerative brain disorders, acquired and congenital.<sup>39,40</sup>

The causative or consequent nature of IGF-1 variations is debated since it was found that specific brain regions connected with stimulus processing and memory (ie: hippocampus, cerebellum, subventricular zone, olfactory bulb) can produce autocrine/paracrine IGF-1 under stimulation by the growth hormone,<sup>41–43</sup> an effect directly correlated with regional potentiation of brain structure and function.

On a pharmacological level, the bulk of circulating IGF-1 is produced by constant hepatic synthesis, amounting to about 10–15 mg daily, which attains a stable plasma concentration around 10–30 nM. IGF-1 naturally has a quite short half-life of 8–16 hours, thus requiring constant release into the blood flow in order to sustain its pharmacological activity. Local brain

production of IGF-1 comes mainly from astrocytes and it appears to be affected by the same compensation and exhaustion dynamics of insulin.<sup>9</sup> Pathologies like Alzheimers' disease seem to be correlated with an initial peak of locally produced IGF-1 that putatively aims to maintain homeostasis in front of IGF-1 receptor hypo-response, eventually leading to functional exhaustion and hypo-production of IGF-1.<sup>44</sup>

Given this mechanism, IGF-1 replacement therapies in various neurodegenerative disorders work like insulin replacement in diabetes, and RTT can be seen as a type of neurodegenerative disorder characterized by deficits on several axes, inside which IGF-1 may provide an alternative source for activation of impaired signaling pathways. Indeed, IGF-1 has shown beneficial effects when given to patients with RTT.<sup>9</sup>

Pathways that are deficient in RTT and can be targeted by IGF-1 may include for instance: cholesterol metabolism since IGF-1 levels are correlated with HDL cholesterol levels in healthy adults,<sup>45</sup> and IGF-1 administration can ameliorate cholesterol metabolism in rodents,<sup>46</sup> oxidative stress, since in cultured cells, IGF-1 administration can prevent the buildup of reactive oxygen species by preserving mitochondrial function;<sup>47</sup> survival and trophism pathways that are interested also by effects from BDNF;<sup>48</sup> IGF-1 can increase mTOR mediated protein translation initiation in cells.<sup>49</sup>

A pilot open-label trial evaluating mecasermin (recombinant human IGF-1 approved for GH deficiency) for RTT patients found that it reduced apneas and behavioral symptoms.<sup>33</sup> A following double-blind placebo-controlled crossover study demonstrated that mecasermin is safe and improves repetitive behaviors and social communication; otherwise, mecasermin treatment did not produce significant improvements.<sup>50</sup> Moreover, electroencephalographic and respiratory parameters worsened and adverse events were reported in 80% of patients, half of whom required hospitalization due to severe adverse events.<sup>50</sup> This unpromising result may hamper further research on mecasermin for RTT.

## GPE: A NMDA Receptor Modulator

Glycine-Proline-Glutamate (GPE) is one of the two degradation products of IGF-1, alongside des(1–3)IGF-1 that are produced by enzymatic cleavage.<sup>1</sup> With respect to previous research, it is likely that IGF-1 actions are mediated both by the whole IGF-1 and by its products des(1-3)-IGF-1 and GPE.<sup>51</sup>

While initial studies attributed the biological activity to the larger portion of IGF-1 it later became clear that also the smaller GPE fragment retained noticeable and distinct effects.<sup>52</sup> While des(1–3)IGF-1 can bind IGF-1 receptors similarly to the parent hormone, the activity proper of GPE was demonstrated to be independent of IGF-1 receptors.<sup>53</sup> It was demonstrated to be mediated by NMDA receptors, which is unsurprising, given that GPE extremities are the known ligands of NMDA receptors glycine and glutamate. Indeed, when competing *in vitro* with the NMDA agonist AP5, only IGF-1 and GPE, but not des(1–3)IGF-1 displayed ligand activity.<sup>54</sup> Moreover, the non-competitive NMDA receptor antagonist MK801 was shown to inhibit the effect of GPE on glial proliferation *in vitro*,<sup>55</sup> and in animal models of NMDA-induced injury, the anti-toxic effect of GPE was similar to that of MK801,<sup>56</sup> overall indicating that NMDA, MK801 and GPE all bind the same receptor. Indeed, GPE also has effects on brain regions that do not respond to IGF-1, like the CA1–2 sub-regions of the hippocampus, rich in NMDA receptors.<sup>57,58</sup>

A role for NMDA receptors in causing RTT symptoms is supported also by the results of a placebo-controlled trial on dextromethorphan for RTT. Dextromethorphan that apparently blocks NMDA receptors caused dose-dependent improvements in seizures, receptive language, and behavioral hyperactivity, but no significant improvement in global clinical severity.<sup>59</sup>

The neuroprotective effects of GPE against ischemia-reperfusion neuronal injury are mediated by a series of mechanisms including inhibition of apoptotic and non-apoptotic death pathways;<sup>60</sup> blood flow increase via NOS induction;<sup>58</sup> better tolerance to glutamate toxicity and increased glutamate clearance by glutamic acid decarboxylase;<sup>58</sup> and maintaining astrocyte integrity.<sup>60</sup> Astrocytes may be key mediators of GPE effects, as human astrocytes bearing RTT mutations can induce phenotypes typical of RTT in co-cultured wild-type neurons, and both IGF-1 and GPE are effective in rescuing some of the caused phenotypes.<sup>61</sup>

Pharmacokinetic considerations on GPE as compared with the whole IGF-1 mainly regard distribution and elimination. It is important to note that most IGF-1 receptors and binding proteins are outside the central nervous system and are involved with metabolic regulations and tissue growth. These peripheral IGF-1 receptors and binding proteins would act as buffers sequestering administered IGF-1, thus lowering its dose-effectiveness inside the central nervous system.

Instead, peripheral tissues lack NMDA receptors, therefore, GPE administration is subject to negligible off-target binding outside the central nervous system. Moreover, the smaller molecular weight of GPE, as compared with IGF-1, would suggest a better distribution to the central nervous system. However, as compared with IGF-1, GPE has lower stability, which is counted in minutes of half-life in the plasma and around half an hour in the brain and cerebrospinal fluid.<sup>62,63</sup>

Although repeated infusion of GPE is required to reach significant concentrations in the brain of rodents, daily intraperitoneal injections of GPE were shown to rescue the characteristic phenotypes of the RTT mouse model, including reduced brain weight, reduced PSD95 expression and spine density in the motor cortex, reduced excitatory postsynaptic currents, and synaptic immaturity.<sup>64</sup> In an effort to find a stable and efficacious analog of GPE, molecules like G-2mPE<sup>65</sup> and cyclic-Gly-Pro<sup>66</sup> and NNZ-2591<sup>67</sup> were developed and tested in vitro and in vivo, all showing different issues and advantages and not reaching a PK-PD profile suitable for a potential human use.<sup>68,69</sup>

## Trofinetide: A Newly Approved GPE-Derivative with Longer Half-Life

### Trofinetide Development and Approval Status

Trofinetide, initially known as NNZ-2566, is a GPE derivative modified with the aim to extend its half-life (Figure 2).


Trofinetide has been extensively studied in vivo for neuroprotection following penetrating brain injury and stroke<sup>70-74</sup> and its mechanism of action was identified in reducing neuroinflammation in these animal models.<sup>73,74</sup>

Trofinetide was then tested in murine models of Fragile X Syndrome, finding a positive effect on behavioral and neuropsychological features, but also on synaptic architecture and signal transduction pathways.<sup>75</sup> This was in line with the effects previously reported for GPE and on the same pathways that are impaired in the context of RTT and related syndromes.

After showing promising efficacy in vivo for Fragile X Syndrome, trofinetide was tested in healthy volunteers to assess its safety and pharmacokinetics as given by oral and injectable administrations, which led to finding a plasma half-life of up to 1.4 hours,<sup>76</sup> which is considerably longer than that of GPE. Trofinetide was then developed for human use with a tentative indication in RTT, until reaching marketing authorization as a first-in-class medication.

The safety, tolerability and potential efficacy of oral trofinetide in RTT were evaluated in two Phase 2 double-blind, placebo-controlled, dose-escalation trials in adolescents and adults (NCT01703533)<sup>77</sup> and then in children and adolescents (NCT02715115).<sup>23</sup>

On 10<sup>th</sup> March 2023, the Food and Drug Administration (FDA) approved oral trofinetide in the US for the treatment of RTT in adult and pediatric patients aged 2 years and older, based on data from two clinical trials performed by Acadia Pharmaceuticals: LAVENDER (NCT04181723)<sup>78</sup> and DAFFODIL (NCT02715115).<sup>79</sup> The US patent for treating RTT with trofinetide has an expiration date in 2032.<sup>1</sup> Alongside granting approval, the FDA considered that the safety database was adequate and the risks associated with trofinetide treatment are acceptable for the indicated population.



**Figure 2** Summary of symptoms and Trofinetide action in Rett Syndrome.

However, the FDA determined that an analysis of spontaneously reported post-marketing adverse events would not be sufficient to prove or disprove neither a serious and unexpected risk of carcinogenicity following exposure to trofinetide nor a serious and unexpected risk of drug–drug interaction. Therefore, the FDA has requested Acadia Pharmaceuticals to conduct two carcinogenicity studies in mice and rats, an *in vitro* drug interaction study focused on CYP2B6, an *in vivo* drug interaction study on OATP1B1 and OATP1B3 and a pharmacokinetics clinical trial to evaluate the effect of moderate renal impairment on the exposure to trofinetide.<sup>1</sup>

### Clinical Studies and Core Efficacy Endpoints

A recent meta-analysis<sup>14</sup> included the three preliminary RCTs conducted by Acadia Pharmaceuticals on trofinetide in RTT patients. Glaze 2017<sup>77</sup> examined adolescents and adults (15–25 years old) and used two fixed doses (35, 70 mg/Kg), measuring efficacy outcomes at day 54. Glaze 2019<sup>23</sup> examined children and adolescents (5–15 years old) and used three fixed doses (50, 100, 200 mg/Kg) measuring efficacy outcomes at day 66. Neul 2023<sup>78</sup> examined patients between the ages of 5 and 20 and used a dose ranging between 200 and 500 mg/Kg per day, measuring efficacy outcomes at 12 weeks. Meta-analytical results indicate that trofinetide improved CGI-I scores and RSBQ scores; particularly improved subscores included impaired communication, compulsive behaviors, and emotional disorders, which improved the lives of patients and their families.<sup>14</sup>

The primary evidence of effectiveness for application of trofinetide in RTT is based on the analysis of clinical efficacy endpoints from the 3 month, multicenter, double-blind, placebo-controlled Phase 3 LAVENDER trial from Acadia Pharmaceuticals.<sup>78</sup> A total of 187 subjects were randomized 1:1 to receive either trofinetide in a weight-based dosing (N = 93) or placebo (N = 94) twice daily. Subjects were required to be between 5 and 20 years of age and have a genetically confirmed diagnosis of RTT. The first co-primary efficacy endpoint was the change from baseline to week 12 on the Rett Syndrome Behavior Questionnaire (RSBQ) total score, as compared to placebo. Regarding this endpoint, the FDA recommended Acadia Pharmaceuticals to include a clinical co-primary endpoint to support their results, because RSBQ scores can be significantly different even when the changes are not clinically meaningful. For this reason, the Clinician's Global Impression of Improvement (CGI-I) was included as a co-primary endpoint. CGI-I is a clinician-completed assessment of how much the individual's illness has improved/worsened relative to a baseline state, scored using a standardized 7-point grid. LAVENDER also evaluated the Communication and Symbolic Behavior Scales Developmental Profile-infant-toddler social composite score (CSBS-DP-IT-SCS) as a secondary endpoint. It is a screening assessment to detect potential communication defects in children, which consists of 24 questions divided into 7 subscales referring to the communicative sphere: emotion and eye gaze, communication, gestures, sounds, words, understanding and object use. After treatment, there was a significant difference between trofinetide-treated and placebo-treated patients favoring trofinetide, on both the co-primary endpoints CGI-I ( $p = 0.003$ ) and RSBQ ( $p = 0.018$ ). For the latter, significant effects were observed on the General Mood subscale ( $p = 0.007$ ) and the Repetitive Face Movement subscale ( $p = 0.047$ ). Nominal improvements were also observed in other areas, such as Breathing problems, Walking/ Standing, Nighttime behaviors and Seizures, but these were not significant. Trofinetide appears to have no effect on the other domains evaluated by the RSBQ.<sup>23</sup>

These results were also supported by a significant improvement in the secondary endpoint CSBS-DP-IT-SCS ( $p = 0.006$ ).<sup>78</sup>

Acadia Pharmaceuticals also conducted an open-label study (DAFFODIL) to evaluate pharmacokinetics and safety of trofinetide in RTT patients between 2 and 4 years of age. The mean CGI-I score at week 12 corresponded to “much improved” with respect to the baseline. The interim PK analysis based on data from 13 children treated with trofinetide for 12 weeks demonstrated a PK exposure to trofinetide and safety profiles similar to those of pediatric patients  $\geq 5$  years of age and of adults, leading to the approval of trofinetide for age 2 and above.<sup>79</sup>

Acadia Pharmaceuticals then conducted two open-label extension trials on patients who concluded LAVENDER, named LILAC and LILAC-2. LILAC demonstrated continued benefits on both RSBQ and CGI-I from open-label treatment with trofinetide, for up to 40 additional weeks after the end of LAVENDER<sup>11</sup> and LILAC-2 for other 32 additional weeks after the end of LILAC.<sup>11</sup> Overall, treatment for 84 weeks with trofinetide demonstrated continued

improvement on both co-primary endpoints, with the same adverse effects of short-term treatment. However, it is worth noting that, at 84 weeks, 59.4% CGI-I and 88% RSBQ evaluations had missing values.

### Adverse Events

The risks associated with trofinetide use did not preclude approval. The most common adverse events were of gastrointestinal nature, including serious diarrhea and vomiting. The FDA recommended that warnings for the clinical significance of diarrhea and weight loss were to be added to the product label.

Based on the three meta-analyzed preliminary studies,<sup>14</sup> diarrhea ( $p = 0.06$ ) and vomiting ( $p = 0.002$ ) were the only adverse events associated with trofinetide treatment. Considering the LAVENDER trial, the most common adverse reactions reported in at least 5% of trofinetide-treated subjects were diarrhea (82%), vomiting (29%), fever (9%), seizure (9%), anxiety (8%), decreased appetite (8%), fatigue (8%), and nasopharyngitis (5%). Diarrhea occurred in nearly 85% of subjects on long-term treatment with trofinetide, while 12% of subjects experienced a loss of greater than 7% of body weight. This is a clinically meaningful risk in a pediatric population.<sup>78</sup>

### Other Therapeutic Strategies: Restoration of MECP2 Expression

These approaches include gene therapy and reactivation of the inactive X chromosome. They have an unclear window of opportunity before which they have to be carried out, in order to produce a significant correction in the degenerative pathway typical of RTT progression. Restoring *MECP2* expression in late life may improve some functional phenotypes but may not improve most structural changes to the brain and/or organ architecture that have already occurred. Gene therapies for RTT are reviewed elsewhere.<sup>4</sup>

### Other Therapeutic Strategies: Restoration of Molecular Pathways Downstream MECP2-Induced Genes in Peripheral Tissues

There is no ascertained common mechanism of action underlying the phenotypes described below that target heart and bone. However, it is likely that a pleiotropic effect of loss of signaling downstream the GH receptor is involved.

#### Cardiac Dysfunction

The main cardiac dysfunction present in people with RTT is the prolonged QT interval, ie the timing between ventricle depolarization and repolarization is delayed in RTT patients as compared to age-matched healthy people. The state-of-the-art treatments for QT prolongation in the general population are beta-blockers, but they were not effective in preventing cardiac alterations in *MECP2* null mice.<sup>4</sup> This suggests that standard treatment is insufficient in the context of cardiac dysfunctions descending from RTT. In view of the lack of efficacy of available treatments, current recommendations are to avoid QT-elongating drugs in people with RTT that have developed prolonged QT intervals, in order to help avoid cardiac dysfunction.<sup>5</sup>

#### Bone Mineral Loss and Fractures

Bone mineral loss and increased likelihood of fractures is an early concern in the RTT population, as opposed to typically developing women, who encounter increased risks of fractures only after menopause. Prophylactic treatment with vitamin D needs to be considered in people with RTT.<sup>20</sup> Another treatment proposed to RTT patients with mineral loss is bisphosphonates, which inhibit bone resorption and therefore prevent the reduction of mineralization in bone.<sup>20</sup> However, in clinical practice bisphosphonates are rarely used due to their adverse effects, such as abdominal pain, inducing osteonecrosis and increasing the risk of non-vertebral fractures with long-term use. This characteristic must be considered in the evaluation of drug–disease interactions considering, for example, that some antiepileptic drugs alter bone metabolism, which increases the risk of fractures. Carbamazepine, phenytoin, and valproic acid also reduce blood calcium levels, further increasing the risk of fractures in RTT patients with epilepsy.<sup>80,81</sup>

## Other Therapeutic Strategies: Restoration of Molecular Pathways Downstream MECP2-Induced Genes in the Brain

Drugs targeting molecular pathways downstream genes induced by *MECP2* aim to restore mainly the excitatory–inhibitory synaptic balance in specific neural circuits. Considered pathways and pharmacological targets include neurotransmitter and neuromodulator systems such as the noradrenergic, serotonergic, glutamatergic, GABAergic, and cholinergic signaling; GH, BDNF and IGF-1 signaling; energy metabolism including the cholesterol biosynthesis pathway; mitochondrial function and oxidative stress; Akt/mTOR mediated regulation of protein translation initiation.<sup>6–8,15,29,82</sup> Recently, a role of the gut microbiota was suggested in the progression and symptoms of RTT, providing an additional pharmacological target.<sup>10</sup>

Overall, the pathways involving GH, IGF-1 and NMDA receptors have been implicated with altered neuronal functions in RTT and related syndromes.

### Breathing

Dysregulation of breathing is a core feature of RTT in up to 93% of patients, it significantly impacts quality of life, and it is thought to contribute to early mortality in some patients.<sup>83</sup> Common issues are hyperventilation, apnea, breath holding, and air swallowing that occurs when waking up. At the origin of breathing dysregulation, there are abnormalities in brain stem connections, with impaired development of the neuronal circuitry responsible for setting the respiratory rhythm, which is secondary to deletion of *MECP2* in GABAergic interneurons<sup>83</sup> and was shown to respond to BDNF administration in *MECP2* mice.<sup>13,84</sup> Several molecules appeared as promising in preclinical and clinical studies to reduce breathing abnormalities and sarizotan (a 5-HT<sub>1a</sub> agonist and a dopamine D<sub>2</sub>-like agonist/partial agonist) reduced apneas in mouse models of RTT, but a study on efficacy, safety, and tolerability of sarizotan in RTT with respiratory symptoms was terminated early during the double-blind period because it did not demonstrate evidence of efficacy on the reduction of respiratory abnormalities, nor on the caregiver-rated impression of change.<sup>13,17,84,85</sup> The noradrenaline uptake inhibitor desipramine<sup>86</sup> was also tested, showing no indication of efficacy.<sup>87</sup> A study to assess the safety and efficacy of fingolimod, which should increase levels of brain-derived neurotrophic factor (BDNF), also had negative results because changes in BDNF levels in the CSF and serum were not found and clinical outcomes were not met.<sup>88</sup>

### Epilepsy

Epilepsy regards approximately 60–80% of RTT individuals, with heterogeneous semiology (focal or generalized), and it appears to be an age-related event, as patients 6 to 12 years of age tend to be at highest risk for seizure development; it is debilitating and often drug-resistant.<sup>89</sup> *MECP2* mutant mice have an imbalance of excitation-to-inhibition in brain regions known to contribute to epilepsy and seizure disorders.<sup>89–91</sup> Epilepsy is considered a major issue in RTT and is managed like in other rare diseases and developmental and epileptic encephalopathies.<sup>92</sup> Tolerability is the main issue with anti-seizure medications, which are associated with many different adverse drug effects. For instance, the use of lamotrigine may cause serious rash or levetiracetam may cause behavioral disturbance,<sup>93–95</sup> therefore, limiting their use to RTT patients with an active seizure disorder would minimize the number of patients with adverse effects, although the neurological best practice would be to prevent seizures with prophylactic treatment.<sup>94,95</sup> Lastly, since RTT patients may be taking many medications, determining drug–drug interactions upfront is also mandatory.<sup>81</sup> For instance, carbamazepine is a known CYP450 inducer and may cause a drug–drug interaction by increasing the clearance and decreasing the blood concentration of other drugs that the patient may be taking.<sup>81,96</sup> Secondly, efficacy is of concern in RTT patients with epilepsy. Sodium valproate has the highest number of positive recommendations for different kinds of epilepsy.<sup>81,94–96</sup> However, no recommendations of a specific treatment were made for RTT-associated epilepsy. As in the general population, 30% of patients with RTT also have drug-resistant epilepsy.<sup>81</sup> In these cases, ketogenic diet or vagus nerve stimulation are considered, alongside innovative drug therapies below.<sup>96,97</sup>

Glatiramer acetate, which is labeled to treat multiple sclerosis by acting on immune cell activation, leads to delayed onset of RTT symptoms and increased BDNF levels.<sup>98</sup> A phase 2 open-label trial to test glatiramer in patients with RTT demonstrated that epileptiform discharges were decreased in four patients, but the trial was terminated due to the development of life-threatening post-injection responses in 3 out of 14 patients.<sup>98,99</sup>

Cannabidiol is an inhibitor of fatty acid aryl hydroxylase, which leads to increased concentrations of endogenous cannabinoids, and also a GPR55 antagonist, overall dampening neuronal excitability.<sup>100</sup> Cannabidiol is efficacious for various syndromic drug-resistant epilepsies, such as Lennox–Gastaut and Dravet, Aicardi, Dup15q, and Doose syndromes, SYNGAP1 encephalopathy, epilepsy with myoclonic absences, and atypical RTT as CDKL5-related encephalopathy, with some reported benefits.<sup>100–102</sup> A long-term safety study of cannabidiol oral solution showed seizure frequency was reduced. However, populations suffered from serious adverse side effects such as diarrhea, vomiting, fatigue, pyrexia and somnolence. The study was terminated early due to the COVID-19 pandemic and recruitment challenges.<sup>103</sup> The available data on cannabinoids used for RTT patients show possible promising effects of cannabidivarin (CBDV), the propyl analog of cannabidiol (CBD). CBDV and CBV are safe and well tolerated.<sup>17,104</sup> They reduce the incidence of seizures, agitation and/or anxiety attacks.<sup>100</sup> These drugs do not alter the safety and effectiveness of diazepam or clobazam, which supports concomitant use in selected patients.<sup>104–108</sup>

Ganaxolone is a synthetic analog of the neuroactive steroid allopregnanolone that acts as a positive allosteric modulator of synaptic and extrasynaptic GABA<sub>A</sub> receptors, thus having antiseizure properties in preclinical and clinical studies.<sup>109</sup> Ganaxolone is now approved for epilepsy associated with the RTT-like CDKL5 syndrome, in patients aged two years and older, which is of immediate interest also for patients with RTT, although there is yet no specific indication of use.<sup>110,111</sup> Ganaxolone might also improve other aspects of children's wellbeing, secondary to a better control of seizures.

### Sleep Disorders

An open-label trial of L-carnitine, an amino acid derivative of methionine and lysine that is required for energy metabolism, showed improved sleep maintenance in a subset of RTT patients.<sup>112</sup>

Additional therapeutic alternatives, not specifically tested for RTT, include the serotonin receptor modulator trazodone,<sup>81</sup> the sedating agent chloral hydrate,<sup>113,114</sup> and melatonin,<sup>115</sup> whereas benzodiazepines are not recommended in RTT, given the respiratory issues of some of these patients.<sup>81,116</sup>

### Behavior Disorders

Disorders of behavior in RTT patients include anxiety, fearful behaviors, various mood disturbances, aggression and self-harm.<sup>117</sup> The scientific literature is poor on this topic, and treatment usually follows what is indicated for disorders associated with autism spectrum disorders and other genetic syndromes, even if RTT presents with peculiar and distinct characteristics, especially with regard to the risk of adverse effects.<sup>118,119</sup> The most successful management of anxiety and mood dysregulations is with selective serotonin modulators, like mirtazapine that appear to protect against disease progression and improve motor, sensory and behavioral symptoms.<sup>120</sup> In support of this clinical evidence, it was demonstrated that serotonin levels are reduced in the hippocampus of *MECP2* null mice, as compared to those of wild-type mice.<sup>5,120</sup> Aggression and self-harm, irritability and impairments in socialization are anecdotally treated using risperidone and other antipsychotic sedating drugs, at the cost of several adverse metabolic and cardiac effects.<sup>120</sup> Moreover, the sigma-1 receptor antagonist blarcamesine appeared to improve RSBQ and CGI-I in a 7-week double-blind, randomized, placebo-controlled clinical trial (NCT03758924) sponsored by Anavex Life Science Corp. In this study, low-dose blarcamesine produced a clinically significant effect in reducing symptoms of RTT ( $p < 0.007$ ) and led to a significant slowing of disease progression ( $p < 0.007$ ).<sup>121</sup>

### Shortcomings of Other Drug Therapies

With the notable exception of gene therapy (yet in early clinical research), the pharmacological approaches described above and summarized (Table 2) are mostly symptomatic, they are used in clinical practice to try and improve the quality of life of families and patients, but they have no impact on the degenerative natural history of the disease. The result is

**Table 2** Treatments for RTT Under Clinical Development with Published Results

Authors (year)	Active Substance	Study Design	RTT Subjects	Results
Djukic A et al (2016) <sup>4</sup>	Glatiramer Acetate	Prospective open label trial – Phase II	10	Gait velocity, memory, and the breath holding index improved significantly. Epileptiform discharges decreased. There was a trend towards improved quality of life, which did not reach statistical significance.
Nissenkorn A et al (2016) <sup>5</sup>	Glatiramer Acetate	Open label trial – phase II	14	4 patients developed an exaggerated immediate postinjection response, which was experienced as life threatening in 3 patients, necessitating arrest of the trial.
Desnous B et al (2023) <sup>6</sup>	Cannabidiol - Clobazam	Longitudinal observational study	10	CBD is well tolerated and may increase the efficacy of clobazam alone. It reduced the incidence of seizures, agitation and/or anxiety attacks. In addition, improvement in spasticity was reported in 4 patients.
Peters JM et al (2023) <sup>7</sup>	Cannabidiol - Diazepam	Long-term safety study – Phase III	21 out of 163	CBD does not alter the safety and effectiveness of diazepam nasal spray and supports concomitant use in appropriate patients.
Tarquino D et al (2023) <sup>8</sup>	Diazepam	Long-term safety study – phase III	16 out of 64	In patients with developmental epileptic encephalopathies, diazepam nasal spray demonstrated a consistent safety profile.
Smith-Hicks et al (2017) <sup>9</sup>	Dextro-methorphan	Prospective randomized open-label trial	38	Dextromethorphan is safe. Statistically significant dose-dependent improvements were seen in clinical seizures, receptive language, and behavioral hyperactivity. There was no significant improvement in global clinical severity.
Flores Gutiérrez J et al (2020) <sup>10</sup>	Mirtazapine	Retrospective study	80	Mirtazapine is well tolerated and it protects from disease progression and improves motor, sensory, and behavioral symptoms.
Hurley EN et al (2022) <sup>11</sup>	Cannabidivarin	Clinical trial – phase I	5	CBDV is safe and well tolerated. The main result was a reduction in seizure frequency. No significant changes were observed in the EEG or in symptoms unrelated to RTT.
Khwaja OS et al (2014) <sup>12</sup>	Mecasermin	Unblinded study - phase I	9 out of 12	Mecasermin is safe and well tolerated in girls with RTT and, as demonstrated in preclinical studies, improves certain breathing and behavioral abnormalities.
Ette et al (2023) <sup>13</sup>	Blarcomesine	Double-blind, randomized study – phase II	25	Low-dose blarcomesine produced a clinically significant effect in reducing symptoms of RTT and it led to a significant slowdown in disease progression.
Mancini et al (2017) <sup>14</sup>	Desipramine	Double-blind, randomized study – phase II	34	Not significant difference between the effects of placebo on the apnea-hypopnea index at 6 months. Neither did we show significant effects for other respiratory parameters.
Ellaway et al (2021) <sup>15</sup>	L-carnitine	Open label trial	21	There was a significant improvement in outcome for the cases compared with the controls in sleep efficiency, energy level, communication skills and expressive speech.

**Table 3** Other Treatments for RTT Under Clinical Development Without Published Results

Drug	Intended Target	Mechanism	Phase
AMO-04 Tianeptine	Glutamate receptors	Modulator	2
Ketamine	GRIN receptors	Modulator	2
VYNT-0126	Not disclosed	Not disclosed	2
NGN-401	<i>MECP2</i> Gene on inactive X chromosome	EXACT transgene regulation technology	I–2
TSHA-102	<i>MECP2</i> Gene	scAAV9 vector carrying recombinant <i>MECP2</i>	I–2
ACP-2591	IGF-1 receptors	Modulator	I
F-15599 NLX-101	5-HT1A cortical receptors	Highly selective agonist	I
Fingolimod	Sphingosine 1-phosphate receptor 1	Modulator	I
Vatiquinone	15-Lipoxygenase	Selective inhibitor	–
GXV001	Not disclosed	Not disclosed	–
Donepezil	AChE	Inhibitor	T
Lovastatin	HMGCR	Inhibitor	T
Sarizotan	5-HT1a and DRD2	Agonist/partial agonist	T

**Note:** – : not clear, T : terminated.

a mostly off-label and heterogeneous use of pharmacological mono- and poly-therapy, which leads to high rates of failure and a severely increased risk of adverse effects and drug–drug and drug–disease interactions. Other treatments are undergoing development, but clinical results are not yet available (Table 3).

## Discussion

### Future Perspectives for Trofinetide

The currently published clinical studies on trofinetide show limitations that could be overcome through future research and with some adjustments: long-term follow-up studies should investigate the real-life impact and sustainability of trofinetide treatment, especially regarding its formulation and dosing, which leads to gastrointestinal adverse events. The duration and economic impact of treatment with trofinetide should also be investigated, possibly with HTA assessments. It would also be necessary to identify the optimal and most tolerable dosing of trofinetide for different types of RTT and related syndromes, as well as different RTT stages and presentations. This is mandatory in order to tailor drug administration to each patient and predict drug response, an important issue with rare diseases. Biomarkers should be determined to pinpoint what physiological, neurocognitive or behavioral mechanisms trofinetide tackles. These aspects would allow for a more patient-specific treatment.

### Current Place in Therapy of Trofinetide

Trofinetide, a first-in-class medication approved by the FDA for the treatment of RTT in people aged 2 years and above, is demonstrated to be relatively tolerable, except for moderate/severe gastrointestinal symptoms. Trofinetide is effective on some, but not all, symptoms of RTT, as shown by the significant improvement only on some sub-scales of the RSBQ, as discussed above (reviewed in a recent meta-analysis).<sup>122</sup> This selectivity of effect likely depends on the absence of binding to IGF-1 receptors. Regarding the methods of administration and management of the drug on the practical side, trofinetide is supplied as 200 mg/mL oral solution in a 450 mL bottle, costing around ten thousand USD and containing enough medication for ten, down to three days, depending on patients' weight. These aspects may constitute a barrier to use, given the chronic nature of RTT. In clinical practice, trofinetide may be suitable as a stand-alone or combined

treatment, depending on each patient's needs. In order to manage these further issues, additional pharmacokinetic and drug–drug interaction studies are under way, as requested by the FDA, to ensure it can be used also in combination with other treatments in complex clinical settings.

It is yet unclear whether trofinetide may be a symptomatic or curative treatment. In theory, these may both be possible, depending on the age of treated patients: ie trofinetide may reduce the progression of RTT for patients below a certain age threshold, acting through the correction of NMDA-mediated neuronal network structuring; trofinetide may only ameliorate RTT symptoms above a certain threshold age, allowing neuronal networks to function better.

Further phase 3 or 4 efficacy studies will also be needed to re-determine the place in therapy of trofinetide when competitors will enter the market and also with respect to other IGF-1 or NMDA targeting therapies. For instance, mecasermin, the recombinant human IGF-1 analog, may be further investigated as a potential alternative to trofinetide given its non-novel status and possibly much lower costs; in this perspective, however, the preliminary unsatisfying results obtained by mecasermin would require scrutiny and proper re-assessment in comparative clinical trials. In view of the significant effect of trofinetide. Overall, trofinetide is a crucial new tool for the pharmacological treatment of RTT and related conditions. Its use is likely to be considerably improved during the first years of marketing.

## Funding

The Authors acknowledge funding from the Italian Ministry of Health (Ministero della Salute, Ricerca Corrente 2024), which had no role in the conception or writing of this work.

## Disclosure

The authors report no conflicts of interest in this work. Simone Pisano and Maria Nobile are joint last authors.

## References

1. Keam SJ. Trofinetide: first Approval. *Drugs*. 2023;83(9):819–824. doi:10.1007/s40265-023-01883-8
2. Hudu SA, Elmigdad F, Qtaitat AA, et al. Trofinetide for rett syndrome: highlights on the development and related inventions of the first USFDA-approved treatment for rare pediatric unmet medical need. *J Clin Med*. 2023;12(15):5114. doi:10.3390/jcm12155114
3. Epistemic AI website. Available from: <https://epistemic.ai/>. Accessed October 14, 2024.
4. Palmieri M, Pozzer D, Landsberger N. Advanced genetic therapies for the treatment of Rett syndrome: state of the art and future perspectives. *Front Neurosci*. 2023;17:1172805. doi:10.3389/fnins.2023.1172805
5. McCauley MD, Wang T, Mike E, et al. Pathogenesis of lethal cardiac arrhythmias in Mecp2 mutant mice: implication for therapy in Rett syndrome. *Sci Transl Med*. 2011;3(113):113ra125. doi:10.1126/scitranslmed.3002982
6. Pack AM, Morrell MJ. Adverse effects of antiepileptic drugs on bone structure: epidemiology, mechanisms and therapeutic implications. *CNS Drugs*. 2001;15(8):633–642. doi:10.2165/00023210-200115080-00006
7. Itoh M, Ide S, Takashima S, et al. Methyl CpG-binding protein 2 (a mutation of which causes Rett syndrome) directly regulates insulin-like growth factor binding protein 3 in mouse and human brains. *J Neuropathol Exp Neurol*. 2007;66(2):117–123. doi:10.1097/nen.0b013e3180302078
8. Buchovecky CM, Hill MG, Borkey JM, Kyle SM, Justice MJ. Evaluation of rett syndrome symptom improvement by metabolic modulators in mecp2-mutant mice. *Curr Protoc Mouse Biol*. 2013;3(4):187–204. doi:10.1002/9780470942390.mo130157
9. Pini G, Scusa MF, Congiu L, et al. IGF1 as a potential treatment for rett syndrome: safety assessment in six rett patients. *Autism Res Treat*. 2012;2012:679801. doi:10.1155/2012/679801
10. Strati F, Cavalieri D, Albanese D, et al. Altered gut microbiota in Rett syndrome. *Microbiome*. 2016;4(1):41. doi:10.1186/s40168-016-0185-y
11. Percy AK, Neul JL, Benke TA, et al. Trofinetide for the treatment of Rett syndrome: results from the open-label extension LILAC study. *Med*. 2024;2024:1.
12. Kyle SM, Vashi N, Justice MJ. Rett syndrome: a neurological disorder with metabolic components. *Open Biol*. 2018;8(2):170216. doi:10.1098/rsob.170216
13. Schmid DA, Yang T, Ogier M, et al. A TrkB small molecule partial agonist rescues TrkB phosphorylation deficits and improves respiratory function in a mouse model of Rett syndrome. *J Neurosci*. 2012;32(5):1803–1810. doi:10.1523/JNEUROSCI.0865-11.2012
14. Abo Zeid M, Elrosasy A, Mohamed RG, et al. A meta-analysis of the efficacy and safety of trofinetide in patients with rett syndrome. *Neurol Sci*. 2024;45:4767–4778. doi:10.1007/s10072-024-07584-8
15. Ricciardi S, Boggio EM, Grosso S, et al. Reduced AKT/mTOR signaling and protein synthesis dysregulation in a Rett syndrome animal model. *Hum Mol Genet*. 2011;20(6):1182–1196. doi:10.1093/hmg/ddq563
16. Liyanage VR, Rastegar M. Rett syndrome and MeCP2. *Neuromolecular Med*. 2014;16(2):231–264. PMID: 24615633; PMCID: PMC5798978. doi:10.1007/s12017-014-8295-9
17. Lopes AG, Loganathan SK, Caliperumal J. Rett Syndrome and the Role of MECP2: signaling to clinical trials. *Brain Sci*. 2024;14(2):120. doi:10.3390/brainsci14020120
18. Kim JA, Kwon WK, Kim JW, Jang JH. Variation spectrum of MECP2 in Korean patients with Rett and Rett-like syndrome: a literature review and reevaluation of variants based on the ClinGen guideline. *J Hum Genet*. 2022;67(10):601–606. doi:10.1038/s10038-022-01044-x

19. Neul JL, Kaufmann WE, Glaze DG, et al. Rett syndrome: revised diagnostic criteria and nomenclature. *Ann Neurol*. 2010;68(6):944–950. doi:10.1002/ana.22124
20. Associazione Italiana Rett. Available from: <https://www.airett.it/sindrome-rett/>. Accessed October 14, 2024.
21. Bernardo P, Cobb S, Coppola A, et al. Neurophysiological Signatures of Motor Impairment in Patients with Rett Syndrome. *Ann Neurol*. 2020;87(5):763–773. doi:10.1002/ana.25712
22. Percy AK, Neul JL, Benke TA, Marsh ED, Glaze DG. A review of the Rett syndrome behaviour questionnaire and its utilization in the assessment of symptoms associated with Rett syndrome. *Front Pediatr*. 2023;11:1229553. doi:10.3389/fped.2023.1229553
23. Glaze DG, Neul JL, Kaufmann WE, et al. Double-blind, randomized, placebo-controlled study of trofinetide in pediatric Rett syndrome. *Neurology*. 2019;92(16):e1912–e1925. doi:10.1212/WNL.0000000000007316
24. Mount R H, Charman T, Hastings R P, Reilly S and Cass H. (2002). The Rett Syndrome Behaviour Questionnaire (RSBQ): refining the behavioural phenotype of Rett syndrome. *Child Psychology Psychiatry*, 43(8), 1099–1110. doi:10.1111/1469-7610.00236
25. Pejhan S, Rastegar M. Role of DNA Methyl-CpG-Binding Protein MeCP2 in rett syndrome pathobiology and mechanism of disease. *Biomolecules*. 2021;11(1):75. doi:10.3390/biom11010075
26. D’Mello SR. Rett and Rett-related disorders: common mechanisms for shared symptoms? *Exp Biol Med (Maywood)*. 2023;248(22):2095–2108. doi:10.1177/15353702231209419
27. Blum WF, Alherbish A, Alsagheir A, et al. The growth hormone-insulin-like growth factor-I axis in the diagnosis and treatment of growth disorders. *Endocr Connect*. 2018;7(6):R212–R222. doi:10.1530/EC-18-0099
28. Wang Y, Wang W, Li D, et al. IGF-1 alleviates NMDA-induced excitotoxicity in cultured hippocampal neurons against autophagy via the NR2B/PI3K-AKT-mTOR pathway. *J Cell Physiol*. 2014;229(11):1618–1629. doi:10.1002/jcp.24607
29. Katz DM, Bird A, Coenraads M, et al. Rett Syndrome: crossing the Threshold to Clinical Translation. *Trends Neurosci*. 2016;39(2):100–113. doi:10.1016/j.tins.2015.12.008
30. Parent H, Ferranti A, Niswender C. Trofinetide: a pioneering treatment for Rett syndrome. *Trends Pharmacol Sci*. 2023;44(10):740–741. doi:10.1016/j.tips.2023.06.008
31. Castro J, Garcia RI, Kwok S, et al. Functional recovery with recombinant human IGF1 treatment in a mouse model of Rett Syndrome. *Proc Natl Acad Sci U S A*. 2014;111(27):9941–9946. doi:10.1073/pnas.1311685111
32. Della Sala G, Putignano E, Chelini G, et al. Dendritic spine instability in a mouse model of CDKL5 disorder is rescued by insulin-like growth factor 1. *Biol Psychiatry*. 2016;80:302–311. doi:10.1016/j.biopsych.2015.08.028
33. Khwaja OS, Ho E, Barnes KV, et al. Safety, pharmacokinetics, and preliminary assessment of efficacy of mecaseprin (recombinant human IGF-1) for the treatment of Rett syndrome. *Proc Natl Acad Sci U S A*. 2014;111(12):4596–4601. doi:10.1073/pnas.1311141111
34. Popken GJ, Hodge RD, Ye P, et al. In vivo effects of insulin-like growth factor-I (IGF-I) on prenatal and early postnatal development of the central nervous system. *Eur J Neurosci*. 2004;19(8):2056–2068. doi:10.1111/j.0953-816X.2004.03320.x
35. Lichtenwalner RJ, Forbes ME, Bennett SA, Lynch CD, Sonntag WE, Riddle DR. Intracerebroventricular infusion of insulin-like growth factor-I ameliorates the age-related decline in hippocampal neurogenesis. *Neuroscience*. 2001;107(4):603–613. doi:10.1016/S0306-4522(01)00378-5
36. Trejo JL, Piriz J, Llorens-Martin MV, et al. Central actions of liver-derived insulin-like growth factor I underlying its pro-cognitive effects. *Mol Psychiatry*. 2007;12(12):1118–1128. doi:10.1038/sj.mp.4002076
37. Markowska AL, Mooney M, Sonntag WE. Insulin-like growth factor-1 ameliorates age-related behavioral deficits. *Neuroscience*. 1998;87(3):559–569. doi:10.1016/S0306-4522(98)00143-2
38. Sonntag WE, Lynch C, Thornton P, Khan A, Bennett S, Ingram R. The effects of growth hormone and IGF-1 deficiency on cerebrovascular and brain ageing. *J Anat*. 2000;197(Pt 4):575–585. doi:10.1046/j.1469-7580.2000.19740575.x
39. Aleman A, Verhaar HJ, De Haan EH, et al. Insulin-like growth factor-I and cognitive function in healthy older men. *J Clin Endocrinol Metab*. 1999;84(2):471–475. doi:10.1210/jcem.84.2.5455
40. Trejo JL, Carro E, Garcia-Galloway E, Torres-Aleman I. Role of insulin-like growth factor I signaling in neurodegenerative diseases. *J Mol Med (Berl)*. 2004;82(3):156–162. doi:10.1007/s00109-003-0499-7
41. Bondy CA, Werner H, Roberts CT, LeRoith D. Cellular pattern of insulin-like growth factor-I (IGF-I) and type I IGF receptor gene expression in early organogenesis: comparison with IGF-II gene expression. *Mol Endocrinol*. 1990;4(9):1386–1398. doi:10.1210/mend-4-9-1386
42. Breese CR, D’Costa A, Rollins YD, et al. Expression of insulin-like growth factor-1 (IGF-1) and IGF-binding protein 2 (IGF-BP2) in the hippocampus following cytotoxic lesion of the dentate gyrus. *J Comp Neurol*. 1996;369(3):388–404. doi:10.1002/(SICI)1096-9861(19960603)369:3<388::AID-CNE5>3.0.CO;2-1
43. Leung KC, Doyle N, Ballesteros M, Waters MJ, Ho KK. Insulin regulation of human hepatic growth hormone receptors: divergent effects on biosynthesis and surface translocation. *J Clin Endocrinol Metab*. 2000;85:4712–4720.
44. Talbot K, Wang HY, Kazi H, et al. Demonstrated brain insulin resistance in Alzheimer’s disease patients is associated with IGF-1 resistance, IRS-1 dysregulation, and cognitive decline. *J Clin Invest*. 2012;122(4):1316–1338. doi:10.1172/JCI59903
45. Succurro E, Arturi F, Grembiale A, et al. Positive association between plasma IGF1 and high-density lipoprotein cholesterol levels in adult nondiabetic subjects. *Eur J Endocrinol*. 2010;163(1):75–80. PMID: 20356932. doi:10.1530/EJE-10-0113
46. Fukunaga K, Imachi H, Lyu J, et al. IGF1 suppresses cholesterol accumulation in the liver of growth hormone-deficient mice via the activation of ABCA1. *Am J Physiol Endocrinol Metab*. 2018;315(6):E1232–E1241. PMID: 30130150. doi:10.1152/ajpendo.00134.2018
47. Pi Y, Goldenthal MJ, Marin-Garcia J. Mitochondrial involvement in IGF-1 induced protection of cardiomyocytes against hypoxia/reoxygenation injury. *Mol Cell Biochem*. 2007;301(1–2):181–189. PMID: 17264981. doi:10.1007/s11010-007-9410-0
48. Reim D, Schmeisser MJ. Neurotrophic Factors in Mouse Models of Autism Spectrum Disorder: focus on BDNF and IGF-1. *Adv Anat Embryol Cell Biol*. 2017;224:121–134. PMID: 28551754. doi:10.1007/978-3-319-52498-6\_7
49. James MJ, Zomerdijk JC. Phosphatidylinositol 3-kinase and mTOR signaling pathways regulate RNA polymerase I transcription in response to IGF-1 and nutrients. *J Biol Chem*. 2004;279(10):8911–8918. PMID: 14688273. doi:10.1074/jbc.M307735200
50. O’Leary HM, Kaufmann WE, Barnes KV, et al. Placebo-controlled crossover assessment of mecaseprin for the treatment of Rett syndrome. *Ann Clin Transl Neurol*. 2018;5(3):323–332. doi:10.1002/acn3.533

51. Vahdatpour C, Dyer AH, Tropea D. Insulin-like growth factor 1 and related compounds in the treatment of childhood-onset neurodevelopmental disorders. *Front Neurosci.* 2016;10:450. doi:10.3389/fnins.2016.00450
52. Sara VR, Carlsson-Sdwirut C, Bergman T, et al. Identification of Gly-Pro-Glu (GPE), the aminoterminal tripeptide of insulin-like growth factor 1 which is truncated in brain, as a novel neuroactive peptide. *Bioch Bioph Res Com.* 1989;165:766–771. doi:10.1016/S0006-291X(89)80032-4
53. Nilsson-Håkansson L, Civalero I, Zhang X, Carlsson-Skwirut C, Sara VR, Nordberg A. Effects of IGF-1, truncated IGF-1 and the tripeptide Gly-Pro-Glu on acetylcholine release from parietal cortex of rat brain. *Neuroreport.* 1993;4(9):1111–1114.
54. Bourguignon JP, Gérard A, Alvarez Gonzalez ML, Franchimont P. Acute suppression of gonadotropin-releasing hormone secretion by insulin-like growth factor I and subproducts: an age-dependent endocrine effect. *Neuroendocrinology.* 1993;58(5):525–530. doi:10.1159/000126586
55. Ikeda T, Waldbillig RJ, Puro DG. Truncation of IGF-I yields two mitogens for retinal Muller glial cells. *Brain Res.* 1995;686:87–92. doi:10.1016/0006-8993(95)00473-4
56. Saura J, Curatolo L, Williams CE, et al. Neuroprotective effects of Gly-Pro-Glu, the N-terminal tripeptide of IGF-1, in the hippocampus in vitro. *Neuroreport.* 1999;10:161–164. doi:10.1097/00001756-199901180-00031
57. Guan J, Williams CE, Skinner SJ, Mallard EC, Gluckman PD. The effects of insulin-like growth factor (IGF)-1, IGF-2, and des-IGF-1 on neuronal loss after hypoxic-ischemic brain injury in adult rats: evidence for a role for IGF binding proteins. *Endocrinology.* 1996;137(3):893–898. doi:10.1210/endo.137.3.8603600
58. Guan J, Waldvogel HJ, Faull RL, Gluckman PD, Williams CE. The effects of the N-terminal tripeptide of insulin-like growth factor-1, glycine-proline-glutamate in different regions following hypoxic-ischemic brain injury in adult rats. *Neuroscience.* 1999;89:649–659. doi:10.1016/S0306-4522(98)00338-8
59. Smith-Hicks CL, Gupta S, Ewen JB, et al. Randomized open-label trial of dextromethorphan in Rett syndrome. *Neurology.* 2017;89(16):1684–1690. doi:10.1212/WNL.0000000000004515
60. Guan J, Thomas GB, Lin H, et al. Neuroprotective effects of the N-terminal tripeptide of insulin-like growth factor-1, glycine-proline-glutamate (GPE) following intravenous infusion in hypoxic-ischemic adult rats. *Neuropharmacology.* 2004;47(6):892–903. doi:10.1016/j.neuropharm.2004.07.002
61. Williams EC, Zhong X, Mohamed A, et al. Mutant astrocytes differentiated from Rett syndrome patients-specific iPSCs have adverse effects on wild-type neurons. *Hum Mol Genet.* 2014;23(11):2968–2980. doi:10.1093/hmg/ddu008
62. Batchelor DC, Lin H, Wen JY, et al. Pharmacokinetics of glycine-proline-glutamate, the N-terminal tripeptide of insulin-like growth factor-1, in rats. *Anal Biochem.* 2003;323(2):156–163. doi:10.1016/j.ab.2003.08.032
63. Baker AM, Batchelor DC, Thomas GB, et al. Central penetration and stability of N-terminal tripeptide of insulin-like growth factor-I, glycine-proline-glutamate in adult rat. *Neuropeptides.* 2005;39(2):81–87. doi:10.1016/j.npep.2004.11.001
64. Tropea D, Giacometti E, Wilson NR, et al. Partial reversal of Rett Syndrome-like symptoms in MeCP2 mutant mice. *Proc Natl Acad Sci USA.* 2009;106:2029–2034. doi:10.1073/pnas.0812394106
65. Harris PWR, Brimble MA, Muir VJ, Lai M, Trotter NS, Callis DJ. Synthesis of proline-modified analogues of the neuroprotective agent glycyl-L-prolyl-glutamic acid (GPE). *Tetrahedron.* 2005;61:10018–10035. doi:10.1016/j.tet.2005.08.026
66. Samonina G, Ashmarin I, Lyapina L. Glyproline peptide family: review on bioactivity and possible origins. *Pathophysiology.* 2002;8:229–234. doi:10.1016/S0928-4680(02)00018-4
67. Guan J, Mathai S, Harris AP, et al. Peripheral administration of a novel diketopiperazine, NNZ 2591, prevents brain injury and improves somatosensory-motor function following hypoxia-ischemia in adult rats. *Neuropharmacology.* 2007;53:749–762. doi:10.1016/j.neuropharm.2007.08.010
68. Guan J, Gluckman PD. IGF-1 derived small neuropeptides and analogues: a novel strategy for the development of pharmaceuticals for neurological conditions. *Br J Pharmacol.* 2009;157(6):881–891. doi:10.1111/j.1476-5381.2009.00256.x
69. Guan J, Harris P, Brimble M, et al. The role for IGF-1-derived small neuropeptides as a therapeutic target for neurological disorders. *Expert Opin Ther Targets.* 2015;19(6):785–793. doi:10.1517/14728222.2015.1010514
70. Lu XC, Chen RW, Yao C, et al. NNZ-2566, a glypromate analog, improves functional recovery and attenuates apoptosis and inflammation in a rat model of penetrating ballistic-type brain injury. *J Neurotrauma.* 2009;26(1):141–154. doi:10.1089/neu.2008.0629
71. Bickerdike MJ, Thomas GB, Batchelor DC, et al. NNZ-2566: a Gly-Pro-Glu analogue with neuroprotective efficacy in a rat model of acute focal stroke. *J Neurol Sci.* 2009;278(1–2):85–90. doi:10.1016/j.jns.2008.12.003
72. Lu XC, Si Y, Williams AJ, Hartings JA, Gryder D, Tortella FC. NNZ-2566, a glypromate analog, attenuates brain ischemia-induced non-convulsive seizures in rats. *J Cereb Blood Flow Metab.* 2009;29(12):1924–1932. doi:10.1038/jcbfm.2009.109
73. Wei HH, Lu XC, Shear DA, et al. NNZ-2566 treatment inhibits neuroinflammation and pro-inflammatory cytokine expression induced by experimental penetrating ballistic-like brain injury in rats. *J Neuroinflammation.* 2009;6:19. doi:10.1186/1742-2094-6-19
74. Cartagena CM, Phillips KL, Williams GL, et al. Mechanism of action for NNZ-2566 anti-inflammatory effects following PBBI involves upregulation of immunomodulator ATF3. *Neuromolecular Med.* 2013;15(3):504–514. doi:10.1007/s12017-013-8236-z
75. Deacon RM, Glass L, Snape M, et al. NNZ-2566, a novel analog of (1-3) IGF-1, as a potential therapeutic agent for fragile X syndrome. *Neuromolecular Med.* 2015;17(1):71–82. doi:10.1007/s12017-015-8341-2
76. Oosterholt SP, Horrigan J, Jones N, Glass L, Della Pasqua O. Population pharmacokinetics of NNZ-2566 in healthy subjects. *Eur J Pharm Sci.* 2017;109:S98–S107. doi:10.1016/j.ejps.2017.05.032
77. Glaze DG, Neul JL, Percy A, et al. A double-blind, randomized, placebo-controlled clinical study of trofinetide in the treatment of rett syndrome. *Pediatr Neurol.* 2017;76:37–46. doi:10.1016/j.pediatrneurol.2017.07.002
78. Neul JL, Percy AK, Benke TA, et al. Trofinetide for the treatment of Rett syndrome: a randomized phase 3 study. *Nat Med.* 2023;29(6):1468–1475. doi:10.1038/s41591-023-02398-1
79. Percy A, Ryther R, Marsh E, et al. Trofinetide for the treatment of Rett syndrome: an open-label study in girls 2 to 4 years of age. *Neurology.* 2023;100:1378. doi:10.1212/WNL.00000000000201897
80. Camm AJ. Clinical trial design to evaluate the effects of drugs on cardiac repolarization: current state of the art. *Heart Rhythm.* 2005;2(2 Suppl): S23–9. doi:10.1016/j.hrthm.2004.09.019

81. Chapleau CA, Lane J, Pozzo-Miller L, Percy AK. Evaluation of current pharmacological treatment options in the management of Rett syndrome: from the present to future therapeutic alternatives. *Curr Clin Pharmacol*. 2013;8(4):358–369. doi:10.2174/15748847113086660069
82. Grosser E, Hirt U, Janic OA, et al. Oxidative burden and mitochondrial dysfunction in a mouse model of Rett syndrome. *Neurobiol Dis*. 2012;48(1):102–114. doi:10.1016/j.nbd.2012.06.007
83. Ramirez JM, Ward CS, Neul JL. Breathing challenges in Rett syndrome: lessons learned from humans and animal models. *Respir Physiol Neurobiol*. 2013;189(2):280–287. doi:10.1016/j.resp.2013.06.022
84. Chao HT, Chen H, Samaco RC, et al. Dysfunction in GABA signalling mediates autism-like stereotypies and Rett syndrome phenotypes. *Nature*. 2010;468(7321):263–269. doi:10.1038/nature09582
85. Abdala AP, Liroy DT, Garg SK, Knopp SJ, Paton JF, Bissonnette JM. Effect of Sarizotan, a 5-HT1a and D2-like receptor agonist, on respiration in three mouse models of Rett syndrome. *Am J Respir Cell Mol Biol*. 2014;50(6):1031–1039.
86. Leonard BE. A comparison of the pharmacological properties of the novel tricyclic antidepressant lofepramine with its major metabolite, desipramine: a review. *Int Clin Psychopharmacol*. 1987;2(4):281–297. doi:10.1097/00004850-198710000-00001
87. Mancini J, Dubus JC, Jouve E, et al. Effect of desipramine on patients with breathing disorders in RETT syndrome. *Ann Clin Transl Neurol*. 2017;5(2):118–127. doi:10.1002/acn3.468
88. Naegelin Y, Kuhle J, Schädelin S, et al. Fingolimod in children with Rett syndrome: the FINGORETT study. *Orphanet J Rare Dis*. 2021;16(1):19. doi:10.1186/s13023-020-01655-7
89. Yıldız N, Serdaroglu E, Kart PÖ, et al. Evaluation of seizure semiology, genetics, magnetic resonance imaging, and electroencephalogram findings in children with Rett syndrome: a multicenter retrospective study. *Epilepsy Res*. 2024;205:107399. doi:10.1016/j.eplesyres.2024.107399
90. Allison K, Maletic-Savatic M, Pehlivan D. MECP2-related disorders while gene-based therapies are on the horizon. *Front Genet*. 2024;15:1332469. doi:10.3389/fgene.2024.1332469
91. Glaze DG, Percy AK, Skinner S, et al. Epilepsy and the natural history of Rett syndrome. *Neurology*. 2010;74(11):909–912. doi:10.1212/WNL.0b013e3181d6b852
92. Guerrini R, Parrini E. Epilepsy in Rett syndrome, and CDKL5- and FOXP1-gene-related encephalopathies. *Epilepsia*. 2012;53(12):2067–2078. doi:10.1111/j.1528-1167.2012.03656.x
93. Specchio N, Balestri M, Striano P, et al. Efficacy of levetiracetam in the treatment of drug-resistant Rett syndrome. *Epilepsy Res*. 2010;88(2–3):112–117. doi:10.1016/j.eplesyres.2009.10.005
94. Pintaudi M, Calevo MG, Vignoli A, et al. Antiepileptic drugs in Rett Syndrome. *Eur J Paediatr Neurol*. 2015;19(4):446–452. doi:10.1016/j.ejpn.2015.02.007
95. Vignoli A, Savini MN, Nowbut MS, et al. Effectiveness and tolerability of antiepileptic drugs in 104 girls with Rett syndrome. *Epilepsy Behav*. 2017;66:27–33. doi:10.1016/j.yebeh.2016.10.006
96. Krajnc N, Župančič N, Oražem J. Epilepsy treatment in Rett syndrome. *J Child Neurol*. 2011;26(11):1429–1433. doi:10.1177/0883073811408422
97. Dolce A, Ben-Zeev B, Naidu S, Kossoff EH. Rett syndrome and epilepsy: an update for child neurologists. *Pediatr Neurol*. 2013;48(5):337–345. doi:10.1016/j.pediatrneurol.2012.11.001
98. Djukic A, Holtzer R, Shinnar S, et al. Pharmacologic treatment of rett syndrome with glatiramer acetate. *Pediatr Neurol*. 2016;61:51–57. doi:10.1016/j.pediatrneurol.2016.05.010
99. Nissenkorn A, Kidon M, Ben-Zeev B. A potential life-threatening reaction to glatiramer acetate in rett syndrome. *Pediatr Neurol*. 2017;68:40–43. doi:10.1016/j.pediatrneurol.2016.11.006
100. Desnoux B, Beretti T, Muller N, et al. Efficacy and tolerance of cannabidiol in the treatment of epilepsy in patients with Rett syndrome. *Epilepsia Open*. 2024;9(1):397–403. doi:10.1002/epi4.12796
101. Devinsky O, Verducci C, Thiele EA, et al. Open-label use of highly purified CBD (Epidiolex®) in patients with CDKL5 deficiency disorder and Aicardi, Dup15q, and Doose syndromes. *Epilepsy Behav*. 2018;86:131–137. doi:10.1016/j.yebeh.2018.05.013
102. Dale T, Downs J, Wong K, Leonard H. The perceived effects of cannabis products in the management of seizures in CDKL5 deficiency disorder. *Epilepsy Behav*. 2021;122:108152. doi:10.1016/j.yebeh.2021.108152
103. Jazz P, Pharmaceuticals GW. A long-term safety study of cannabidiol oral solution (GWP42003-P, CBD-OS) in patients with rett syndrome; Available from: <https://classic.clinicaltrials.gov/show/NCT04252586>. Accessed July 31, 2023.
104. Hurlley EN, Ellaway CJ, Johnson AM, et al. Efficacy and safety of cannabidivarin treatment of epilepsy in girls with Rett syndrome: a Phase 1 clinical trial. *Epilepsia*. 2022;63(7):1736–1747. doi:10.1111/epi.17247
105. Karimzadeh A, Baradaran-Salimi K, Voges B, et al. Short post-injection seizure duration is associated with reduced power of ictal brain perfusion SPECT to lateralize the seizure onset zone. *EJNMMI Res*. 2024;14(1):40. doi:10.1186/s13550-024-01095-5
106. D’Onofrio G, Kuchenbuch M, Hachon-Le Camus C, et al. Slow Titration of Cannabidiol Add-On in Drug-Resistant Epilepsies Can Improve Safety With Maintained Efficacy in an Open-Label Study. *Front Neurol*. 2020;11:829. doi:10.3389/fneur.2020.00829
107. Peters JM, Puri V, Segal E, Misra SN, Rabinowicz AL, Carrazana E. Concomitant cannabidiol does not impact safety and effectiveness of diazepam nasal spray for seizure clusters: post hoc analysis of a phase 3 safety study. *Epilepsy Behav*. 2023;144:109248. doi:10.1016/j.yebeh.2023.109248
108. Tarquinio D, Wheless JW, Segal EB, Misra SN, Rabinowicz AL, Carrazana E. Safety of diazepam nasal spray in pediatric patients with developmental epileptic encephalopathies: results from a long-term phase 3 safety study. *J Child Neurol*. 2023;38(6–7):389–393. doi:10.1177/08830738231185424
109. Downs J, Jacoby P, Specchio N, et al. Effects of ganaxolone on non-seizure outcomes in CDKL5 deficiency disorder: double-blind placebo-controlled randomized trial. *Eur J Paediatr Neurol*. 2024;51:140–146. doi:10.1016/j.ejpn.2024.06.005
110. Pong AW, Ross J, Tyrlikova I, et al. Epilepsy: expert opinion on emerging drugs in phase 2/3 clinical trials. *Expert Opin Emerg Drugs*. 2022;27(1):75–90. doi:10.1080/14728214.2022.2059464
111. Knight EMP, Amin S, Bahi-Buisson N, et al.; Marigold Trial Group. Safety and efficacy of ganaxolone in patients with CDKL5 deficiency disorder: results from the double-blind phase of a randomised, placebo-controlled, phase 3 trial. *Lancet Neurol*. 2022;21(5):417–427. doi:10.1016/S1474-4422(22)00077-1

112. Ellaway CJ, Peat J, Williams K, Leonard H, Christodoulou J. Medium-term open label trial of L-carnitine in Rett syndrome. *Brain Dev.* 2001;23 (Suppl 1):S85–9. doi:10.1016/S0387-7604(01)00346-1
113. Cappuccio G, Bernardo P, Raiano E, et al. Pain and sleep disturbances in Rett syndrome and other neurodevelopmental disorders. *Acta Paediatr.* 2019;108(1):171–172. doi:10.1111/apa.14576
114. Miyamoto A, Oki J, Takahashi S, Okuno A. Serum melatonin kinetics and long-term melatonin treatment for sleep disorders in Rett syndrome. *Brain Dev.* 1999;21:59–62. doi:10.1016/S0387-7604(98)00072-2
115. McArthur AJ, Budden SS. Sleep dysfunction in Rett syndrome: a trial of exogenous melatonin treatment. *Dev Med Child Neurol.* 1998;40:186–192. doi:10.1111/j.1469-8749.1998.tb15445.x
116. Liou DT, Wu WW, Bissonnette JM. Autonomic dysfunction with mutations in the gene that encodes methyl-CpG-binding protein 2: insights into Rett syndrome. *Auton Neurosci.* 2011;161:55–62. doi:10.1016/j.autneu.2011.01.006
117. Gemelli T, Berton O, Nelson ED, Perrotti LI, Jaenisch R, Monteggia LM. Postnatal loss of methyl-CpG binding protein 2 in the forebrain is sufficient to mediate behavioral aspects of Rett syndrome in mice. *Biol Psychiatry.* 2006;59:468–476. doi:10.1016/j.biopsych.2005.07.025
118. Hesketh S, Jessop DS, Hogg S, Harbuz MS. Differential actions of acute and chronic citalopram on the rodent hypothalamic-pituitary-adrenal axis response to acute restraint stress. *J Endocrinol.* 2005;185(3):373–382. doi:10.1677/joe.1.06074
119. Venkat A, Jauch E, Russell WS, Crist CR, Farrell R. Care of the patient with an autism spectrum disorder by the general physician. *Postgrad Med J.* 2012;88(1042):472–481. doi:10.1136/postgradmedj-2011-130727
120. Flores Gutiérrez J, De Felice C, Natali G, et al. Protective role of mirtazapine in adult female Mecp2<sup>±</sup> mice and patients with Rett syndrome. *J Neurodev Disord.* 2020;12(1):26. doi:10.1186/s11689-020-09328-z
121. Ette EI, Fadiran EO, Missling C, Hammond E. The new big is small: leveraging knowledge from small trials for rare disease drug development: blarcamesine for Rett syndrome. *Br J Clin Pharmacol.* 2023. doi:10.1111/bcp.15843
122. Mohammed HE, Bady Z, Haseeb ME, Aboeldahab H, Sharaf-Eldin WE, Zaki MS. Is trofinetide a future treatment for Rett syndrome? A comprehensive systematic review and meta-analysis of randomized controlled trials. *BMC Med.* 2024;22(1):299. doi:10.1186/s12916-024-03506-9

## Drug Design, Development and Therapy

Dovepress

### Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>