

Childhood Maltreatment and Adolescent Eating Disorders' Symptoms: A Moderated Mediation Model of Social Anxiety and Physical Activity

Sihong Li, Xuerong Luo, Lintong Song, Xueping Gao, Yanmei Shen

Department of Psychiatry, National Clinical Research Center for Mental Disorders, and National Center for Mental Disorders, The Second Xiangya Hospital of Central South University, Changsha, Hunan, 410011, People's Republic of China

Correspondence: Yanmei Shen; Xueping Gao, Department of Psychiatry, National Clinical Research Center for Mental Disorders, and National Center for Mental Disorders, The Second Xiangya Hospital of Central South University, Changsha, Hunan, 410011, People's Republic of China, Email ymshen@csu.edu.cn; xuepinggao@csu.edu.cn

Background: Childhood maltreatment contributes to an increased risk of eating disorders in adolescents. However, less is known regarding the underlying mechanism between childhood maltreatment and eating disorders' symptoms. This study tested the mediation effect of social anxiety in the association between childhood maltreatment and eating disorders' symptoms and whether this mediation was moderated by physical activity.

Methods: A longitudinal study was conducted among 848 students in grades 7–8 from a middle school in Changsha, China. The students completed measurements on childhood maltreatment (Childhood Trauma Questionnaire) at the baseline and measurements on social anxiety (The Social Anxiety Scale-Adolescents), physical activity, and eating disorders' symptoms (The Eating Attitudes Test) after six months. Mediation and moderation analyses were carried out in SPSS macro-PROCESS.

Results: Emotional abuse was significantly related to eating disorders' symptoms, and the association was mediated by social anxiety (indirect effect: $\beta = 0.03$, 95% CI: 0.01 to 0.05, $p < 0.05$). Furthermore, physical activities moderated the path of emotional abuse to social anxiety ($\beta = 0.32$, 95% CI: 0.25 to 0.39, $p < 0.01$) and social anxiety to eating disorders' symptoms ($\beta = 0.18$, 95% CI: 0.10 to 0.26, $p < 0.01$).

Conclusion: This research emphasizes the importance of social anxiety in the relationship between childhood maltreatment and eating disorders' symptoms. Additionally, these associations are weaker for adolescents with more physical activities, addressing the importance of physical activity in the prevention and management of eating disorders' symptoms.

Keywords: childhood maltreatment, social anxiety, eating disorders, adolescent, physical activity, moderated mediation model

Introduction

Eating disorders (ED) are severe psychiatric disorders characterized by maladaptive eating behaviors or excessive attention to food and body weight and size.¹ ED includes anorexia nervosa (AN), bulimia nervosa (BN), binge eating disorder (BED) and so on.¹ The incidence of ED peaks during the early stages of life, especially in youth and adolescence, a critical phase for the development of eating patterns.^{2,3} Additionally, recent research has shown a strong correlation between ED and a variety of behavioral and psychosocial problems, including substance abuse, depression, anxiety, and even suicidality.^{4–10} Moreover, there has been a 25% rise in the occurrence of eating disorders globally, while only roughly 20% of those impacted seek out medical help.^{11,12} In China, ED symptoms are common problems among middle school students, with a prevalence of 8.9%.¹³ Given the early onset, high prevalence, and heavy disease burden of ED, investigating the influencing factors and underlying mechanisms of ED symptoms is critical in informing future prevention and intervention efforts.

Research has established a correlation between childhood maltreatment (CM) and the development of ED.¹⁴ CM encompasses various subtypes, such as physical and emotional neglect, as well as physical and emotional abuse (EA).

Two systematic reviews and meta-analyses pointed out that individuals suffering from ED had more exposure to CM compared to other mental disorders, regardless of the specific CM type.^{15,16} Additionally, a prospective study revealed that exposure to CM increased the likelihood of developing ED, such as AN and BN, in adulthood.¹⁷ Notably, among all forms of CM, EA has been linked explicitly to symptoms of ED, such as the desire for thinness, body dissatisfaction, and bingeing and purging behaviors.^{18,19} A network analysis study also showed that various types of CM were associated with ED core symptoms solely via EA.²⁰ These results suggest that CM, especially EA, may lead to the emergence of ED.

Previous studies have established a robust connection between CM and ED. However, the specific mechanisms through which CM affects ED symptoms are still unclear. Moderated mediation analysis is a valuable method for evaluating the associations between various factors. Two variables establish the relationship through mediators, whereas moderators affect the triangular relationship.²¹ Therefore, identifying mediating and moderating factors from CM to ED symptoms is essential for the interventions of ED symptoms and the minimization of the adverse effects of CM.

The Mediating Role of Social Anxiety (SA)

Social anxiety (SA) denotes elevated levels of fear in social situations, resulting in notable distress and functional impairment at a clinical level.¹⁷ The association between CM and SA has been extensively documented in scholarly literature. According to Arrindell (1983 and 1989), experiencing inadequate care and rejection of family in childhood have distinct associations with difficulties in interpersonal social interactions.^{22,23} Subsequent research has further underscored the predictive role of CM in SA. For instance, several surveys conducted both in clinical and non-clinical populations discovered that CM was linked to more severe SA symptoms and impaired functioning, particularly in cases of emotional abuse and neglect.^{24–26}

Furthermore, abundant evidence reveals that SA is a potential risk factor for ED.^{27–32} A meta-analysis cited SA as one of the most common comorbid conditions in ED, indicating that higher levels of SA were correlated with more severe ED psychopathology.³³ Moreover, it is reported that SA onset preceded ED in the majority of those with both disorders.^{34,35} In addition, according to attachment theory, the way in which primary caregivers respond to children during early childhood (eg, CM) influences an individual's interpersonal relationships or attachment styles in later life.^{36,37} The attachment style characterized by avoidance and anxiety may negatively affect interpersonal closeness and intimacy (eg, SA),³⁸ which have been consistently associated with symptoms of ED.^{33,39,40} Therefore, drawing from the findings of the literature review and attachment theory, it is reasonable to hypothesize that SA mediates the relation between CM and ED symptoms.

The Moderating Role of Physical Activity

The response of adolescents to CM varies, with some able to achieve healthy development despite experiencing maltreatment. However, there has been limited research addressing this topic. Therefore, it's important to explore the potential moderating variables in the pathway from CM to ED symptoms to minimize the adverse outcomes of CM.

Physical activity (PA) refers to any physical movement that raises caloric expenditure beyond the baseline resting level.⁴¹ The World Health Organization (WHO) guidelines recommend that kids and teenagers aged 5 to 17 should participate in at least 60 minutes of daily PA with moderate to vigorous intensity.⁴² However, a school-based survey including 11,110 European adolescents found that only 17.9% of boys and 10.7% of girls met the standard for WHO-recommended sufficient PA (60 min/day).⁴³ PA is known to have a stress-buffering effect, which can mitigate the negative psychological impacts of stressors such as CM.⁴⁴ According to the self-decision theory, engaging in behaviors such as PA can satisfy psychological needs, potentially reducing SA to a certain extent.^{45,46} Furthermore, PA contributes to greater well-being and lower levels of anxiety in both sexes. A randomized controlled trial (RCT) showed that exercise intervention combined with general cognitive behavior treatment (CBT) was more effective in reducing SA than CBT combined with education.⁴⁷ In addition, previous studies also showed that elevated levels of PA were associated with the initiation and continuation of ED.^{48,49} One RCT even proposed that physical exercise and dietary therapy could serve as alternative treatments of CBT for ED.⁵⁰ However, it remains unknown whether WHO-recommended sufficient PA could moderate the indirect link between CM and ED symptoms through SA in adolescents.

The Present Study

To sum up, we tested our proposed model among Chinese adolescents using a longitudinal study design. We aimed to explore whether social anxiety mediated the relationship between CM and ED symptoms and whether this triangular association was moderated by WHO-recommended sufficient PA. Collectively, these two inquiries form a model of moderated mediation. The proposed model is depicted in Figure 1.

Methods

Participants and Procedure

From November to December 2020, 1162 students in grades 7–8 were recruited from a middle school in Changsha City, Hunan, China, using the clustered sampling method. Inclusion criteria included: (a) aged between 11 to 16 years old; (b) enrolled in middle school full-time; (c) willing to participate with informed consent; (d) able to understand and complete the questionnaire survey. We excluded students who were absent due to sickness or other reasons during the research period, as well as those who were not able to complete the questionnaire due to severe physical and mental illness.

Ethical approval was granted by the Ethics Committee of the Second Xiangya Hospital at Central South University. Prior to the survey, we trained school teachers on the procedures for distributing the questionnaire, offering guidance, and assisting students experiencing difficulties in completing the questionnaire. The questionnaires were distributed via WeChat, one of the largest online social media platforms in China. Permission was acquired from both parents and students. Prior to data collection, teachers notified parents and obtained their consent for their children's participation. Then, students accessed the survey using their parents' mobile phones and provided electronic informed consent on the first page of the survey.

A total of 1084 students completed the baseline assessment for CM; among them, 848 completed assessments on ED symptoms and SA six months later, from May to June 2021. Therefore, 848 students were included in the final analysis.

Measures

Sociodemographic Characteristics

Sociodemographic variables included gender, age, nationality, and only child.

CM

CM was evaluated using the Childhood Trauma Questionnaire (CTQ) to explore adverse childhood experiences before the age of 16 by prompting participants to recall and report on their past or present experiences. The CTQ consists of 23 items, categorized into four subscales: emotional abuse, emotional neglect, physical abuse, and physical neglect. Each item utilizes a 5-point frequency scale, ranging from “never happened” (scored as 1) to “always happened” (scored as 5).⁵¹ Higher scores indicated a greater degree of trauma experienced in childhood. The CTQ has demonstrated good internal consistency in the




Figure 1 The proposed model.

current study.^{52,53} The previous studies demonstrated that the Chinese adaptation of the CTQ exhibits strong reliability and validity for evaluating childhood trauma.⁵⁴

ED Symptoms

ED symptoms were assessed using the 26-item Eating Attitudes Test (EAT-26). The scale has demonstrated good reliability, validity, sensitivity, and specificity for measuring ED symptoms, with higher scores indicating the more number of ED symptoms including disturbed eating attitudes and behaviors.^{55–57} It is commonly employed in various populations, including students from middle high school and college and individuals with specific risk factors, such as athletes.⁵⁸ The Chinese version of the EAT-26 serves as a reliable and valid self-assessment tool for detecting individuals with ED symptoms in mainland China.⁵⁹

PA

The guidelines established by the World Health Organization recommend that children and adolescents engage in a cumulative time of 60 minutes for moderate to vigorous physical activity each day to promote their overall health.⁶⁰ Thus, PA was assessed by one self-reported question:

Over the past week, on how many days did you engage in a minimum of 60 minutes of moderate to vigorous physical activity, resulting in an elevated heart rate and occasional heavy breathing?

They could choose from response options ranging from 0 to 7 days per week. This question was selected from the Youth Risk Behavior Survey (YRBS).⁶¹

SA

SA was evaluated using the simplified version of the Social Anxiety Scale for Adolescents (SAS-A).⁶² The SAS-A includes 13 items, which are divided into three subscales: Fear of Negative Evaluation, Social Avoidance and Distress in new social situations or with unfamiliar peers, and Social Avoidance and Distress that is more general or pervasive (SAD-G).^{62,63} Each item is rated from “not applicable” (scored as 1) to “applicable to me very much” (scored as 5). The SAS-A has demonstrated good internal consistency in the current study.⁶⁴

Statistical Analysis

SPSS software (Version 26.0; IBM, Inc., Chicago, IL) was used for data analyses, with $p < 0.05$ indicating statistical significance. A one-sample Kolmogorov–Smirnov test was performed to assess the normality of continuous. Due to the non-normality of CM, SA, and ED scores, Spearman correlation analyses were conducted to assess their correlations. To examine the hypothesized moderated mediation model, we utilized Model 59 of SPSS PROCESS 4.0 macro. Firstly, Harman’s single-factor analysis was conducted to evaluate any potential common method deviation. Afterwards, the influence of PA on the model was analyzed with Model 59, as depicted in Figure 1. We standardized all the continuous variables and calculated interaction terms. Furthermore, the moderation and mediation effect was tested using a bootstrapping method with 5000 samples, with a 95% confidence interval (CI) not including zero indicating a significant effect. All models were adjusted for confounding variables, including age, gender, nationality, and single child.

Results

After testing Model 59 on each subscale of CTQ and SAS-A, we discovered a significant moderated mediation model, which is depicted in Figure 2.

Common Method Deviation Test

Harman’s single-factor analysis was utilized to examine the presence of common method bias. An exploratory factor analysis was conducted on all the items in the EA, SAD-G, and ED symptoms’ scales. The findings indicated that the first factor had an interpretation rate of 30.02%, which was less than 40% of the reference value. As a result, we could infer that the common method deviation was not statistically significant.




Figure 2 The final moderated mediation model.

Note: *** $p < 0.001$.

Abbreviation: SAD-G, Social Avoidance and Distress that is more general or pervasive.

Preliminary Analysis

Table 1 presents the descriptive statistics of all primary study variables, and zero-order correlations for EA, ED symptoms, PA and SAD-G. Results showed that EA was positively correlated with ED symptoms and SAD-G and negatively correlated with PA. Besides, PA was negatively correlated with SAD-G.

Mediation Analyses

Table 2 displays the outcomes of the mediation analyses. The total effect of EA on ED symptoms (c) was 0.17 (95% CI: 0.11 to 0.24), while the direct effect (c') was 0.14 (95% CI: 0.07 to 0.21). SAD-G was associated with both EA ($\beta=0.27$, 95% CI: 0.20 to 0.33) and ED symptoms ($\beta=0.12$, 95% CI: 0.05 to 0.19). The bootstrapping index for an indirect impact ($\beta = 0.03$) as statistically significant with the inclusion of SAD-G as a mediating factor, as the 95% CI did not encompass zero (0.01 to 0.05). SA accounted for 18% of the total effect of EA on ED symptoms.

Table 1 Descriptive Statistics and Correlations of the Main Study Variables

Variables	M, SD	EA	ED	SAD-G	PA
Age	12.59,0.64				
Gender, n(%)					
male	449(52.9)				
female	399(47.1)				
Nationality, (%)					
Han	803(94.7)				
Others	45(5.3)				
Only Child, (%)					
Yes	505(59.6)				
No	343(40.4)				
EA	7.70, 3.57	1.00	0.15**	0.27**	-0.01
ED	8.26, 10.17	0.15**	1.00	0.19**	0.03
SAD-G	4.05, 2.43	0.27**	0.19**	1.00	-0.11**
PA	4.89, 2.35	-0.01	0.36	-0.11**	1.00

Note: ** $p < 0.01$.

Abbreviations: N =848; SD, standard deviation; EA, emotional abuse; ED, eating disorders' symptoms; SAD-G, Social Avoidance and Distress that is more general or pervasive; PA, physical activity.

Table 2 Mediation Effect of SA on the Association Between EA and ED

Outcome	SAD-G			ED		
	β	LCI	UCI	β	LCI	UCI
EA	0.27***	0.20	0.33	0.14***	0.07	0.21
SAD-G				0.12***	0.05	0.19
F	R-sq	0.11***		R-sq	0.22***	
		21.31			7.43	

Note: *** $p < 0.001$.

Abbreviations: N =848; EA, emotional abuse; ED, eating disorders' symptoms; SAD-G, Social Avoidance and Distress that is more general or pervasive; PA, physical activity; LCI, lower 95% confidence interval; UCI, 95% upper confidence interval.

Moderated Mediation Analyses

The moderated mediation analysis (Table 3) indicated that PA concurrently moderated the effect of EA on SAD-G ($\beta = 0.32$, 95% CI: 0.25 to 0.39) and the effect of SAD-G on ED symptoms ($\beta = 0.18$, 95% CI: 0.10 to 0.26). Nevertheless, PA did not show any significant moderating effect in the direct route from EA to ED symptoms ($\beta = 0.12$, 95% CI 0.04 to 0.20). The results of moderated mediation for SAD-G are displayed in Figures 3 and 4.

To further investigate the impact of EA on ED symptoms across various PA levels, we categorized PA into two levels based on the WHO guidelines: low PA (<60 min/day) and sufficient PA (≥ 60 min/day). Table 4 displays the findings. The direct effect of EA on ED symptoms was notably significant in both the low PA group (effect = 0.12, 95% CI: 0.04 to 0.20) and the sufficient PA group (effect = 0.16, 95% CI: 0.01 to 0.30). Furthermore, in the low PA group, the indirect impact of EA on ED symptoms was significant (effect = 0.06, 95% CI: 0.03 to 0.09). However, in the sufficient PA group, the indirect impact of EA on ED symptoms was nonsignificant (effect = -0.01, 95% CI: 0.03 to 0.01), suggesting that the association between EA and ED symptoms was attenuated for adolescents with sufficient PA. The index of moderated mediation is -0.05 with the 95% CI of -0.09 to -0.02.

Discussion

Our study showed that SA mediated the relationship between EA and adolescent ED symptoms. Moreover, PA moderated the indirect effect of EA on ED symptoms. Specifically, the effects of EA on SAD-G and of SAD-G on ED symptoms were weaker in the WHO-recommended sufficient PA group, emphasizing the possible protective effect of PA against ED symptoms.

Table 3 Moderated Mediation Analysis Results for the Relationship Between EA and ED

Outcome	SAD-G			ED		
	β	LCI	UCI	β	LCI	UCI
EA	0.32***	0.25	0.39	0.12**	0.04	0.20
SAD-G				0.18***	0.10	0.26
ED						
EA*PA	-0.24**	-0.40	-0.08	0.04	-0.13	0.20
SA*PA				-0.26**	-0.43	-0.09
F	R-sq	0.36***		R-sq	0.26***	
		17.37			6.95	

Notes: * $p < 0.05$. ** $p < 0.01$. *** $p < 0.001$.

Abbreviations: N =848; EA, emotional abuse; ED, eating disorders' symptoms; SA/SAD-G, Social Avoidance and Distress that is more general or pervasive; PA, physical activity; LCI, lower 95% confidence interval; UCI, 95% upper confidence interval.




Figure 3 PA as a moderator between EA and SAD-G.

Abbreviations: SAD-G, Social Avoidance and Distress that is more general or pervasive.




Figure 4 PA as a moderator between SAD-G and ED symptoms.

Note: SAD-G: Social Avoidance and Distress that is more general or pervasive.

The Mediating Role of SAD-G

Our findings indicated that SAD-G mediated the relationship between EA and ED symptoms. This suggests that adolescents who have experienced EA have a higher risk of SAD-G, which further leads to an increased risk of ED symptoms. Previous studies in both clinical and non-clinical samples showed that EA was positively related to ED.^{65,66}

Table 4 Conditional Direct and Indirect Effects of EA on ED by PA Groups

	Paths	EA→ED			EA→SA→ED		
		Effect	LCI	UCI	Effect	LCI	UCI
Low		0.12**	0.04	0.20	0.06*	0.03	0.09
Sufficient		0.16*	0.01	0.30	−0.01	−0.03	0.01

Notes: * $p < 0.05$. ** $p < 0.01$.

Abbreviations: N =848; EA, emotional abuse; ED, eating disorders' symptoms; SA/ SAD-G, Social Avoidance and Distress that is more general or pervasive; PA, physical activity; LCI, lower 95% confidence interval; UCI, 95% upper confidence interval; EA→ED, the direct path from EA to ED symptoms; EA→SA→ED, the indirect path from EA to ED symptoms through SA.

Our study advanced previous understanding by further exploring the underlying mechanism in the association between EA and ED symptoms among adolescents. Previous studies have identified various mediators between CM and ED symptoms, such as emotion regulation,¹⁸ alexithymia,⁶⁷ attachment insecurity,⁶⁸ internal shame, and psychological distress.⁶⁹ Our study identified a new mediator— SAD-G, which is in line with the attachment theory. This theory pointed out that insufficient care and family rejection during childhood are distinctly linked to challenges in social interactions with others, which further leads to ED.^{40,70} EA, as opposed to other forms of abuse, has a stronger tendency to predict the diagnosis of SA.⁷¹ Moreover, EA is closely related to the severity and worse function of SA.^{25,72} A systematic review and meta-analysis also revealed the connection between SA and ED.³³ SA is one of the most common comorbid anxiety disorders among ED and a risk factor for ED. Our study extends this conclusion by showing that SAD-G seems to be more closely related to ED symptoms. This may be explained by the attachment theory, in which challenges in social interactions with others caused by EA are more applicable to closer relationships.^{70,73} Another explanation may be that SAD-G and ED share some common vulnerability factors, like alexithymia.^{74,75} Thus, our integrated model suggests that risk factors in the environmental context (eg childhood EA) may enhance some risk traits (eg, SAD-G), which may in turn increase their ED symptoms tendency.

These findings emphasized the importance of early identification and intervention for adolescents who have experienced EA, as they are at greater risk of developing both SAD-G and ED. Clinicians should consider screening for social anxiety symptoms in patients with a history of emotional abuse and eating disorders, and interventions should target both emotional regulation and social interaction difficulties. Moreover, addressing SAD-G in adolescents could potentially reduce the risk of developing ED symptoms.

The Moderating Role of PA

This study revealed that PA moderated the path between EA and SAD-G (first-stage moderation) and the path between SAD-G and ED symptoms (second-stage moderation).

For the first stage of moderation, PA weakens the association between EA and SAD-G. Previous studies have demonstrated that PA has the potential to effectively treat SA or alleviate SA symptoms in both clinical and non-clinical populations.⁷⁶ RCT studies also showed that compared to usual treatment, the PA intervention was more effective in reducing SA symptoms.^{77,78} Longitudinal studies also proved that SA symptoms were markedly reduced after PA treatment.^{79–84}

For the second stage of moderation, PA attenuates the association between SA and ED symptoms. Although excessive and compulsive PA was considered a symptom of ED,⁸⁵ some studies also pointed out that moderate PA could decrease the frequency of BED episodes.⁸⁶ In the case of AN and BN, incorporating PA into treatment may enhance the quality of life, social interactions, body composition, and overall physical health.^{85,87,88} Our findings indicate that PA could be considered as a key component of interventions aimed at adolescents with a history of EA, SA and ED symptoms. Clinicians could encourage physical activity as a comprehensive treatment plan for adolescents with SA and ED symptoms. Additionally, the significance of conducting additional research on different types, intensities, frequencies,

durations, and timing of PA is highlighted by these studies. It is important to identify the specific parameters that have the greatest influence on patients with ED.

Limitations

Our study has several limitations. First, all data were collected based on self-reported scales, which may result in recall bias. Future studies should consider using multiple informants when collecting data to replicate our findings further. Second, the model was tested on a sample of middle school students, who may not represent other populations. Future studies are needed to test the model in different populations, such as a clinical sample. Third, PA and ED symptoms were assessed based on one self-reported item from the YRBS and EAT-26. Future studies should use more detailed questions and more objective measurement tools to get a more accurate assessment and diagnosis. Additionally, the subgroup analysis with different type of eating disorder (ie, anorexia nervosa or bulimia nervosa) should be considered in further studies. Fourth, CM was assessed using the CTQ, which may capture maltreatment experiences from several years ago and potentially impact the interpretation of the study results. Thus, we will consider longer observation periods in future research designs to better understand the persistence of these effects. Lastly, this study was carried out during the COVID-19 pandemic, which may potentially impact PA, ED symptoms, and SA. Thus, future research should more comprehensively consider the influence of pandemic factors on study outcomes and further explore this association.

Conclusion

Our findings showed that the CM was associated with ED symptoms both directly and indirectly through SA, this highlights the importance of addressing social anxiety as a key factor in understanding the pathways from EA to ED symptoms. Additionally, PA further moderated the mediation effect, with higher levels of PA buffering the negative effects of EA on SA and, subsequently, on ED symptoms.

These findings carry significant implications and open up possibilities for potential interventions. First, interventions targeting decreasing adolescent SA, particularly in those with a history of CM, may help prevent or mitigate the development of ED. Second, PA could be an accessible, low-cost intervention that can be integrated into both clinical and community settings. Incorporating PA into intervention programs may not only improve psychological outcomes such as SA but also reduce the overall risk of developing ED symptoms. Furthermore, our findings highlight the need for prioritizing prevention and intervention efforts among adolescents with high levels of SA and low PA level. By focusing on this at-risk population, interventions can be more tailored and effective in reducing the incidence of ED symptoms.

Abbreviations

CM, Childhood maltreatment; ED, Eating disorders; SA, Social anxiety; EA, Emotional abuse; SAD-G, Social Avoidance and Distress that is more general or pervasive; PA, Physical activities; CTQ, Childhood Trauma Questionnaire; YRBS, Youth Risk Behavior Survey; EAT-26, The Eating Attitudes Test; SD, Standard deviations; CBT, Cognitive behavior treatment; BED, Binge eating disorder; BN, Bulimia nervosa; AN, Anorexia nervosa; WHO, World Health Organization.

Data Sharing Statement

The datasets used or analyzed during the current study are available from the corresponding author upon reasonable request.

Ethics Approval and Consent to Participate

This study protocol was approved by the Ethics Committee of the Second Xiangya Hospital of Central South University. All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2000. Informed consent was obtained from all patients and their parents or legal guardians to be included in the study.

Acknowledgments

We would like to show our great gratitude to the school personnel who have offered invaluable support. We would also like to show our great appreciation to all of the students who participated in this project.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This work was supported by the National Natural Science Foundation of China (Grant No. 82201703) and the Scientific Research Launch Project for new employees of the Second Xiangya Hospital of Central South University. The funders had no role in the study design, in the data collection and analysis, in the writing of the report, and in the decision to submit the paper for publication.

Disclosure

The authors report no conflicts of interest in this work.

References

- Treasure J, Duarte TA, Schmidt U. Eating disorders. *Lancet*. 2020;395(10227):899–911. doi:10.1016/S0140-6736(20)30059-3
- Schaumburg K, Zerwas S, Goodman E, Yilmaz Z, Bulik CM, Micali N. Anxiety disorder symptoms at age 10 predict eating disorder symptoms and diagnoses in adolescence. *J Child Psychol Psychiatr*. 2019;60(6):686–696. doi:10.1111/jcpp.12984
- Harrison F. Eating disorders. *Lancet*. 2003;361:407–416.
- Stice EO, Marti ZA, N C. A meta-analytic review of trials that tested whether eating disorder prevention programs prevent eating disorder onset. *Clin Psychol Rev*. 2021;87:102046. doi:10.1016/j.cpr.2021.102046
- Friborg O, Martinussen M, Kaiser S, Overgrd KT, Rosenvinge JH. Personality disorders in eating disorder not otherwise specified and binge eating disorder. *J Nerv Mental Dis*. 2014;202(2):119–125. doi:10.1097/NMD.0000000000000080
- Gruza RA, Przybeck TR, Cloninger CR. Prevalence and correlates of binge eating disorder in a community sample. *Comprehensive Psychiatry*. 2007;48(2):124–131. doi:10.1016/j.comppsy.2006.08.002
- Micali N, Martini MG, Thomas JJ, et al. Lifetime and 12-month prevalence of eating disorders amongst women in mid-life: a population-based study of diagnoses and risk factors. *BMC Med*. 2017;15(1):12. doi:10.1186/s12916-016-0766-4
- Hudson JI, Hiripi E, Pope HG Jr, Kessler RC. The prevalence and correlates of eating disorders in the national comorbidity survey replication. *Biol Psychiatry*. 2007;61(3):348–358. doi:10.1016/j.biopsych.2006.03.040
- Swanson SA, Crow SJ, Le Grange D, Swendsen J, Merikangas KR. Prevalence and correlates of eating disorders in adolescents. Results from the national comorbidity survey replication adolescent supplement. *Arch Gen Psychiatry*. 2011;68(7):714–723. doi:10.1001/archgenpsychiatry.2011.22
- Comerci GD. Medical complications of anorexia nervosa and bulimia nervosa. *Med Clin North Am*. 1990;74(5):1293–1310. doi:10.1016/S0025-7125(16)30517-X
- Mitchison D, Mond J, Bussey K, et al. DSM-5 full syndrome, other specified, and unspecified eating disorders in Australian adolescents: prevalence and clinical significance. *Psychol Med*. 2020;50(6):981–990. doi:10.1017/S0033291719000898
- Solmi F, Hotopf M, Hatch SL, Treasure J, Micali N. Eating disorders in a multi-ethnic inner-city UK sample: prevalence, comorbidity and service use. *Soc Psychiatry Psychiatr Epidemiol*. 2016;51(3):369–381. doi:10.1007/s00127-015-1146-7
- Li S, Song L, Twayigira M, Fan T, Luo X, Shen Y. Eating disorders among middle school students in a Chinese population: prevalence and associated clinical correlates. *J Psychiatr Res*. 2022;154:278–285. doi:10.1016/j.jpsychires.2022.07.062
- Eielsen HP, Ulvenes P, Hoffart A, Ø R, Rosenvinge JH, Vrabel K. Childhood trauma and outcome trajectories in patients with longstanding eating disorders across 17 years. *Int J Eat Disord*. 2024;57(1):81–92. doi:10.1002/eat.24067
- Caslini M, Bartoli F, Crocamo C, Dakanalis A, Clerici M, Carrà G. Disentangling the association between child abuse and eating disorders: a systematic review and meta-analysis. *Psychosomatic Med*. 2016;78(1):79–90. doi:10.1097/PSY.0000000000000233
- Molendijk ML, Hoek HW, Brewerton TD, Elzinga BM. Childhood maltreatment and eating disorder pathology: a systematic review and dose-response meta-analysis. *Psychol Med*. 2017;47(8):1402–1416. doi:10.1017/S0033291716003561
- Talmon A, Widom CS. Childhood maltreatment and eating disorders: a prospective investigation. *Child Maltreat*. 2022;27(1):88–99. doi:10.1177/1077559520988786
- Burns EE, Fischer S, Jackson JL, Harding HG. Deficits in emotion regulation mediate the relationship between childhood abuse and later eating disorder symptoms. *Child Abuse Negl*. 2012;36(1):32–39. doi:10.1016/j.chiabu.2011.08.005
- Dawson D, Strodl E, Kitamura H. Childhood maltreatment and disordered eating: the mediating role of emotion regulation. *Appetite*. 2022;172:105952. doi:10.1016/j.appet.2022.105952
- Monteleone AM, Cascino G, Pellegrino F, et al. The association between childhood maltreatment and eating disorder psychopathology: a mixed-model investigation. *Eur Psychiatry*. 2019;61:111–118. doi:10.1016/j.eurpsy.2019.08.002

21. Edwards JR, Lambert LS. Methods for integrating moderation and mediation: a general analytical framework using moderated path analysis. *Psychol Methods*. 2007;12(1):1–22. doi:10.1037/1082-989X.12.1.1
22. Arrindell W, Emmelkamp P, Monsma A, Brilman E. The role of perceived parental rearing practices in the aetiology of phobic disorders: a controlled study. *British J Psychiatry*. 1983;143(2):183–187. doi:10.1192/bjp.143.2.183
23. Arrindell W, Kwee M, Methorst G, van der Ende J, Pol E, Moritz B. Perceived parental rearing styles of agoraphobic and socially phobic in-patients. *British J Psychiatry*. 1989;155(4):526–535. doi:10.1192/bjp.155.4.526
24. Simon N, Herlands N, Marks E, et al. Childhood maltreatment linked to greater symptom severity and poorer quality of life and function in social anxiety disorder. *Depression Anxiety*. 2009;26(11):1027–1032. doi:10.1002/da.20604
25. Bruce L, Heimberg R, Blanco C, Schneier F, Liebowitz M. Childhood maltreatment and social anxiety disorder: implications for symptom severity and response to pharmacotherapy. *Depression Anxiety*. 2012;29(2):131–138. doi:10.1002/da.20909
26. Zhao J, Sun R, Shangguan M. Childhood psychological maltreatment and social anxiety in college students: the roles of parasymphathetic nervous system activity and parent-child separation experience. *Child Abuse Negl*. 2024;151:106723. doi:10.1016/j.chiabu.2024.106723
27. Hinrichsen H, Sheffield A, Waller G. The role of parenting experiences in the development of social anxiety and agoraphobia in the eating disorders. *Eating Behav*. 2007;8(3):285–290. doi:10.1016/j.eatbeh.2006.11.003
28. Bulik CM. Anxiety, depression, and eating disorders. *Eating Disord Obesity*. 2002;2(1):193–198.
29. Schwalberg M, Barlow D, David H, Howard LJ. Comparison of bulimics, obese binge eaters, social phobics, and individuals with panic disorder. *J Abnormal Psychol*. 1992;101(4):675–681. doi:10.1037/0021-843X.101.4.675
30. Schaefer LM, Thompson JK. Self-objectification and disordered eating: a meta-analysis. *Int J Eating Disorders*. 2018;51(6):483–502. doi:10.1002/eat.22854
31. Kathleen M, Griffiths B, Jones A. Self-objectification and depression: an Integrative systematic review. *J Affective Disord*. 2014;171:22–32.
32. Sabiston CM, Pila E, Pinsonnault-Bilodeau G, Cox AE. Social physique anxiety experiences in physical activity: a comprehensive synthesis of research studies focused on measurement, theory, and predictors and outcomes. *Int Rev Sport Exercise Psychol*. 2014;7(1):158–183. doi:10.1080/1750984X.2014.904392
33. Kerr-Gaffney J, Harrison A, Tchanturia K. Social anxiety in the eating disorders: a systematic review and meta-analysis. *Psychol Med*. 2018;48(15):2477–2491. doi:10.1017/S0033291718000752
34. Bulik CM, Sullivan PF, Fear JL, Joyce PR. Eating disorders and antecedent anxiety disorders: a controlled study. *Acta Psychiatr Scand*. 1997;96(2):101–107. doi:10.1111/j.1600-0447.1997.tb09913.x
35. Godart NT, Flament MF, Lecrubier Y, Jeammet P. Anxiety disorders in anorexia nervosa and bulimia nervosa: co-morbidity and chronology of appearance. *Eur Psychiatry*. 2000;15(1):38–45. doi:10.1016/S0924-9338(00)00212-1
36. Tasca GA. Attachment and eating disorders: a research update. *Curr Opin Psychol*. 2019;25:59–64. doi:10.1016/j.copsyc.2018.03.003
37. Cooper ML, Shaver PR, Collins NL. Attachment styles, emotion regulation, and adjustment in adolescence. *J Pers Soc Psychol*. 1998;74(5):1380–1397. doi:10.1037/0022-3514.74.5.1380
38. Fraley RC, Shaver PR. Adult romantic attachment: theoretical developments, emerging controversies, and unanswered questions. *Rev General Psychol*. 2000;4(2):132–154. doi:10.1037/1089-2680.4.2.132
39. Dakanalis A, Timko CA, Zanetti MA, et al. Attachment insecurities, maladaptive perfectionism, and eating disorder symptoms: a latent mediated and moderated structural equation modeling analysis across diagnostic groups. *Psychiatry Res*. 2014;215(1):176–184. doi:10.1016/j.psychres.2013.10.039
40. Monteleone AM, Cardi V, Volpe U, et al. Attachment and motivational systems: relevance of sensitivity to punishment for eating disorder psychopathology. *Psychiatry Res*. 2018;260:353–359. doi:10.1016/j.psychres.2017.12.002
41. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep*. 1985;100(2):126–131.
42. World Health Organisation. Global recommendations on physical activity for health. 2010.
43. McMahon EM, Corcoran P, O'Regan G, et al. Physical activity in European adolescents and associations with anxiety, depression and well-being. *Eur Child Adolesc Psychiatry*. 2017;26(1):111–122. doi:10.1007/s00787-016-0875-9
44. Campbell TS, Donoghue KM, Ghosh U, Nelson CM, Roth TL. Early life stress affects bdnf regulation: a role for exercise interventions. *Int J Mol Sci*. 2022;23(19):11729. doi:10.3390/ijms231911729
45. Ren Y, Li M. Influence of physical exercise on social anxiety of left-behind children in rural areas in China: the mediator and moderator role of perceived social support. *J Affect Disord*. 2020;266:223–229. doi:10.1016/j.jad.2020.01.152
46. Brunet J, Sabiston CM. Social physique anxiety and physical activity: a self-determination theory perspective. *Psychol Sport Exercise*. 2009;10(3):329–335. doi:10.1016/j.psychsport.2008.11.002
47. Merom D, Phongsavan P, Wagner R, et al. Promoting walking as an adjunct intervention to group cognitive behavioral therapy for anxiety disorders—a pilot group randomized trial. *J Anxiety Disord*. 2008;22(6):959–968. doi:10.1016/j.janxdis.2007.09.010
48. Rizk M, Mattar L, Kern L, et al. Physical activity in eating disorders: a systematic review. *Nutrients*. 2020;12(1). doi:10.3390/nu12010183
49. Wang H, He X, Tang Y, Tang J, Yang J. Unveiling the links between physical activity, self-identity, social anxiety, and emotional eating among overweight and obese young adults. *Front Psychol*. 2023;14:1255548. doi:10.3389/fpsyg.2023.1255548
50. Mathisen TF, Rosenvinge JH, Friberg O, et al. Is physical exercise and dietary therapy a feasible alternative to cognitive behavior therapy in treatment of eating disorders? A randomized controlled trial of two group therapies. *Int J Eat Disord*. 2020;53(4):574–585. doi:10.1002/eat.23228
51. Bernstein DP, Stein JA, Newcomb MD, et al. Development and validation of a brief screening version of the childhood trauma questionnaire. *Child Abuse Negl*. 2003;27(2):169–190. doi:10.1016/S0145-2134(02)00541-0
52. Bernstein DP, Ahluvalia T, Pogge D, Handelsman L. Validity of the Childhood Trauma Questionnaire in an adolescent psychiatric population. *J Am Acad Child Adolesc Psychiatry*. 1997;36(3):340–348. doi:10.1097/00004583-199703000-00012
53. Yao SQ. Initial reliability and validity of childhood trauma questionnaire(CTQ-SF) applied in Chinese college students. *Chin J Clin Psychol*. 2005.
54. He J, Zhong X, Gao Y, Xiong G, Yao S. Psychometric properties of the Chinese version of the childhood trauma questionnaire-short form (CTQ-SF) among undergraduates and depressive patients. *Child Abuse Negl*. 2019;91:102–108. doi:10.1016/j.chiabu.2019.03.009
55. Garfinkel P, Newman A. The eating attitudes test: twenty-five years later. *Eating Weight Disord*. 2001;6(1):1–24. doi:10.1007/BF03339747

56. Wang J, Hao QH, Peng W, Tu Y, Zhang L, Zhu TM. Relationship between smartphone addiction and eating disorders and lifestyle among Chinese college students. *Front Public Health*. 2023;11:1111477.
57. Garner DM, Olmsted MP, Bohr Y, Garfinkel PE. The eating attitudes test: psychometric features and clinical correlates. *Psychol Med*. 1982;12(4):871–878. doi:10.1017/S0033291700049163
58. EAT-26 Self-Test; n.d. Available from: <https://www.eat-26.com>. Accessed November 05, 2024.
59. Kang Q, Chan RCK, Li X, et al. Psychometric properties of the Chinese version of the eating attitudes test in young female patients with eating disorders in Mainland China. *Eur Eat Disord Rev*. 2017;25(6):613–617. doi:10.1002/erv.2560
60. Bull FC, Al-Ansari SS, Biddle S, et al. World Health Organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med*. 2020;54(24):1451–1462. doi:10.1136/bjsports-2020-102955
61. Underwood J, Brener N, Thornton J, et al. Overview and methods for the youth risk behavior surveillance system - United States, 2019. *MMWR Supplements*. 2020;69(1):1–10. doi:10.15585/mmwr.su6901a1
62. H Zhu. A study on the relationship of attachment to social anxiety among adolescents. In *Edited by Chongqing CSU*. 2008.
63. Greca A, Lopez N. Social anxiety among adolescents: linkages with peer relations and friendships. *J Abnormal Child Psychol*. 1998;26(2):83–94. doi:10.1023/A:1022684520514
64. Hou S, Twayigira M, Luo X, et al. The relationship between emotional neglect and non-suicidal self-injury among middle school students in China: the mediating role of social anxiety symptoms and insomnia. *BMC Psychiatry* 2023, 23(1):248.
65. Czaia J, Rief W, Hilbert A. Emotion regulation and binge eating in children. *Int J Eat Disord*. 2009;42(4):356–362. doi:10.1002/eat.20630
66. Mills P, Newman EF, Cossar J, Murray G. Emotional maltreatment and disordered eating in adolescents: testing the mediating role of emotion regulation. *Child Abuse Negl*. 2015;39:156–166. doi:10.1016/j.chiabu.2014.05.011
67. Barone E, Carfagno M, Cascino G, et al. Childhood maltreatment, alexithymia and eating disorder psychopathology: a mediation model. *Child Abuse Negl*. 2023;146:106496. doi:10.1016/j.chiabu.2023.106496
68. Tasca GA, Ritchie K, Zachariades F, et al. Attachment insecurity mediates the relationship between childhood trauma and eating disorder psychopathology in a clinical sample: a structural equation model. *Child Abuse Negl*. 2013;37(11):926–933. doi:10.1016/j.chiabu.2013.03.004
69. O’Loughlin E, Galligan R, Grant S. Childhood maltreatment, shame, psychological distress, and binge eating: testing a serial mediational model. *J Eat Disord*. 2023;11(1):96. doi:10.1186/s40337-023-00819-7
70. Jewell T, Collyer H, Gardner T, et al. Attachment and mentalization and their association with child and adolescent eating pathology: a systematic review. *Int J Eat Disord*. 2016;49(4):354–373. doi:10.1002/eat.22473
71. Gibb BE, Chelminski I, Zimmerman M. Childhood emotional, physical, and sexual abuse, and diagnoses of depressive and anxiety disorders in adult psychiatric outpatients. *Depression Anxiety*. 2007;24(4):256–263. doi:10.1002/da.20238
72. Kuo JR, Goldin PR, Werner K, Heimberg RG, Gross JJ. Childhood trauma and current psychological functioning in adults with social anxiety disorder. *J Anxiety Disord*. 2011;25(4):467–473. doi:10.1016/j.janxdis.2010.11.011
73. Hazan C, Shaver P. Romantic love conceptualized as an attachment process. *J Pers Soc Psychol*. 1987;52(3):511–524. doi:10.1037/0022-3514.52.3.511
74. Courty A, Godart N, Lalanne C, Berthoz S. Alexithymia, a compounding factor for eating and social avoidance symptoms in anorexia nervosa. *Compr Psychiatry*. 2015;56:217–228. doi:10.1016/j.comppsy.2014.09.011
75. Levinson CA, Brosos LC, Vanzhula I, et al. Social anxiety and eating disorder comorbidity and underlying vulnerabilities: using network analysis to conceptualize comorbidity. *Int J Eat Disord*. 2018;51(7):693–709. doi:10.1002/eat.22890
76. Zika M, Becker L. Physical activity as a treatment for social anxiety in clinical and non-clinical populations: a systematic review and three meta-analyses for different study designs. *Front Human Neurosci*. 2021;15:653108. doi:10.3389/fnhum.2021.653108
77. Jazaieri H, Goldin P, Werner K, Ziv M, Gross J. A randomized trial of MBSR versus aerobic exercise for social anxiety disorder. *J Clin Psychol*. 2012;68(7):715–731. doi:10.1002/jclp.21863
78. Yu H, Li F, Hu Y, et al. Improving the metabolic and mental health of children with obesity: a school-based nutrition education and physical activity intervention in Wuhan, China. *Nutrients*. 2020;12(1):194. doi:10.3390/nu12010194
79. Shin YJ, Lee JH. Effects of circuit training according to the feedback type on psychological and physical health of workers with social anxiety disorder. *Iran J Public Health*. 2018;47(Suppl 1):65–73.
80. ten Have M, de Graaf R, Monshouwer K. Physical exercise in adults and mental health status: findings from the Netherlands mental health survey and incidence study (NEMESIS). *J Psychosomatic Res*. 2011;71(5):342–348. doi:10.1016/j.jpsychores.2011.04.001
81. Zink J, Belcher B, Kechter A, Stone M, Leventhal A. Reciprocal associations between screen time and emotional disorder symptoms during adolescence. *Preventive Med Reports*. 2019;13:281–288. doi:10.1016/j.pmedr.2019.01.014
82. Lamarche L, Gammage KL. The effects of leader gender on self-presentational concerns in exercise. *Psychol Health*. 2010;25(7):769–781. doi:10.1080/08870440902866886
83. Lamarche L, Gammage KL, Strong HA. The effect of mirrored environments on self-presentational efficacy and social anxiety in women in a step aerobics class. *Psychol Sport Exercise*. 2009;10(1):67–71. doi:10.1016/j.psychsport.2008.06.006
84. Nahm J-W, Kim M-J, Kim Y-A, Kim T. The development of resistance exercise prescription program algorithm for adult health management on IT environment. *J Korean Soc Living Environm Syst*. 2015;22(5):777. doi:10.21086/ksles.2015.10.22.5.777
85. Bratland-Sanda S, Rosenvinge J, Vrabel K, et al. Physical activity in treatment units for eating disorders: clinical practice and attitudes. *Eating Weight Disord*. 2009;14(2–3):e106–112. doi:10.1007/BF03327807
86. Deboer L, Tart C, Presnell K, Powers M, Baldwin A, Smits J. Physical activity as a moderator of the association between anxiety sensitivity and binge eating. *Eating Behav*. 2012;13(3):194–201. doi:10.1016/j.eatbeh.2012.01.009
87. Sundgot-Borgen J, Rosenvinge JH, Bahr R, Schneider LS. The effect of exercise, cognitive therapy, and nutritional counseling in treating bulimia nervosa. *Med Sci Sports Exerc*. 2002;34(2):190–195. doi:10.1097/00005768-200202000-00002
88. Duesund L, Skårderud F. Use the body and forget the body: treating anorexia nervosa with adapted physical activity. *Clin Child Psychol Psychiatry*. 2003;8(1):53–72. doi:10.1177/135910450300800100

Psychology Research and Behavior Management

Dovepress

Publish your work in this journal

Psychology Research and Behavior Management is an international, peer-reviewed, open access journal focusing on the science of psychology and its application in behavior management to develop improved outcomes in the clinical, educational, sports and business arenas. Specific topics covered in the journal include: Neuroscience, memory and decision making; Behavior modification and management; Clinical applications; Business and sports performance management; Social and developmental studies; Animal studies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/psychology-research-and-behavior-management-journal>