

Supplementary Table S1. Non-ocular AEs and SAEs

n (%)	All patients N=144
Non-ocular AE	28 (19.4)
Device-related AE	0
Surgery-related AE	0
SAE	6 (4.2)
Most common AEs	
COVID-19	3 (2.1)
Nasopharyngitis	3 (2.1)
Odontopathy	3 (2.1)

AE, adverse event; SAE, serious adverse event.

Supplementary Table S2. Severe visual disturbances reported preoperatively and at 6 and 12 months postoperative^a

Visual disturbance, n (%)	Preoperative (n=142)	Postoperative	
		6 months (n=135)	12 months (n=135)
Starbursts	14 (9.9)	13 (9.6)	21 (15.6)
Halos	12 (8.5)	9 (6.7)	6 (4.4)
Glare	8 (5.6)	6 (4.4)	3 (2.3)
Hazy vision	16 (11.3)	3 (2.2)	1 (0.7)
Blurred vision	18 (12.8)	3 (2.2)	2 (1.5)
Double vision	8 (5.7)	0	0

^aData were reported from the Questionnaire for Visual Disturbances.

Supplementary Table S3. Spectacle independence preoperatively and at 6 and 12 months postoperative in dim and bright conditions^a

Distance	Preoperative, n (%) (n=140)	Postoperative	
		6 months, n (%) (n=142)	12 months, n (%) (n=142)
Near, 40 cm			
Overall	46 (32.9)	116 (86.6)	115 (86.5)
Bright conditions	48 (34.3)	121 (90.3)	117 (88.0)
Dim conditions	53 (37.9)	117 (87.3)	116 (87.2)
Intermediate, 80 cm			
Overall	53 (37.9)	126 (94.0)	120 (90.2)
Bright conditions	51 (36.4)	126 (94.0)	122 (91.7)
Dim conditions	52 (37.1)	123 (91.8)	121 (91.0)
Far, 5 m			
Overall	66 (47.1)	125 (93.3)	125 (94.0)
Bright conditions	64 (45.7)	128 (95.5)	125 (94.0)
Dim conditions	62 (44.3)	121 (90.3)	121 (91.0)

Data were reported from the Intraocular Lens Satisfaction Questionnaire.

^aProportion of patients answering “never” to the following: In the past 7 days, how often did you need to wear eyeglasses to see?

Supplementary Figure S1. Mean \pm SD monocular DCIVA (A), UIVA (B), DCNVA (C), and UNVA (D) after surgery. DCIVA, distance corrected intermediate visual acuity; DCNVA, distance corrected near visual acuity; logMAR, logarithm of the minimum angle of resolution; UIVA, uncorrected intermediate visual acuity; CNVA, uncorrected near visual acuity.

