

No associations between aromatase gene polymorphisms and breast cancer risk in Saudi patients

Mohammed Alanazi¹
 Huda A Alabdulkarim²
 Jilani P Shaik¹
 Abdulrahman Al Naeem³
 Mohammad Elrobh¹
 Abdullah Al Amri¹
 Fatimah Basil al-Mukaynizi¹
 Abdelhabib Semlali¹
 Arjumand Warsy¹
 Narasimha Reddy Parine¹

¹Department of Biochemistry, College of Science, King Saud University,
²Comprehensive Cancer Center at King Fahad Medical City, ³Department of Women's Imaging, King Fahad Medical City, Riyadh, Kingdom of Saudi Arabia

Background: Cytochrome P450 (CYP)19A1 encodes aromatase, the enzyme responsible for the conversion of androgens to estrogens, and may play a role in variation in outcomes among women with breast cancer. The aim of this study was to analyze the genetic association of rs4646 (A > C) and rs700518 (Val > Val) in the CYP19A1 gene with the risk of breast cancer.

Methods: These two single nucleotide polymorphisms (SNPs) were analyzed in a primary study group of breast cancer patients and healthy control subjects. Genotypes were determined by the TaqMan SNP analysis technique. The study data were analyzed using the chi-square or *t*-test and logistic regression analysis by Statistical Package for the Social Sciences version 16 software.

Results: rs4646 and rs700518 had no association with susceptibility to breast cancer. There was no significant association for either of these SNPs overall in breast cancer samples when compared with healthy control samples. Our data do not support a relationship between the CYP19A1 rs4646 and rs700518 SNPs and risk of breast cancer. It may be that there are ethnic differences with regard to this relationship.

Conclusion: This study demonstrated that CYP19A1 rs4646 and rs700518 SNPs may not be involved in the etiology of breast cancer in the Saudi population. Confirmation of our findings in larger populations of other ethnicities could provide evidence for the role of the CYP19A1 gene in breast carcinomas.

Keywords: CYP19A1, rs4646, rs700518, breast cancer, genetic polymorphisms

Introduction

Breast cancer is one of the most common types of malignancy and is a leading cause of death worldwide, accounting for nearly one million new cases diagnosed and half a million deaths annually.¹ Incidence rates vary widely according to region, being high in developed countries and lower in developing countries.² This type of cancer is obviously a significant public health problem in Saudi Arabia.

Cytochrome P450 (CYP)19A1 encodes the enzyme aromatase, which catalyzes the conversion of the C19 androgens, androstenedione and testosterone, to estrone and estradiol, respectively.^{3,4} Specific single nucleotide polymorphisms (SNPs) in the intronic regions of CYP19A1 have been shown to play a role in altering regulation of transcription and/or splicing of CYP19A1, producing different enzyme products with variable enzymatic activity when compared with the normal gene product.^{5,6} Studies have identified SNPs in CYP19A1 that are associated with an increased risk of cancer, primarily in European American, North Indian, and Chinese populations.^{7,8} Several studies have been done on polymorphisms in the CYP19 gene in an attempt to identify

Correspondence: Narasimha Reddy Parine
 Genome Research Chair, Department of Biochemistry, College of Science, PO Box 2455, King Saud University, Riyadh 11451, Kingdom of Saudi Arabia
 Tel +966 11467 5802
 Fax +966 11467 5802
 Email redyparine@gmail.com

an association between genetic variations and breast cancer risk, but few studies have found such an association. One report suggested that a tetra-nucleotide repeat polymorphism in intron 4 (TTTA)_n was strongly associated with the risk of breast cancer.⁹ However, studies of other genetic variations have not shown a clear association with risk of breast cancer, thus generating a situation of inconsistent results.^{10–13} Hence, in the present population-based, case–control study, we investigated the genotype distribution of rs4646 and rs700518 SNPs in patients with breast cancer.

Materials and methods

Study population

The study population comprised 148 females (median age 48 years) suffering from breast cancer and attending the outpatient clinics of the clinical co-investigators at King Fahad Medical City Hospital, Riyadh, Saudi Arabia, and 154 age-matched normal healthy controls also attending King Fahad Medical City Hospital for minor illnesses and recruited following physical examinations after exclusion of a breast cancer diagnosis and history of cancer or cancer-related diseases. The patients and controls were of Saudi Arabian ethnicity. Demographic data, age at diagnosis, tumor grade, and immunohistochemical determination of estrogen receptor (ER), progesterone receptor (PR), and HER2 receptor status were recorded. The study was approved by the institutional review board of King Khalid University Hospital. Written informed consent was obtained from all participants.

Genotyping

Genomic DNA was extracted from blood samples taken from breast cancer cases and controls using a QIAmp DNA blood mini kit (Qiagen, Valencia, CA, USA) following the manufacturer's instructions. The rs1130409 SNP in the APE1

gene was genotyped using the TaqMan allelic discrimination assay as described previously.¹⁴ Ten percent of the samples were subjected to repeated analysis for verification of the genotyping procedures.¹⁵

Statistical analysis

Genotype and allelic frequencies were compared using Fisher's Exact test (two-tailed) as described by Alanazi et al¹⁵ to describe the χ^2 test and odds ratios, and 95% confidence intervals to know the variation between cancer cases and controls. The statistical analysis was performed using Statistical Package for the Social Sciences version 21.0 (SPSS Inc, Chicago, IL).¹⁶

Results

One hundred and forty-eight breast cancer cases and 154 healthy controls were included in this study. The clinical characteristics of the breast cancer cases and the healthy controls are given in Table 1. Of 148 confirmed cases of breast cancer, 67 were ER-positive, 78 were ER-negative, 89 were PR-positive, 59 were PR-negative, 57 were HER2-positive, and 89 were HER2-negative.

All the genotypic distributions were consistent with that expected in the Hardy–Weinberg model. The homozygous wild-type allele was used as a reference to calculate the odds of acquiring breast cancer in comparison with the other two genotypes. The genotype frequencies of the analyzed SNPs along with the resulting odds ratio and significance levels are shown in Table 1. We did not find any significant association between the two SNPs and risk of breast cancer.

In the present study, we found no association with CYP19A1 rs4646 and rs700518 genotypes between breast cancer cases and matched healthy controls. The frequencies of rs4646 (A > C) genotypes in breast cancer cases were 8 (0.05), 46 (0.31), and 94 (0.64), respectively, whereas in

Table 1 Genotype frequencies of *Aromatase* gene polymorphism in breast cancer cases and controls

SNP	Variant	Cases	Controls	OR	CI	χ^2 value	P-value
rs4646	AA	8 (0.05)	8 (0.05)	Ref			
	AC	46 (0.31)	47 (0.31)	0.979	0.339–2.828	0.002	0.96831
	CC	94 (0.64)	99 (0.64)	0.949	0.350–2.625	0.01	0.93490
	AC + CC	140 (0.95)	146 (0.95)	0.959	0.350–2.625	0.01	0.93490
	A	62 (0.20)	63 (0.20)	Ref			
	C	234 (0.80)	245 (0.80)	0.971	0.655–1.439	0.02	0.88154
rs700518 Val > Val	AA	60 (0.41)	55 (0.36)	Ref			
	AG	66 (0.45)	71 (0.46)	0.852	0.519–1.400	0.40	0.52715
	GG	22 (0.15)	28 (0.18)	0.720	0.369–1.404	0.93	0.33451
	AG + GG	88 (0.59)	99 (0.64)	0.815	0.512–1.297	0.75	0.38789
	A	186 (0.63)	181 (0.59)	Ref			
	G	110 (0.37)	127 (0.41)	0.843	0.608–1.169	1.05	0.30561

Abbreviations: CI, confidence interval; OR, odds ratio; Ref, reference; SNP, single nuclear polymorphism.

healthy controls the frequencies were 8 (0.05), 47 (0.30), and 99 (0.65), respectively. Breast cancer patients did not show any risk when compared with healthy individuals (Table 1). As shown in Table 1, the frequency of the rs700518, (A > G) A/A, A/G, and G/G genotypes were 60 (0.41), 66 (0.45), and 22 (0.15), respectively, in breast cancer patients and 55 (0.36), 71 (0.46), and 28 (0.18), respectively, in controls. Breast cancer patients did not show any risk when compared with healthy individuals (Table 1).

The correlation between CYP19A1 rs4646 and rs700518 SNP status and clinicopathological characteristics was also analyzed, but interestingly none of these parameters showed an association with breast cancer in this Saudi population. Breast cancer patients did not show any association between age and ER-positive, ER-negative, PR-positive, HER2-positive, and HER2-negative status (Tables 2–5).

We also generated a regional linkage disequilibrium (LD) plot using SNAP (SNP Annotation and Proxy Search, <http://www.broadinstitute.org/mpg/snap/ldplot.php>) for CYP19A1 rs4646 and rs700518. The LD plot indicated that there are multiple loci near rs4646 and rs700518 with high

LD ($r^2 > 0.8$), which suggests that fine mapping is necessary to evaluate the genetic effect of CYP19A1 on cancer as well as functional studies (Figures 1 and 2).

Discussion

Numerous genetic variations of CYP19A1 have been reported, even though the possible functional significance of most of these polymorphisms is still undefined. Various studies conducted in specific breast cancer populations have analyzed a few common CYP19A1 polymorphisms and generated inconsistent results with regard to their possible association with cancer risk, sex hormone levels, HER2 status, and survival.^{9,17–20} In the present study, our data do not support any appreciable association between CYP19A1 rs4646 and rs700518 genotypes and breast cancer risk in the Saudi population (Table 1). We analyzed the association of both these SNPs with various clinical parameters and found no evidence of heterogeneity for either rs4646 or rs700518 in Saudi breast cancer patients, and found no association except with rs4646 in PR-negative patients (Tables 2–5).

Table 2 Genotype frequencies of Aromatase gene polymorphism in breast cancer cases (above 48 years old versus below 48 years old)

SNP	Variant	Cases	Controls	OR	CI	χ^2 value	P-value	
rs4646		Above 48						
	AA	4 (0.06)	1 (0.01)	Ref				
	AC	21 (0.30)	20 (0.29)	0.263	0.027–2.554	1.49	0.22256	
	CC	45 (0.64)	47 (0.69)	0.239	0.026–2.224	1.83	0.17573	
	AC + CC	66 (0.94)	67 (0.99)	0.246	0.027–2.262	1.78	0.18228	
	AA	29 (0.21)	22 (0.16)	Ref				
rs4646	CC	111 (0.79)	114 (0.84)	0.739	0.400–1.363	0.94	0.33150	
		Below 48						
	AA	4 (0.49)	7 (0.08)	Ref				
	AC	25 (0.44)	27 (0.31)	1.620	0.423–6.210	0.50	0.47887	
	CC	49 (0.07)	52 (0.60)	1.649	0.454–5.984	0.59	0.44337	
	AC + CC	74 (0.51)	79 (0.92)	1.639	0.461–5.830	0.58	0.44136	
rs700518	AA	33 (0.71)	41 (0.24)	Ref				
	CC	123 (0.29)	131 (0.76)	1.167	0.693–1.963	0.34	0.56148	
	Val > Val		Above 48					
		AA	34 (0.33)	24 (0.35)	Ref			
		AG	31 (0.45)	36 (0.53)	0.608	0.299–1.236	1.90	0.16804
		GG	5 (0.22)	8 (0.12)	0.441	0.129–1.515	1.74	0.18673
AG + GG		36 (0.67)	44 (0.65)	0.578	0.292–1.144	2.50	0.11416	
Val > Val	AA	99 (0.56)	84 (0.62)	Ref				
	GG	41 (0.44)	52 (0.38)	0.669	0.405–1.105	2.47	0.11581	
	rs700518		Below 48					
		AA	26 (0.0)	31 (0.36)	Ref			
		AG	35 (0.0)	39 (0.45)	1.070	0.535–2.139	0.04	0.84815
GG		17 (0.0)	16 (0.19)	1.267	0.537–2.990	0.29	0.58913	
AG + GG		52 (0.0)	55 (0.64)	1.127	0.592–2.148	0.13	0.71557	
Val > Val	AA	87 (0.0)	101 (0.59)	Ref				
	GG	69 (0.0)	71 (0.41)	1.128	0.728–1.749	0.29	0.58937	

Abbreviations: CI, confidence interval; OR, odds ratio; Ref, reference; SNP, single nuclear polymorphism.

Table 3 Genotype frequencies of *Aromatase* gene polymorphism in breast cancer cases (ER-positive versus ER-negative)

SNP	Variant	Cases	Controls	OR	CI	χ^2 value	P-value
rs4646	ER-positive						
	AA	2 (0.03)	8 (0.05)	Ref			
	AC	18 (0.27)	47 (0.31)	1.532	0.297–7.913	0.26	0.60859
	CC	47 (0.70)	99 (0.64)	1.899	0.388–9.293	0.65	0.42165
	AC + CC	65 (0.97)	146 (0.95)	1.781	0.368–8.618	0.53	0.46758
	AA	22 (0.16)	63 (0.20)	Ref			
rs4646	ER-negative						
	CC	112 (0.84)	245 (0.80)	1.309	0.767–2.234	0.98	0.32231
	AA	6 (0.08)	8 (0.05)	Ref			
	AC	26 (0.33)	47 (0.31)	0.738	0.231–2.357	0.26	0.60680
	CC	46 (0.59)	99 (0.64)	0.620	0.203–1.889	0.72	0.39647
	AC + CC	72 (0.92)	146 (0.95)	0.658	0.220–1.966	0.57	0.45044
rs700518	ER-positive						
	Val > Val	30 (0.45)	55 (0.36)	Ref			
	AG	32 (0.48)	71 (0.46)	0.775	0.418–1.435	0.66	0.41646
	GG	5 (0.07)	28 (0.18)	0.458	0.179–1.173	2.72	0.09915
	AG + GG	37 (0.55)	99 (0.64)	0.685	0.382–1.228	1.62	0.20312
	AA	92 (0.69)	181 (0.59)	Ref			
rs700518	ER-negative						
	Val > Val	42 (0.31)	127 (0.41)	0.697	0.455–1.067	2.78	0.09567
	AA	28 (0.36)	55 (0.36)	Ref			
	AG	33 (0.42)	71 (0.46)	0.913	0.494–1.688	0.08	0.77148
	GG	17 (0.22)	28 (0.18)	1.193	0.560–2.538	0.21	0.64739
	AG + GG	50 (0.64)	99 (0.64)	0.992	0.562–1.751	0.001	0.97807
	AA	89 (0.57)	181 (0.59)	Ref			
	GG	67 (0.43)	127 (0.41)	1.073	0.727–1.584	0.13	0.72348

Abbreviations: CI, confidence interval; ER, estrogen receptor; OR, odds ratio; Ref, reference; SNP, single nuclear polymorphism.

Table 4 Genotype frequencies of *Aromatase* gene polymorphism in breast cancer cases (PR-positive versus PR-negative)


SNP	Variant	Cases	Controls	OR	CI	χ^2 value	P-value
rs4646	PR-positive						
	AA	5 (0.06)	8 (0.05)	Ref			
	AC	28 (0.31)	47 (0.31)	0.953	0.284–3.201	0.01	0.93817
	CC	56 (0.63)	99 (0.64)	0.905	0.282–2.900	0.03	0.86660
	AC + CC	84 (0.94)	146 (0.95)	0.921	0.292–2.905	0.02	0.88768
	AA	38 (0.21)	63 (0.20)	Ref			
rs4646	PR-negative						
	CC	140 (0.79)	245 (0.80)	0.947	0.602–1.490	0.05	0.81502
	AA	3 (0.05)	8 (0.05)	Ref			
	AC	18 (0.31)	47 (0.31)	1.021	0.243–4.284	0.0001	0.97704
	CC	38 (0.64)	99 (0.64)	1.024	0.258–4.063	0.0012	0.97358
	AC + CC	56 (0.95)	146 (0.95)	1.023	0.262–3.994	0.001	0.97409
rs700518	PR-positive						
	Val > Val	24 (0.20)	63 (0.20)	Ref			
	AG	94 (0.80)	245 (0.80)	1.007	0.595–1.706	0.001	0.97888
	AA	39 (0.44)	55 (0.36)	Ref			
	AG	38 (0.43)	71 (0.46)	0.755	0.427–1.333	0.94	0.33189
	GG	12 (0.13)	28 (0.18)	0.604	0.274–1.333	1.57	0.21003
rs700518	PR-negative						
	Val > Val	50 (0.66)	99 (0.64)	0.712	0.418–1.214	1.56	0.21131
	AA	116 (0.65)	181 (0.59)	Ref			
	GG	62 (0.35)	127 (0.41)	0.762	0.519–1.117	1.95	0.16306
	AA	21 (0.36)	55 (0.36)	Ref			
	AG	28 (0.47)	71 (0.46)	1.033	0.530–2.011	0.01	0.92423
	GG	10 (0.17)	28 (0.18)	0.935	0.388–2.255	0.02	0.88168
	AG + GG	38 (0.64)	99 (0.64)	1.005	0.537–1.881	0.0012	0.98683
	AA	70 (0.59)	181 (0.59)	Ref			
	GG	48 (0.41)	127 (0.41)	0.977	0.635–1.505	0.01	0.91689

Abbreviations: CI, confidence interval; PR, progesterone receptor; OR, odds ratio; Ref, reference; SNP, single nuclear polymorphism.

Table 5 Genotype frequencies of *Aromatase* gene polymorphism in breast cancer cases (HER2-positive versus HER2-negative)

SNP	Variant	Case	Control	OR	CI	χ^2 value	P-value
rs4646	HER2-positive						
	AA	3 (0.05)	8 (0.05)	Ref			
	AC	21 (0.37)	47 (0.31)	1.191	0.287–4.945	0.06	0.80916
	CC	33 (0.58)	99 (0.64)	0.889	0.223–3.548	0.03	0.86748
	AC + CC	54 (0.95)	146 (0.95)	0.986	0.252–3.855	0.0001	0.98418
	AA	27 (0.24)	63 (0.20)	Ref			
rs4646	HER2-negative						
	AA	4 (0.04)	8 (0.05)	Ref			
	AC	25 (0.28)	47 (0.31)	1.064	0.292–3.882	0.01	0.92536
	CC	60 (0.67)	99 (0.64)	1.212	0.350–4.198	0.09	0.76122
	AC + CC	85 (0.96)	146 (0.95)	1.164	0.340–3.982	0.06	0.80817
	AA	33 (0.19)	63 (0.20)	Ref			
rs700518	HER2-positive						
	Val > Val	24 (0.42)	55 (0.36)	Ref			
	AG	25 (0.44)	71 (0.46)	0.807	0.416–1.564	0.40	0.52475
	GG	8 (0.14)	28 (0.18)	0.655	0.261–1.644	0.82	0.36534
	AG + GG	33 (0.58)	99 (0.64)	0.764	0.411–1.421	0.73	0.39436
	AA	73 (0.64)	181 (0.59)	Ref			
rs700518	HER2-negative						
	Val > Val	41 (0.36)	127 (0.41)	0.800	0.513–1.249	0.96	0.32619
	AA	35 (0.39)	55 (0.36)	Ref			
	AG	41 (0.46)	71 (0.46)	0.907	0.512–1.608	0.11	0.73934
	GG	13 (0.15)	28 (0.18)	0.730	0.334–1.596	0.63	0.42891
	AG + GG	54 (0.61)	99 (0.64)	0.857	0.500–1.468	0.32	0.57434
rs700518	HER2-negative						
	Val > Val	111 (0.62)	181 (0.59)	Ref			
	AA	67 (0.38)	127 (0.41)	0.860	0.589–1.256	0.61	0.43580

Abbreviations: CI, confidence interval; OR, odds ratio; Ref, reference; SNP, single nuclear polymorphism.

**Figure 1** Regional linkage disequilibrium plot for the single nuclear polymorphism rs4646.




Figure 2 Regional linkage disequilibrium plot for the rs700518 single nucleotide polymorphism.

Some strengths and limitations of our study should be noted. Its strengths include the population-based and prospective study design in the central region of Saudi Arabia, thereby minimizing selection bias, and the detailed review of the cancer diagnosis, thereby minimizing disease misclassification. However, our nested case-control sample was relatively small in size, which hampered our ability to evaluate specific gene-disease association. Confirmation of our findings in larger populations of women of different ethnicities could provide evidence for the role of the CYP19A1 gene in breast carcinoma.

Acknowledgment

The authors extend their appreciation to the Deanship of Scientific Research at King Saud University for funding this work through research group project RGPVPP-081.

Disclosure

The authors declare that they have no competing interests in this work.

References

1. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. GLOBOCAN 2008 v2.0, Cancer Incidence and Mortality Worldwide: IARC Cancer Base No. 10 Internet. Lyon, France: International Agency for Research on Cancer; 2010. Available from: <http://globocan.iarc.fr>. Accessed July 22, 2013.
2. Center MM, Jemal A, Smith RA, Ward E. Worldwide variations in colorectal cancer. *CA Cancer J Clin*. 2009;59(6):366–378.
3. Simpson ER, Clyne C, Rubin G, Boon WC, Robertson K, et al. Aromatase – a brief overview. *Annu Rev Physiol*. 2002;64:93–127.
4. Olson SH, Bandera EV, Orlov I. Variants in estrogen biosynthesis genes, sex steroid hormone levels, and endometrial cancer: a HuGE review. *Am J Epidemiol*. 2007;165:235–245.
5. Millar DS, Horan M, Chuzhanova NA, Cooper DN. Characterisation of a functional intronic polymorphism in the human growth hormone (GH1) gene. *Hum Genomics*. 2010;4:289–301.
6. Cooper DN. Functional intronic polymorphisms: buried treasure awaiting discovery within our genes. *Hum Genomics*. 2010;4:284–288.
7. Cai Q, Kataoka N, Li C, et al. Haplotype analyses of CYP19A1 gene variants and breast cancer risk: results from the Shanghai Breast Cancer Study. *Cancer Epidemiol Biomarkers Prev*. 2008;17:27–32.
8. Chattopadhyay S, Siddiqui S, Akhtar MS, et al. Genetic polymorphisms of ESR1, ESR2, CYP17A1, and CYP19A1 and the risk of breast cancer: a case control study from North India. *Tumour Biol*. 2014;35:4517–4527.
9. Fasching PA, Loehberg CR, Strissel PL, et al. Single nucleotide polymorphisms of the aromatase gene (CYP19A1), HER2/neu status, and prognosis in breast cancer patients. *Breast Cancer Res Treat*. 2008;112: 89–98.
10. Kristensen VN, Harada N, Yoshimura N, et al. Genetic variants of CYP19 (aromatase) and breast cancer risk. *Oncogene*. 2000;19:1329–1333.
11. Haiman CA, Hankinson SE, Spiegelman D, Brown M, Hunter DJ. No association between a single nucleotide polymorphism in CYP19 and breast cancer risk. *Cancer Epidemiol Biomarkers Prev*. 2002;11: 215–216.
12. Ralph DA, Zhao LP, Aston CE, et al. Age specific association of steroid hormone pathway gene polymorphisms with breast cancer risk. *Cancer*. 2007;109:1940–1948.
13. Zhang L, Gu L, Qian B, et al. Association of genetic polymorphisms of ER-alpha and the estradiol-synthesizing enzyme genes CYP17 and CYP19 with breast cancer risk in Chinese women. *Breast Cancer Res Treat*. 2009;114:327–338.
14. Livak KJ. Allelic discrimination using fluorogenic probes and the 5' nuclease assay. *Genet Anal*. 1999;14:143–149.
15. Alanazi M, Pathan AA, Ajaj SA, et al. DNA repair genes XRCC1, XRCC3, XPD, and OGG1 polymorphisms among the central region population of Saudi Arabia. *Biol Res*. 2013;46:161–167.
16. Alanazi M, Pathan AA, Abduljaleel Z, et al. Association between PARP-1 V762A polymorphism and breast cancer susceptibility in Saudi population. *PLoS One*. 2013;8:e85541.

17. Dunning AM, Dowsett M, Healey CS, et al. Polymorphisms associated with circulating sex hormone levels in postmenopausal women. *J Natl Cancer Inst.* 2004;96:936–945.
18. Kristensen VN, Andersen TI, Lindblom A, Erikstein B, Magnus P, Borresen-Dale AL. A rare CYP19 (aromatase) variant may increase the risk of breast cancer. *Pharmacogenetics.* 1998;8:43–48.
19. Probst-Hensch NM, Ingles SA, Diep AT, et al. Aromatase and breast cancer susceptibility. *Endocr Relat Cancer.* 1999;6:165–173.
20. Long JR, Kataoka N, Shu XO, et al. Genetic polymorphisms of the CYP19A1 gene and breast cancer survival. *Cancer Epidemiol Biomarkers Prev.* 2006;15:2115–2122.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress