



Open Access Full Text Article

ERRATUM

Erratum

Barrese V, Miceli F, Soldovieri MV, Ambrosino P, Iannotti FA, Cilio MR, Tagliatela M. Neuronal potassium channel openers in the management of epilepsy: role and potential of retigabine. *Clinical Pharmacology: Advances and Applications*. 2010;2:225–236.

On page 233, left-hand column, second paragraph, line 17 forward, it says:

“The most common reasons for discontinuation were dizziness, confusion, somnolence, and asthenia, with the highest proportion of withdrawals in patients receiving retigabine 1200 mg/kg.⁷² In RESTORE 1, where retigabine was used at 1200 mg/kg, urinary or renal disorders were observed in 12% of patients.”

In this passage, both instances of “retigabine 1200 mg/kg” should be “retigabine 1200 mg/day.”

Clinical Pharmacology: Advances and Applications

Publish your work in this journal

Clinical Pharmacology: Advances and Applications is an international, peer-reviewed, open access journal publishing original research, reports, reviews and commentaries on all areas of drug experience in humans. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.

Submit your manuscript here: <http://www.dovepress.com/clinical-pharmacology-advances-and-applications-journal>

Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.