

# Triazole derivatives with improved in vitro antifungal activity over azole drugs

Shichong Yu<sup>1,\*</sup>  
Xiaoyun Chai<sup>1,\*</sup>  
Yanwei Wang<sup>1</sup>  
Yongbing Cao<sup>2</sup>  
Jun Zhang<sup>3</sup>  
Qiuye Wu<sup>1</sup>  
Dazhi Zhang<sup>1</sup>  
Yuanying Jiang<sup>2</sup>  
Tianhua Yan<sup>4</sup>  
Qingyan Sun<sup>1</sup>

<sup>1</sup>Department of Organic Chemistry, School of Pharmacy, Second Military Medical University, Shanghai, People's Republic of China; <sup>2</sup>Drug Research Center, School of Pharmacy, Second Military Medical University, Shanghai, People's Republic of China; <sup>3</sup>Overseas Education Faculty of the Second Military Medical University, Shanghai, People's Republic of China; <sup>4</sup>Department of Pharmacology, School of Pharmacy, China Pharmaceutical University, Nanjing, People's Republic of China

\*These authors contributed equally to this work

Correspondence: Qingyan Sun  
Department of Organic Chemistry,  
School of Pharmacy, Second Military  
Medical University, Shanghai,  
People's Republic of China  
Tel +86 21 8187 1228-15  
Fax +86 21 8187 1225  
Email sqy\_2000@163.com

Tianhua Yan  
Department of Pharmacology, School of  
Pharmacy, China Pharmaceutical  
University, Majia Road 49, Nanjing 210009,  
People's Republic of China  
Email thyan2012@163.com

**Abstract:** A series of triazole antifungal agents with piperidine side chains was designed and synthesized. The results of antifungal tests against eight human pathogenic fungi in vitro showed that all the compounds exhibited moderate-to-excellent activities. Molecular docking between 8d and the active site of *Candida albicans* CYP51 was provided based on the computational docking results. The triazole interacts with the iron of the heme group. The difluorophenyl group is located in the S3 subsite and its fluorine atom (2-F) can form H-bonds with Gly307. The side chain is oriented into the S4 subsite and formed hydrophobic and van der Waals interactions with the amino residues. Moreover, the phenyl group in the side chain interacts with the phenol group of Phe380 through the formation of  $\pi$ - $\pi$  face-to-edge interactions.

**Keywords:** synthesis, CYP51, molecular docking, azole agents

## Introduction

Over the past two decades, a dramatically significant increase in fungal infections has been observed. Among these, the widespread diffusion of topical and systemic infectious diseases caused by the opportunistic pathogen *Candida albicans* is often related to the use of broad-spectrum antibiotics, immunosuppressive agents, anticancer, and anti-AIDS drugs.<sup>1,2</sup> One of the principal problems in the treatment of *Candida albicans* infections is the spread of antifungal drug resistance, mainly in patients chronically subjected to antimycotic therapy such as HIV-infected individuals.<sup>3,4</sup>

More recently, there has been an expansion in the number of antifungal drugs available. Major classes of antifungal compounds are currently in clinical use: polyenes, azole derivatives, allylamines, thiocarbamates, echinocandins, and fluoropyrimidines.<sup>5-8</sup> Despite this growing list of antifungal agents, treatment of fungal diseases remains unsatisfactory. In a word, the limitations of current antifungal drugs, increased incidence of systemic fungal infections, and rapid development of drug resistance have highlighted the need for new antifungal agents with a new structure of compounds and with fewer side effects.<sup>9-13</sup>

The current standard of therapies is the fungicidal polyenes, amphotericin B (very good antifungal activity, but liver and kidney toxicity), and the safe azoles (not fungicidal, but fungistatic). In particular, the azoles are important antifungal agents widely used in clinics.<sup>14</sup> Azoles exert antifungal activity through the inhibition of cytochrome P450 14 $\alpha$ -demethylase (CYP51), which is crucial in the process of ergosterol biosynthesis. The CYP51 enzyme contains an iron protoporphyrin unit located in its active site, which catalyzes the oxidative removal of the 14 $\alpha$ -methyl group of lanosterol by

typical monooxygenase activity.<sup>15</sup> Azole antifungal agents bind to the iron of the porphyrin and cause the blockade of the fungal ergosterol biosynthesis pathway by preventing the access of the natural substrate lanosterol to the active site of the enzyme.<sup>16</sup> The depletion of ergosterol and accumulation of 14 $\alpha$ -methylated sterols alter membrane fluidity, with concomitant reduction in activity of membrane-associated enzymes and increased permeability. The net effect is to inhibit fungal growth and replication.<sup>17</sup>

In our previous research, numerous studies on the structure-activity relationships (SAR) of antifungal azoles had been developed, and these studies had led to new compounds endowed with better biological and pharmacological properties.<sup>18–23</sup> Molecular docking studies indicated that the triazole ring, the difluorophenyl group, and the hydroxyl group were the pharmacophores of antifungal agents; the side chains can be oriented into substrate access channel 2 (FG loop) and form hydrophobic and van der Waals interactions.<sup>20,21</sup> In view of the importance of the piperazinyl for antifungal activity in our previous study,<sup>20,21</sup> we sought to investigate this further with the piperidine ring, which contains an oxygen atom. This oxygen atom could improve the flexibility of the molecule, making the side chain more easily lock into its proper position. Also, it may interact with the amino acid residues through hydrogen bonding.

We focused our attention on installing various substituted acids to form piperidinol ester. In order to study the SAR of the new compounds, we inserted aromatic groups (6a–c) and alkyl groups (compounds 7a–m) into the side chains. We also inserted halogens, and alkoxy, nitro, and cyano groups into the phenyl group (compounds 8a–r) of the side chains, respectively. Almost all of the azole antifungal drugs have halogens present, which play a very important role in these drugs. They are very useful to modulate the electronic effects on the phenyl rings of drugs. For example, chlorine has strong inductive electron-attracting effects, while those of fluorine are very weak. Moreover, these atoms may also influence the steric characteristics and the hydrophilic–hydrophobic balance of the molecules.

## Chemistry

The general synthetic methodology for the preparation of the title compounds (6a–c, 7a–m, and 8a–r) is outlined in Figure 1. As a key intermediate of our designed triazole antifungals, the oxirane compound 4 was synthesized using the reported procedure.<sup>24</sup> The intermediate 5 was synthe-

sized by ring-open reaction of oxirane 4 with piperidinol. The title compounds were synthesized and a good yield was obtained when the reaction was performed in CH<sub>2</sub>Cl<sub>2</sub> in the presence of 4-dimethylaminopyridine (DMAP) and 3-(3-dimethylaminopropyl)-1-ethylcarbodiimide hydrochloride (EDCI). All the new compounds (6a–c, 7a–m, and 8a–r) described above were characterized by infra-red (IR), high-resolution electrospray ionization mass spectra (HR-MS), and nuclear magnetic resonance (NMR) spectroscopic analysis.


## Pharmacology

The in vitro antifungal activities of all the title compounds were evaluated against eight human pathogenic fungi: *Candida albicans* (*C. alb.*), *Candida parapsilosis* (*C. para.*), *Candida tropicalis* (*C. tro.*), *Cryptococcus neoformans* (*C. neo.*), *Fonsecaea compacta* (*F. com.*), *Trichophyton rubrum* (*T. rub.*), *Microsporum gypseum* (*M. gyp.*), and *Aspergillus fumigatus* (*A. fum.*), which are often encountered clinically, and were compared with fluconazole (FCZ), itraconazole (ICZ), voriconazole (VCZ). *C. alb.* and *C. neo.* were provided by Shanghai Changzheng Hospital (Shanghai, People's Republic of China); *C. par.*, *C. tro.*, *F. com.*, *T. rub.*, *M. gyp.*, and *A. fum.* were provided by Shanghai Changhai Hospital. *C. alb.* (ATCCY0109) and *C. neo.* (ATCCBLS108) were used as the quality control strains, and tested in each assay. FCZ, ICZ, and VCZ served as the positive control, and were obtained from their respective manufacturers.

The antifungal potency was evaluated by means of the minimum inhibitory concentrations (MICs) using the serial test in the broth microdilution modification method published by the Clinical and Laboratory Standards Institute (CLSI) method M 27-A2.<sup>25</sup> The MIC was defined as the first well with an approximate 80% reduction in growth compared to the growth of the drug-free well. For assays, the title compounds to be tested were dissolved in dimethyl sulfoxide (DMSO), serially diluted in growth medium, inoculated, and incubated at 35°C. Growth MICs were determined at 24 hours for *C. alb.* and at 72 hours for *C. neo.* The data points were calculated from the mean of replicates (three times). All of our susceptibility tests were performed in triplicate using each antifungal agent.

## Materials and methods

In our studies, the 3D structures of the designed azoles were built using the Builder module within the Insight II 2000 software package of Discovery Studio 2.5 from Accelrys (San



**Figure 1** Conditions: (a)  $\text{ClCH}_2\text{COCl}$ ,  $\text{AlCl}_3$ ,  $50^\circ\text{C}$ , 5 hours, with 87.0% yield; (b)  $\text{C}_6\text{H}_5\text{CH}_2$ ,  $\text{NaHCO}_3$ , 1*H*-1,2,4-triazole, reflux, 5 hours, with 43.2% yield; (c)  $\text{C}_6\text{H}_5\text{CH}_2$ ,  $(\text{CH}_3)_3\text{SOI}$ ,  $\text{NaOH}$ , centylmethylammonium bromide,  $60^\circ\text{C}$ , 3 hours, with 54.3% yield; (d)  $\text{CH}_3\text{SO}_3\text{H}$ ,  $0^\circ\text{C}$ , 1 hour, with 89.5% yield; (e)  $\text{CH}_3\text{CH}_2\text{OH}$ ,  $\text{Et}_3\text{N}$ , 4-piperidinol, reflux, 6 hours, with 70.2% yield; (f) various acids, DMAP, EDCl,  $\text{CH}_2\text{Cl}_2$ , reflux, 8 hours, with 67.0%–85.0% yield.

Diego, CA, USA). The flexible ligand-docking procedure in the Affinity module within Insight II was used to define the lowest energy position for the substrate using a Monte Carlo docking protocol.

Melting points (Mp) were measured on a Yamato MP-21 melting-point apparatus (Tokyo, Japan) and are uncorrected. IR spectra were recorded in potassium bromide disks on a Hitachi 270-50 spectrophotometer (Tokyo, Japan).  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were recorded in  $\text{CDCl}_3$  unless otherwise indicated with a Bruker Avance II 500 spectrometer (Billerica, MA, USA), using tetramethylsilane (TMS) as the internal standard. HR-MS was performed on an Agilent 6538 Q-TOF mass spectrometer (Santa Clara, CA, USA). Thin layer chromatography (TLC) analysis was carried out on silica gel plates GF254 (Qingdao Haiyang Chemical, Qingdao, People's Republic of China). Column chromatography was performed with silica gel 60 G (Qindao Haiyang Chemical).

The solvents and reagents were used as received or dried prior to use as needed.

## Preparation of compounds

### 1-(2-[2,4-difluorophenyl]-2-hydroxy-3-[1*H*-1,2,4-triazol-1-yl]propyl)piperidin-4-ol (5)

To a stirred mixture of: 1-(2-[2,4-difluorophenyl]-2,3-epoxypropyl)-1*H*-1,2,4-triazole methanesulfonate (4) (1.65 g, 0.005 mol),  $\text{C}_2\text{H}_5\text{OH}$  (30 mL),  $\text{N}(\text{C}_2\text{H}_5)_3$  (3 mL), 4-piperidinol (2) (0.70g, 0.006 mol) were added and heated at  $70^\circ\text{C}$ – $80^\circ\text{C}$  for 5 hours. The reaction was monitored by TLC. After filtration, the filtrate was evaporated under reduced pressure. Water (30 mL) was added to the residue, which was then extracted with ethyl acetate (80 mL  $\times$  3). The extract was washed with saturated NaCl solution (20 mL  $\times$  3), dried over anhydrous  $\text{Na}_2\text{SO}_4$ , and evaporated. The residue was separated and purified readily

by chromatography on silica gel to afford compound 5 (1.23g), with a 70.2% yield. Mp: 79°C–82°C; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ: 8.12 (1H, s, triazole-C<sub>3</sub>-H), 7.79 (1H, s, triazole-C<sub>5</sub>-H), 6.65–7.23 (3H, m, Ar-H), 5.25 (1H, br, OH), 3.76–3.91 (1H, m, CH-O), 4.56 (1H, d, *J*=14.0 Hz, triazole-CHa), 4.52 (1H, d, *J*=14.0 Hz, triazole-CHb), 3.15 (1H, d, *J*=13.6 Hz, NCHa), 2.78 (1H, d, *J*=13.6 Hz, NCHb), 1.50–2.33 (8H, m, piperidine-H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ: 162.5, 161.8, 150.8, 143.1, 130.2, 115.8, 110.4, 104.6, 69.8, 63.7, 55.5, 51.7, 47.1, 46.8, 31.8, 31.4; IR (KBr): 3,420, 3,084, 2,952, 1,414, 1,363, 1,320, 1,266, 1,175, 1,134, 1,082, 1,012, 965, 885, 864, 794,

714, 672, 651, 631, 560, 514, 465; HR-MS (ESI): *m/z* calculated for C<sub>16</sub>H<sub>20</sub>F<sub>2</sub>N<sub>4</sub>O<sub>2</sub>, [M+H]<sup>+</sup> 339.1554; found: 339.1567.

### General procedure for the target compounds 6a–c, 7a–m, and 8a–r

To a stirred mixture of: 1-(2-[2,4-difluorophenyl]-2-hydroxy-3-[1*H*-1,2,4-triazol-1-yl] propyl)-piperidin-4-ol (5) (0.338 g, 0.001 mol), DMAP (100 mg) and EDCI (200 mg) in 50 mL dichloromethane at 0°C and substituted acid (0.001 mol) were added and stirred for 8–12 hours. The reaction was monitored by TLC. After filtration, the filtrate was evaporated under

**Table I** Antifungal activities of the title compounds in vitro<sup>a</sup> (MIC<sub>80</sub><sup>b</sup>, nmol/mL)

Compound	<i>C. alb.</i> Y0109	<i>C. par.</i> 0306392	<i>C. tro.</i>	<i>C. neo.</i> BLS108	<i>F. com.</i>	<i>T. rub.</i> 0501124	<i>M. gyp.</i> 0310388	<i>A. fum.</i> 0504656
6a	0.034	0.137	0.137	0.137	0.548	0.137	0.137	2.192
6b	0.126	0.032	0.126	2.019	0.505	0.505	0.126	8.078
6c	0.479	0.479	0.479	0.479	7.660	1.915	0.479	1.915
7a	0.164	2.630	0.658	0.658	2.630	0.658	0.658	10.521
7b	0.159	0.634	0.159	0.634	0.634	0.634	0.634	10.147
7c	0.153	0.153	0.153	0.153	0.612	0.612	0.153	2.450
7d	0.153	0.612	0.612	0.612	0.612	0.612	0.153	9.799
7e	0.148	0.592	0.148	0.592	0.592	0.592	0.592	9.474
7f	0.036	0.573	0.573	0.573	0.573	0.573	0.143	9.170
7g	0.009	0.139	0.139	0.139	0.555	0.555	0.139	2.221
7h	0.034	0.538	8.615	0.034	2.154	0.135	0.135	2.154
7i	0.508	2.031	0.508	0.508	2.031	2.031	0.508	2.031
7j	0.480	1.922	1.922	0.480	7.688	1.922	0.480	30.751
7k	2.304	9.214	2.304	36.858	36.858	9.214	2.304	147.431
7l	2.222	35.548	2.222	35.548	35.548	8.887	35.548	142.191
7m	0.604	9.660	2.415	9.660	9.660	9.660	9.660	9.660
8a	0.033	0.008	0.033	0.033	0.133	0.133	0.133	0.534
8b	0.032	0.129	0.129	0.129	0.514	0.129	0.129	2.057
8c	0.005	0.129	0.129	0.129	0.514	0.129	0.129	2.057
8d	0.005	0.129	0.129	0.129	0.129	0.032	0.032	0.514
8e	0.124	0.498	0.124	0.498	7.965	0.031	0.498	1.991
8f	0.124	0.124	0.498	0.498	7.965	0.498	0.498	1.991
8g	0.124	0.124	0.124	0.498	0.498	0.124	0.124	1.991
8h	0.114	0.458	0.458	0.458	1.831	0.458	0.458	1.831
8i	0.114	0.458	0.458	0.458	1.831	0.458	0.458	1.831
8j	0.007	0.458	0.458	0.458	0.458	0.458	0.114	1.831
8k	0.117	0.466	0.466	0.466	1.865	0.466	0.466	1.865
8l	0.032	0.130	0.032	0.130	0.518	0.518	0.130	0.518
8m	0.031	0.125	0.125	0.502	0.502	0.502	0.125	2.007
8n	0.030	0.118	0.118	0.473	0.473	1.893	0.473	1.893
8o	0.030	0.487	0.122	0.122	0.487	0.122	0.122	1.949
8p	0.122	0.122	0.122	0.122	0.487	0.122	0.122	0.487
8q	0.030	0.122	0.122	0.122	0.122	0.030	0.030	0.487
8r	0.032	0.507	0.127	0.127	0.507	0.127	0.127	2.028
ICZ	0.089	0.354	0.089	0.354	0.354	0.089	0.354	1.417
VCZ	0.011	0.045	0.045	0.045	0.179	0.045	0.045	0.716
FCZ	0.816	0.816	3.265	3.265	52.241	13.060	3.265	208.966

**Notes:** <sup>a</sup>*C. alb.*, *Candida albicans*; *C. par.*, *Candida parapsilosis*; *C. tro.*, *Candida tropicalis*; *C. neo.*, *Cryptococcus neoformans*; *F. com.*, *Fonsecaea compacta*; *T. rub.*, *Trichophyton rubrum*; *M. gyp.*, *Microsporum gypseum*; *A. fum.*, *Aspergillus fumigatus*, ICZ, itraconazole; VCZ, voriconazole; FCZ, fluconazole; <sup>b</sup>MIC<sub>80</sub>: 80% minimal inhibitory concentration; FCZ, fluconazole; ICZ, itraconazole; VCZ, voriconazole.

reduced pressure. The residue was then extracted with ethyl acetate (60 mL ×3). The extract was washed with saturated NaCl solution (20 mL ×3), dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, and evaporated. The residue was crystallized from ethyl acetate to afford the title compounds, with 67.0%–85.0% yield.

### 1-(2-[2,4-difluorophenyl]-2-hydroxy-3-[1H-1,2,4-triazol-1-yl]propyl)piperidin-4-yl-2-phenylacetate (6a)

Mp: 93°C–95°C; <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ: 8.13 (1H, s, triazole-C<sub>3</sub>-H), 7.78 (1H, s, triazole-C<sub>5</sub>-H), 6.80–7.54 (8H, m, Ar-H), 5.20 (1H, br, OH), 4.72–4.74 (1H, m, CH-O), 4.55 (1H, d, *J*=14.0 Hz, triazole-CHa), 4.50 (1H, d, *J*=14.0 Hz, triazole-CHb), 3.57 (2H, s, ArCH<sub>2</sub>N), 3.10 (1H, d, *J*=13.6 Hz, NCHa), 2.70 (1H, d, *J*=13.6 Hz, NCHb), 1.52–2.40 (8H, m, piperidine-H); <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ: 167.6, 152.8, 146.5, 135.7, 131.2, 130.8, 130.6, 130.2, 129.8, 128.6, 119.8, 112.2, 105.8, 105.5, 105.2, 73.8, 69.5, 63.4, 55.2, 51.9, 46.8, 32.2; IR (KBr): 3,426, 3,082, 2,957, 2,827, 1,723, 1,617, 1,499, 1,469, 1,416, 1,364, 1,320, 1,270, 1,219, 1,170, 1,137, 1,082, 1,042, 1,010, 962, 882, 861, 818, 795, 760, 710, 677, 657, 631, 577, 515, 470, 439; HR-MS (ESI): *m/z* calculated for C<sub>24</sub>H<sub>26</sub>F<sub>2</sub>N<sub>4</sub>O<sub>3</sub>, [M+H]<sup>+</sup> 457.1973; found: 457.2034.

## Results and discussion

The in vitro antifungal activities of newly synthesized azole derivatives 6a–c, 7a–m, and 8a–r against eight strains of important pathogenic fungi are reported in Table 1, together with the positive controls FCZ, ICZ, and VCZ. From the in vitro antifungal activity assay, we can see that all the synthesized compounds show moderate-to-excellent activity against nearly all the tested fungal pathogens. The MIC values of compounds 7k–7m against *C. alb.* were decreased from 2.304 nmol/mL to 0.604 nmol/mL, while compounds 7a–7j ranged from 0.009 nmol/mL to 0.480 nmol/mL. From that, we can see that the activity of type 7 would be weakened when the H atom of the alkyl group side chain is substituted by a halogen atom. Furthermore, the activity got lower when the halogen electron-withdrawing action ability was enhanced. Tested against *C. alb.*, compounds 8c and 8d showed MICs about 18-fold greater than that of ICZ (with a MIC value of 0.089 nmol/mL), and was also comparable to that of VCZ (with a MIC value of 0.011 nmol/mL). Additionally, compounds 7g and 8j showed MIC values of 0.009 nmol/mL and 0.007 nmol/mL, respectively. The activity of all the title compounds except


compounds 7k and 7l exhibited moderate activity against *A. fum.*, and showed higher activity than FCZ (with a MIC value of 208.97 nmol/mL). VCZ is an antifungal agent, which is characterized by a broad spectrum of activity in vitro and which shows fungistatic activity against fungi such as *A. fum.* Compounds 8a, 8d, 8l, 8p, and 8q exhibited similar activity against *A. fum.* when compared to VCZ (with a MIC value of 0.716 nmol/mL). Compound 8a was about six-fold more active in MIC values against *C. par.* than VCZ (with a MIC value of 0.045 nmol/mL); in addition, it was about 100-fold more potent than FCZ. Lastly, compound 6b showed a MIC value of 0.032 nmol/mL, which is comparable to the positive control VCZ, and was

**Table 2** GOLD docking scores of the compounds

Compd.	Fitness <sup>[a]</sup>	S(hb_ext) <sup>[b]</sup>	S(vdw_ext) <sup>[c]</sup>	S(int) <sup>[d]</sup>
6a	78.17	10.00	50.36	-4.36
6b	72.14	10.00	46.61	-3.64
6c	65.10	10.00	44.94	-5.94
7a	62.71	10.00	39.59	-2.45
7b	61.46	10.00	42.43	-4.92
7c	59.81	10.00	36.66	-5.01
7d	58.73	10.00	41.64	-3.54
7e	46.65	10.00	33.37	-2.72
7f	70.35	10.00	45.98	-8.14
7g	73.25	9.99	45.02	-9.02
7h	75.95	10.02	48.93	-8.30
7i	67.84	9.91	42.52	-9.97
7j	57.35	0.98	59.92	-12.74
7k	45.01	0.99	34.76	-3.06
7l	55.36	10.00	35.51	-6.25
7m	57.96	5.98	39.04	-4.22
8a	67.40	10.00	45.59	-7.14
8b	73.21	10.00	46.72	-10.98
8c	71.77	10.00	45.73	-5.86
8d	72.29	8.49	44.33	-7.96
8e	81.00	10.00	53.18	-10.62
8f	75.74	10.00	51.74	-13.38
8g	72.74	10.00	53.99	-6.20
8h	78.32	10.00	53.18	-9.25
8i	72.40	9.95	49.20	-11.49
8j	79.29	10.00	50.50	-8.60
8k	76.98	10.00	52.48	-8.07
8l	70.81	10.00	47.47	-6.72
8m	64.76	10.00	43.77	-13.54
8n	76.32	10.00	54.23	-13.53
8o	74.95	9.21	50.87	-15.00
8p	66.78	10.00	49.55	-10.18
8q	70.70	9.98	44.58	-12.44
8r	61.20	0.03	50.35	-7.03

**Notes:** [a] Fitness, the negative of the sum of the component energy terms; [b] S(hb\_ext), protein-ligand hydrogen bond energy; [c] S(vdw\_ext), protein-ligand van der Waals (vdw) energy; [d] S(int), the sum of ligand internal vdw energy and ligand torsional strain energy.





**Figure 2** Binding of compound 8d to the active site of CYP51 from *C. albicans*.

found to be a potent inhibitor of fungal growth even on fungi strains resistant to FCZ.

In this study, we docked all the compounds to clarify their binding modes with *Candida albicans* CYP51 (CACYP51) using the flexible docking in the Affinity module within the Insight II 2000 software package and which provided information for further optimization. As shown in Table 2, compounds 6a, 7g, 7h, 8b, 8c, 8d, 8j, 8n, and 8o have relatively higher fitness scores, which are consistent with their antifungal activity. In contrast, weakly active compounds such as 7i, 7j, 7k, 7l, and 7m have relatively lower fitness scores. The good correlation between antifungal activity and docking score indicates that a suitable side chain has a great impact on antifungal activity.

Figure 2 shows the binding mode of the designed compound 8d to the active site of CYP51. The active site of CYP51 was constructed in previous studies.<sup>20,21</sup> Both the *R* and *S* isomers of the compounds were docked into the active site and the interaction energies of the *R* and *S* isomers were calculated, respectively. The interaction energies of both the *S* and *R* isomers show that the two conformations are both in good qualitative agreement with in vitro antifungal activities. However, the interaction energies of the *R* isomers of the compounds were lower than that of the corresponding *S* isomers. Note that only

the binding mode of the *R* conformation is displayed in Figure 2. Clearly, the regions in the active site of CYP51 for ligand non-covalent binding can be divided into four subsites, S1–S4.<sup>26,27</sup> Molecular docking results revealed that the compound 8d binds to the active site of CACYP51 through the formation of a coordination bond with the iron of the heme group. The difluorophenyl group is located in the S3 subsite, which is a narrow hydrophobic cleft formed by Phe126, Leu121, Tyr118, Met306, and Gly307. The fluorine atom (2-F) can form H-bonds with the residue Gly307. The side chain of the compound 8d is oriented in the S4 subsite and forms hydrophobic and van der Waals interactions with Tyr64, Met92, Ala117, Tyr118, Phe228, Pro230, Leu376, Ile379, Phe380, and Met508. Moreover, the phenyl group attached to the piperidinol of the side chain interacts with the phenol group of Phe380 through the formation of  $\pi$ – $\pi$  face-to-edge interactions.

## Conclusion

In conclusion, a series of triazoles was successfully synthesized and characterized by IR, HR-MS, and NMR spectroscopic analysis. This study was undertaken to evaluate the effects on the antifungal properties of the designed compounds. The data of the in vitro antifungal activity indicates that all the synthesized compounds show

moderate-to-excellent activity against nearly all the tested fungal pathogens. The activity of type 7 is weakened when the alkyl group is substituted by a halogen atom. Moreover, the activity got lower when the halogen electron-withdrawing action ability was enhanced. Halogens are common in antifungal agents and, together with alkyl, alkoxy, nitro, and cyano groups, were inserted into the phenyl of the side chains. However, in our research, the introduction of the halogens on the phenyl group of the designed compounds did not improve activity. Conversely, antifungal activity to some other fungi strains was decreased and the antifungal spectrum was narrowed to *C. par.*, *C. tro.*, *C. neo.*, and *F. com.* Molecular docking results revealed that the compound 8d binds to the active site of CACYP51 through the formation of a coordination bond with the iron of the heme group. The difluorophenyl group is located in the S3 subsite. The fluorine (2-F) can form H-bonds with the residue Gly307. The side chain of the compound 8d is oriented in the S4 subsite and formed hydrophobic and van der Waals interactions with the amino residues. Moreover, the phenyl group attached to the piperidinol of the side chain interacts with the phenol group of Phe380 through the formation of  $\pi$ - $\pi$  face-to-edge interactions. More insightful observations of the active site would give us some key information about the SAR for further design of broad-spectrum antifungal agents.

## Acknowledgments

This work was supported by the National Natural Science Foundation of China (Grant No 30300437, 81273556), the Pharmaceutical Education research project of Chinese Association of Higher Medical Education Professional Committee (Grant No ZD201220), and the Creativity and Innovation Training Program of Second Military Medical University (Grant No ZD2013029).

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Singh N. Trends in the epidemiology of opportunistic fungal infections: predisposing factors and the impact of antimicrobial use practices. *Clin Infect Dis.* 2001;33(10):1692–1696.
- Ablordepey SY, Fan P, Ablordepey JH, Mardenborough L. Systemic antifungal agents against AIDS-related opportunistic infections: current status and emerging drugs in development. *Curr Med Chem.* 1999;6(12):1151–1195.
- Wildfeuer A, Seidl HP, Paule I, Haberleiter A. In vitro evaluation of voriconazole against clinical isolates of yeasts, moulds and dermatophytes in comparison with itraconazole, ketoconazole, amphotericin B and griseofulvin. *Mycoses.* 1998;41(7–8):309–319.
- Georgopapadakou NH. Antifungals: mechanism of action and resistance, established and novel drugs. *Curr Opin Microbiol.* 1998;1(5):547–557.
- Georgopapadakou NH, Walsh TJ. Antifungal agents: chemotherapeutic targets and immunologic strategies. *Antimicrob Agents Chemother.* 1996;40(2):279–291.
- Kontoyiannis DP, Mantadakis E, Samonis G. Systemic mycoses in the immunocompromised host: an update in antifungal therapy. *J Hosp Infect.* 2003;53(4):243–258.
- White TC, Marr KA, Bowden RA. Clinical, cellular, and molecular factors that contribute to antifungal drug resistance. *Clin Microbiol Rev.* 1998;11(2):382–402.
- Wong-Beringer A, Kriengkauykiat J. Systemic antifungal therapy: new options, new challenges. *Pharmacotherapy.* 2003;23(11):1441–1462.
- Ghannoum MA, Rice LB. Antifungal agents: mode of action, mechanisms of resistance, and correlation of these mechanisms with bacterial resistance. *Clin Microbiol Rev.* 1999;12(4):501–517.
- Sheehan DJ, Hitchcock CA, Sibley CM. Current and emerging azole antifungal agents. *Clin Microbiol Rev.* 1999;12(1):40–79.
- Sheng CQ, Zhang WN, Ji HT, et al. Structure-based optimization of azole antifungal agents by CoMFA, CoMSIA, and molecular docking. *J Med Chem.* 2006;49(8):2512–2525.
- Pfaller MA, Messer SA, Hollis RJ, Jones RN. In vitro activities of posaconazole (Sch 56592) compared with those of itraconazole and fluconazole against 3,685 clinical isolates of *Candida* spp. and *Cryptococcus neoformans*. *Antimicrob Agents Chemother.* 2001;45(10):2862–2864.
- Lepesheva GI, Hargrove TY, Ott RD, Nes WD, Waterman MR. Biodiversity of CYP51 in trypanosomes. *Biochem Soc Trans.* 2006;34(Pt 6):1161–1164.
- Lewis RE. Current concepts in antifungal pharmacology. *Mayo Clin Proc.* 2011;86(8):805–817.
- Di Santo R, Tafi A, Costi R, et al. Antifungal agents. 11. N-substituted derivatives of 1-[(aryl)(4-aryl-1 H-pyrrol-3-yl) methyl]-1 H-imidazole: synthesis, anti-candida activity, and QSAR studies. *J Med Chem.* 2005;48(16):5140–5153.
- Odds FC, Brown AJ, Gow NA. Antifungal agents: mechanisms of action. *Trends Microbiol.* 2003;11(6):272–279.
- Shyadehi AZ, Lamb DC, Kelly SL, et al. The mechanism of the acyl-carbon bond cleavage reaction catalyzed by recombinant sterol 14 alpha-demethylase of *Candida albicans* (other names are: lanosterol 14 alpha-demethylase, P-45014DM, and CYP51). *J Biol Chem.* 1996;271(21):12445–12450.
- Chai X, Zhang J, Hu H, et al. Design, synthesis, and biological evaluation of novel triazole derivatives as inhibitors of cytochrome P450 14 alpha-demethylase. *Eur J Med Chem.* 2009;44(5):1913–1920.
- Chai X, Zhang J, Yu S, et al. Design, synthesis, and biological evaluation of novel 1-(1H-1,2,4-triazol-1-yl)-2-(2,4-difluorophenyl)-3-substituted benzylamino-2-propanols. *Bioorg Med Chem Lett.* 2009;19(6):1811–1814.
- Chai X, Zhang J, Cao Y, et al. New azoles with antifungal activity: Design, synthesis, and molecular docking. *Bioorg Med Chem Lett.* 2011;21(2):686–689.
- Chai X, Zhang J, Cao Y, et al. Design, synthesis and molecular docking studies of novel triazole as antifungal agent. *Eur J Med Chem.* 2011;46(7):3167–3176.
- Jiang Y, Cao Y, Zhang J, et al. Design, synthesis and antifungal evaluation of 1-(2-(2,4-difluorophenyl)-2-hydroxy-3-(1H-1,2,4-triazol-1-yl) propyl)-1H-1,2,4-triazol-5(4H)-one. *Eur J Med Chem.* 2011;46(7):3135–3141.
- Xu J, Cao Y, Zhang J, et al. Design, synthesis and antifungal activities of novel 1,2,4-triazole derivatives. *Eur J Med Chem.* 2011;46(7):3142–3148.
- Aoyama Y, Yoshida Y, Nishino T, et al. Isolation and characterization of an altered cytochrome P-450 from a yeast mutant defective in lanosterol 14 alpha-demethylation. *J Biol Chem.* 1987;262(29):14260–14264.

25. Clinical and Laboratory Standards Institute (CLSI). Reference Method for Broth Dilution Antifungal Susceptibility Testing of Yeasts Approved standard. Document M38-A3, Wayne, PA; 2002.
26. Ji HT, Zhang WN, Zhang M, et al. Structure-based de novo design, synthesis, and biological evaluation of non-azole inhibitors specific for lanosterol 14 alpha-demethylase of fungi. *J Med Chem.* 2003;46(4):474–485.
27. Ji H, Zhang W, Zhou Y, et al. A three-dimensional model of lanosterol 14 $\alpha$ -demethylase of *Candida albicans* and its interaction with azole antifungals. *J Med Chem.* 2000;43(13):2493–2505.

## Drug Design, Development and Therapy

Dovepress

### Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which

has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/drug-design-development-and-therapy-journal>