

Sex differences in stroke: a socioeconomic perspective

Ahmad Delbari¹
 Farzane Keyghobadi²
 Yadollah Abolfathi
 Momtaz^{1,3}
 Fariba Keyghobadi²
 Reza Akbari²
 Hومان Kamranian²
 Mohammad Shouride Yazdi²
 Sayed Shahaboddin
 Tabatabaei¹
 Seyed-Mohammad
 Fereshtehnejad⁴

¹Iranian Research Center on Aging, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran; ²Research Center on Healthy Aging, Sabzevar University of Medical Sciences, Sabzevar, Khorasan, Iran; ³Malaysian Research Institute on Ageing (MyAgeing™), Universiti Putra Malaysia, Serdang, Selangor, Malaysia; ⁴Department of Neurobiology, Division of Clinical Geriatrics, Care Sciences and Society (NVS), Karolinska Institutet, Stockholm, Sweden

Correspondence: Yadollah Abolfathi Momtaz
 Iranian Research Center on Aging, University of Social Welfare and Rehabilitation Sciences, Kodakpar Ave, Daneshjo Blvd, Evin 1985713834, Tehran, Iran
 Email yabolfathi@gmail.com

Background: A number of studies have explored the issue of sex differences in stroke from biomedical perspective; however, there are still large gaps in the existing knowledge. The purpose of this study was to assess whether the differences in socioeconomic status and living conditions between men and women may explain the part of the sex differences in incidence and outcomes of stroke.

Methods: All stroke participants aged ≥ 60 years admitted in Vaseie Hospital in Sabzevar, Iran, from March 21, 2013, until March 20, 2014, were included in this study. Computerized tomography and magnetic resonance imaging were used to confirm stroke. A series of χ^2 tests were performed and Statistical Program for Social Sciences, Version 21.0, was used to investigate the potential differences between older men and women in stroke incidence and outcomes.

Results: A total of 159 incident stroke cases were documented during 1 year. The annual rate of stroke was statistically significantly higher in elderly women than in elderly men (401 vs 357 per 100,000; $P < 0.001$). Female elderly participants had significantly lower socioeconomic status, poorer living conditions, and higher lifetime history of depression, hypertension, and diabetes mellitus than their male counterparts.

Conclusion: The findings from this study showed that elderly women are more adversely affected by stroke in terms of incidence and outcomes of stroke than elderly men. The most noticeable result is that sex differences in socioeconomic status and living conditions may result in increased incidence of stroke and poorer outcomes in elderly women. Therefore, it is imperative to identify vulnerable elderly women and provide them appropriate treatment and services.

Keywords: aged, incidence, mortality, outcome, sex differences, socioeconomic disparities, vascular disease

Introduction

Currently, stroke is an emerging major public health problem and the cause of 9% of overall mortality.¹ It is estimated that by 2020 stroke will be the second leading cause of death.² The results from several studies show that incidence and burden of stroke are mostly different for women and men,³⁻⁵ and elderly women are more likely affected by stroke than men. Some studies show an increased incidence of stroke and poorer outcomes, such as disability and quality of life, in elderly women than in elderly men.^{1,4,6-9} Although sex differences in stroke have increasingly been explored from biomedical perspectives, and several explanations such as pregnancy complications, genetic factors, clinical presentation, steroid hormones, hemoglobin levels, oral contraceptives, and menopause have been provided to explain the observed sex differences in stroke,^{4,10-12} there are still large gaps in the existing knowledge.^{4,7,13} It seems that the reasons for sex disparities in stroke are multifactorial, and the mechanisms that increase the risk of stroke in elderly women are not well understood.^{4,14} Therefore, a

greater understanding of the differences in stroke between men and women may enhance the development of effective prevention and treatment strategies to improve stroke care and quality of life in the rapidly aging populations. As mentioned earlier, some studies reported sex differences in stroke from physiological perspectives and found that intrinsic sex differences may explain some of the differences in stroke between men and women, but there are still considerable gaps in our understanding of sex differences in stroke.⁴

Some studies have found that socioeconomic status contributes to stroke;^{2,15,16} however, only limited studies that investigated the differences in stroke between elderly men and women from a socioeconomic perspective are available. There are significant differences in socioeconomic status and living conditions between elderly men and women, wherein older women have lower socioeconomic status and living conditions.^{11,17–19} A growing body of literature indicates that lower socioeconomic status and living conditions contribute to some risk factors for stroke such as hypertension, diabetes mellitus, and depression.^{20–24} Participants with low socioeconomic status are less likely to receive specific processes of care after stroke compared with participants with high socioeconomic status.^{22,25} Some studies have found that older adults living alone, being unmarried, and living in deprived areas are less likely to receive stroke care.^{26,27} Some other studies have found that access to health care may affect both incidence and recovery from stroke. Low access to health care for various prevention and therapeutic services is related to higher rates of stroke incidence and mortality.^{2,21} In addition, low socioeconomic status and poor living conditions decrease the willingness to seek medical care among older adults.^{28–30} Figure 1 presents the model of hypothesized relationships. Elderly women, compared to elderly men, have

lower socioeconomic status and living conditions before stroke, which result in differences in risk factor distribution, availability of stroke prevention interventions, and provision of care that consequently increase the incidence of stroke and result in poorer outcomes of stroke.

According to the aforementioned possible underlying mechanism, the present study aimed to assess whether the differences in socioeconomic status and living conditions between men and women may explain part of the sex differences in incidence and outcomes of stroke.

Methodology

This hospital-based study was conducted in Sabzevar (a region in the northeast of Iran, with a population of ~500,000) from March 21, 2013, until March 20, 2014. All the participants aged ≥ 60 years admitted in Vaseie Hospital in Sabzevar, Iran, were included in this study. Computerized tomography scan and magnetic resonance imaging were used to confirm stroke. All participants were followed up until a definite outcome was determined. Permission was obtained from hospital administration and Ethical Committee, and then, data such as sociodemographic characteristics and prestroke risk factors were collected.

Ethical approval

This study was conducted in accordance with the 1964 Declaration of Helsinki and its later amendments and was approved by the Ethics Committee of the Sabzevar University of Medical Sciences, Iran. Written informed consent was obtained from all participants or from their relatives; the objectives of this study were explained, and confidentiality of information was assured to them.

Statistical Program for Social Sciences for Windows, Version 21.0 (IBM Corporation, Armonk, NY, USA), was used for data analysis. Descriptive statistics including percentages, measures of central tendency, and measures of variability were used to present the sample. A series of χ^2 tests were conducted to investigate the potential differences between older men and women in stroke incidence and outcomes. A two-tailed P -value ≤ 0.05 was used to determine the statistical significance.

Results

A total of 159 incident strokes among participants aged ≥ 60 years were observed during a period of 12 months from January 2013 to January 2014. The mean age of the participants was 76.38 years (standard deviation = 8.50; range 60–95 years), 52.2% were female, 69.2% were unemployed,




Figure 1 Socioeconomic mechanism underlying sex differences in stroke.

Table 1 Descriptive characteristics of the sample

Variable	Category	n	%	M	SD
Age	Young-old (60–74 years)	65	40.9	76.38	8.50
	Old-old (75–84 years)	57	35.8		
	Oldest-old (≥ 85 years)	37	23.3		
Sex	Male	76	47.8		
	Female	83	52.2		
Marital status	Unmarried	43	27.0		
	Married	116	73.0		
Education	No formal education	123	77.4		
	Formal education	36	22.6		
Employment status	Unemployed	110	69.2		
	Employed	49	30.8		
Stratum	Urban	57	35.8		
	Rural	102	64.2		
Living arrangement	Spouse	79	49.7		
	Family/children	52	32.7		
	Living alone	28	17.6		
Medical conditions	Hypertension	126	79.2		
	Diabetes mellitus	42	26.4		
	Depression	25	15.7		
HAT				22.31	29.16
	Within 4 hours	66	41.5		
	Within 24 hours	61	38.4		
Mortality	After 24 hours	32	20.1		
		8	5.0		

Abbreviations: SD, standard deviation; HAT, hospital arrival time after the onset of stroke.

and 77.4% reported not having formal education. The annual rate of stroke was 378 cases per 100,000 persons. There was a significantly higher incidence of stroke in elderly women than in elderly men (401 vs 357 per 100,000; $P < 0.001$); ~41% of stroke victims were in young-old (60–74) group. As shown in Table 1, ~16% of participants reported a history of depression, 26.4% diabetes, and 79.2% hypertension. The mean of hospital arrival time after the onset of stroke was 22.31 hours (standard deviation = 29.16).

Table 2 presents the sociodemographic and health characteristics of the sample with respect to sex. There was a strong association between marital status and sex ($\chi^2[1] = 17.05$; $P < 0.001$), with 88.2% of elderly male participants with stroke compared to 59.0% of elderly female participants with stroke being married. Slightly more than one-fourth of elderly women were living alone, whereas only a minority of the men (7.9%) were living alone. This sex difference was highly significant ($\chi^2[2] = 17.22$; $P < 0.001$). Majority (77.3%) of stroke participants in this sample had no formal education. Male stroke participants had significantly higher level of education than their female counterparts ($\chi^2[2] = 13.8$; $P < 0.001$).

As shown in Table 2, statistically significant differences were observed in risk factors for stroke between men and

Table 2 Sociodemographic and health characteristics of the sample with respect to sex

Variable	Category	Male		Female		χ^2
		n	%	n	%	
Age	Young-old (60–74 years)	26	34.2	39	47.0	3.64
	Old-old (75–84 years)	28	36.8	29	34.9	
	Oldest-old (≥ 85 years)	22	28.9	15	18.1	
Marital status	Unmarried	9	11.8	34	41.0	17.05***
	Married	67	88.2	49	59.0	
Education	No formal education	49	64.5	74	89.2	13.8***
	Formal education	27	35.5	9	10.8	
Employment status	Unemployed	43	56.6	67	80.7	10.85**
	Employed	33	43.4	16	19.3	
Stratum	Urban	34	44.7	23	27.7	5.00*
	Rural	42	55.3	60	72.3	
Living arrangement	With spouse	50	65.8	29	34.9	17.22***
	With family/children	20	26.3	32	38.6	
	Living alone	6	7.9	22	26.5	
Medical conditions	Hypertension	55	72.4	71	85.5	4.19*
	Diabetes mellitus	14	18.4	28	33.7	
	Depression	7	9.2	18	21.7	
HAT	Within 4 hours	32	42.1	34	41.0	7.22*
	Within 24 hours	35	46.1	26	31.3	
	After 24 hours	9	11.8	23	27.7	
Death	Yes	2	2.6	6	7.2	1.76
	No	74	97.4	77	92.8	

Notes: * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

Abbreviation: HAT, hospital arrival time after the onset of stroke.

women. Significantly more number of women reported lifetime history of depression ($\chi^2[1] = 4.66$; $P \leq 0.05$), hypertension ($\chi^2[1] = 4.19$; $P \leq 0.05$), and diabetes mellitus ($\chi^2[1] = 4.79$; $P \leq 0.05$) than men. In addition, hospital arrival time after the onset of stroke was longer in women ($\chi^2[1] = 7.22$; $P \leq 0.05$) than in men. Women were more likely to be unemployed than men ($\chi^2[1] = 10.85$; $P \leq 0.05$). Although no significant association was observed between sex and mortality rate, elderly women had higher mortality rate after stroke than men. In female participants, in-hospital mortality rate was 7.2%, while in male participants it was 2.6%. No significant difference in age was observed between elderly male and female survivors with stroke.

Discussion

In order to identify sex differences in stroke and to explore sex disparities in terms of incidence and outcome of stroke in later life, the present study was conducted from a socioeconomic perspective. Although some epidemiological studies show that stroke is more common among men,^{31,32} our finding that is in line with some studies showing increased incidence of stroke among postmenopausal women^{1,7,33} revealed a significantly higher rate of stroke in older women than in their age-matched male counterparts. Although age as a nonmodifiable risk factor may explain higher rate of stroke in older women,^{31,34} the finding from the current study showed no significant difference between elderly men and women in age.

In-hospital mortality rate among elderly women was three times more than that among men, which supports a growing body of evidence showing a greater in-hospital mortality in women than in men after stroke.^{4,35} It seems that elderly women are not evaluated as elderly men and possibly not treated appropriately. The findings from a Swedish registry showed that older women were less likely to be treated with antithrombotic drugs and less likely to receive anticoagulation.³⁶ Similarly, another study showed that older women were less likely to have echocardiography or any carotid investigation after adjusting for other confounding factors such as age and initial stroke severity.³⁷

Since the main aim of the present study was to investigate sex disparities in stroke, we explored these differences from a socioeconomic perspective. The findings from the current study showed that older women, compared to older men, had lower socioeconomic status and were more likely being unmarried and living alone. These findings are consistent with the results of previous studies showing sex differences in socioeconomic status^{38,39} and living conditions.^{11,17,40}

Multiple studies have found that people in low socioeconomic status and poor living conditions are more susceptible to acquiring chronic medical conditions and poorer health status.^{18,41} In parallel with previous studies,^{23,27} older women reported higher rates of past history of hypertension, diabetes mellitus, and depression, which might be associated with increased incidence of stroke.⁴²⁻⁴⁴ Older women are more vulnerable to social risk factors for depression than men.⁴⁵ Some studies show that low socioeconomic status is more closely associated with diabetes in women than in men.⁴⁶ The prevalence of hypertension is greater in women after menopause than in men, especially those with a history of hypertension during pregnancy. In addition, socioeconomic determinants of hypertension are more prevalent among women.^{47,48}

Other findings from the current study showed a sex disparity in delayed hospital arrival, that is, elderly women had longer prehospital delay. This finding is consistent with the results of previous studies that have demonstrated elderly women, compared to men, are less likely to be admitted to the acute stroke care unit within the eligible time that may potentially delay treatment and results in poor stroke outcomes.^{4,27,49,50} In addition, some elderly female stroke participants might be managed at home by traditional techniques⁹ that may lead to poor stroke outcomes.

In sum, in line with some previous studies demonstrating that low socioeconomic status and poor living conditions are associated with increased incidence and mortality from stroke,^{25,41,51} this study hypothesized that differences in socioeconomic status and living conditions between elderly men and women may explain part of sex disparities in stroke in terms of incidence rate and outcomes. Our findings showing lower socioeconomic status and poorer living conditions for older female participants with stroke may account for part of sex disparities in stroke.

Although, to our knowledge, this study is one of the first studies exploring sex differences in stroke from a socioeconomic perspective and found interesting results, it has some limitations. First, no data on household income status of participants were available. However, it is noteworthy to mention that level of education and employment status can reflect income effect.⁵² Second, the cross-sectional nature of the study precludes any definitive conclusion; therefore, a longitudinal study with a large sample size is required to confirm the proposed mechanism. The third limitation of the current study is that a multivariate analysis was not feasible, which may affect the results.

Conclusion

The findings from the current study supported some previous research studies showing that elderly women are more adversely affected by stroke in terms of incidence and in-hospital mortality of stroke than elderly men. The most noticeable result of our study was that sex differences in socioeconomic status and living conditions might result in higher rate of stroke incidence and poorer outcomes in elderly women. Therefore, it is imperative to identify vulnerable elderly women with stroke and provide them appropriate treatment and services.

Acknowledgments

This study was supported by Sabzevar University of Medical Sciences, Iran. The authors would like to thank Associate Professor Dr Robab Sahaf and anonymous reviewers for their helpful comments and suggestions for the revision and improvement in this manuscript.

Disclosure

The authors report no conflicts of interest in this work.

References

- Haast RA, Gustafson DR, Kiliaan AJ. Sex differences in stroke. *J Cereb Blood Flow Metab*. 2012;32(12):2100–2107.
- Cox AM, McKeivitt C, Rudd AG, Wolfe CD. Socioeconomic status and stroke. *Lancet Neurol*. 2006;5(2):181–188.
- Petrea RE, Beiser AS, Seshadri S, Kelly-Hayes M, Kase CS, Wolf PA. Gender differences in stroke incidence and poststroke disability in the Framingham Heart Study. *Stroke*. 2009;40(4):1032–1037.
- Turtzo LC, McCullough LD. Sex differences in stroke. *Cerebrovasc Dis*. 2008;26(5):462–474.
- Morvic S, Rundek T, Demarin V. Gender differences in stroke. *Period Biol*. 2012;114(3):267–268.
- Ahangar AA, Ashraf Vaghefi SB, Ramaezani M. Epidemiological evaluation of stroke in Babol, northern Iran (2001–2003). *Eur Neurol*. 2005;54(2):93–97.
- Roth DL, Haley WE, Clay OJ, et al. Race and gender differences in 1-year outcomes for community-dwelling stroke survivors with family caregivers. *Stroke*. 2011;42(3):626–631.
- Gargano JW, Reeves MJ. Sex differences in stroke recovery and stroke-specific quality of life: results from a Statewide Stroke Registry. *Stroke*. 2007;38(9):2541–2548.
- Foerch C, Ghandehari K, Xu G, Kaul S. Exploring gender distribution in patients with acute stroke: a multi-national approach. *J Res Med Sci*. 2013;18(1):10–16.
- Kimberly WT, Lima FO, O'Connor S, Furie KL. Sex differences and hemoglobin levels in relation to stroke outcomes. *Neurology*. 2013; 80(8):719–724.
- Reeves MJ, Bushnell CD, Howard G, et al. Sex differences in stroke: epidemiology, clinical presentation, medical care, and outcomes. *Lancet Neurol*. 2008;7(10):915–926.
- Bushnell CD, Hurn P, Colton C, et al. Advancing the study of stroke in women: summary and recommendations for future research from an NINDS-Sponsored Multidisciplinary Working Group. *Stroke*. 2006; 37(9):2387–2399.
- Reid JM, Dai D, Gubituz GJ, Kapral MK, Christian C, Phillips SJ. Gender differences in stroke examined in a 10-Year cohort of patients admitted to a Canadian teaching hospital. *Stroke*. 2008;39(4):1090–1095.
- Tsai KL, Lin CL, Hsu C. Is gender-specific therapy necessary for patients of hemorrhagic stroke? *J Exp Clin Med*. 2012;4(6):309–312.
- Addo J, Ayerbe L, Mohan KM, et al. Socioeconomic status and stroke an updated review. *Stroke*. 2012;43(4):1186–1191.
- Marshall IJ, Wang Y, Crichton S, McKeivitt C, Rudd AG, Wolfe CD. The effects of socioeconomic status on stroke risk and outcomes. *Lancet Neurol*. 2015;14(12):1206–1218.
- Yahaya N, Abdullah SS, Momtaz YA, Hamid TA. Quality of life of older Malaysians living alone. *Educ Gerontol*. 2010;36(10–11):893–906.
- Momtaz YA, Ibrahim R, Hamid TA, Yahaya N. Sociodemographic predictors of elderly's psychological well-being in Malaysia. *Aging Ment Health*. 2011;15(4):437–445.
- Hamid TA, Momtaz YA, Abdul Rashid SNS. Older women and lower self-rated health. *Educ Gerontol*. 2010;36(6):521–528.
- Winkleby MA, Jatulis DE, Frank E, Fortmann SP. Socioeconomic status and health: how education, income, and occupation contribute to risk factors for cardiovascular disease. *Am J Public Health*. 1992;82(6):816–820.
- Kapral MK, Wang H, Mamdani M, Tu JV. Effect of socioeconomic status on treatment and mortality after stroke. *Stroke*. 2002;33(1):268–275.
- Arrich J, Müllner M, Lalouschek W, Greisenegger S, Crevenna R, Herkner H. Influence of socioeconomic status and gender on stroke treatment and diagnostics. *Stroke*. 2008;39(7):2066–2072.
- Momtaz YA, Hamid TA, Yahaya N, Ibrahim R. Effects of chronic comorbidity on psychological well-being among older persons in Northern Peninsular Malaysia. *Appl Res Qual Life*. 2010;5(2):133–146.
- Kautzky-Willer A, Dorner T, Jensby A, Rieder A. Women show a closer association between educational level and hypertension or diabetes mellitus than males: a secondary analysis from the Austrian HIS. *BMC Public Health*. 2012;12:392.
- Langagergaard V, Palnum KH, Mehnert F, et al. Socioeconomic differences in quality of care and clinical outcome after stroke: a nationwide population-based study. *Stroke*. 2011;42(10):2896–2902.
- Lazzarino AI, Palmer W, Bottle A, Aylin P. Inequalities in stroke patients' management in English public hospitals: a survey on 200,000 patients. *PLoS One*. 2011;6(3):e17219.
- Bushnell C, McCullough LD, Awad IA, et al. Guidelines for the prevention of stroke in women: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014;45(5):1545–1588.
- Momtaz YA, Hamid TA, Ibrahim R, Yahaya N, Abdullah SS. Moderating effect of Islamic religiosity on the relationship between chronic medical conditions and psychological well-being among elderly Malays. *Psychogeriatrics*. 2012;12(1):43–53.
- Wiltshire JC, Roberts V, Brown R, Sarto GE. The effects of socioeconomic status on participation in care among middle-aged and older adults. *J Aging Health*. 2009;21(2):314–335.
- Ahmed SM, Tomson G, Petzold M, Kabir ZN. Socioeconomic status overrides age and gender in determining health-seeking behaviour in rural Bangladesh. *Bull World Health Organ*. 2005;83(2):109–117.
- Lloyd-Jones D, Adams R, Carnethon M, et al. Heart disease and stroke statistics – 2009 update a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation*. 2009;119(3):e21–e181.
- Appelros P, Stegmayr B, Terént A. Sex differences in stroke epidemiology: a systematic review. *Stroke*. 2009;40(4):1082–1090.
- Das SK, Banerjee TK, Biswas A, et al. A prospective community-based study of stroke in Kolkata, India. *Stroke*. 2007;38(3):906–910.
- Caso V, Paciaroni M, Agnelli G, et al. Gender differences in patients with acute ischemic stroke. *Womens Health (Lond)*. 2010;6(1):51–57.
- Persky RW, Turtzo LC, McCullough LD. Stroke in women: disparities and outcomes. *Curr Cardiol Rep*. 2010;12(1):6–13.
- Glader E-L, Stegmayr B, Norrving B, et al. Sex differences in management and outcome after stroke a Swedish national perspective. *Stroke*. 2003;34(8):1970–1975.
- Williams JE, Chimowitz MI, Cotsonis GA, Lynn MJ, Waddy SP, WASID Investigators. Gender differences in outcomes among patients with symptomatic intracranial arterial stenosis. *Stroke*. 2007;38(7):2055–2062.

38. Hamid TA, Momtaz YA, Ibrahim R. Predictors and prevalence of successful aging among older Malaysians. *Gerontology*. 2011;58(4):366–370.
39. Momtaz YA, Hamid TA, Yusoff S, et al. Loneliness as a risk factor for hypertension in later life. *J Aging Health*. 2012;24(4):696–710.
40. Momtaz YA, Haron SA, Ibrahim R, Hamid TA. Social embeddedness as a mechanism for linking social cohesion to well-being among older adults: moderating effect of gender. *Clin Interv Aging*. 2014;9:863.
41. Arrich J, Lalouschek W, Müllner M. Influence of socioeconomic status on mortality after stroke retrospective cohort study. *Stroke*. 2005;36(2):310–314.
42. O'Donnell MJ, Xavier D, Liu L, et al. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case-control study. *Lancet*. 2010;376(9735):112–123.
43. Yong H, Foody JA, Linong J, et al. A systematic literature review of risk factors for Stroke in China. *Cardiol Rev*. 2013;21(2):77–93.
44. Delbari A, Salman Roghani R, Tabatabaei SS, Rahgozar M, Lokk J. Stroke epidemiology and one-month fatality among an urban population in Iran. *Int J Stroke*. 2011;6(3):195–200.
45. Carayanni V, Stylianopoulou C, Koulierakis G, Babatsikou F, Koutis C. Sex differences in depression among older adults: are older women more vulnerable than men in social risk factors? The case of open care centers for older people in Greece. *Eur J Ageing*. 2012;9(2):177–186.
46. Rathmann W, Haastert B, Icks A, et al. Sex differences in the associations of socioeconomic status with undiagnosed diabetes mellitus and impaired glucose tolerance in the elderly population: the KORA Survey 2000. *Eur J Public Health*. 2005;15(6):627–633.
47. Bybee KA, Dew ML, Lawhorn SL, Stevens TL. *Cardiovascular Disease in Women Essentials*. Kansas City, MO: Jones & Bartlett Learning; 2012.
48. Warrell DA, Cox TM, Firth JD. *Oxford Textbook of Medicine*. Vol 1. Cambridge, UK: Oxford University Press; 2003.
49. Smith MA, Lisabeth LD, Bonikowski F, Morgenstern LB. The role of ethnicity, sex, and language on delay to hospital arrival for acute ischemic stroke. *Stroke*. 2010;41(5):905–909.
50. Lutfiyya MN, Ng L, Asner N, Lipsky MS. Disparities in stroke symptomatology knowledge among US midlife women: an analysis of population survey data. *J Stroke Cerebrovasc Dis*. 2009;18(2):150–157.
51. Putman K, De Wit L, Schoonacker M, et al. Effect of socioeconomic status on functional and motor recovery after stroke: a European multicentre study. *J Neurol Neurosurg Psychiatry*. 2007;78(6):593–599.
52. Litwak E, Kulis S. Technology, proximity, and measures of kin support. *J Marriage Fam*. 1987;49(3):649–661.

Clinical Interventions in Aging

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine,

CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>

Dovepress