

Exploration of sex differences in Rhes effects in dopamine mediated behaviors

Gabriel C Quintero^{1,2}
Daniela Spano³

¹INDICASAT AIP, Centro de Neurociencias, Panamá, ²Department of Psychology, University of New Orleans, N Orleans, Louisiana, USA; ³CEINGE Biotechnologie Avanzate, Naples, Italy

Abstract: Studies have shown that Ras homolog enriched in striatum (Rhes) proteins are highly expressed in areas of the central nervous system that have high dopaminergic innervation. In this study, we used Rhes mutant mice (Wild type, Rhes KO, Rhes Heterozygous) of both sexes to explore differences in the effects of Rhes protein levels in basal levels of activity, anxiety, and stereotypy, in relation to sex. Adult male and female mice were evaluated in an open field test for measuring basal levels of activity and anxiety for 5 consecutive days, and they were tested in the apomorphine-induced stereotypy paradigm. Rhes protein levels affected basal levels of activity but it was not found to be related to sex differences. Moreover, a decrease in Rhes protein levels was linked to a nonsignificant anxiolytic effect, mainly in female mice. Finally, a decrease in Rhes protein levels does not affect dopamine D₁ and D₂ receptor (D₁/D₂) synergism in female or male mice. Together, these results suggest that Rhes protein levels affect locomotion activity, and have an influence in anxiety depending on sex; Rhes protein levels do not affect D₁/D₂ synergism in both sexes.

Keywords: behavior, dopamine receptor, Rhes protein, mutant mice, sexes

Introduction

Dopamine (DA) is a catecholamine that serves as a major neurotransmitter, mediating diverse behaviors like movement, attention, motivation, and cognition.¹ DA receptors are normally divided in D1 and D2 subfamilies. The D1 subfamily is composed of the D₁ and D₅ subtypes, and the D2 subfamily composed of the D₂, D₃, and D₄ subtypes.² Under normal conditions, the majority of motor behaviors are mediated by the combined action of the D₁ and D₂ subtypes rather than the other subtypes.^{3,4} This combined action of D₁ and D₂ subtypes is known as D₁/D₂ receptor synergism.^{5,6} The D₁/D₂ receptor synergism has been supported empirically by studies at different levels: behavioral, electrophysiological, and gene expression in the striatum.^{6–10} However, the mechanism subjacent to this synergism has not been discovered. In order to explain this, it is necessary to elucidate the distribution of D₁ and D₂ receptors at the level of the striatum, but this has remained theoretically controversial; in effect, information on this topic is complicated. Theories range from complete colocalization to total segregation in striatal neurons.¹¹ Interaction of D₁ and D₂ receptors subserving synergism could be intercellular, intracellular, or combination of both.

Furthermore, DA receptors are functionally dynamic, altering their sensitivity to agonists in response to changes in the extracellular environment. For instance, following depletion of endogenous DA by means of selective destruction of mesostriatal DA projection with 6-hydroxydopamine (6-OHDA), there is a consequent

Correspondence: Gabriel C Quintero
INDICASAT AIP, Centro de Neurociencias, #219 Clayton, Ciudad del Saber, Panamá
Tel +507 397 9772
Fax +507 397 9772
Email gquintero2008@gmail.com

breakdown in D_1/D_2 synergism.^{12,13} This breakdown in D_1/D_2 synergism can also be generated by treatment with reserpine, a catecholamine depleting agent.^{14,15} Under both of these conditions, independent stimulation of either D_1 or D_2 receptors can elicit the full expression of DA-induced behaviors. This breakdown of synergism is invariably associated with profound supersensitivity of both D_1 and D_2 receptors. It is necessary to consider that the increase in DA receptor sensitivity is not necessarily correlated to an increase in DA receptor density, a phenomenon known as DA receptor dissociation. Moreover, the D_1/D_2 synergism is independent of D_1 and D_2 receptor density.¹⁵

Ras homolog enriched in striatum (Rhes) protein is a novel striatal specific Ras-like small G protein very similar to Dexas-1.¹⁶ Rhes proteins are expressed in different areas of the central nervous system, but mainly in the striatum and olfactory tubercle.^{16–18} In addition, *Rhes* mRNA is particularly enriched in brain regions that receive dopaminergic inputs, such as the Striatum and Nucleus Accumbens.¹⁷ *Rhes* mRNA is also found in low to moderate levels in other areas of the nervous system, such as the cerebellum, thalamus, hippocampus, and olfactory bulb during the early stages of postnatal development.^{17,19} *Rhes* mRNA in mammalian brains is localized in D_1R and D_2R bearing striatal projection neurons.²⁰ In rodents, the modulation of Rhes protein expression in the basal ganglia depends mainly on thyroid hormone levels during early stages of development, but depends on dopamine levels and DA innervations during the adult stages.^{16,19}

In general, there is a consensus that Rhes proteins could directly modulate the activation of the heterotrimeric G proteins to which G protein-coupled receptors (GPCRs) are coupled.^{21,22} Vargiu et al proposed possible explanations for the mechanism of interaction between Rhes and GPCR, and the inhibitory effect of activated D_2 receptor on cyclic adenosine monophosphate (cAMP).¹⁸ This study also reported that Rhes proteins exert their effects by interacting with G-alpha inhibitory proteins (G*α*i).

Previous studies have shown that decreases in mRNA levels occur after manipulations of the dopaminergic system of the basal ganglia in adult animals (eg, DA depletion by 6-OHDA or reserpine treatment), which results in consequent dopamine receptor supersensitivity and breakdown of the D_1/D_2 receptor synergism.¹⁷ Rhes protein levels are also decreased under similar conditions of dopamine receptor supersensitivity and the breakdown of D_1/D_2 synergism.¹⁷

In striatum, there is a relationship among the expression of Rhes proteins, the degree of DA receptor sensitivity, D_1/D_2

synergism, and the levels of dopamine neurotransmitters. Conditions that result in the breakdown of D_1/D_2 synergism and DA receptor supersensitivity (ie, DA depletion by 6-OHDA or reserpine treatment) invariably cause a decrease in *Rhes* mRNA levels in the striatum.¹⁷ Possible mechanisms by which Rhes protein levels affect DA receptor signaling have been proposed. Errico et al reported that in mutant mice, a decrease in Rhes proteins results in D_1R increased cAMP/protein kinase A (PKA) signaling mediated by G-olfactory (olf) proteins, and decreased coupling efficiency of D_2 receptor to its downstream Gi/o proteins.²⁰ Another study by Harrison et al added that Rhes proteins physically interact with G*α*i and can interfere with adenylyl cyclase (AC) activation by the G-olf coupled D_1 receptor in a pertussis toxin (PTX)-sensitive manner.²³ Furthermore, Thapliyal et al reported that Rhes proteins inhibit N-type calcium channels, but decrease the ability of ligands for Gi/o-coupled receptors to inhibit these channels, an effect mediated by β - γ subunits released from PTX-sensitive G proteins.²⁴

At a more behavioral level, the striatal Rhes protein has been linked to the inhibition of dopamine mediated behaviors in Rhes knockout (KO) mice. For instance, Rhes KO males denoted enhanced stereotypy after dopamine drug agonism.²⁵ Furthermore, Rhes knockout mice exhibited a higher D_2 receptor antagonist-induced catalepsy than Rhes wild type (WT) mice.²⁰

Other behavioral studies by Errico et al reported that Rhes KO mice displayed hyperactivity and increased locomotor response in the novelty-induced exploration task and open field test, respectively.²⁰ However, no differences were found between KO and WT mice in prepulse inhibition on their startle reflex, nor in motor coordination in the accelerating rotarod paradigm test. Furthermore, SKF 81297, a full D_1R agonist, produced a higher enhancement of basal levels of activity in Rhes KO compared to their WT counterparts at doses of 1.5 mg/kg and 5 mg/kg.²⁰ On the other hand, D_1R agonist SKF 38393 induces less of an increase in grooming behavior in Rhes mutant mice compared to their WT counterparts.²⁵

In addition, the postsynaptic D_2R -like antagonist haloperidol induced higher cataleptic behavior in Rhes KO compared to WT mice at all doses (0.25, 0.5, and 1.5 mg/kg). However, it seems that Rhes protein levels do not affect presynaptic D_2R functioning as evaluated by locomotor responses to quinpirole at different doses as there were no differences between WT and KO mice. In summary, Errico et al's behavioral studies concluded that Rhes protein levels act by modulating the motor output associated with postsynaptic

dopamine receptors transmission, and that decreasing Rhes protein levels facilitates the motor stimulation induced by a D_1 R agonist and strengthens the cataleptic response induced by the administration of a D_2 R antagonist.²⁰

Another preliminary study by Spano et al showed that Rhes KO mice displayed a sex dependent increase in anxiety levels, deficits in motor coordination, while learning and memory processes remained unchanged.²⁶ The present study evaluated whether or not there were sex related differences in the behavioral effects of Rhes protein levels on dopamine mediated behaviors, and assesses in both sexes the hypothesis that Rhes protein levels are causally related to the state of D_1/D_2 synergism and the degree of D_1 and D_2 receptor sensitivity.

Materials and methods

Subjects

In the present study, we used Rhes mutant mice of both sexes that had been backcrossed for ten generations on C57/Bl6 background, and bred at the University of New Orleans.²⁶ Mice employed were 2 to 3 months old, and divided in 3 genotypes: WT, KO, and heterozygous (Het). The number of mice totaled 58; 26 females ($n \geq 8$, per genotype group), and 32 males ($n \geq 9$, per genotype group). The mice were kept in same sex cages, in groups, with free access to food and water. Artificial lighting was provided from 7:00 am to 7:00 pm. The present study followed the guidelines of the NIH regarding the care and use of animals for experimental procedures, and the guidelines of the Institutional Animal Care and Use Committee (IACUC) of the University of New Orleans. Every effort was performed to minimize animal discomfort and to reduce the number of animals used. For controlling pain levels, mice were deeply anesthetized with intraperitoneal (IP) injections of tribromoethanol (100–150 mg per 10 g of body weight) for the tail biopsy procedures.

Open field (basal levels of activity and anxiety)

To determine whether the presence or absence of Rhes protein levels plays a role in basal levels of activity and anxiety, mice were placed in an open field arena (43.2 cm long by 33 cm wide by 32 cm high) for 30 minutes, for 5 consecutive days. During the test, the distance traveled in the arena field and the time spent in the periphery of the open field (thigmotaxis) were recorded by a digital video camera (Sony Corporation of America, New York, NY) and the video signal was sent directly to a computerized behavior analysis system (SMART System, Harvard Apparatus, Holliston, MA). The time the

mice spent in thigmotaxis was used as an index of anxiety. The periphery and center of the arena was defined as: the total area of the arena (1426 cm²) divided in two concentric rectangles (two areas of 713 cm²) with sides parallel to the arena limits. Thus, the total area of the arena was divided into 2 equal parts (center and outside) with the same probability of being occupied.

Stereotypy test

We tested whether Rhes protein levels affected D_1/D_2 synergism and DA receptor sensitivity by measuring stereotypy in response to dopaminergic drug administration. Stereotypy was used since it is a well characterized behavioral measure of D_1/D_2 synergism.²⁷

Species-specific stereotyped behavior was recorded under each of the four following conditions: (1) stimulation of neither receptor subtype (two saline injections at the scheduled time points), (2) stimulation of both D_1 and D_2 receptors [injection of saline 30 minutes prior to injection of the mixed D_1/D_2 agonist apomorphine (3 mg/kg)], (3) stimulation of D_2 receptors alone [injection of the D_1 antagonist SCH 23390 (0.1 mg/kg) 30 minutes prior to an injection of apomorphine (3 mg/kg)], (4) stimulation of D_1 receptors alone [injection of the selective D_2 antagonist eticlopride (0.3 mg/kg) 30 minutes prior to injection of apomorphine (3.0 mg/kg)]. All injections were IP, and every mouse was subjected to all conditions using a Latin square design in four sessions, each separated by 72–96 hours. This treatment regimen was used rather than selective agonists in order to eliminate the effects of endogenous DA at the heterotypic receptor. Each mouse was placed in a Plexiglas cylinder (measuring 22 cm high and 10.2 cm in diameter) with a thin layer of wood chip bedding on the floor. After 30 minutes of habituation, the appropriate pretreatment injection was given, followed 30 minutes later by an injection of apomorphine or saline. The testing session was terminated 60 minutes later. Behavior was recorded with a Sony digital video camera for subsequent analysis and rated on a scale of 0 to 5 every 5 minutes during a 30 second observation period. The behavioral analyst was blind to the genotype or pharmacological treatment. This rating scale, which was modified for mice from a similar study in rats, has been previously employed in other mutant mouse studies and is defined as follows: 0 = still; 1 = grooming or normal exploration; 2 = discontinuous unfocused stereotypy (eg, brief episodes of strong sniffing); 3 = continuous unfocused stereotypy behavior (ie, stereotypy directed to multiple objects and/or

surfaces); 4 = continuous focused sniffing (ie, sniffing of one object or within a highly circumscribed area); 5 = continuous focused oral stereotypy (ie, licking/chewing of one object or within a highly circumscribed area).^{15,27} Evaluation of the D₁ receptor behavior was described in a previous related work.²⁵ Specifically, grooming in response to the D₁ agonist SKF 38393 was observed. Male mice were placed in Plexiglas cylinders for 30 minutes of habituation, then mice received an injection of SKF 38393 (10 mg/kg, IP), and returned to the testing cylinders, and their behavior was recorded for the next hour. The amount of grooming (total seconds spent grooming every 5 minutes during a 60 second observation period) was recorded by an experimenter unaware of genotype.

Statistical analysis

Statistical analysis was performed using Analysis of Variance (ANOVA) factorial with post hoc tests. Significance was set at *P* value of <0.05. IBM SPSS Statistics (v 19; Armonk, NY) for Windows program was employed for statistical analysis. The numerical values described in the Results section represent the Mean \pm Standard Error of the Mean (SE).

Results

The data for all the behavioral experiments were analyzed by a mixed repeated measures design with the IBM SPSS Statistics 19 for Windows. Table 1 shows a general summary of experimental data for all behavioral tests.

Basal levels of activity

Figure 1 shows locomotion traveled distance across 5 days (30 minutes per day). Mixed repeated measures ANOVA of traveled distance showed significant main effect of days (Wilks' Lambda: 0.8, *F* (4, 64) = 5.2, *P* < 0.01), a significant main effect of genotype (Wilks' Lambda: 0.8, *F* (8, 128) = 2.3, *P* < 0.02), and significant main effect of sex (Wilks' Lambda: 0.9, *F* (4, 64) = 2.6, *P* < 0.05), but a nonsignificant days \times genotype \times sex interaction (Wilks'

Lambda: 0.9, *F* (8, 128) = 1.1, *P* = 0.4). Subsequent analysis of the main effect of days with *t* student comparisons revealed that the distance traveled on day 1 (7425.7 cm \pm 184.6) was significantly different from distances traveled on day 2 (7167.0 cm \pm 172.3), day 4 (7296.2 cm \pm 165.4), and day 5 (7594.8 cm \pm 170.9). The distance traveled on day 3 (6986.9 cm \pm 143.2) also differed from the distance traveled on day 5 (7594.8 cm \pm 170.9) (*P* < 0.001, all contrasts). Further analysis of the main effect of genotype with *t* student comparisons revealed that WT mice (7848.8 cm \pm 269.2) displayed higher levels of activity compared to Het mice (7053.7 cm \pm 312.6) on day 1 (*t* (16) = 2.6, *P* < 0.02); however, WT mice (6489.7 cm \pm 236.9) showed lower levels of activity than KO mice (7220.1 cm \pm 182.4) on day 3 [*t* (20) = -2.2, *P* < 0.05]. On day 5, KO mice (8160.1 cm \pm 372.6) exhibited higher levels of activity than Het mice (7280.9 cm \pm 156.7) (*t* (22) = -2.8, *P* < 0.02). On the other hand, further analysis of the main effect of sex with *t* student comparisons was unable to reveal significant differences between males and females on any of the five days (*P* > 0.05). In general, the locomotion tests showed that variation in Rhes protein levels could be related to variation in the levels of activity, but this trend could vary across days of exposition to the open field.

Figure 2 shows locomotion-traveled distance in the first 15 minutes of day 1. Mixed repeated measures design of traveled distance showed a significant effect of time (Wilks' Lambda: 0.8, *F* (2, 55) = 8.8, *P* < 0.001), a nonsignificant effect of genotype (Wilks' Lambda: 0.9, *F* (4, 110) = 0.7, *P* = 0.6), a nonsignificant effect of sex (Wilks' Lambda: 0.9, *F* (2, 55) = 0.3, *P* = 0.7), and a significant time \times genotype \times sex interaction (Wilks' Lambda: 0.8, *F* (4, 110) = 2.8, *P* = 0.03). Subsequent analysis of the effect of time with *t* student comparisons did not find a significant difference between groups. Compared to Spano et al's results, our results did not show significant differences between genotypes; however, Rhes Het mice exhibited lower levels of activity compared to WT mice in the first 10 minutes, and Spano et al's results

Table 1 Data summary

Locomotion activity			Anxiety			Stereotypy		
Distance traveled (cm)			Thigmotaxis (sec)			Stereotypy scores		
Day	Mean	SE	Day	Mean	SE	Treatment	Mean	SE
1	7425.7	184.6	1	1105.9	27.3	Condition		
2	7167	172.3	2	1198.3	26.2	Saline	0.7	0.0
3	6986.9	143.2	3	1162.2	26.9	D1	0.2	0.0
4	7296.2	165.4	4	1230.6	29.4	D2	0.6	0.1
5	7594.8	170.9	5	1205.5	28.3	D1 + D2	2.0	0.1

Abbreviations: sec, seconds; SE, standard error.




Figure 1 Profile of locomotor activity in female and male wild type (WT), knockout (KO), and heterozygous (Het) mice (day X sex X genotype) in the open field.

Note: Each point of the graph represents the average distance traveled (cm) in the open field (\pm SE) per day.

Abbreviations: WT, wild type; KO, knockout; Het, heterozygous; SE, standard error.

reported that KO mice showed lower levels of activity than their WT counterparts in the first 5 minutes (however, the main effects in that were not significant).²⁶ Taking together this study's and Spano et al's previous results, it could be

interpreted as Rhes protein level reduction is related to some decrease in basal levels of activity. Also, differences between both studies could be explained by differences in sampling.




Figure 2 Profile of locomotor activity in WT, KO, and Het mice during the first 15 minutes of day 1 in the open field.

Notes: Each point of the graph represents the average distance traveled (m) in the open field (\pm SE) per time interval. Data for males and females were pooled ($n = 18$ to 23 per genotype).

Abbreviations: WT, wild type; KO, knockout; Het, heterozygous; SE, standard error.

Anxiety

Figure 3 shows time spent in thigmotaxis across 5 days (30 minutes per day). Mixed repeated measures design for thigmotaxis time showed a significant main effect of days (Wilks' Lambda: 0.7, $F(4, 64) = 5.6$, $P < 0.01$), a nonsignificant main effect of genotype (Wilks' Lambda: 0.8, $F(8, 128) = 1.7$, $P = 0.09$), a nonsignificant main effect of sex (Wilks' Lambda: 0.9, $F(4, 64) = 0.9$, $P = 0.4$), and a nonsignificant day \times genotype \times sex interaction (Wilks' Lambda: 0.8, $F(8, 128) = 1.6$, $P = 0.1$). Subsequent analysis of the main effect of days with *t* student comparisons revealed that thigmotaxis in day 1 (1105.9 sec \pm 27.3) was significantly different from the rest of the days: day 2 (1198.3 sec \pm 26.2), day 3 (1162.2 sec \pm 26.9), day 4 (1230.6 sec \pm 29.4), and day 5 (1205.5 sec \pm 28.3) ($P < 0.001$, all contrasts). Despite the fact that the genotype main effect was not significant, there was a marked trend ($P = 0.095$) that a decrease in Rhes protein levels was related to lower levels of anxiety (lower time in thigmotaxis) (WT: 1198.5 sec \pm 42.8, KO: 1133.0 sec \pm 46.4, Het: 1136.8 sec \pm 41.8).

Stereotypy

Figure 4 shows stereotypy scores averaged across the 60 minute observation period. Mixed repeated measures design of stereotypy scores showed significant main effects of treatment (Wilks' Lambda: 0.1, $F(3, 51) = 240.3$, $P < 0.001$),

genotype (Wilks' Lambda: 0.7, $F(6, 102) = 4.1$, $P < 0.001$) and sex (Wilks' Lambda: 0.8, $F(3, 51) = 3.1$, $P = 0.04$), but a nonsignificant treatment \times genotype \times sex interaction (Wilks' Lambda: 0.9, $F(6, 102) = 0.7$, $P = 0.6$).

Concomitant analysis of the main effect of treatment with *t* student comparisons revealed that $D_1 + D_2$ treatment (2.0 ± 0.1) revealed significantly higher stereotypy scores compared to the rest of the treatments [saline (0.7 ± 0.0), D_1 (0.2 ± 0.0), and D_2 (0.6 ± 0.1)]; moreover, *t* student's comparisons also revealed that D_1 treatment (0.2 ± 0.0) was significantly different from saline (0.7 ± 0.0) and D_2 treatment (0.6 ± 0.1) ($P < 0.001$, all contrasts). This suggests that $D_1 + D_2$ treatment varied significantly from saline, D_1 , and D_2 treatment conditions, signifying that Rhes protein levels do not influence D_1/D_2 synergism.

Subsequent analysis of the main effect of genotype with *t* student comparisons revealed that under D_1 agonism treatment conditions, the WT group had (0.3 ± 0.0) higher stereotypy scores compared to the KO (0.1 ± 0.0) and Het groups (0.1 ± 0.0) ($P < 0.001$, all contrasts); furthermore, *t* students comparisons also revealed that under $D_1 + D_2$ treatment condition the WT group (1.7 ± 0.1) showed lower levels of stereotypy compared to KO (2.2 ± 0.2) and Het (2.2 ± 0.2) groups, but these differences did not reach significant levels ($P = 0.06$ and $P = 0.11$, respectively). Furthermore, the analysis of the main effect of sex with *t* student comparisons




Figure 3 Profile of anxiety behavior in female and male WT, KO, and Het mice in the open field.

Notes: Every bar represents the mean time in seconds spent in the periphery of the open field (\pm SE). The total amount of time for screening the behavior per day was 30 minutes (1800 seconds).

Abbreviations: WT, wild type; KO, knockout; Het, heterozygous; SE, standard error.




Figure 4 Profile of stereotypy behavior in female and male WT, KO, and Het mice.

Notes: Every bar represents the average score of all the time points (\pm SE). The higher the score, the higher the level of stereotypy. There was significant treatment effect ($P < 0.001$), genotypic effect ($P < 0.001$), and sex effect ($P < 0.05$). Specifically, D2 + D1 Treatment differed significantly from the rest of the Treatments ($P < 0.001$). Moreover, D1 Treatment differed significantly from the rest of the Treatments ($P < 0.001$). ** $P < 0.001$ WT groups differs from KO and HET; * $P < 0.01$ male WT groups differs from male KO and HET.

Abbreviations: WT, wild type; KO, knockout; Het, heterozygous; SE, standard error.

revealed that under $D_1 + D_2$ agonism treatment, the male group (2.2 ± 0.1) showed higher levels of stereotypy compared to the female group (1.7 ± 0.1) ($P = 0.05$).

In synthesis some trends were evident in the effects of treatments: under D_1 treatment condition, both KO and Het mice showed significantly lower stereotypy scores compared to WT mice. Furthermore, under $D_1 + D_2$ treatment, KO and Het mice showed a higher (nonsignificant) response than WT mice.

Discussion

The present results of the effects of Rhes protein levels in basal levels of activity are in partial agreement with previous similar behavioral studies. For instance, Spano et al reported in the open field that KO mice showed lower levels of activity in the first 5 minutes of the open field (15 minute test) compared to WT mice.²⁶ In our study, the open field test was performed for 30 minutes for 5 consecutive days, however, we included an additional analysis of the data as performed by Spano et al (15 minutes of day 1, Figure 2); this analysis denoted that Het mice showed lower levels of activity than their WT counterpart, but the tendency was not significant. It is important to consider that Spano et al's study reported a significant post hoc effect despite the absence of a main significant effect.²⁶ However, both studies agreed that a decrease in Rhes protein levels could be associated with a decrease in basal levels of activity.

Another study by Errico et al explored basal levels of activity in Rhes mutant mice. They reported that KO mice expressed higher rather than lower levels of activity compared to WT mice in the last 20 minutes of the open field test (30 minute session).²⁰ In the present study, KO mice also showed higher levels of activity than WT mice but only on the third day of the test. In this sense, both studies agree that a reduction in Rhes protein levels is related to an increase in basal levels of activity. The hyperactivity of the Rhes mutant mice can be explained by the control exerted by Rhes protein levels on G-olf protein levels, which are expressed higher in striatum of Rhes KO mice compared to their WT counterparts (KO mice expressed 114% to 124% of WT mice levels), according to Errico et al.²⁰

The present study found a marked but not significant association between a decrease in Rhes protein levels and anxiolysis; KO and Het mice showed lower levels of anxiety compared to their WT counterparts, a tendency that was more marked in female mice (shown in Figure 3). Our results in anxiety differ from those obtained by Spano et al.²⁶ Those authors found higher levels of anxiety in the female KO mice compared to their WT counterpart, while our study found the opposite tendency: KO mice displayed significantly lower levels of anxiety. This discrepancy could be explained by differences in the number of subjects employed and the type of anxiety test. Spano et al's study used the elevated plus maze test rather than the open field test. As a reference, other

mutant models of anxiety, such as the Galanin R1 receptor (GAL-R1), have also found inconsistent results between the elevated plus maze and open field tests.²⁸ In general, despite the apparent discrepancies between our study and Spano et al's study, one commonality is that female KO mice differ from WT and Het mice on anxiety levels, an effect that should be explored further.²⁶

Different authors have pointed out relationships between DA receptor dynamics and anxiety states. For instance, different authors have reported that lower doses of apomorphine (an agonist of D_1 and D_2 receptors) have sedating, relaxing and anxiolytic effects.^{29–33} Moreover, imaging studies (PET) have shown marked relationships between central D_2 receptor density, DA synthesis capacity in the striatum, and neuroticism anxiety (an intense feeling of anxiety, disproportionate to the trigger and can actually interfere with an individual's ability to function normally).^{34,35} In general, other studies have proposed a relevant role of DA in the genesis and treatment of anxiety states.³⁶ A possible explanation that can be drawn from the study is that because Rhes protein levels influence DA receptor supersensitivity, then an increase in DA receptor sensitivity is related to an anxiolytic effect, mainly in female mice.¹⁷ Moreover, a clinical study reported that dopamine receptor density could affect anxiety conditions.³⁷ In effect, that study reached a consensus that sex differences of neuroticism-anxiety related to variations in D_2 receptor density exist; it reported a sex specific (male but not female) association between lower neuroticism-anxiety and the expression of the A1 + allele of the DRD2 Taq 1 polymorphism (lower levels of anxiety correlated to lower levels of D_2 receptor density). Then, if there were associations between dopamine receptor density and anxiety, even sex specific relationships, then the existence of relationships between dopamine receptor sensitivity and anxiety should also be possible.

Our present results in the stereotypy test suggest a sex difference in the response to apomorphine; males showed higher levels of stereotypy response compared to females. Other similar studies in rodents have found different results; for instance, it has been reported that females display higher levels of apomorphine and cocaine-induced stereotypy than males.^{38,39} A possible explanation for this difference is due to levels of estrogen in females.⁴⁰ Female rats in estrus (high estrogen level) display low levels of stereotypy; furthermore, males treated with estrogen display a lower level of stereotypy compared to their nontreated counterparts.⁴⁰

The ovarian hormones effect on stimulants response is complex, and research results are contradictory.⁴¹ Some studies

found no effect of ovariectomy in amphetamine-induced stereotypy.^{38,42} However, other studies have found larger effects of cocaine and amphetamine in stereotypy under the estrus phase.^{43,44}

The decrease in Rhes protein levels does not seem to affect D_1/D_2 synergism, because D_1 agonism and D_2 agonism treatments induced lower stereotypic responses compared to $D_1 + D_2$ agonism treatment in KO and Het mice. If D_1/D_2 synergism were broken, then D_1 or D_2 agonism treatments should generate a higher stereotypic response, similar to $D_1 + D_2$ agonism treatment. Furthermore, KO and Het mice showed lower stereotypic responses under D_1 treatment compared to their WT counterparts, suggesting that Rhes protein levels are facilitatory of D_1 mediated behaviors. This is also supported by previous results published by our group reporting that Rhes protein levels appear to facilitate D_1 -specific grooming behaviors in males.²⁵ Our results suggest that Rhes protein levels are inhibitory of behaviors mediated by $D_1 + D_2$ receptor co-stimulation. A possible explanation for the difference between the facilitatory effect of Rhes proteins on D_1 agonism behavior and the inhibitory effect on $D_1 + D_2$ agonism behavior could be linked to a previous result by Errico et al. They reported that the mutant mice model that decreases Rhes protein levels results in a D_1R increased cAMP/PKA signaling which is mediated by G-olf proteins, and a decreased coupling efficiency of the D_2 receptor to its downstream Gi/o proteins.²⁰ Our previous work with a group of male mice reported a significant difference between KO and Het mice compared to WT mice under $D_1 + D_2$ treatment. In the present study, female groups did not show the same trend as male groups, and this could explain the differences in trends when both sexes were combined.²⁵

The present results suggest that Rhes proteins do not seem to be indispensable for DA D_1/D_2 receptor synergism in both sexes. However, Rhes proteins exert intricate effects on D_1 and D_2 receptor sensitivity. Specifically, Rhes proteins suppress behaviors generated by combined agonism of D_1 and D_2 receptors, mainly in male mice (Figure 4). The modulatory effects on D_1 receptors are more intricate as well. Although it has been shown previously to be inhibitory to $G\alpha_S$ activation and $G\alpha$ -olf expression, its presence is necessary for full expression of the D_1 -mediated grooming behavior. Despite the fact that Rhes protein levels do not seem to affect DA synergism, they do affect the sensitivity of the response to DA receptor activation. The increased response to $D_1 + D_2$ receptor co-activation by apomorphine in male but not in female KO mice suggests that Rhes proteins are repressive of this response and a potential sex difference. Other studies

performed in Rhes mutant mice suggest a repressive role for Rhes proteins in DA-mediated events. For example, Rhes KO mice show an increased locomotor response to the D₁ agonist SKF 81297, and increased G α -olf levels and Glutamate R1 phosphorylation.²⁰ In a heterologous expression system, Rhes protein levels were shown to inhibit the activation of G α S-coupled receptors at a point proximal to G protein activation.¹⁸ The results for D₂ receptor activation are less clear, with KO mice showing reduced D₂-mediated G protein activation and increased sensitivity to D₂ antagonists.²⁰ Our results indicate that for stereotypy, a behavior elicited only by concomitant stimulation of D₁ and D₂ receptors, Rhes proteins are inhibitory mainly in male mice rather than female mice.

To elucidate the effects of Rhes on activation of D₁ and D₂ receptors independently, we tested behavior induced by each receptor subtype alone. Agonism of D₁ receptors did not induce stereotypy, but the scores on the rating scale were reduced in KO and Het mice compared to their WT counterparts in both sexes. This effect is probably linked to an enhanced sensitivity to the D₂ antagonist eticlopride, which was administered prior to apomorphine in the D₁ group, and is in agreement with the findings of Errico et al, who reported increased catalepsy to haloperidol treatment in KO mice.²⁰ Activation of D₂ receptors did not induce significant differences in stereotypy among experimental groups for both sexes.

As in all experiments using mice in which a gene is knocked-out from the single cell stage, interpretations of the present results require caution. This model of mutant mice is limited by potential genetic compensatory responses that could lead to anatomical and physiological compensations during development. Subsequent studies could explore the present findings with a knockout model in which the knocking out condition is restricted in time, for example, to adulthood (temporal conditional knockout), or restricted in space; for instance, limited to the striatum.

Conclusion

In conclusion, the present research shows that changes in Rhes protein levels affect basal levels of activity and stereotypy response under DA receptor agonism conditions; furthermore, a decrease in Rhes protein levels could be linked to anxiolysis in females. A decrease in Rhes protein levels does not affect D₁/D₂ synergism in female and males.

Acknowledgments

The authors thank Dr G J LaHoste for research support and lab facilities. This study was supported by NIH grant

RR016816, and also by SNI funding (SENACYT-PANAMA) awarded to GQ.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Goldman-Rakic PS, Selemon LD. New frontiers in basal ganglia research. Introduction. *Trends Neurosci.* 1990;13(7):241–244.
2. Sibley DR, Monsma FJ Jr. Molecular biology of dopamine receptors. *Trends Pharmacol Sci.* 1992;13(2):61–69.
3. LaHoste GJ, Henry BL, Marshall JF. Dopamine D1 receptors synergize with D2, but not D3 or D4, receptors in the striatum without the involvement of action potentials. *J Neurosci.* 2000;20(17):6666–6671.
4. O'Sullivan GJ, Kinsella A, Grandy DK, Tighe O, Croke DT, Waddington JL. Ethological resolution of behavioral topography and D2-like vs D1-like agonist responses in congenic D4 dopamine receptor “knockouts”: identification of D4:D1-like interactions. *Synapse.* 2006;59(2):107–118.
5. Braun AR, Chase TN. Obligatory D-1/D-2 receptor interaction in the generation of dopamine agonist related behaviors. *Eur J Pharmacol.* 1986;131(2–3):301–306.
6. LaHoste GJ, Marshall JF. Dopamine receptor interactions in the brain. In: Stone TW, editor. *CNS Neurotransmitters and Neuromodulators: Dopamine*. Boca Raton, FL: CRC Press; 1996:107–119.
7. Bordi F, Meller E. Enhanced behavioral stereotypes elicited by intrastriatal injection D1 and D2 dopamine agonists in intact rats. *Brain Res.* 1989;504(2):276–283.
8. Walters JR, Bergstrom DA, Carlson JH, Chase TN, Braun AR. D1 dopamine receptor activation required for postsynaptic expression of D2 agonist effects. *Science.* 1987;236(4802):719–722.
9. White FJ, Bednars LM, Wachtel SR, Hjorth S, Brooderson RJ. Is stimulation of both D1 and D2 receptors necessary for the expression of dopamine-mediated behaviors? *Pharmacol Biochem Behav.* 1988;30(1):189–193.
10. LaHoste GJ, Yu J, Marshall JF. Striatal Fos expression is indicative of dopamine D1/D2 synergism and receptor supersensitivity. *Proc Natl Acad Sci U S A.* 1993;90(16):7451–7455.
11. Deng YP, Lei WL, Reiner A. Differential perikaryal localization in rats of D1 and D2 dopamine receptors on striatal projection neuron types identified by retrograde labeling. *J Chem Neuroanat.* 2006;32(2–4):101–116.
12. Arnt J. Hyperactivity induced by stimulation of separate dopamine D-1 and D-2 receptors in rats with bilateral 6-OHDA lesions. *Life Sci.* 1985;37(8):717–723.
13. Arnt J, Hyttel J. Differential inhibition by dopamine D-1 and D-2 antagonists of circling behaviour induced by dopamine agonists in rats with unilateral 6-hydroxydopamine lesions. *Eur J Pharmacol.* 1984;102(2):349–354.
14. Hu XT, Wachtel SR, Galloway MP, White FJ. Lesions of the nigrostriatal dopamine projection increase the inhibitory effects of D1 and D2 dopamine agonists on caudate-putamen neurons and relieve D2 receptors from the necessity of D1 receptor stimulation. *J Neurosci.* 1990;10(7):2318–2329.
15. LaHoste GJ, Marshall JF. Dopamine supersensitivity and D1/D2 synergism are unrelated to changes in striatal receptor density. *Synapse.* 1992;12(1):14–26.
16. Falk JD, Vargiu P, Foye PE, et al. Rhes: A striatal-specific Ras homolog related to Dexas1. *J Neurosci Res.* 1999;57(6):782–788.
17. Harrison LM, LaHoste GJ. Rhes, the Ras homolog enriched in striatum, is reduced under conditions of dopamine supersensitivity. *Neuroscience.* 2006;137(2):483–492.
18. Vargiu P, De Abajo R, Garcia-Ranea JA, et al. The small GTP-binding protein, Rhes, regulates signal transduction from G protein-coupled receptors. *Oncogene.* 2004;23(2):559–568.

19. Harrison LM, Lahoste GJ, Ruskin DN. Ontogeny and dopaminergic regulation in brain of ras homolog enriched in striatum (Rhes). *Brain Res.* 2008;1245:16–25.
20. Errico F, Santini E, Migliarini S, et al. The GTP-binding protein Rhes modulates dopamine signalling in striatal medium spiny neurons. *Mol Cell Neurosci.* 2008;37(2):335–345.
21. Cismowski MJ, Ma C, Ribas C, et al. Activation of heterotrimeric G-protein signaling by a ras-related protein. Implications for signal integration. *J Biol Chem.* 2000;275(31):23421–23424.
22. Graham TE, Prossnitz ER, Dorin RI. Dexas1/AGS-1 inhibits signal transduction from the Gi-coupled formyl peptide receptor to Erk-1/2 MAP kinases. *J Biol Chem.* 2002;277(13):10876–10882.
23. Harrison LM, He Y. Rhes and AGS1/Dexas1 affect signaling by dopamine D1 receptors through adenylyl cyclase. *J Neurosci Res.* 2011; 89(6):874–882.
24. Thapliyal A, Bannister RA, Hanks C, Adams BA. The monomeric G proteins AGS1 and Rhes selectively influence Galphai-dependent signaling to modulate N-type (CaV2.2) calcium channels. *Am J Physiol Cell Physiol.* 2008;295(5):C1417–C1426.
25. Quintero GC, Spano D, Lahoste GJ, Harrison LM. The Ras homolog Rhes affects dopamine D1 and D2 receptor-mediated behavior in mice. *Neuroreport.* 2008;19(16):1563–1566.
26. Spano D, Branchi I, Rosica A, et al. Rhes is involved in striatal function. *Mol Cell Biol.* 2004;24(13):5788–5796.
27. Nolan EB, Harrison LM, Lahoste GJ, Ruskin DN. Behavioral synergism between D(1) and D(2) dopamine receptors in mice does not depend on gap junctions. *Synapse.* 2007;61(5):279–287.
28. Holmes A, Kinney JW, Wrenn CC, et al. Galanin GAL-R1 receptor null mutant mice display increased anxiety-like behavior specific to the elevated plus-maze. *Neuropsychopharmacology.* 2003;28(6): 1031–1044.
29. Angrist B, Thompson H, Shopsin B, Gershon S. Clinical studies with dopamine-receptor stimulants. *Psychopharmacologia.* 1975;44(3): 273–280.
30. Corsini GU, Del Zompo M, Manconi S, Piccardi MP, Onali PL, Mangoni A. Evidence for dopamine receptors in the human brain mediating sedation and sleep. *Life Sci.* 1977;20(9):1613–1618.
31. Douglas CJ. Delirium tremens. *New York Medical Journal.* 1900:72.
32. Douglas CJ. A prompt emergency hypnotic. *New York Medical Journal.* 1917;106:1032.
33. Lal S, De la Vega CE. Apomorphine and psychopathology. *J Neurol Neurosurg Psychiatry.* 1975;38(7):722–726.
34. Kestler LP, Malhotra AK, Finch C, Adler C, Breier A. The relation between dopamine D2 receptor density and personality: preliminary evidence from the NEO personality inventory-revised. *Neuropsychiatry Neuropsychol Behav Neurol.* 2000;13(1):48–52.
35. Laakso A, Wallius E, Kajander J, et al. Personality traits and striatal dopamine synthesis capacity in healthy subjects. *Am J Psychiatry.* 2003; 160(5):904–910.
36. Goldberg HL, Finnerty RJ. The comparative efficacy of buspirone and diazepam in the treatment of anxiety. *Am J Psychiatry.* 1979;136(9): 1184–1187.
37. Wacker J, Reuter M, Hennig J, Stemmler G. Sexually dimorphic link between dopamine D2 receptor gene and neuroticism-anxiety. *Neuroreport.* 2005;16(6):611–614.
38. Savageau MM, Beatty WW. Gonadectomy and sex differences in the behavioral responses to amphetamine and apomorphine of rats. *Pharmacol Biochem Behav.* 1981;14(1):17–21.
39. Walker QD, Cabassa J, Kaplan KA, et al. Sex differences in cocaine-stimulated motor behavior: disparate effects of gonadectomy. *Neuropsychopharmacology.* 2001;25(1):118–130.
40. Miller JC. Sex differences in dopaminergic and cholinergic activity and function in the nigro-striatal system of the rat. *Psychoneuroendocrinology.* 1983;8(2):225–236.
41. Van Hartesveldt C, Joyce JN. Effects of estrogen on the basal ganglia. *Neurosci Biobehav Rev.* 1986;10(1):1–14.
42. Camp DM, Robinson TE. Susceptibility to sensitization. II. The influence of gonadal hormones on enduring changes in brain monoamines and behavior produced by the repeated administration of D-amphetamine or restraint stress. *Behav Brain Res.* 1988;30(1):69–88.
43. Quiñones-Jenab V, Ho A, Schlussman SD, Franck J, Kreek MJ. Estrous cycle differences in cocaine-induced stereotypic and locomotor behaviors in Fischer rats. *Behav Brain Res.* 1999;101(1):15–20.
44. Becker JB, Cha JH. Estrous cycle-dependent variation in amphetamine-induced behaviors and striatal dopamine release assessed with microdialysis. *Behav Brain Res.* 1989;35(2):117–125.

Neuropsychiatric Disease and Treatment

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS.

Submit your manuscript here: <http://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>

Dovepress

The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.