

# Emerging targets in cancer management: role of the CXCL12/CXCR4 axis

Monica Cojoc<sup>1</sup>  
Claudia Peitzsch<sup>1</sup>  
Franziska Trautmann<sup>1</sup>  
Leo Polishchuk<sup>2</sup>  
Gennady D Telegeev<sup>2</sup>  
Anna Dubrovskaja<sup>1</sup>

<sup>1</sup>OncoRay National Center for Radiation Research in Oncology, Medical Faculty Carl Gustav Carus, Dresden University of Technology, Dresden, Germany; <sup>2</sup>Institute of Molecular Biology and Genetics, National Academy of Sciences of Ukraine, Kyiv, Ukraine

**Abstract:** The chemokine CXCL12 (SDF-1) and its cell surface receptor CXCR4 were first identified as regulators of lymphocyte trafficking to the bone marrow. Soon after, the CXCL12/CXCR4 axis was proposed to regulate the trafficking of breast cancer cells to sites of metastasis. More recently, it was established that CXCR4 plays a central role in cancer cell proliferation, invasion, and dissemination in the majority of malignant diseases. The stem cell concept of cancer has revolutionized the understanding of tumorigenesis and cancer treatment. A growing body of evidence indicates that a subset of cancer cells, referred to as cancer stem cells (CSCs), plays a critical role in tumor initiation, metastatic colonization, and resistance to therapy. Although the signals generated by the metastatic niche that regulate CSCs are not yet fully understood, accumulating evidence suggests a key role of the CXCL12/CXCR4 axis. In this review we focus on physiological functions of the CXCL12/CXCR4 signaling pathway and its role in cancer and CSCs, and we discuss the potential for targeting this pathway in cancer management.

**Keywords:** epithelial-to-mesenchymal transition, cancer stem cells, metastasis

## Introduction to the CXCL12/CXCR4 axis

Chemokines are a class of small (8–10 kDa) inflammatory or homeostatic cytokines sharing a common biological activity in stimulating the migration of different types of cells including lymphocytes, monocytes, neutrophils, endothelial cells, mesenchymal stem cells, and malignant epithelial cells.<sup>1,2</sup> Chemokines are classified into four conserved groups – CXC, CC, C, and CX3C – based on the number and spacing of their N-terminal cysteine residues: CXC chemokines have a single nonconserved amino acid residue (X) between the first N-terminal cysteine residues (C); CC chemokines have these two cysteine residues adjacent; C chemokines have only one N-terminal cysteine; whereas CX3C chemokines contain three nonconserved amino acid residues separating the N-terminal cysteine pair. More than 50 chemokines have been discovered so far.<sup>1,3</sup> The classification of the chemokine receptors is based on the type of their ligands. For example, CXC receptors bind CXC ligands, while CC receptors bind CC ligands, etc.<sup>3</sup>

To date, over 20 chemokine receptors have been identified.<sup>1,3–5</sup> Chemokine receptors belong to a family of G protein-coupled receptors (GPCRs) containing seven transmembrane-spanning  $\alpha$ -helix domains. One of the intracellular loops of the chemokine receptors couples with heterotrimeric G proteins that mediate a cascade of intracellular signaling following ligand binding.<sup>4</sup> The heterotrimeric G protein is composed of the G $\alpha$ , G $\beta$ , and G $\gamma$  subunits. Both G $\alpha$  and G $\beta$  subunits have covalently

Correspondence: Anna Dubrovskaja  
OncoRay National Center for Radiation  
Research in Oncology, Medical Faculty  
Carl Gustav Carus, TU Dresden,  
Fetscherstr 74/PF41,  
01307 Dresden, Germany  
Email [anna.dubrovskaja@oncoray.de](mailto:anna.dubrovskaja@oncoray.de)


Gennady D Telegeev  
Institute of Molecular Biology and  
Genetics, NAS of Ukraine, Zabolotnogo  
Str, 150 Kyiv 03143, Ukraine  
Email [g.d.telegeev@imbg.org.ua](mailto:g.d.telegeev@imbg.org.ua)

attached lipid tails that anchor G proteins to the plasma membrane. In the inactive or basal state, the  $G\alpha$  subunit contains the guanine nucleotide diphosphate (GDP).

Upon activation, GPCR acts as a GEF (guanine nucleotide exchange factor) and promotes the conformational change of the  $G\alpha$  subunit and replacement of the bound GDP by guanine nucleotide triphosphate (GTP). This exchange triggers the further conformation changes within the  $G\alpha$  subunit, which allows the trimeric G protein to be released from the receptor, and to dissociate into the GTP-bound  $G\alpha$  subunit and  $G\beta/\gamma$  dimer. Both the activated components interact with various effector proteins and initiate unique intracellular signaling cascades, such as activation of phospholipase C (PLC), regulation of adenylyl cyclase, triggering of different kinase cascades including mitogen-activated protein kinase (MAPK), c-Jun N-terminal kinase

(JNK), p38, and the phosphoinositide 3-kinase (PI3K) routes (Figure 1).<sup>5-9</sup>

The distinct routes of the GPCRs signaling depend on the coupled  $G\alpha$  subunits, which are classified into four families;  $G\alpha_s$ ,  $G\alpha_i$ ,  $G\alpha_q$  and  $G\alpha_{12}$ . GPCRs coupled to the  $G\alpha_s$  stimulate adenylyl cyclase whereas  $G\alpha_i$  bound GPCRs inhibit it. The adenylyl cyclase serves as an effector enzyme that catalyzes 5'adenosine triphosphate into cyclic adenosine monophosphate (cAMP) and thereby activates cAMP-dependent protein kinase, which regulates a host of other downstream effectors including MAPK signaling pathway.<sup>5,10</sup> Activated  $G\alpha_i$  is also able to activate the Src family of tyrosine kinases, which play an important role in signal integration.<sup>3,6</sup> GPCRs coupled to  $G\alpha_q$  act through PLC $\beta$ , which cleaves phosphatidylinositol 4,5-bisphosphate to form the second messenger molecules called diacylglycerol and inositol-1,4,5-trisphosphate (IP<sub>3</sub>).



**Figure 1** A schematic of the CXCL12/CXCR4 signaling pathways.

**Abbreviations:** ER, endoplasmic reticulum; GTP, Guanosine-5'-triphosphate; RTK, receptor tyrosine kinase; CXCL, chemokine (C-X-C motif) ligand; CXCR, C-X-C chemokine receptor; cAMP, cyclic adenosine monophosphate; PKA, protein kinase A; JAK, Janus kinase; STAT, signal transducer and activator of transcription; Cdc42, Cell division control protein 42 homolog; Rac, Ras-related C3 botulinum toxin substrate; Rho, Ras homolog gene family; GRK, G protein-coupled receptor kinase; FOXO, Forkhead box protein; NF $\kappa$ B, nuclear factor kappa-light-chain-enhancer of activated B cells; PIP2, phosphatidylinositol bisphosphate; FAK, focal adhesion kinase; PLC, phospholipase C; PKC, protein kinase C; Ras, Rat sarcoma protein family; ERK, extracellular-signal-regulated kinase.

Diacylglycerol activates another enzyme called protein kinase C (PKC), whereas  $IP_3$  diffuses to the endoplasmic reticulum where it opens calcium channels and triggers the release of calcium from intracellular stores into the cytoplasm. This intracellular calcium mobilization is frequently used for analysis of chemokine receptor activity.<sup>11</sup> GPCRs coupling to  $G\alpha_{12}$  alternatively act via the Rho-GEF, which in turn activates the small G protein RhoA (Figure 1).<sup>5,12</sup>

Activation of PI3K by GPCRs is thought to be dependent on the direct binding of  $G\beta\gamma$  subunits.<sup>13</sup> PI3K activation triggers a signaling cascade leading to the activation of AKT (also called protein kinase B) and its downstream targets including phosphoinositide-dependent kinase 1 (PDK1), glycogen synthase kinase 3 (GSK3), mammalian target of rapamycin (mTOR), p70 ribosomal protein S6 kinase (p70<sup>S6K</sup>), forkhead family transcription factors (FOXO), and other signaling proteins. Notably, PI3K activation in response to the GPCR-mediated signaling results in the activation of focal adhesion kinase (FAK), which induces migratory activity in different types of cells, including tumor cells.<sup>6,14,15</sup>

The duration of the GPCR signaling depends on the  $G\alpha$  subunit lifespan in the GTP-bound state. Hydrolysis of the GTP of  $G\alpha$ -GTP to GDP leads to the inactivation of the  $G\alpha$  subunit and to its reassociation with the  $G\beta/\gamma$  dimer, which terminates all effector interactions.<sup>16,17</sup> In addition, chemokine receptor signaling is tightly regulated by the process of internalization and lysosomal degradation. Upon GPCR signaling activation, intracellular domains of receptors are phosphorylated by the second messenger kinases such as G protein-coupled receptor kinases (GRKs), followed by the binding of the phosphorylated receptors with regulatory proteins called arrestins. Arrestins impair communication of GPCRs with the G proteins and target them for lysosomal degradation following protein internalization and trafficking (Figure 1).<sup>18,19</sup>

Interestingly, it has been reported that several chemokine receptors including CCR2, CCR5, CXCR1, CXCR2, CXCR4, and CXCR7 can undergo homo- or hetero-dimerization upon ligand binding; a process that was proposed to regulate distinct intracellular signaling pathways.<sup>7,20,21</sup> Chemokines and their receptors display a high degree of redundancy in that most chemokines bind to multiple receptors and vice versa. The chemokine stromal cell-derived growth factor 1 (SDF-1), also known as CXCL12, binds primarily to its cognate receptor CXCR4, which is also a coreceptor for the entry of the Human immunodeficiency virus (HIV) into the target immune cells (T helper cells) besides the CD4 receptor.<sup>22</sup>

The assumption that CXCR4 is the only receptor for CXCL12 was recently challenged, since it was demonstrated

that this chemokine also binds to the orphan receptor called CXCR7, which is a receptor for the interferon-inducible T cell chemoattractant CXCL11/I-TAC. Moreover, CXCR7 constitutively forms heterodimers with the CXCR4 receptor. Growing evidence indicates that binding of CXCL12 to CXCR7 does not result in activation of signaling pathways typical of G proteins. It has been proposed that CXCR7 serves as a ligand scavenger or acts as a “decoy” receptor.<sup>23,24</sup> Very recently, this receptor has been described as an activator of various signaling pathways in a CXCL12-dependent manner.<sup>23,24</sup> CXCR7 is broadly expressed in normal tissues including the heart, brain, spleen, kidney, lung, testis, ovary, thyroid, and human placenta.<sup>23</sup> A germline deletion of CXCR7 resulted in perinatal lethality, and expression of CXCR7 was associated with cardiac development.<sup>25</sup> Moreover, CXCR7 is upregulated in many malignant cells, including breast, lung, cervical, pancreatic, and prostate cancer cells, and found to be involved in tumor cell growth, survival, and metastasis.<sup>21,22,24</sup>

## Physiological function of the CXCL12/CXCR4 signaling

CXCL12 is a small (8 kDa) homeostatic chemokine that was originally described as an efficacious lymphocyte chemoattractant and regulator of hematopoiesis, and was soon after also characterized as a modulator of multiple physiological processes.<sup>26–29</sup> CXCL12 is a pleiotropic chemokine that is widely expressed in different organs including the brain, lung, colon, heart, kidney, and liver where it acts as a chemoattractant for immature and mature hematopoietic cells; it thus plays an important role in inflammation and immune surveillance of tissues. Additionally, CXCL12 serves as an emergent salvage signal for initiating tissue regeneration and repair. Of note, various tissues respond to the chemical or physical insults such as toxic agents, hypoxia, and irradiation by increasing expression and secretion of CXCL12, which is important for recruitment of CXCR4 positive stem and progenitor cells to a site requiring tissue regeneration.<sup>29</sup> CXCL12 is expressed from a single gene in six splice variant isoforms known as SDF-1 $\alpha$ , SDF-1 $\beta$ , SDF-1 $\gamma$ , SDF-1 $\delta$ , SDF-1 $\epsilon$ , and SDF-1 $\phi$ .<sup>30–32</sup> These CXCL12 isoforms share the same first three exons but contain different fourth exons. Different splice variants are characterized by distinct properties such as stability and tissue of origin. SDF-1 $\alpha$  is constitutively produced in many organs but tends to undergo rapid degradation in the blood. In contrast, SDF-1 $\beta$  displays high proteolytic stability and is expressed in highly vascularized organs such as the liver, spleen, and kidney. SDF-1 $\gamma$  is present in less vascularized organs such as the heart and brain.<sup>30–32</sup>



and metastasis development in more than 75% of all cancers including breast, ovarian, lung, colon, prostate, kidney, melanoma, brain, esophageal, pancreatic, and many forms of leukemia.<sup>56–60</sup> Whereas CXCR4 expression is low or absent in many normal tissues, it is highly expressed in more than 23 different types of tumors including prostate, brain, breast, kidney, pancreas, ovarian cancer, and melanoma (Figure 2).<sup>56</sup> CXCL12 protein levels are highest in organs that are known to be the common sites of metastasis including the liver, bone marrow, and lungs, suggesting that tumor cells may use chemokine-mediated trafficking patterns, which are normally utilized during organogenesis, vascularization, and tissue regeneration.<sup>30–32,55</sup>

The CXCR4/CXCL12 axis also indirectly promotes tumor metastasis by mediating invasion and proliferation of tumor cells, and enhancing tumor-associated neoangiogenesis.<sup>61–63</sup> The stromal fibroblasts found within carcinomas and termed carcinoma-associated fibroblasts (CAFs) constitute the major cellular component of the tumor microenvironment. For a long period of time, neither the source of CAFs nor the differences between CAFs and fibroblasts from nonneoplastic tissue have been defined. Recent studies have shown that CAFs can arise from multiple origins including resident tissue fibroblasts, epithelial cells, human mesenchymal stem cells, and HSC.<sup>64–66,67,68</sup>

Analogous to their normal counterparts, which are present in inflammatory environments and aid in wound healing through tissue remodeling and repair, CAFs promote angiogenesis and matrix degradation through elevated secretion of CXCL12 and matrix metalloproteinase-1 (MMP-1).<sup>66,69</sup> During the past 25 years, the accumulation of evidence supports a critical and complex role of MMPs in tumor development. For example, MMPs can increase genomic instability in normal epithelium, which may result in tumor formation. MMPs are the key enzymes responsible for degradation of the extracellular matrix, which is the physical barrier for cancer cell invasion.<sup>70</sup> A recent study has identified *CXCR4* as a CAF-associated gene, implying an existence of an autocrine feedback loop (Figure 2).<sup>71</sup> The CXCR4/CXCL12 signaling in CAFs results in a multitude of cellular functions, including migration within the tumor microenvironment, adhesion, proliferation, and secretion of MMPs.<sup>72–75</sup>

A large number of studies demonstrated that CAF-derived CXCL12 not only stimulates carcinoma cell growth directly through the CXCR4 receptor displayed on tumor cells, or indirectly through MMP-mediated tissue remodeling, but also serves to recruit endothelial progenitor cells into tumors and thereby promotes neoangiogenesis.<sup>64,65</sup>

Moreover, CAF-mediated CXCL12 promotes an epithelial-to-mesenchymal transition (EMT) in primary tumors.<sup>76</sup> The acquisition of the EMT program is a critical process for the progression of cancers from local carcinomas to invasive malignancies, which is often associated with the loss of epithelial differentiation and gain of mesenchymal phenotype. Recent studies have shown a molecular link between EMT and self-renewal and demonstrated that cancer cells undergoing EMT gain the cancer stem cell (CSC) phenotype and tumorigenicity.<sup>77–80</sup>

A growing body of evidence indicates that a subset of cancer cells referred to as CSCs play a critical role in tumor initiation and resistance to anticancer therapy.<sup>77</sup> Similarly to normal stem cells, CSCs possess the ability to self-renew and to differentiate into all cell populations within the tumor mass.<sup>81</sup> This stem cell concept of tumorigenesis was proven the first time in 1994 by John Dick et al who demonstrated that CD34<sup>+</sup>CD38<sup>-</sup> acute myeloid leukemia cell subset is a cell population initiating the malignant disease in immunodeficient mice.<sup>82</sup> Clarke et al introduced the CSC concept in solid tumors in 2003.<sup>83</sup> In this study they identified CD44<sup>+</sup>CD24<sup>-/low</sup> breast cancer CSCs. As few as 100 cells with the CD44<sup>+</sup>CD24<sup>-/low</sup> phenotype were able to form tumors in mice, whereas tens of thousands of cells with alternate phenotypes were unable to initiate tumor growth.<sup>83</sup> During the last few years, similar discoveries were made in other tumor types. Moreover, this population has been implicated to therapy resistance and tumor recurrence.<sup>84–86</sup>

Several lines of evidence suggest a link between the EMT and CSC characteristics in various types of cancer. Alterations in genes associated with developmental pathways such as Wnt, SHH and Notch are common in CSC populations, and could facilitate the EMT process. Analogous to differentiated somatic cells, which can be reset to a pluripotent stage by the process of induced differentiation, non-CSC tumor cells can be reprogrammed by the activation of developmental pathways and EMT programs that change their self-renewal and differentiation potency.<sup>79,80,85,87,88</sup>

The positive correlation between EMT and CSC properties could lead to the concept of “migrating cancer stem cells” as the basis of metastatic colonization.<sup>89,90</sup> Recent findings demonstrated that a distinct subpopulation of CSCs can initiate tumor growth at secondary sites. The features of CSCs such as invasion, attachment-independent survival, and the ability to interact with micromilieu at the extravasation site support their involvement in metastatic dissemination.<sup>84,91</sup> Moreover, recent studies have revealed striking similarities of the signaling pathways regulating

CSC and driving metastasis formation. For example, Liu and colleagues reported a 186-gene “invasiveness” gene signature (IGS) that was identified by comparing the gene expression profiles of normal breast epithelium and breast CSCs with CD44<sup>+</sup>/CD24<sup>-low</sup> phenotype. Among 295 breast cancer patients, there was a significant association between the IGS signature and metastasis-free survival. The IGS was also applied to discriminate low- and high-risk patients with medulloblastoma, lung, and prostate cancers, demonstrating the prognostic value of markers that define CSCs.<sup>92</sup> More specifically, several reports demonstrated that CSCs are involved in metastatic dissemination in

xenograft models of breast, pancreatic, and colorectal carcinoma.<sup>83,93,94</sup>

In support of the hypothesis that metastatic tumor cells could have both EMT and CSC phenotypes, clinical studies have shown that the majority (>80%) of the circulating tumor cells (CTCs) in patients with metastatic castration-resistant prostate cancer coexpress stem cell marker CD133 and mesenchymal proteins including vimentin, N-cadherin, and O-cadherin.<sup>95</sup> Study of CTCs from metastatic breast cancer patients revealed that 62% of the cells were positive for at least one of three EMT markers – Twist1, AKT2, PI3K $\alpha$  – and 69% of the total

**Table I** CXCR4 as a marker for putative cancer stem cell populations in solid tumors

Tumor type	Additional CSC markers analyzed in combination with CXCR4	Biological functions of the CXCL12-CXCR4 axis	Methods used	References
Renal	N/A	Clonogenicity, tumorigenicity, drug resistance	<ul style="list-style-type: none"> <li>– Analysis of the FASC sorted patient-derived cells for their spherogenicity and tumorigenicity;</li> <li>– Inhibition of CXCR4 expression by siRNA;</li> <li>– Chemical inhibition of CXCR4 functions in vitro;</li> <li>– Combination treatment with AMD3100 and pazopanib, sunitinib and sorafenib</li> </ul>	161
Prostate	CD133, CD44	Spherogenicity, tumorigenicity, differentiation potential, drug resistance	<ul style="list-style-type: none"> <li>– Analysis of the FASC sorted cells from the established cell lines for their differentiation potential, cell adhesion, clonal growth, spherogenicity, and tumorigenicity;</li> <li>– Inhibition of CXCR4 in vitro and in vivo by using chemical inhibitor and neutralizing antibody;</li> <li>– Combination treatment with AMD3100 and Taxotere (Sanofi SA) (docetaxel)</li> </ul>	110
Colon	CD133	Migration and metastasis	<ul style="list-style-type: none"> <li>– Analysis of the FASC sorted patient-derived cells for their migratory potential in vitro and metastatic properties in vivo</li> </ul>	204
Pancreas	CD133	Migration, tumorigenicity, drug resistance and metastasis	<ul style="list-style-type: none"> <li>– Analysis of the FASC sorted patient-derived cells for their migratory properties in vitro, tumorigenic and metastatic properties in vivo;</li> <li>– Analysis of the FASC sorted patient-derived cells for their resistance to the standard chemotherapeutic agent gemcitabine in vitro and in vivo</li> </ul>	205
Glioma	CD133	Clonogenicity, spherogenicity, tumorigenicity, differentiation potential, chemo- and radioresistance	<ul style="list-style-type: none"> <li>– Analysis of the FASC sorted patient-derived cells for their clonogenic properties, spherogenicity, and differentiation potential in vitro, and tumorigenicity in vivo;</li> <li>– Analysis of the FASC sorted patient-derived cells for their resistance to the standard chemotherapeutic agent temozolomide and radiation treatment</li> </ul>	206
Gefitinib-resistant non-small cell lung cancer	N/A	Spherogenicity, chemo- and radioresistance, tumorigenicity	<ul style="list-style-type: none"> <li>– Analysis of the FASC sorted A549/GR cells for their sphere forming and self-renewal capacity in vitro and tumorigenic properties in vivo;</li> <li>– Analysis of the FASC sorted A549/GR cells for their resistance to irradiation and to the standard chemotherapeutic agent gefitinib and cisplatin in vitro</li> </ul>	176

**Abbreviations:** CXCR, CXC chemokine receptor; CXCL, chemokine (C-X-C motif) ligand; CSC, cancer stem cell; FACS, fluorescence-activated cell sorting; siRNA, small interfering RNA.

CTC population were positive for the CSC marker aldehyde dehydrogenase (ALDH).<sup>96</sup>

Although the signals generated by the metastatic niche that regulate CSCs are not fully understood, recent studies provide strong evidence for the important role of the chemokine receptor CXCR4 for CSC maintenance, dissemination, and consequent metastatic colonization. The role of CXCR4 has been examined in the CSC context in various types of cancer including pancreatic, colon, renal, brain, lung, and prostate cancer (Table 1 and Figure 2). Moreover, Fusi et al showed that CTCs from patients with metastatic carcinoma or melanoma are positive for CXCR4 expression.<sup>97</sup>

All together, these findings demonstrate that activation of the CXCL12/CXCR4 signaling can be indicative of the metastatic CSC population and suggest that therapeutic modulation of the CXCR4/CXCL12 axis may be essential for inhibition of metastatic and tumorigenic potential of CSCs.

## Regulation and biological effect of CXCR4

The CXCR4/CXCL12 signaling in tumor cells is regulated at several levels.<sup>49</sup> First, expression of the CXCR4 and CXCL12 genes are regulated at the transcriptional level by hypoxia. Several reports suggest that low tumor oxygenation and other signals from the tumor microenvironment such as growth factors collaborate to promote EMT associated with high invasiveness and resistance to chemo- and radiotherapy.<sup>98</sup> In hypoxia, the lack of oxygen leads to the hypoxia inducible factor 1 $\alpha$  (HIF1 $\alpha$ )-dependent activation of the CXCR4 and CXCL12 expression.<sup>99,100</sup> In addition, tumor fibroblasts and macrophages within the irradiated tumor niche start to produce growth factors and cytokines including CXCL12 which may lead to the invasive behavior of CSCs.<sup>101,102</sup> The overlapping signaling mechanisms that govern tumor resistance to conventional treatment and invasiveness could explain why, in some cases, cancer, which relapses after treatment, can develop into more aggressive metastatic disease, which is difficult to treat, and is associated with poor clinical prognosis.<sup>103–106</sup>

Studies of the molecular interaction between stroma and tumor cells suggest that CSCs can acquire resistance to chemo- and radiotherapy by adhesion to extracellular matrix or accessory cells via integrin signaling and, thus, may be responsible for residual disease and relapses. At the molecular level, CXCR4 is an important mediator of the interaction of tumor cells with extracellular matrix proteins such as laminin, fibronectin, and collagen, which contributes to metastatic spread.<sup>107–109</sup> Our recent study demonstrated that CXCL12/

CXCR4 signaling pathway regulates the adhesion of CD133<sup>+</sup>/CD44<sup>+</sup> prostate cancer progenitors to the extracellular protein fibronectin that is important for metastatic process. Moreover, the expression of  $\alpha$ 5 and  $\beta$ 3 integrin subunits, which form receptors for fibronectin, are strongly upregulated in CD133<sup>+</sup>/CD44<sup>+</sup> progenitor cells compared to CD133<sup>-</sup>/CD44<sup>-</sup> prostate cancer cells.<sup>110</sup> Despite the fact that CXCR4 does not directly modulate cell attachment, CXCR4 receptor engagement by CXCL12 plays an essential role in managing cell adhesion by modulation of integrin expression, FAK phosphorylation, and activation of p38 MAPK and ROCK kinases.<sup>108,111</sup> The disruption of the interaction of cancer cells with their microenvironmental milieu by CXCR4 inhibition leads to their sensitization to the cytotoxic therapeutic agents.<sup>112,113</sup> These findings are consistent with data of high CXCR4 expression by nasopharyngeal carcinoma cells in postradiotherapy patients.<sup>114</sup> Moreover, elevated CXCR4 expression shows prognostic value for patients with renal, colorectal, and breast carcinoma.<sup>115–119</sup>

In addition to HIF-1 $\alpha$ , some other transcription factors can influence CXCR4 transcription, including v-ets erythroblastosis virus E26 oncogene homolog 1 and NF- $\kappa$ B nuclear factor kappa-light-chain enhancer of activated B cells, which mediate CXCR4-dependent tumor invasion upon stimulation with hepatocyte growth factor.<sup>37,120,121</sup> Furthermore, a novel vesnarinone-responsive molecule Krüppel-like factor 2 and histone deacetylase 3-interacting protein CREB3 were also shown to activate the transcription of the CXCR4 and, therefore, contribute to cell migration.<sup>122,123</sup> CXCR4 expression and function are positively regulated by the developmental signaling pathways Wnt, SHH and Notch and the oncogenic pathways PI3K/AKT, NF- $\kappa$ B, and JAK/STAT that are also strongly implicated as CSC regulators.<sup>124–128</sup> In turn, activation of the CXCL12/CXCR4 signaling may affect these pathways, suggesting a positive feedback loop between CXCR4 and the signaling routes regulating self-renewal capacity and tumorigenicity of cancer cells.<sup>126,127,129–133</sup> At the intracellular level, CXCL12/CXCR4 signaling triggers several phosphorylation cascades controlled by Src and AKT.

The PI3K/AKT axis serves as the central route in the CXCL12/CXCR4 signaling cascade.<sup>134–135</sup> Recently, we showed that activation of the CXCL12/CXCR4 and PI3K/AKT signaling pathways is important for self-renewal and tumorigenicity of prostate cancer cells with stem cell characteristics.<sup>136,137</sup> PI3K/AKT signaling regulates transcription through the FOXO by phosphorylating conserved serine/threonine residues. Transcriptionally active FOXOs affect a wide range of biological processes, including cell survival,

DNA repair, oxidative stress response, and longevity.<sup>138</sup> Among the members of the FOXO family, FOXO3A has been shown to be important for the maintenance of neural, hematopoietic, endothelial stem cells,<sup>139–141</sup> and cancer stem-like cell populations.<sup>136,137</sup> The chromatin immunoprecipitation assay demonstrated that FOXO3A binds to the CXCR4 promoter.<sup>110</sup> These data suggested that the CXCR4/AKT positive feedback system may play a role in the maintenance and dissemination of the prostate cancer progenitors.

In addition, CXCL12/CXCR4 signaling may promote tumor growth through transactivation of receptor tyrosine kinases such as EGFR, IGF-1R, and FGFR, which contributes to enhanced invasive signals and metastatic growth of breast, prostate, and ovarian tumors.<sup>142–144</sup>

CXCR4 has also been demonstrated to elicit intracellular signals through interaction with the scaffolding proteins independently of heterotrimeric G-protein coupling.<sup>145</sup> For example, CXCR4 signaling can be modulated by  $\beta$ -arrestin that induces CXCR4 internalization and attenuates CXCR4-mediated G protein activation.  $\beta$ -arrestin can be recruited to the CXCR4/CXCR7 heterodimeric complex resulting in potentiation of downstream  $\beta$ -arrestin-dependent cell signaling pathways, including ERK1/2, p38 MAPK, and SAPK/JNK, which enhances cell migration in response to CXCL12 stimulation.<sup>21,24,25,146–148</sup>

Finally, recent studies support a new view of CXCR7 as a signaling receptor independent of G proteins.<sup>149</sup> CXCL12 binding to CXCR7 activates the PI3K/AKT, PLC/MAPK, and protein kinase C pathways and promotes tumor growth, neovascularization, and dissemination.<sup>24,25,146</sup>

In summary, activation of CXCL12/CXCR4 axis may be critical for different aspects of tumor initiation, progression, metastasis, and therapy resistance, and targeting CXCR4 signaling might be beneficial in cancer treatment.

## Critical analysis of the potential for targeting the CXCL12/CXCR4 axis in cancer management

Multiple agents are currently being developed to target CXCL12/CXCR4 signaling in cancer. Among these inhibitors is anti-CXCR4 drug AMD3100, also known as plerixafor (Mozobil; Sanofi SA, Paris, France), which is approved for stem cell mobilization in patients with non-Hodgkin's lymphoma and multiple myeloma, while the CXCL12 analog CTCE-9908 (Chemokine Therapeutics Corp, Vancouver, BC, Canada) is approved for clinical use in patients with osteosarcoma. Novel CXCR4 antagonists BKT140 (Emory University), POL6326 (Polyphor Ltd,

Allschwill, Switzerland), and TG-0054 (ChemoCentryx, Inc, Mountain View, CA, USA), which were characterized as powerful human stem cell mobilizers, are currently in clinical trials for multiple myeloma, leukemia, and lymphoma. NOX-A12 (Noxxon Pharma AG, Berlin, Germany) is the only anticancer agent in active clinical development that neutralizes CXCL12 resulting in a complete block of CXCL12 signaling through its two receptors, CXCR4 and CXCR7. The anti-CXCL12 aptamer NOX-A12 is in clinical trial for the treatment of chronic lymphocytic leukemia and multiple myeloma. CXCR4 inhibitor MSX-122 (Altiris Therapeutics Inc, Tucker, GA, USA) is in Phase I trials for advanced malignant diseases that are metastatic or unresectable and that are resistant to standard therapy, while CXCR7-specific inhibitor CCX2066 (ChemoCentryx, Inc) is in preclinical studies.<sup>134,150</sup>

CXCR4 antagonist AMD3100 is the most studied among the agents that inhibit CXCL12/CXCR4 signaling. AMD3100 was initially studied as an anti-HIV agent and then it was discovered that this compound increases white blood cell counts in the blood and is able to mobilize stem cells from the bone marrow. This observation led to the examination of its anticancer activity.<sup>151</sup> AMD3100 has already been shown to decrease metastatic potential in animal models for different types of tumors, including breast, ovarian and colorectal cancer, melanoma, and oral squamous cell carcinoma.<sup>152–158</sup> Similarly, blocking CXCR4 receptor function by a monoclonal antibody or polypeptide inhibits cancer cell proliferation, motility, and invasion in multiple preclinical models both in vitro and in vivo.<sup>110,113,159,160</sup>

The fact that CXCR4 is present in normal and cancer stem-like cells in various tissues suggests that this molecule could be essential for maintenance and viability of tumor progenitor cell population. Indeed, recent data suggest that inactivation of the CXCL12/CXCR4 axis by neutralizing antibody or with the CXCR4-specific small molecule antagonist AMD3100 inhibits glioma, renal, colon, pancreas, and prostate cancer progenitors as well as tumor initiating population within gefitinib-resistant lung cancer and tamoxifen-resistant breast cancer cells in vitro and in animal models (Table 1).<sup>107,161–163</sup>

Preclinical and clinical data demonstrated that tumor cells can be protected from the effect of ionizing radiation by hypoxia, and determination of microenvironmental parameters such as tumor hypoxic fraction, vasculature, and perfusion may have a prognostic value for the response to radiotherapy.<sup>164–166</sup> Moreover, low oxygen tension is a critical microenvironmental factor in regulating tumor initiating

cells.<sup>98,167</sup> Hypoxia promotes expansion of glioma and colon CSCs and converts non-stem cancer cells into CSC populations with increased self-renewal capacity.<sup>168,169</sup> The effects of reduced oxygen tension on CSCs are mediated at least in part through the activation of the HIF signaling pathway.<sup>167</sup> As described above, CXCR4 expression is also induced under hypoxic stress via activation of the HIF pathway.<sup>99,100</sup>

Pharmacologic inhibition of the CXCL12/CXCR4 interaction by AMD3100 or neutralizing antibody prevents the recurrence of glioblastoma after irradiation in mice by inhibition of vasculogenesis.<sup>170</sup> Preclinical studies have shown that radiation upregulates HIF-1 expression level and activity in vivo.<sup>171,172</sup> This induces the *CXCL12* gene expression and promotes the mobilization of CD11b<sup>+</sup> monocytes from the BM, recruitment of these BM derived cells into the tumors, and development of functional tumor vasculature,<sup>170,173,174</sup> thereby supporting all remaining viable tumor cells. Similarly, concomitant treatment with local irradiation and AMD3100 induced a significant tumor growth delay and increased radiocurability in lung tumors by retention of BM derived cells.<sup>175</sup>

Another study demonstrated the role of CXCR4 in tumor radioresistance more definitively and showed that activation of CXCR4-mediated STAT3 signaling in non-small cell lung cancer cells (NSCLC) is functionally crucial for the maintenance of stemness and resistance to radiotherapy.<sup>176</sup> Another molecular route that can underlie the CXCR4-mediated radioprotection is the integrin signaling pathway. Accumulating evidence suggests that CXCR4 route enhances integrin-mediated adhesion and cooperates with integrin signaling in mediating chemoresistance.<sup>177–179</sup> In fact, CXCR4 engagement by CXCL12 induces expression of the integrin receptors such as  $\alpha 3$ ,  $\alpha 5$ ,  $\beta 1$ , and  $\beta 3$  subunits, activation of FAK, and integrin-linked kinase, which is accompanied by the up-regulation of ERK1/2, JNK, and p38 phosphorylation.<sup>108,179–182</sup> Previous findings showed that integrins might induce tumor radioresistance via activating SAPK/JNK, MEK1/2, PI3K/AKT, and NF- $\kappa$ B signaling pathways.<sup>183–185</sup>

Collectively, these clinical and preclinical data are consistent with the CXCL12/CXCR4 pathway being a potential target to inhibit tumor growth and neovascularization, metastatic dissemination, and therapy resistance. Furthermore, the availability of pharmacologic inhibitors impinging CXCL12/CXCR4 signaling pathway opens novel opportunities for translational and clinical studies. However, important challenges remain prior to the clinical use of CXCR4 inhibitors in patients with solid tumors. First, CXCR4 is expressed by

numerous types of healthy tissues.<sup>26–29,33–39,183–188</sup> Interference with CXCL12/CXCR4 signaling leads to deficiencies in hematopoiesis and organ homeostatic functions as well as in tissue repair after various stresses and insults, including cytotoxic drugs and radiation injury. Given the ubiquitous expression of CXCR4 and the functional importance of the CXCL12/CXCR4 signaling axis, this may impede the use of the CXCR4-targeted therapeutic tools in the clinic. Clinical studies of AMD3100 as a mobilizer of HSC in non-Hodgkin's lymphoma and multiple myeloma patients have demonstrated that AMD3100 has minimal side effects. However, in the clinic, AMD3100 is given to the patients as a daily subcutaneous injection with granulocyte colony-stimulating factor for a limited time (1–7 days),<sup>189</sup> whereas in preclinical studies for evaluating the antitumor efficacy of AMD3100, it is often delivered continuously via a subcutaneous osmotic infusion pump.<sup>110,162,190,191</sup> In clinical study for HIV treatment, AMD3100 was administered as a daily intravenous infusion or subcutaneous injection for a period of 11–14 days and, despite its efficacy for the treatments for HIV patients, the trials were discontinued due to cardiac toxicity.<sup>192,193</sup>

Accumulating experimental evidence suggests that combinatorial strategies based on bulk tumor reduction and CSC-specific pathway inhibition offer a promising treatment modality and are predicted to have a greater efficacy in tumor reduction and prevention of relapse than monotherapies.<sup>88,110,137</sup> If these conditions can be met, the use of two types of therapy in low-dose combination can provide better therapeutic effects with less side-effect toxicity. Recent prostate tumor xenograft studies in mice showed that a combination of AMD3100, which targets prostate cancer stem-like cells, and the conventional chemotherapeutic drug Taxotere (Sanofi SA), which targets the bulk tumor, is significantly more effective in eradicating tumors as compared to monotherapy.<sup>110,136,137,194</sup> However, efficacy of CXCR4 inhibitors against CSC function in cancer patients remains to be determined. The ongoing clinical trials for CXCR4 inhibitors as chemosensitizers in acute myeloid leukemia and other hematological malignancies will help to elucidate this question.

## Conclusion

Although the collective evidence from the preclinical and clinical studies support the potential efficacy of CXCR4 inhibitors for development of innovative approaches to cancer treatment, several significant challenges remain before translation of these inhibitors into the clinic. A major

factor that can prevent a successful clinical use of the CXCR4-targeting anticancer therapy is a potential side effect on the stem cell compartment in normal tissues. This may be especially important when this treatment is combined with radiotherapy and other cytotoxic therapy associated with depletion of normal tissue progenitors. Thus, biology-driven rational design of novel combination therapies will be critical for the development of low-side-effect cancer treatment and can be based on the synergistic antitumor effect of CXCR4 inhibition and conventional therapy. There is also a need for the evaluation of the relationship between CXCR4 and tumor initiating cells in cancer patients. In fact, evaluation of the CXCR4 and CXCL12 expression level may have significant prognostic value in various types of cancer, including glioma, prostate, breast, colon, ovarian, pancreatic, and lung cancer where high expression of CXCR4 or CXCL12 predicts poor patient outcome.<sup>195–203</sup> However, direct proof for stemness of CXCR4<sup>+</sup> cells from primary human tumor tissues is still missing. A better understanding of the role of CXCR4 pathway for the maintenance of tumor progenitor population may be necessary for the development of screening tests to identify the patients who are likely to respond to CXCR4 inhibition. In addition, recent discovery of the cancer stem cell plasticity and heterogeneity can make the CXCR4<sup>+</sup> tumor cell population a moving target that could be hard to track and eradicate.<sup>85</sup> Nevertheless, the key role of CXCR4 in tumor initiation, vascularization, dissemination, and therapy resistance underscore the importance of CXCR4 inhibition for the optimization of current anticancer treatment strategies.

## Acknowledgment

The authors wish to acknowledge funding from the German Federal Ministry of Education and Research (Grant-No 03Z1NN11).

## Disclosure

The authors report no conflicts of interest in this work.

## References

- Viola A, Luster AD. Chemokines and their receptors: drug targets in immunity and inflammation. *Annu Rev Pharmacol Toxicol.* 2008;48:171–197.
- Smith H, Whittall C, Weksler B, Middleton J. Chemokines stimulate bidirectional migration of human mesenchymal stem cells across bone marrow endothelial cells. *Stem Cells Dev.* 2012;21(3):476–486.
- Balkwill F. Cancer and the chemokine network. *Nat Rev Cancer.* 2004;4(7):540–550.
- Gilman AG. G proteins: transducers of receptor-generated signals. *Annu Rev Biochem.* 1987;56:615–649.
- Pierce KL, Premont RT, Lefkowitz RJ. Seven-transmembrane receptors. *Nat Rev Mol Cell Biol.* 2002;3(9):639–650.
- New DC, Wong YH. Molecular mechanisms mediating the G protein-coupled receptor regulation of cell cycle progression. *J Mol Signal.* 2007;2:2.
- Mellado M, Rodríguez-Frade JM, Mañes S, Martínez-A C. Chemokine signaling and functional responses: the role of receptor dimerization and TK pathway activation. *Annu Rev Immunol.* 2001;19:397–421.
- Epstein RJ. The CXCL12-CXCR4 chemotactic pathway as a target of adjuvant breast cancer therapies. *Nat Rev Cancer.* 2004;4(11):901–909.
- Lattin J, Zidar DA, Schroder K, Kellie S, Hume DA, Sweet MJ. G-protein-coupled receptor expression, function, and signaling in macrophages. *J Leukoc Biol.* 2007;82(1):16–32.
- Gerits N, Kostenko S, Shiryayev A, Johannessen M, Moens U. Relations between the mitogen-activated protein kinase and the cAMP-dependent protein kinase pathways: comradeship and hostility. *Cell Signal.* 2008;20(9):1592–1607.
- Princen K, Hatse S, Vermeire K, De Clercq E, Schols D. Evaluation of SDF-1/CXCR4-induced Ca<sup>2+</sup> signaling by fluorometric imaging plate reader (FLIPR) and flow cytometry. *Cytometry A.* 2003;51(1):35–45.
- Zhang R, Xie X. Tools for GPCR drug discovery. *Acta Pharmacol Sin.* 2012;33(3):372–384.
- Schwindinger WF, Robishaw JD. Heterotrimeric G-protein betagamma-dimers in growth and differentiation. *Oncogene.* 2001;20(13):1653–1660.
- Rozenfurt E. Signal transduction pathways in the mitogenic response to G protein-coupled neuropeptide receptor agonists. *J Cell Physiol.* 1998;177(4):507–517.
- Liang J, Slingerland JM. Multiple roles of the PI3K/PKB (Akt) pathway in cell cycle progression. *Cell Cycle.* 2003;2(4):339–345.
- McCudden CR, Hains MD, Kimple RJ, Siderovski DP, Willard FS. G-protein signaling: back to the future. *Cell Mol Life Sci.* 2005;62(5):551–577.
- Teicher BA, Fricker SP. CXCL12 (SDF-1)/CXCR4 pathway in cancer. *Clin Cancer Res.* 2010;16(11):2927–2931.
- Ferguson SS. Evolving concepts in G protein-coupled receptor endocytosis: the role in receptor desensitization and signaling. *Pharmacol Rev.* 2001;53(1):1–24.
- Luttrell LM, Gesty-Palmer D. Beyond desensitization: physiological relevance of arrestin-dependent signaling. *Pharmacol Rev.* 2010;62(2):305–330.
- Mellado M, Rodríguez-Frade JM, Vila-Coro AJ, et al. Chemokine receptor homo- or heterodimerization activates distinct signaling pathways. *EMBO J.* 2001;20(10):2497–2507.
- Décaillot FM, Kazmi MA, Lin Y, Ray-Saha S, Sakmar TP, Sachdev P. CXCR7/CXCR4 heterodimer constitutively recruits beta-arrestin to enhance cell migration. *J Biol Chem.* 2011;286(37):32188–32197.
- Berger EA, Murphy PM, Farber JM. Chemokine receptors as HIV-1 coreceptors: roles in viral entry, tropism, and disease. *Annu Rev Immunol.* 1999;17:657–700.
- Sun X, Cheng G, Hao M, et al. CXCL12/CXCR4/CXCR7 chemokine axis and cancer progression. *Cancer Metastasis Rev.* 2010;29(4):709–722.
- Singh AK, Arya RK, Trivedi AK, et al. Chemokine receptor trio: CXCR3, CXCR4 and CXCR7 crosstalk via CXCL11 and CXCL12. *Cytokine Growth Factor Rev.* 2013;24(1):41–49.
- Yu S, Crawford D, Tsuchihashi T, Behrens TW, Srivastava D. The chemokine receptor CXCR7 functions to regulate cardiac valve remodeling. *Dev Dyn.* 2011;240(2):384–393.
- Aiuti A, Webb IJ, Bleul C, Springer T, Gutierrez-Ramos JC. The chemokine SDF-1 is a chemoattractant for human CD34<sup>+</sup> hematopoietic progenitor cells and provides a new mechanism to explain the mobilization of CD34<sup>+</sup> progenitors to peripheral blood. *J Exp Med.* 1997;185:111–120.
- Ma Q, Jones D, Borghesani PR, et al. Impaired B-lymphopoiesis, myelopoiesis and derailed cerebellar neuron migration in CXCR4- and SDF-1-deficient mice. *Proc Natl Acad Sci U S A.* 1998;95:9448–9453.
- Zou YR, Kottmann AH, Kuroda M, Taniuchi I, Littman DR. Function of the chemokine receptor CXCR4 in haematopoiesis and in cerebellar development. *Nature.* 1998;393(6685):595–599.

29. Kucia M, Jankowski K, Reza R, et al. CXCR4-SDF-1 signalling, locomotion, chemotaxis and adhesion. *J Mol Histol.* 2004;35(3):233–245.
30. Janowski M. Functional diversity of SDF-1 splicing variants. *Cell Adh Migr.* 2009;3(3):243–249.
31. Yu L, Cecil J, Peng SB, et al. Identification and expression of novel isoforms of human stromal cell-derived factor 1. *Gene.* 2006;374: 174–179.
32. Ho TK, Shiwen X, Abraham D, Tsui J, Baker D. Stromal-cell-derived factor-1 (SDF-1)/CXCL12 as potential target of therapeutic angiogenesis in critical leg ischaemia. *Cardiol Res Pract.* 2012;2012:143209.
33. Ratajczak MZ, Majka M, Kucia M, et al. Expression of functional CXCR4 by muscle satellite cells and secretion of SDF-1 by muscle-derived fibroblasts is associated with the presence of both muscle progenitors in bone marrow and hematopoietic stem/progenitor cells in muscles. *Stem Cells.* 2003;21(3):363–371.
34. Kucia M, Reza R, Miekus K, et al. Trafficking of normal stem cells and metastasis of cancer stem cells involve similar mechanisms: pivotal role of the SDF-1-CXCR4 axis. *Stem Cells.* 2005;23(7):879–894.
35. Day CE, Guillen C, Willars GB, Wardlaw AJ. Characterization of the migration of lung and blood T cells in response CXCL12 in a three-dimensional matrix. *Immunology.* 2010;130(4):564–571.
36. Agle KA, Vongsa RA, Dwinell MB. Calcium mobilization triggered by the chemokine CXCL12 regulates migration in wounded intestinal epithelial monolayers. *J Biol Chem.* 2010;285(21):16066–16075.
37. Esencay M, Newcomb EW, Zagzag D. HGF upregulates CXCR4 expression in gliomas via NF-kappaB: implications for glioma cell migration. *J Neurooncol.* 2010;99(1):33–40.
38. Burger JA, Kipps TJ. CXCR4: a key receptor in the crosstalk between tumor cells and their microenvironment. *Blood.* 2006;107(5):1761–1767.
39. Tiveron MC, Cremer H. CXCL12/CXCR4 signalling in neuronal cell migration. *Curr Opin Neurobiol.* 2008;18(3):237–244.
40. Schober A, Zernecke A. Chemokines in vascular remodeling. *Thromb Haemost.* 2007;97(5):730–737.
41. Petit I, Jin D, Rafii S. The SDF-1-CXCR4 signaling pathway: a molecular hub modulating neo-angiogenesis. *Trends Immunol.* 2007;28(7): 299–307.
42. Karshovska E, Zagorac D, Zernecke A, Weber C, Schober A. A small molecule CXCR4 antagonist inhibits neointima formation and smooth muscle progenitor cell mobilization after arterial injury. *J Thromb Haemost.* 2008;6(10):1812–1815.
43. Zernecke A, Schober A, Bot I, et al. SDF-1alpha/CXCR4 axis is instrumental in neointimal hyperplasia and recruitment of smooth muscle progenitor cells. *Circ Res.* 2005;96(7):784–791.
44. Rehimi R, Khalida N, Yusuf F, Morosan-Puopolo G, Brand-Saberi B. A novel role of CXCR4 and SDF-1 during migration of cloacal muscle precursors. *Dev Dyn.* 2010;239(6):1622–1631.
45. Nervi B, Link DC, DiPersio JF. Cytokines and hematopoietic stem cell mobilization. *J Cell Biochem.* 2006;99(3):690–705.
46. Yu J, Li M, Qu Z, Yan D, Li D, Ruan Q. SDF-1/CXCR4-mediated migration of transplanted bone marrow stromal cells toward areas of heart myocardial infarction through activation of PI3K/Akt. *J Cardiovasc Pharmacol.* 2010;55(5):496–505.
47. Gambaryan N, Perros F, Montani D, et al. Targeting of c-kit+ haematopoietic progenitor cells prevents hypoxic pulmonary hypertension. *Eur Respir J.* 2011;37(6):1392–1399.
48. Yu L, Hales CA. Effect of chemokine receptor CXCR4 on hypoxia-induced pulmonary hypertension and vascular remodeling in rats. *Respir Res.* 2011;12:21.
49. Ratajczak MZ, Zuba-Surma E, Kucia M, et al. The pleiotropic effects of the SDF-1-CXCR4 axis in organogenesis, regeneration and tumorigenesis. *Leukemia.* 2006;20(11):1915–1924.
50. Bleul CC, Fuhlbrigge RC, Casasnovas JM, Aiuti A, Springer TA. A highly efficacious lymphocyte chemoattractant, stromal cell-derived factor 1 (SDF-1). *J Exp Med.* 1996;184(3):1101–1109.
51. Lazarini F, Tham TN, Casanova P, Arenzana-Seisdedos F, Dubois-Dalcq M. Role of the alpha-chemokine stromal cell-derived factor (SDF-1) in the developing and mature central nervous system. *Glia.* 2003;42(2):139–148.
52. Vagima Y, Lapid K, Kollet O, Goichberg P, Alon R, Lapidot T. Pathways implicated in stem cell migration: the SDF-1/CXCR4 axis. *Methods Mol Biol.* 2011;750:277–289.
53. Burger JA, Burger M, Kipps TJ. Chronic lymphocytic leukemia B cells express functional CXCR4 chemokine receptors that mediate spontaneous migration beneath bone marrow stromal cells. *Blood.* 1999;94(11):3658–3667.
54. Möhle R, Failenschmid C, Bautz F, Kanz L. Overexpression of the chemokine receptor CXCR4 in B cell chronic lymphocytic leukemia is associated with increased functional response to stromal cell-derived factor-1 (SDF-1). *Leukemia.* 1999;13(12):1954–1959.
55. Müller A, Homey B, Soto H, et al. Involvement of chemokine receptors in breast cancer metastasis. *Nature.* 2001;410(6824):50–56.
56. Balkwill F. The significance of cancer cell expression of the chemokine receptor CXCR4. *Semin Cancer Biol.* 2004;14(3):171–179.
57. Vandercappellen J, Van Damme J, Struyf S. The role of CXC chemokines and their receptors in cancer. *Cancer Lett.* 2008;267(2):226–244.
58. Zlotnik A. New insights on the role of CXCR4 in cancer metastasis. *J Pathol.* 2008;215(3):211–213.
59. Furusato B, Mohamed A, Uhlén M, et al. CXCR4 and cancer. *Pathol Int.* 2010;60(7):497–505.
60. Darash-Yahana M, Pikarsky E, Abramovitch R, et al. Role of high expression levels of CXCR4 in tumor growth, vascularization, and metastasis. *FASEB J.* 2004;18(11):1240–1242.
61. Sun X, Wei L, Chen Q, Terek RM. CXCR4/SDF1 mediate hypoxia induced chondrosarcoma cell invasion through ERK signaling and increased MMP1 expression. *Mol Cancer.* 2010;9:17.
62. Eck SM, Blackburn JS, Schmucker AC, Burrage PS, Brinckerhoff CE. Matrix metalloproteinase and G protein coupled receptors: co-conspirators in the pathogenesis of autoimmune disease and cancer. *J Autoimmun.* 2009;33(3–4):214–221.
63. Orimo A, Weinberg RA. Stromal fibroblasts in cancer: a novel tumor-promoting cell type. *Cell Cycle.* 2006;5(15):1597–1601.
64. Xu J, Clark RA. Extracellular matrix alters PDGF regulation of fibroblast integrins. *J Cell Biol.* 1996;132:239–249.
65. Dunphy JE. The Fibroblast – A Ubiquitous Ally for the Surgeon. *N Engl J Med.* 1963;268:1367–1377.
66. Mishra PJ, Mishra PJ, Humeniuk R, et al. Carcinoma-associated fibroblast-like differentiation of human mesenchymal stem cells. *Cancer Res.* 2008;68(11):4331–4339.
67. Orimo A, Gupta PB, Sgroi DC, et al. Stromal fibroblasts present in invasive human breast carcinomas promote tumor growth and angiogenesis through elevated SDF-1/CXCL12 secretion. *Cell.* 2005;121(3):335–348.
68. Shangguan L, Ti X, Krause U, et al. Inhibition of TGF-β/Smad signaling by BAMBI blocks differentiation of human mesenchymal stem cells to carcinoma-associated fibroblasts and abolishes their protumor effects. *Stem Cells.* 2012;30(12):2810–2819.
69. McDonald LT, LaRue AC. Hematopoietic stem cell derived carcinoma-associated fibroblasts: a novel origin. *Int J Clin Exp Pathol.* 2012;5(9):863–873.
70. Shuman Moss LA, Jensen-Taubman S, Stetler-Stevenson WG. Matrix metalloproteinases: changing roles in tumor progression and metastasis. *Am J Pathol.* 2012;181(6):1895–1899.
71. Eck SM, Cote AL, Winkelman WD, et al. CXCR4 and matrix metalloproteinase-1 are elevated in breast carcinoma-associated fibroblasts and in normal mammary fibroblasts exposed to factors secreted by breast cancer cells. *Mol Cancer Res.* 2009;7(7): 1033–1044.
72. Alsayed Y, Ngo H, Runnels J, et al. Mechanisms of regulation of CXCR4/SDF-1 (CXCL12)-dependent migration and homing in multiple myeloma. *Blood.* 2007;109:2708–2717.
73. Busillo JM, Benovic JL. Regulation of CXCR4 signaling. *Biochim Biophys Acta.* 2007;1768(4):952–963.
74. Zhou Y, Larsen PH, Hao C, Yong VW. CXCR4 is a major chemokine receptor on glioma cells and mediates their survival. *J Biol Chem.* 2002;277(51):49481–49487.

75. Luker KE, Luker GD. Functions of CXCL12 and CXCR4 in breast cancer. *Cancer Lett.* 2006;238(1):30–41.
76. Jung Y, Kim JK, Shiozawa Y, et al. Recruitment of mesenchymal stem cells into prostate tumours promotes metastasis. *Nat Commun.* 2013;4:1795.
77. Scheel C, Weinberg RA. Cancer stem cells and epithelial-mesenchymal transition: concepts and molecular links. *Semin Cancer Biol.* 2012; 22(5–6):396–403.
78. Gupta PB, Onder TT, Jiang G, et al. Identification of selective inhibitors of cancer stem cells by high-throughput screening. *Cell.* 2009;138(4):645–659.
79. Mani SA, Guo W, Liao MJ, et al. The epithelial-mesenchymal transition generates cells with properties of stem cells. *Cell.* 2008;133(4): 704–715.
80. Asiedu MK, Ingle JN, Behrens MD, Radisky DC, Knutson KL. TGF $\beta$ /TNF $\alpha$ -mediated epithelial-mesenchymal transition generates breast cancer stem cells with a claudin-low phenotype. *Cancer Res.* 2011;71(13):4707–4719.
81. Shackleton M, Quintana E, Fearon ER, Morrison SJ. Heterogeneity in cancer: cancer stem cells versus clonal evolution. *Cell.* 2009;138(5): 822–829.
82. Lapidot T, Sirard C, Vormoor J, et al. A cell initiating human acute myeloid leukaemia after transplantation into SCID mice. *Nature.* 1994;367(6464):645–648.
83. Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF. Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci U S A.* 2003;100(7):3983–3988.
84. Baccelli I, Trumpp A. The evolving concept of cancer and metastasis stem cells. *J Cell Biol.* 2012;198(3):281–293.
85. Peitzsch C, Kurth I, Kunz-Schughart L, Baumann M, Dubrovskaya A. Discovery of the cancer stem cell related determinants of radioresistance. *Radiother Oncol.* Epub July 3, 2013.
86. Baumann M, Krause M, Hill R. Exploring the role of cancer stem cells in radioresistance. *Nat Rev Cancer.* 2008;8(7):545–554.
87. Tang DG. Understanding cancer stem cell heterogeneity and plasticity. *Cell Res.* 2012;22(3):457–472.
88. Scaffidi P, Misteli T. In vitro generation of human cells with cancer stem cell properties. *Nat Cell Biol.* 2011;13(9):1051–1061.
89. Brabletz T, Jung A, Spaderna S, Hlubek F, Kirchner T. Opinion: migrating cancer stem cells – an integrated concept of malignant tumour progression. *Nat Rev Cancer.* 2005;5(9):744–749.
90. Hindriksen S, Bijlsma MF. Cancer Stem Cells, EMT, and Developmental Pathway Activation in Pancreatic Tumors. *Cancers.* 2012;4(4): 989–1035.
91. Dalerba P, Cho RW, Clarke MF. Cancer stem cells: models and concepts. *Annu Rev Med.* 2007;58:267–284.
92. Liu R, Wang X, Chen GY, et al. The prognostic role of a gene signature from tumorigenic breast-cancer cells. *N Engl J Med.* 2007;356(3): 217–226.
93. Liu H, Patel MR, Prescher JA, et al. Cancer stem cells from human breast tumors are involved in spontaneous metastases in orthotopic mouse models. *Proc Natl Acad Sci U S A.* 2010;107(42):18115–18120.
94. Pang R, Law WL, Chu AC, et al. A subpopulation of CD26+ cancer stem cells with metastatic capacity in human colorectal cancer. *Cell Stem Cell.* 2010;6(6):603–615.
95. Armstrong AJ, Marengo MS, Oltean S, et al. Circulating tumor cells from patients with advanced prostate and breast cancer display both epithelial and mesenchymal markers. *Mol Cancer Res.* 2011;9(8): 997–1007.
96. Aktas B, Tewes M, Fehm T, Hauch S, Kimmig R, Kasimir-Bauer S. Stem cell and epithelial-mesenchymal transition markers are frequently overexpressed in circulating tumor cells of metastatic breast cancer patients. *Breast Cancer Res.* 2009;11:R46.
97. Fusi A, Liu Z, Kümmerlen V, Nonnemacher A, Jeske J, Keilholz U. Expression of chemokine receptors on circulating tumor cells in patients with solid tumors. *J Transl Med.* 2012;10:52.
98. Hill RP, Marie-Egyptienne DT, Hedley DW. Cancer stem cells, hypoxia and metastasis. *Semin Radiat Oncol.* 2009;19(2):106–111.
99. Ishikawa T, Nakashiro K, Klosek SK, et al. Hypoxia enhances CXCR4 expression by activating HIF-1 in oral squamous cell carcinoma. *Oncol Rep.* 2009;21(3):707–712.
100. Zagzag D, Lukyanov Y, Lan L, et al. Hypoxia-inducible factor 1 and VEGF upregulate CXCR4 in glioblastoma: implications for angiogenesis and glioma cell invasion. *Lab Invest.* 2006;86(12): 1221–1232.
101. Nguyen DH, Oketch-Rabah HA, Illa-Bochaca I, et al. Radiation acts on the microenvironment to affect breast carcinogenesis by distinct mechanisms that decrease cancer latency and affect tumor type. *Cancer Cell.* 2011;19(5):640–651.
102. Sofia Vala I, Martins LR, Imaizumi N, et al. Low doses of ionizing radiation promote tumor growth and metastasis by enhancing angiogenesis. *PLoS One.* 2010;5:e11222.
103. Fagundes H, Perez CA, Grigsby PW, Lockett MA. Distant metastases after irradiation alone in carcinoma of the uterine cervix. *Int J Radiat Oncol Biol Phys.* 1992;24(2):197–204.
104. Anderson P, Dische S. Local tumor control and the subsequent incidence of distant metastatic disease. *Int J Radiat Oncol Biol Phys.* 1981;7(12):1645–1648.
105. Fuks Z, Leibel SA, Wallner KE, et al. The effect of local control on metastatic dissemination in carcinoma of the prostate: long-term results in patients treated with 125I implantation. *Int J Radiat Oncol Biol Phys.* 1991;21(3):537–547.
106. von Essen CF. Radiation enhancement of metastasis: a review. *Clin Exp Metastasis.* 1991;9(2):77–104.
107. Engl T, Relja B, Marian D, et al. CXCR4 chemokine receptor mediates prostate tumor cell adhesion through alpha5 and beta3 integrins. *Neoplasia.* 2006;8(4):290–301.
108. Sun YX, Fang M, Wang J, Cooper CR, Pienta KJ, Taichman RS. Expression and activation of alpha v beta 3 integrins by SDF-1/CXCL12 increases the aggressiveness of prostate cancer cells. *Prostate.* 2007;67(1):61–73.
109. Goel HL, Li J, Kogan S, Languino LR. Integrins in prostate cancer progression. *Endocr Relat Cancer.* 2008;15(3):657–664.
110. Dubrovskaya A, Elliott J, Salamone RJ, et al. CXCR4 Expression in Prostate Cancer Progenitor Cells. *PLoS ONE.* 2012;7(2):e31226.
111. Struckhoff AP, Vitko JR, Rana MK, et al. Dynamic regulation of ROCK in tumor cells controls CXCR4-driven adhesion events. *J Cell Sci.* 2010;123(Pt 3):401–412.
112. Azab AK, Runnels JM, Pitsillides C, et al. CXCR4 inhibitor AMD3100 disrupts the interaction of multiple myeloma cells with the bone marrow microenvironment and enhances their sensitivity to therapy. *Blood.* 2009;113(18):4341–4351.
113. Zeng Z, Samudio IJ, Munsell M, et al. Inhibition of CXCR4 with the novel RCP168 peptide overcomes stroma-mediated chemoresistance in chronic and acute leukemias. *Mol Cancer Ther.* 2006;5(12): 3113–3121.
114. Ou DL, Chen CL, Lin SB, Hsu CH, Lin LI. Chemokine receptor expression profiles in nasopharyngeal carcinoma and their association with metastasis and radiotherapy. *J Pathol.* 2006;210(3):363–373.
115. Parker CC, Kim RH, Li BD, Chu QD. The chemokine receptor CXCR4 as a novel independent prognostic marker for node-positive breast cancer patients. *J Surg Oncol.* 2012;106(4):393–398.
116. Hiller DJ, Meschonat C, Kim R, Li BD, Chu QD. Chemokine receptor CXCR4 level in primary tumors independently predicts outcome for patients with locally advanced breast cancer. *Surgery.* 2011;150(3): 459–465.
117. Yopp AC, Shia J, Butte JM, et al. CXCR4 expression predicts patient outcome and recurrence patterns after hepatic resection for colorectal liver metastases. *Ann Surg Oncol.* 2012;19 Suppl 3:S339–S346.
118. D'Alterio C, Consales C, Polimeno M, et al. Concomitant CXCR4 and CXCR7 expression predicts poor prognosis in renal cancer. *Curr Cancer Drug Targets.* 2010;10(7):772–781.

119. Chu QD, Panu L, Holm NT, Li BD, Johnson LW, Zhang S. High chemokine receptor CXCR4 level in triple negative breast cancer specimens predicts poor clinical outcome. *J Surg Res*. 2010;159(2): 689–695.
120. Maroni P, Bendinelli P, Matteucci E, Desiderio MA. HGF induces CXCR4 and CXCL12-mediated tumor invasion through Ets1 and NF-kappaB. *Carcinogenesis*. 2007;28(2):267–279.
121. Matteucci E, Ridolfi E, Maroni P, Bendinelli P, Desiderio MA. c-Src/histone deacetylase 3 interaction is crucial for hepatocyte growth factor dependent decrease of CXCR4 expression in highly invasive breast tumor cells. *Mol Cancer Res* 2007;5(8):833–845.
122. Kim HC, Choi KC, Choi HK, et al. HDAC3 selectively represses CREB3-mediated transcription and migration of metastatic breast cancer cells. *Cell Mol Life Sci*. 2010;67(20):3499–3510.
123. Uchida D, Onoue T, Begum NM, et al. Vesnarinone downregulates CXCR4 expression via upregulation of Krüppel-like factor 2 in oral cancer cells. *Mol Cancer*. 2009;8:62.
124. Jin Z, Zhao C, Han X, Han Y. Wnt5a promotes Ewing sarcoma cell migration through upregulating CXCR4 expression. *BMC Cancer*. 2012;12:480.
125. Tamura M, Sato MM, Nashimoto M. Regulation of CXCL12 expression by canonical Wnt signaling in bone marrow stromal cells. *Int J Biochem Cell Biol*. 2011;43(5):760–767.
126. Han Y, He T, Huang DR, Pardo CA, Ransohoff RM. TNF-alpha mediates SDF-1 alpha-induced NF-kappa B activation and cytotoxic effects in primary astrocytes. *J Clin Invest*. 2001;108(3):425–435.
127. Kukreja P, Abdel-Mageed AB, Mondal D, Liu K, Agrawal KC. Up-regulation of CXCR4 expression in PC-3 cells by stromal-derived factor-1alpha (CXCL12) increases endothelial adhesion and transendothelial migration: role of MEK/ERK signaling pathway-dependent NF-kB activation. *Cancer Res*. 2005;65(21):9891–9898.
128. Soldevila G, Licona I, Salgado A, Ramírez M, Chávez R, García-Zepeda E. Impaired chemokine-induced migration during T-cell development in the absence of Jak 3. *Immunology*. 2004;112(2): 191–200.
129. Wang J, Cai J, Han F, et al. Silencing of CXCR4 blocks progression of ovarian cancer and depresses canonical Wnt signaling pathway. *Int J Gynecol Cancer*. 2011;21(6):981–987.
130. Fareh M, Turchi L, Virolle V, et al. The miR 302-367 cluster drastically affects self-renewal and infiltration properties of glioma-initiating cells through CXCR4 repression and consequent disruption of the SHH-GLI-NANOG network. *Cell Death Differ*. 2012;19(2):232–244.
131. Peng SB, Peek V, Zhai Y, et al. Akt activation, but not extracellular signal-regulated kinase activation, is required for SDF-1alpha/CXCR4-mediated migration of epitheloid carcinoma cells. *Mol Cancer Res*. 2005;3(4):227–236.
132. Helbig G, Christopherson KW 2nd, Bhat-Nakshatri P, et al. NF-kappaB promotes breast cancer cell migration and metastasis by inducing the expression of the chemokine receptor CXCR4. *J Biol Chem*. 2003;278(24):21631–21638.
133. Vila-Coro AJ, Rodríguez-Frade JM, Martín De Ana A, Moreno-Ortiz MC, Martínez-AC, Mellado M. The chemokine SDF-1alpha triggers CXCR4 receptor dimerization and activates the JAK/STAT pathway. *FASEB J*. 1999;13(13):1699–1710.
134. Ramsey DM, McAlpine SR. Halting metastasis through CXCR4 inhibition. *Bioorg Med Chem Lett*. 2013;23(1):20–25.
135. Chinni SR, Sivalogan S, Dong Z, et al. CXCL12/CXCR4 signaling activates Akt-1 and MMP-9 expression in prostate cancer cells: the role of bone microenvironment-associated CXCL12. *Prostate*. 2006;66(1):32–48.
136. Dubrovskaya A, Kim S, Salamone RJ, et al. The role of PTEN/Akt/PI3K signaling in the maintenance and viability of prostate cancer stem-like cell populations. *Proc Natl Acad Sci U S A*. 2009;106(1):268–273.
137. Dubrovskaya A, Elliott J, Salamone RJ, et al. Combination therapy targeting both tumor-initiating and differentiated cell populations in prostate carcinoma. *Clin Cancer Res*. 2010;16(23):5692–5702.
138. Huang H, Tindall DJ. FOXO factors: a matter of life and death. *Future Oncol*. 2006;2(1):83–89.
139. Miyamoto K, Araki KY, Naka K, et al. Foxo3a is essential for maintenance of the hematopoietic stem cell pool. *Cell Stem Cell*. 2007;1(1): 101–112.
140. Zhu S, Evans S, Yan B, et al. Transcriptional regulation of Bim by FOXO3a and Akt mediates scleroderma serum-induced apoptosis in endothelial progenitor cells. *Circulation*. 2008;118(21): 2156–2165.
141. Wu Y, Peng H, Cui M, et al. CXCL12 increases human neural progenitor cell proliferation through Akt-1/FOXO3a signaling pathway. *J Neurochem*. 2009;109(4):1157–1167.
142. Porcile C, Bajetto A, Barbieri F, et al. Stromal cell-derived factor-1alpha (SDF-1alpha/CXCL12) stimulates ovarian cancer cell growth through the EGF receptor transactivation. *Exp Cell Res*. 2005;308(2):241–253.
143. Chinni SR, Yamamoto H, Dong Z, Sabbota A, Bonfil RD, Cher ML. CXCL12/CXCR4 transactivates HER2 in lipid rafts of prostate cancer cells and promotes growth of metastatic deposits in bone. *Mol Cancer Res*. 2008;6(3):446–457.
144. Cabioglu N, Summy J, Miller C, et al. CXCL-12/stromal cell-derived factor-1alpha transactivates HER2-neu in breast cancer cells by a novel pathway involving Src kinase activation. *Cancer Res*. 2005;65(15): 6493–6497.
145. Defea K. Beta-arrestins and heterotrimeric G-proteins: collaborators and competitors in signal transduction. *Br J Pharmacol*. 2008; 153 Suppl 1:S298–S309.
146. Heinrich EL, Lee W, Lu J, Lowy AM, Kim J. Chemokine CXCL12 activates dual CXCR4 and CXCR7-mediated signaling pathways in pancreatic cancer cells. *J Transl Med*. 2012;10:68.
147. Sun Y, Cheng Z, Ma L, Pei G. Beta-arrestin 2 is critically involved in CXCR4-mediated chemotaxis, and this is mediated by its enhancement of p38 MAPK activation. *J Biol Chem*. 2002;277(51): 49212–49219.
148. Cheng ZJ, Zhao J, Sun Y, et al. Beta-arrestin differentially regulates the chemokine receptor CXCR4-mediated signaling and receptor internalization, and this implicates multiple interaction sites between beta-arrestin and CXCR4. *J Biol Chem*. 2000;275(4):2479–2485.
149. Sánchez-Martín L, Sánchez-Mateos P, Cabañas C. CXCR7 impact on CXCL12 biology and disease. *Trends Mol Med*. 2013;19(1):12–22.
150. de Nigris F, Schiano C, Infante T, Napoli C. CXCR4 inhibitors: tumor vasculature and therapeutic challenges. *Recent Pat Anticancer Drug Discov*. 2012;7(3):251–264.
151. Peled A, Wald O, Burger J. Development of novel CXCR4-based therapeutics. *Expert Opin Investig Drugs*. 2012;21(3):341–353.
152. Smith MC, Luker KE, Garbow JR, et al. CXCR4 regulates growth of both primary and metastatic breast cancer. *Cancer Res*. 2004;64(23):8604–8612.
153. Kim M, Koh YJ, Kim KE, et al. CXCR4 signaling regulates metastasis of chemoresistant melanoma cells by a lymphatic metastatic niche. *Cancer Res*. 2010;70(24):10411–10421.
154. D'Alterio C, Barbieri A, Portella L, et al. Inhibition of stromal CXCR4 impairs development of lung metastases. *Cancer Immunol Immunother*. 2012;61(10):1713–1720.
155. Uchida D, Onoue T, Kuribayashi N, et al. Blockade of CXCR4 in oral squamous cell carcinoma inhibits lymph node metastases. *Eur J Cancer*. 2011;47(3):452–459.
156. Kajiyama H, Shibata K, Terauchi M, Ino K, Nawa A, Kikkawa F. Involvement of SDF-1alpha/CXCR4 axis in the enhanced peritoneal metastasis of epithelial ovarian carcinoma. *Int J Cancer*. 2008;122(1): 91–99.
157. Matsusue R, Kubo H, Hisamori S, et al. Hepatic stellate cells promote liver metastasis of colon cancer cells by the action of SDF-1/CXCR4 axis. *Ann Surg Oncol*. 2009;16(9):2645–2653.
158. Ehteshami M, Min E, Issar NM, Kasl RA, Khan IS, Thompson RC. The role of the CXCR4 cell surface chemokine receptor in glioma biology. *J Neurooncol*. 2013;113(2):153–162.

159. Pan J, Mestas J, Burdick MD, et al. Stromal derived factor-1 (SDF-1/CXCL12) and CXCR4 in renal cell carcinoma metastasis. *Mol Cancer*. 2006;5:56.
160. O'Boyle G, Swidenbank I, Marshall H, Barker CE, Armstrong J, White SA, Fricker SP, Plummer R, Wright M, Lovat PE. Inhibition of CXCR4-CXCL12 chemotaxis in melanoma by AMD11070. *Br J Cancer*. 2013;108(8):1634–1640.
161. Gassenmaier M, Chen D, Buchner A, et al. CXC chemokine receptor 4 is essential for maintenance of renal cell carcinoma-initiating cells and predicts metastasis. *Stem Cells*. Epub April 30, 2013.
162. Dubrovskaja A, Hartung A, Bouchez LC, et al. CXCR4 activation maintains a stem cell population in tamoxifen-resistant breast cancer cells through AhR signalling. *Br J Cancer*. 2012;107(1):43–52.
163. Redjal N, Chan JA, Segal RA, Kung AL. CXCR4 inhibition synergizes with cytotoxic chemotherapy in gliomas. *Clin Cancer Res*. 2006;12(22):6765–6771.
164. Overgaard J. Hypoxic radiosensitization: adored and ignored. *J Clin Oncol*. 2007;25(26):4066–4074.
165. Yaromina A, Krause M, Thames H, et al. Pre-treatment number of clonogenic cells and their radiosensitivity are major determinants of local tumour control after fractionated irradiation. *Radiother Oncol*. 2007;83(3):304–310.
166. Zips D, Böke S, Kroeber T, et al. Prognostic value of radiobiological hypoxia during fractionated irradiation for local tumor control. *Strahlenther Onkol*. 2011;187(5):306–310.
167. Li Z, Rich JN. Hypoxia and hypoxia inducible factors in cancer stem cell maintenance. *Curr Top Microbiol Immunol*. 2010;345:21–30.
168. Hedderston JM, Li Z, McLendon RE, Hjelmeland AB, Rich JN. The hypoxic microenvironment maintains glioblastoma stem cells and promotes reprogramming towards a cancer stem cell phenotype. *Cell Cycle*. 2009;8(20):3274–3284.
169. Soeda A, Park M, Lee D, et al. Hypoxia promotes expansion of the CD133-positive glioma stem cells through activation of HIF-1 $\alpha$ . *Oncogene*. 2009;28(45):3949–3959.
170. Kioi M, Vogel H, Schultz G, Hoffman RM, Harsh GR, Brown JM. Inhibition of vasculogenesis, but not angiogenesis, prevents the recurrence of glioblastoma after irradiation in mice. *J Clin Invest*. 2010;120(3):694–705.
171. Chen FH, Chiang CS, Wang CC, et al. Radiotherapy decreases vascular density and causes hypoxia with macrophage aggregation in TRAMP-C1 prostate tumors. *Clin Cancer Res*. 2009;15(5):1721–1729.
172. Moeller BJ, Cao Y, Li CY, Dewhirst MW. Radiation activates HIF-1 to regulate vascular radiosensitivity in tumors: role of reoxygenation, free radicals, and stress granules. *Cancer Cell*. 2004;5(5): 429–441.
173. Lyden D, Hattori K, Dias S, et al. Impaired recruitment of bone-marrow-derived endothelial and hematopoietic precursor cells blocks tumor angiogenesis and growth. *Nat Med*. 2001;7(11):1194–1201.
174. Ceradini DJ, Kulkarni AR, Callaghan MJ, et al. Progenitor cell trafficking is regulated by hypoxic gradients through HIF-1 induction of SDF-1. *Nat Med*. 2004;10(8):858–864.
175. Kozin SV, Kamoun WS, Huang Y, Dawson MR, Jain RK, Duda DG. Recruitment of myeloid but not endothelial precursor cells facilitates tumor regrowth after local irradiation. *Cancer Res*. 2010;70(14): 5679–5685.
176. Jung MJ, Rho JK, Kim YM, et al. Upregulation of CXCR4 is functionally crucial for maintenance of stemness in drug-resistant non-small cell lung cancer cells. *Oncogene*. 2013;32(2):209–221.
177. Shen W, Bendall LJ, Gottlieb DJ, Bradstock KF. The chemokine receptor CXCR4 enhances integrin-mediated in vitro adhesion and facilitates engraftment of leukemic precursor-B cells in the bone marrow. *Exp Hematol*. 2001;29(12):1439–1447.
178. Hartmann TN, Burger JA, Glodek A, Fujii N, Burger M. CXCR4 chemokine receptor and integrin signaling co-operate in mediating adhesion and chemoresistance in small cell lung cancer (SCLC) cells. *Oncogene*. 2005;24(27):4462–4471.
179. Jones J, Marian D, Weich E, et al. CXCR4 chemokine receptor engagement modifies integrin dependent adhesion of renal carcinoma cells. *Exp Cell Res*. 2007;313(19):4051–4065.
180. Kuonen F, Secondini C, Rüegg C. Molecular pathways: emerging pathways mediating growth, invasion, and metastasis of tumors progressing in an irradiated microenvironment. *Clin Cancer Res*. 2012;18(19):5196–5202.
181. Bettink SI, Werner C, Chen CH, et al. Integrin-linked kinase is a central mediator in angiotensin II type 1- and chemokine receptor CXCR4 signaling in myocardial hypertrophy. *Biochem Biophys Res Commun*. 2010;397(2):208–213.
182. Kijowski J, Baj-Krzyworzeka M, Majka M, et al. The SDF-1-CXCR4 axis stimulates VEGF secretion and activates integrins but does not affect proliferation and survival in lymphohematopoietic cells. *Stem Cells*. 2001;19(5):453–466.
183. Eke I, Koch U, Hehlhans S, et al. PINCH1 regulates Akt1 activation and enhances radioresistance by inhibiting PP1 $\alpha$ . *J Clin Invest*. 2010;120(7):2516–2527.
184. Ahmed KM, Zhang H, Park CC. NF- $\kappa$ B Regulates Radioresistance Mediated By  $\beta$ 1-Integrin in Three-Dimensional Culture of Breast Cancer Cells. *Cancer Res*. 2013;73(12):3737–3748.
185. Ou J, Luan W, Deng J, Sa R, Liang H.  $\alpha$ V integrin induces multicellular radioresistance in human nasopharyngeal carcinoma via activating SAPK/JNK pathway. *PLoS One*. 2012;7(6):e38737.
186. Jordan NJ, Kolios G, Abbot SE, et al. Expression of functional CXCR4 chemokine receptors on human colonic epithelial cells. *J Clin Invest*. 1999;104(8):1061–1069.
187. Zhu B, Xu D, Deng X, et al. CXCL12 enhances human neural progenitor cell survival through a CXCR7- and CXCR4-mediated endocytotic signaling pathway. *Stem Cells*. 2012;30(11):2571–2583.
188. Dong F, Harvey J, Finan A, Weber K, Agarwal U, Penn MS. Myocardial CXCR4 expression is required for mesenchymal stem cell mediated repair following acute myocardial infarction. *Circulation*. 2012;126(3):314–324.
189. De Clercq E. The bicyclam AMD3100 story. *Nat Rev Drug Discov*. 2003;2(7):581–587.
190. Righi E, Kashiwagi S, Yuan J, et al. CXCL12/CXCR4 blockade induces multimodal antitumor effects that prolong survival in an immunocompetent mouse model of ovarian cancer. *Cancer Res*. 2011;71(16):5522–5534.
191. Rubin JB, Kung AL, Klein RS, et al. A small-molecule antagonist of CXCR4 inhibits intracranial growth of primary brain tumors. *Proc Natl Acad Sci U S A*. 2003;100(23):13513–13518.
192. Scholten DJ, Canals M, Maussang D, et al. Pharmacological modulation of chemokine receptor function. *Br J Pharmacol*. 2012;165(6): 1617–1643.
193. Hendrix CW, Flexner C, MacFarland RT, et al. Pharmacokinetics and safety of AMD-3100, a novel antagonist of the CXCR-4 chemokine receptor, in human volunteers. *Antimicrob Agents Chemother*. 2000; 44(6):1667–1673.
194. Domanska UM, Timmer-Bosscha H, Nagengast WB, et al. CXCR4 inhibition with AMD3100 sensitizes prostate cancer to docetaxel chemotherapy. *Neoplasia*. 2012;14(8):709–718.
195. Ramos EA, Grochowski M, Braun-Prado K, et al. Epigenetic changes of CXCR4 and its ligand CXCL12 as prognostic factors for sporadic breast cancer. *PLoS One*. 2011;6(12):e29461.
196. Wu Y, Jin M, Xu H, et al. Clinicopathologic Significance of HIF-1 $\alpha$ , CXCR4, and VEGF Expression in Colon Cancer. *Clin Dev Immunol*. 2010;2010:537531.
197. Popple A, Durrant LG, Spendlove I, et al. The chemokine, CXCL12, is an independent predictor of poor survival in ovarian cancer. *Br J Cancer*. 2012;106(7):1306–1313.
198. Maréchal R, Demetter P, Nagy N, et al. High expression of CXCR4 may predict poor survival in resected pancreatic adenocarcinoma. *Br J Cancer*. 2009;100(9):1444–1451.

199. Wang M, Chen GY, Song HT, Hong x, Yang ZY, Sui GJ. Significance of CXCR4, phosphorylated STAT3 and VEGF-A expression in resected non-small cell lung cancer. *Exp Ther Med*. 2011;2(3):517–522.
200. Zhang NH, Li J, Li Y, et al. Co-expression of CXCR4 and CD133 proteins is associated with poor prognosis in stage II–III colon cancer patients. *Exp Ther Med*. 2012;3(6):973–982.
201. Jung SJ, Kim C, Park CH, et al. Correlation between Chemokine Receptor CXCR4 Expression and Prognostic Factors in Patients with Prostate Cancer. *Korean J Urol*. 2011;52(9):607–611.
202. Zhang L, Ye SB, Ma G, et al. The expressions of MIF and CXCR4 protein in tumor microenvironment are adverse prognostic factors in patients with esophageal squamous cell carcinoma. *J Transl Med*. 2013;11:60.
203. Bennani-Baiti IM, Cooper A, Lawlor ER, et al. Intercohort Gene Expression Co-analysis Reveals Chemokine Receptors as Prognostic Indicators in Ewing's Sarcoma. *Clin Cancer Res*. 2010;16(14):3769–3778.
204. Zhang SS, Han ZP, Jing YY, et al. CD133(+)/CXCR4(+) colon cancer cells exhibit metastatic potential and predict poor prognosis of patients. *BMC Med*. 2012;10:85.
205. Hermann PC, Huber SL, Herrler T, et al. Distinct populations of cancer stem cells determine tumor growth and metastatic activity in human pancreatic cancer. *Cell Stem Cell*. 2007;1(3):313–323.
206. Zheng X, Xie Q, Li S, Zhang W. CXCR4-positive subset of glioma is enriched for cancer stem cells. *Oncol Res*. 2011;19(12):555–561.

### OncoTargets and Therapy

## Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress