

The association between the migration inhibitory factor –173G/C polymorphism and cancer risk: a meta-analysis

Xiao Zhang¹
 Wenhao Weng¹
 Wen Xu²
 Yulan Wang¹
 Wenjun Yu¹
 Xun Tang¹
 Lifang Ma¹
 Qihui Pan³
 Jiayi Wang¹
 Fenyong Sun¹

¹Department of Clinical laboratory medicine, Shanghai Tenth People's Hospital of Tongji University,

²Department of Clinical laboratory medicine, Zhongshan Hospital, Fudan University, ³Department of Central Laboratory, Shanghai Tenth People's Hospital of Tongji University, Shanghai, People's Republic of China

Correspondence: Jiayi Wang; Fenyong Sun
 Number 301 Middle Yanchang Road, Shanghai Tenth People's Hospital of Tongji University, Shanghai 200072, People's Republic of China
 Tel +86 21 6630 0588
 Fax +86 21 6630 3643; +86 21 6630 0588
 Email karajan2@163.com; sunfenyongtongji@126.com

Abstract: Previous studies have suggested that macrophage migration inhibitory factor (MIF) –173G/C polymorphism may be associated with cancer risk. However, previous research has demonstrated conflicting results. Therefore, we followed the preferred reporting items for systematic reviews and meta-analyses (PRISMA) guidelines and the meta-analysis on genetic association studies checklist, and performed a meta-analysis to investigate the association between MIF –173G/C polymorphisms and the risk of cancer. Odds ratios (ORs) and corresponding 95% confidence intervals (CIs) were combined to measure the association between MIF promoter polymorphisms and cancer risk. The pooled ORs were performed for the dominant model, recessive model, allelic model, homozygote comparison, and heterozygote comparison. The publication bias was examined by Begg's funnel plots and Egger's test. A total of ten studies enrolling 2,203 cases and 2,805 controls met the inclusion criteria. MIF (–173G/C) polymorphism was significantly associated with increased cancer risk under the dominant model (OR=1.32, 95%, CI=1.00–1.74, $P=0.01$) and the heterozygote comparison (OR=1.38, CI=1.01–1.87, $P=0.04$). In subgroup analysis, MIF polymorphism and prostate were related to increased risk of prostate and non-solid cancer. In conclusion, MIF polymorphism was significantly associated with cancer risk in heterozygote comparison. The MIF –173G/C polymorphism may be associated with increased cancer risk.

Keywords: MIF, SNP, systematic review, cancer susceptibility

Introduction

Macrophage migration inhibitory factor (MIF) was first identified nearly 50 years ago and has been used as a cytokine and an enzyme.^{1,2} MIF is a member of the transferring growth factor- β (TGF- β) super family, which is expressed by a broad variety of cells, including B- and T-lymphocytes as well as endocrine, endothelial, and epithelial cells of diverse histogenetic origin.³ Presently, MIF is considered to play an important role in the pro- and anti-inflammatory response to infection since it is constitutively expressed and acts as an upstream regulator of many other inflammatory cytokines.^{4,5}

Recently, several studies have shown that MIF can promote tumor growth and viability by modulating immune responses and supporting tumor-associated angiogenesis.⁶ A few experiments suggested that MIF mRNA and MIF protein are overexpressed in a number of cancers.⁷ Tan et al reported that MIF is upregulated in patients with pancreatic cancer and causes dysfunction of insulin secretion in β -cells.⁸ Krockenberger et al reported that MIF is clearly overexpressed on the protein level in invasive cervical cancer compared to cervical dysplasia.⁹ Two polymorphisms in the promoter region of MIF have been reported in the past. One is a single nucleotide polymorphism (SNP)

at the nucleotide position –173 (G to C)¹⁰ and the other is a tetranucleotide CATT repeat beginning at position –794.¹¹ The association between these two polymorphisms and diseases has been extended to several inflammatory conditions including Graves' disease,¹² idiopathic thrombocytopenic purpura,¹³ and Vogt-Koyanagi-Harada (VKH) syndrome.¹⁴ These studies indicate that these two polymorphisms of MIF are associated with inflammatory diseases. Similarly, some studies have reported that the polymorphism of MIF resulted in an increased risk of cancer. With new studies about the polymorphism of MIF and the risk of cancer emerging, there has been no meta-analysis conducted regarding the association between MIF promoter polymorphism and the risk of cancer in recent times. The aim of this study is to perform a meta-analysis of all available studies that analyze the association between the polymorphism of MIF promoter and the risk of cancer.

Materials and methods

Literature search

The preferred reporting items for systematic reviews and meta-analyses (PRISMA) statement (Figure S1) and the meta-analysis on genetic association studies checklist (Figure S2) were followed in our meta-analysis. A comprehensive search of EMBASE, PubMed, Web of Science, OVID, Cochrane Library, and China National Knowledge Infrastructure (CNKI) was done from database inception to July 22, 2014 without language restriction. The search strategy was “macrophage migration inhibitory factor or MIF” and “polymorphism or variant or mutation or genotype.” To complete our research, we also studied the review articles and references of retrieved articles manually. The literature review was performed independently by X Zhang and J Wang and the disagreements were resolved through consensus by all the authors.^{15,16}

Selection criteria

Studies were included in the meta-analysis if the following inclusion criteria were satisfied: 1) case-control studies focused on association between the MIF promoter polymorphism and cancer risk, 2) studies enrolled more than 30 patients, 3) studies provided sufficient data to estimate the odds ratio (OR) and 95% confidence intervals (CIs) according to MIF promoter polymorphism, and 4) when study patients overlapped with patients in other included studies, we selected the first study published. The two researchers (J Wang and X Zhang) independently read the titles and abstracts and excluded the uncorrelated studies; then the

full-texts were examined by our review team. The studies were selected according to the inclusion criteria.^{15,16}

Data abstraction

Two independent reviewers (X Zhang and J Wang) extracted the following information: authors, year of publication, country, tumor type, number of cases and controls analyzed, mean value of age, source of controls (hospital-based controls or population-based controls), and genotyping method. If both univariate and multivariate analyses were reported, we utilized the multivariate analysis because it involves observation and analysis of more than one statistical outcome variable at a time thus is more accurate. If articles provided insufficient data (missing data, inconsistencies, or any other uncertainties), we attempted to contact the first and corresponding authors for necessary information via telephone or email.^{15,16}

Statistical analysis

ORs and corresponding 95% CIs were combined to measure the association between MIF promoter polymorphisms and cancer risk. Hardy–Weinberg equilibrium (HWE) for each study was determined by the chi-square test. The pooled ORs were calculated for the allelic model (mutation [M] allele versus [vs] wild [W] allele), dominant model (WM + MM vs WW), recessive model (MM vs WM + WW), homozygote comparison (MM vs WW), and heterozygote comparison (WM vs WW) respectively, and $P < 0.05$ denoted statistical significance. Statistical heterogeneity among the studies was evaluated using the Q -test and I^2 -test. When heterogeneity among the studies was observed, the pooled OR was calculated by random-effect models. Sensitivity analyses were performed to identify the potential influence of the individual data set to the pooled ORs. Subgroup analyses were conducted with respect to cancer type and source of controls. The statistical significance was analyzed by Student's t -test. These analyses were performed by Review Manager Version 5.1 software (<http://ims.cochrane.org/revman>). Both Begg's and Egger's tests were performed using R (<http://cran.r-project.org/bin/windows/base>).^{15,16}

Results

Characteristics of identified studies

Following an initial search, 166 studies were retrieved from PubMed; 233 studies from EMBASE; 313 studies from OVID; 266 studies from Web of Science; 50 studies from Cochrane Library; 532 studies from CNKI; and five additional review articles were added to make our search comprehensive. After duplicated records were removed,

878 published studies were identified. We excluded 780 unrelated studies by reading the titles and abstracts. Next, we downloaded the full-text of the remaining 98 studies and excluded 65 unrelated studies. Of the remaining 33 studies considered for performing the meta-analysis, some studies were found to report incomplete data or report other associations between MIF and cancer. We tried our best to communicate with the first and corresponding authors to get the necessary data. Some authors were able to provide the necessary data for our study, while others did not. Ultimately, after further reviewing in detail, ten studies were included in our meta-analysis.^{17–26} Figure 1 shows in detail the selection process. These ten studies were published between 2005 and 2014. There were 2,203 cases and 2,805 controls included in our meta-analysis. Studies were carried out in People's Republic of China, Taiwan, Japan, Iran, Italy, and USA. Polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) was used in seven studies.^{17,18,20,21,23,25,26} One study used polymerase chain reaction-single strand conformation polymorphism (PCR-SSCP).²⁴ The other two studies employed denaturing high-performance liquid chromatography (DHLPC) wave analysis¹⁹ and a Genetic

Analyzer,²² respectively. Three studies assessed prostate cancer;^{20,22,26} three studies assessed leukemia^{17,19,25} and one each for gastric cancer,²⁴ cervical cancer,¹⁸ colorectal cancer,²¹ and bladder cancer.²³ The genotype distribution in one study deviated from HWE.²⁶ The main characteristics of all the included studies are listed in Table 1.

Meta-analysis

Overall, ten prospective studies enrolling 2,203 cases and 2,805 controls were included in our meta-analysis. A statistically significant association between MIF (–173G/C) polymorphism and cancer risk was found under the dominant model (OR=1.32, CI=1.00–1.74, $P=0.01$) (Figure 2) and the heterozygote comparison (OR=1.38, CI=1.01–1.87, $P=0.04$) (Figure S3). There was no statistical significant association under the recessive model (OR=0.98, 95% CI 0.67–1.45, $P=0.93$) (Figure S4), homozygote comparison (OR=1.02, 95% CI 0.64–1.63, $P=0.93$) (Figure S5), and allelic model (OR=1.32, 95% CI 1.00–1.74, $P=0.05$) (Figure S6). Furthermore, in our subgroup analysis, a significant association was found in the prostate group under the dominant model (OR=3.34, 95% CI 2.24–4.97, $P<0.001$), allelic model




Figure 1 Flow diagram summarizing the selection of eligible studies.

Table 1 Baseline characteristics of studies included in the meta-analysis

Study	Year	Country	Tumor Type	Cases	Controls	Age	Source of controls	Genotyping method	HWE
Ramireddy et al ¹⁷	2014	Taiwan	Acute myeloid leukemia	256	256	Mean age: cases: 53.4 controls: 55.8	HB	PCR-RFLP	0.06
Wu et al ¹⁸	2011	People's Republic of China	Cervical cancer	250	147	Mean age: cases: 49.08±9.405 controls: 47.99±10.750	PB	PCR-RFLP	0.28
Ziino et al ¹⁹	2005	Italy	Acute lymphoblastic leukemia	151	355	NR	PB	PCR and DHLPC	0.05
Razzaghi et al ²⁰	2012	Iran	Prostate cancer	61	71	NR	PB	Wave analysis	0.88
Ramireddy et al ²¹	2014	Taiwan	Colorectal cancer	192	256	Mean age: cases: 62.1 controls: 55.8	PB	PCR-RFLP	0.13
Meyer-Siegler et al ²²	2007	USA	Prostate cancer	131	128	Mean age: cases: 70.16±0.89 controls: 64.39±1.09	PB	PCR and ABI 310	–
Yuan et al ²³	2012	People's Republic of China	Bladder cancer	325	345	Cases: ≤55 years: 66 persons, >55 years: 259 persons; controls: ≤55 years: 83 persons, >55 years: 262 persons	PB	Genetic analyzer	0.94
Arisawa et al ²⁴	2007	Japan	Gastric cancer	232	430	Mean age: cases: 62.99±10.73 controls: 54.72±18.84	HB	PCR-SSCP	0.81
Xue et al ²⁵	2010	People's Republic of China	Acute lymphoblastic leukemia	346	516	Cases: <6 years: 156 persons, ≥6 years: 190 persons; controls: <6 years: 251 persons, ≥6 years: 265 persons	PB	PCR-RFLP	0.8
Ding et al ²⁶	2009	People's Republic of China	Prostate cancer	259	301	Cases: ≤70 years: 123 persons, >70 years: 136 persons; controls: ≤70 years: 153 persons, >70 years: 148 persons	HB	PCR-RFLP	0.01

Abbreviations: HB, hospital-based; PB, population-based; HWE, Hardy-Weinberg equilibrium; PCR-RFLP, polymerase chain reaction-restriction fragment length polymorphism; DHLPC, denaturing high-performance liquid chromatography; PCR-SSCP, polymerase chain reaction-single strand conformation polymorphism; NR, no report.




Figure 2 Forest plot of MIF –173G/C polymorphism and cancer risk in dominant model.
Abbreviation: CI, confidence interval.

(OR=2.94, 95% CI 1.91–4.54, $P<0.001$), and heterozygote comparison (OR=2.39, 95% CI 1.65–3.47, $P<0.001$). MIF (–173G/C) polymorphism was also significantly associated with non-solid cancer risk under the dominant model (OR=1.27, 95% CI 1.03–1.56, $P=0.03$) and heterozygote comparison (OR=1.32, 95% CI 1.06–1.63, $P=0.01$). Table S1 presents the results of overall and subgroup analyses.

Sensitivity analysis

We performed sensitivity analysis by omitting one study at a time and calculating the pooled ORs again. However, the results did not show any significant statistical differences when studies were omitted. Therefore, the stability of the study was not influenced by any individual study. Table S2 presents the sensitivity analysis in the dominant model.

Publication bias

Both Begg's funnel plot and Egger's test were carried out to evaluate the publication bias of the studies. The results are presented in Figure 3 and Table 2. Publication bias was found under the dominant model ($P=0.0286$) according to Begg's funnel plot. When Egger's test was performed, publication bias was found under the recessive model ($P=0.0075$) and homozygote comparison ($P=0.03$). Results indicate that there may be publication bias existing in our meta-analysis. Table 2 presents the results of Begg's funnel plot and Egger's test under the five genetic models.

Discussion

In our meta-analysis, ten studies enrolling 2,203 cases and 2,805 controls were included. The results indicated that MIF –173G/C polymorphism was significantly associated with cancer risk.

MIF is known as a major regulator of inflammation and a central upstream mediator of innate immune response, and functions as a key mediator to counter-regulate the inhibitory effects of glucocorticoids within the immune system.²⁷ There are numerous studies suggesting that MIF polymorphism might be associated with the risk of immune disease. Liu et al reported that MIF polymorphism is associated with new-onset Graves' disease in a Taiwanese Chinese population.¹² Hao et al carried out a meta-analysis to investigate the association between MIF polymorphism and the risk of inflammatory bowel disease (IBD).²⁸ They found that MIF –173G/C polymorphism contributed to the susceptibility of IBD.

MIF is also involved in cancer growth and progression. The elevated MIF and mRNA levels have been observed in many tumor cells and pre-tumor states. Krockenberger et al found that MIF was significantly overexpressed on both the protein level and the mRNA level in invasive cervical cancer and MIF protein was overexpressed in SiHA and CaSki cervical cancer cell lines.⁹ Huang et al reported that MIF expression levels in hepatocellular carcinoma tissues and cell lines were significantly up-regulated compared with adjacent normal tissues or a normal liver cell line.²⁹ Moreover, several studies suggested that MIF polymorphism might be associated with the risk of cancer. Only one study reported that MIF –173G/C polymorphism is associated with a decreased risk of cancer.²³ All the other studies reported the opposite conclusion. We also found a meta-analysis that investigated the association between the MIF –173G/C polymorphism and cancer risk.³⁰ However, there were only five studies included in that meta-analysis, and the result was only under the dominant model. In recent times, some new studies have been emerging; for instance, Yuan et al reported that MIF –173G/C polymorphism is associated with




Figure 3 Publication bias in this meta-analysis.

Notes: (A) Begg's funnel plots of MIF -173G/C polymorphism in dominant model. (B) Egger's test of MIF -173G/C polymorphism in dominant model.

Abbreviation: MIF, migration inhibitory factor.

decreased cancer risk.²³ This conclusion contradicted with the conclusion in the previous meta-analysis. Therefore, we added new studies in our meta-analysis and calculated ORs in the dominant model, recessive model, homozygote comparison, heterozygote comparison, and allelic model. In our meta-analysis, we found that MIF -173G/C polymorphism is significantly associated with cancer risk in the dominant model (OR=1.32, 95% CI 1.00–1.74, $P=0.01$) and heterozygote comparison (OR=1.38, 95% CI 1.01–1.87, $P=0.04$). There were no significant associations between

MIF -173G/C polymorphism and cancer risk in the recessive model (OR=0.98, 95% CI 0.67–1.45, $P=0.93$), homozygote comparison (OR=1.02, 95% CI 0.64–1.63, $P=0.93$), and allelic model (OR=1.32, 95% CI 1.00–1.74, $P=0.05$). Drawing from these results, we conclude from our meta-analysis that MIF -173G/C polymorphism might increase the risk of cancer.

There are several limitations in our meta-analysis. First, publication bias exists in the current meta-analysis. If the future studies find that MIF polymorphism was not associated with cancer risk, then publication bias might cause false outcomes. Second, there were some studies lacking in necessary data to calculate ORs under different genetic models. Although we had tried our best to communicate with the first and corresponding authors, some were unable to reply. Third, the patients included in the meta-analysis were limited. It was difficult for us to perform subgroup analyses and obtain specific results. Additionally, only papers published in English or Chinese were included in our meta-analysis, and

Table 2 A summary of P -values for Begg's funnel plot and Egger's test in five genetic models

	Begg's funnel plot	Egger's test
Dominant model	0.0286	0.1128
Recessive model	0.1361	0.0075
Homozygote comparison	0.1361	0.03
Heterozygote comparison	0.4767	0.2992
Allelic model	0.7614	0.2373

eligible studies written in other languages that could have fulfilled our study criterion were not included.

Conclusion

Our meta-analysis concluded that MIF -173G/C polymorphism might increase the risk of cancer. Given the above limitations, more studies are needed to confirm the association between MIF polymorphism and the risk of cancer.

Acknowledgment

This study was supported by Natural Science Foundation of People's Republic of China (Grant Nos: 81201363 and 81301689), and the Climbing training program (assigned to Jiayi Wang) from Shanghai Tenth People's Hospital.

Disclosure

The authors report no conflicts of interest in this work.

References

- Bloom BR, Bennett B. Mechanism of a reaction in vitro associated with delayed-type hypersensitivity. *Science*. 1966;153(3731):80–82.
- Lolis E, Bucala R. Macrophage migration inhibitory factor. *Expert Opin Ther Targets*. 2003;7(2):153–164.
- He M, Metz C, Cheng KF, et al. Novel arylazoarylmethane as potential inhibitor of macrophage migration inhibitory factor. *Arch Pharm*. 2014;347(2):104–107.
- Grieb G, Merk M, Bernhagen J, Bucala R. Macrophage migration inhibitory factor (MIF): a promising biomarker. *Drug News Perspect*. 2010;23(4):257–264.
- Vujicic M, Senerovic L, Nikolic I, Saksida T, Stosic-Grujicic S, Stojanovic I. The critical role of macrophage migration inhibitory factor in insulin activity. *Cytokine*. 2014;69(1):39–46.
- Yasasever V, Camlica H, Duranyildiz D, Oguz H, Tas F, Dalay N. Macrophage migration inhibitory factor in cancer. *Cancer Invest*. 2007;25(8):715–719.
- Armstrong ME, Gantier M, Li L, et al. Small interfering RNAs induce macrophage migration inhibitory factor production and proliferation in breast cancer cells via a double-stranded RNA-dependent protein kinase-dependent mechanism. *J Immunol*. 2008;180(11):7125–7133.
- Tan L, Ye X, Zhou Y, et al. Macrophage migration inhibitory factor is overexpressed in pancreatic cancer tissues and impairs insulin secretion function of beta-cell. *J Transl Med*. 2014;12:92.
- Krockenberger M, Engel JB, Kolb J, et al. Macrophage migration inhibitory factor expression in cervical cancer. *J Cancer Res Clin Oncol*. 2010;136(5):651–657.
- Amoli MM, Donn RP, Thomson W, et al. Macrophage migration inhibitory factor gene polymorphism is associated with sarcoidosis in biopsy proven erythema nodosum. *J Rheumatol*. 2002;29(8):1671–1673.
- Baugh JA, Chitnis S, Donnelly SC, et al. A functional promoter polymorphism in the macrophage migration inhibitory factor (MIF) gene associated with disease severity in rheumatoid arthritis. *Genes Immun*. 2002;3(3):170–176.
- Liu YH, Chen CC, Yang CM, Chen YJ, Tsai FJ. Dual effect of a polymorphism in the macrophage migration inhibitory factor gene is associated with new-onset Graves disease in a Taiwanese Chinese population. *PLoS One*. 2014;9(3):e92849.
- Lao W, Xiang Y, Fang M, Yang X. Polymorphism and expression of macrophage migration inhibitory factor does not contribute to glucocorticoid resistance in idiopathic thrombocytopenic purpura. *Pharmazie*. 2013;68(10):846–849.
- Zhang C, Liu S, Hou S, et al. MIF gene polymorphisms confer susceptibility to Vogt-Koyanagi-Harada syndrome in a Han Chinese population. *Invest Ophthalmol Vis Sci*. 2013;54(12):7734–7738.
- Zhang X, Weng W, Xu W, et al. Role of Bcl-2-938 C>A polymorphism in susceptibility and prognosis of cancer: a meta-analysis. *Sci Rep*. 2014;4:7241.
- Zhang X, Weng W, Xu W, et al. Prognostic significance of interleukin 17 in cancer: a meta-analysis. *Int J Clin Exp Med*. 2014;7(10):3258–3269.
- Ramireddy L, Lin CY, Liu SC, et al. Association study between macrophage migration inhibitory factor -173 polymorphism and acute myeloid leukemia in Taiwan. *Cell Biochem Biophys*. 2014;70(2):1159–1165.
- Wu S, Lian J, Tao H, Shang H, Zhang L. Correlation of macrophage migration inhibitory factor gene polymorphism with the risk of early-stage cervical cancer and lymphatic metastasis. *Oncol Lett*. 2011;2(6):1261–1267.
- Ziino O, D'Urbano LE, De Benedetti F, et al. The MIF -173G/C polymorphism does not contribute to prednisone poor response in vivo in childhood acute lymphoblastic leukemia. *Leukemia*. 2005;19(12):2346–2347.
- Razzaghi M, Mazloomfard M, Lotfi B. Macrophage migration inhibitory factor gene -173*C polymorphism and risk of prostate cancer. *Eur Urol Suppl*. 2012;19(12):142.
- Ramireddy L, Chen WT, Peng CT, et al. Association between genetic polymorphism of the MIF gene and colorectal cancer in Taiwan. *J Clin Lab Anal*. 2014.
- Meyer-Siegler KL, Vera PL, Iczkowski KA, et al. Macrophage migration inhibitory factor (MIF) gene polymorphisms are associated with increased prostate cancer incidence. *Genes Immun*. 2007;8(8):646–652.
- Yuan Q, Wang M, Wang M, Zhang Z, Zhang W. Macrophage migration inhibitory factor gene -173G>C polymorphism and risk of bladder cancer in southeast China: a case-control analysis. *Mol Biol Rep*. 2012;39(3):3109–3115.
- Arisawa T, Tahara T, Shibata T, et al. Functional promoter polymorphisms of the macrophage migration inhibitory factor gene in gastric carcinogenesis. *Oncol Rep*. 2008;19(1):223–228.
- Xue Y, Xu H, Rong L, et al. The MIF -173G/C polymorphism and risk of childhood acute lymphoblastic leukemia in a Chinese population. *Leuk Res*. 2010;34(10):1282–1286.
- Ding GX, Zhou SQ, Xu Z, et al. The association between MIF -173 G>C polymorphism and prostate cancer in southern Chinese. *J Surg Oncol*. 2009;100(2):106–110.
- Xu L, Li Y, Sun H, et al. Current developments of macrophage migration inhibitory factor (MIF) inhibitors. *Drug Discov Today*. 2013;18(11–12):592–600.
- Hao NB, He YF, Luo G, Yong X, Zhang Y, Yang SM. Macrophage migration inhibitory factor polymorphism and the risk of ulcerative colitis and Crohn's disease in Asian and European populations: a meta-analysis. *BMJ Open*. 2013;3(12):e003729.
- Huang XH, Jian WH, Wu ZF, et al. Small interfering RNA (siRNA)-mediated knockdown of macrophage migration inhibitory factor (MIF) suppressed cyclin D1 expression and hepatocellular carcinoma cell proliferation. *Oncotarget*. 2014;5(14):5570–5580.
- Vera PL, Meyer-Siegler KL. Association between macrophage migration inhibitory factor promoter region polymorphism (-173 G/C) and cancer: a meta-analysis. *BMC Res Notes*. 2011;4:395.

Supplementary materials

Table S1 A summary of ORs for the overall and subgroup analyses of MIF polymorphism and cancer risk

Subgroups	Dominant model (ORs)	95% CI	P-value	Recessive model (ORs)	95% CI	P-value	Allelic model (ORs)	95% CI	P-value
Overall	1.57	1.1–2.24	0.01	0.98	0.67–1.45	0.93	1.32	1.00–1.74	0.05
Prostate cancer	3.34	2.24–4.97	<0.001	–	–	–	2.94	1.91–4.54	<0.001
Other cancer	1.2	0.9–1.59	0.21	0.98	0.67–1.45	0.93	1.12	0.92–1.36	0.27
Solid cancer	1.78	1.04–3.04	0.04	1.04	0.64–1.69	0.88	1.44	0.94–2.22	0.1
Non-solid cancer	1.27	1.03–1.56	0.03	0.81	0.40–1.66	0.57	1.17	0.98–1.40	0.07
Asian	1.41	0.97–2.06	0.07	0.98	0.67–1.45	0.93	1.32	0.96–1.81	0.1
Caucasian	2.13	0.78–5.81	0.14	–	–	–	1.34	0.67–2.71	0.41
HB	1.8	1.06–3.04	0.03	0.8	0.45–1.44	0.46	1.67	0.90–3.12	0.1
PB	1.49	0.93–2.37	0.1	1.06	0.64–1.75	0.82	1.15	0.87–1.52	0.32
Subgroups	Homozygote comparison (ORs)	95% CI	P-value	Heterozygote comparison (ORs)	95% CI	P-value			
Overall	1.02	0.64–1.63	0.93	1.38	1.01–1.87	0.04			
Prostate cancer	–	–	–	2.39	1.65–3.47	<0.001			
Other cancer	1.02	0.64–1.63	0.93	1.23	0.90–1.68	0.19			
Solid cancer	1.05	0.56–2.00	0.87	1.44	0.88–2.35	0.15			
Non-solid cancer	0.9	0.47–1.75	0.76	1.32	1.06–1.63	0.01			
Asian	1.02	0.64–1.63	0.93	1.4	0.97–2.01	0.07			
Caucasian	–	–	–	1.23	0.77–1.98	0.23			
HB	0.88	0.50–1.56	0.67	1.75	1.22–2.51	0.002			
PB	1.08	0.56–2.10	0.82	1.2	0.81–1.79	0.35			

Abbreviations: ORs, odds ratios; MIF, migration inhibitory factor; CI, confidence interval; HB, hospital-based; PB, population-based.

Table S2 The influence of individual study on ORs in dominant model

Study omitted	Year	OR	95% CI	P-value	Heterogeneity	
					I ²	P-value
None		1.57	1.10–2.24	0.01	87	P<0.001
Ramireddy et al ²	2014	1.60	1.07–2.39	0.02	88	P<0.001
Leukemia						
Wu et al ³	2011	1.55	1.05–2.27	0.03	88	P<0.001
Ziino et al ⁴	2005	1.65	1.12–2.43	0.01	88	P<0.001
Razzaghi et al ⁵	2012	1.54	1.06–2.24	0.02	88	P<0.001
Ramireddy et al ⁶	2014	1.60	1.07–2.37	0.02	88	P<0.001
CRC						
Meyer-Siegler et al ⁷	2007	1.40	1.01–1.93	0.04	83	P<0.001
Yuan et al ⁸	2012	1.75	1.31–2.35	0.0002	77	P<0.001
Arisawa et al ⁹	2007	1.61	1.07–2.42	0.02	88	P<0.001
Xue et al ¹⁰	2010	1.62	1.07–2.44	0.02	88	P<0.001
Ding et al ¹¹	2009	1.44	1.03–2.03	0.04	84	P<0.001

Abbreviations: OR, odds ratio; CI, confidence interval.



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
Title			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	Title
Abstract			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	Abstract
Introduction			
Rationale	3	Describe the rationale for the review in the context of what is already known.	Introduction
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	Introduction
Methods			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (eg, Web address), and, if available, provide registration information including registration number.	NA
Eligibility criteria	6	Specify study characteristics (eg, PICOS, length of follow-up) and report characteristics (eg, years considered, language, publication status) used as criteria for eligibility, giving rationale.	Literature search and selection criteria
Information sources	7	Describe all information sources (eg, databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	Literature search and selection criteria
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Literature search and selection criteria
Study selection	9	State the process for selecting studies (ie, screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	Data abstraction
Data collection process	10	Describe method of data extraction from reports (eg, piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	Data abstraction
Data items	11	List and define all variables for which data were sought (eg, PICOS, funding sources) and any assumptions and simplifications made.	Statistical analysis
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	Data abstraction
Summary measures	13	State the principal summary measures (eg, risk ratio, difference in means).	Statistical analysis
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (eg, <i>I</i> ²) for each meta-analysis.	Statistical analysis
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (eg, publication bias, selective reporting within studies).	Statistical analysis

(Continued)

Section/topic	#	Checklist item	Reported on page #
Additional analyses	16	Describe methods of additional analyses (eg, sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	Statistical analysis
Results			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	Figure 1
Study characteristics	18	For each study, present characteristics for which data were extracted (eg, study size, PICOS, follow-up period) and provide the citations.	Table S1
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	Table 2 and Figure 3
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	Figure 2 Figures S1–S4
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	Table S2
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	Figure 3 and Table 2
Additional analysis	23	Give results of additional analyses, if done (eg, sensitivity or subgroup analyses, meta-regression [see Item 16]).	Tables 1 and S2
Discussion			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (eg, healthcare providers, users, and policy makers).	Discussion
Limitations	25	Discuss limitations at study and outcome level (eg, risk of bias), and at review-level (eg, incomplete retrieval of identified research, reporting bias).	Discussion
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	Discussion
Funding			
Funding	27	Describe sources of funding for the systematic review and other support (eg, supply of data); role of funders for the systematic review.	Acknowledgment

Figure S1 Preferred reporting items for systematic reviews and meta-analyses (PRISMA) checklist

Notes: Data from Moher D, Liberati A, Tetzlaff J, Altman DG. The PRIS MA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRIS MA Statement. *PLoS Med.* 6(6):e1000097. For more information, visit: www.prisma-statement.org.

#	Item	Section name and paragraph number within manuscript
Introduction		
1	Provide a detailed justification for the polymorphism studied; if a single polymorphism was analyzed, give details as to why others were not included in the meta-analysis.	Para 2 of Introduction
2	Provide a detailed justification for the population(s) and clinical condition studied.	Para 2 of Introduction
Methods		
3	Provide full details of the search strategy employed; outline the full electronic search strategy – specific combination of keywords and any limits applied- for at least one database. Specify whether synonyms of polymorphisms/genes (eg, SNP number) were searched.	Para 1 of Materials and methods
4	Report full details on the inclusion and exclusion criteria applied for selecting studies. <i>Please list the excluded articles and the reasons for exclusion of each article in a supplementary file.</i>	Para 1 of Materials and methods, Para 1 of Results
5	Provide details on how the quality of the studies included in the analyses was assessed.	Para 2 of Materials and methods
6	Describe steps taken to contact study authors to identify additional studies and to request missing data.	Para 3 of Materials and methods
7	Describe how environmental effects were adjusted for, if this adjustment was not conducted, outline the reasons for this.	Para 4 of Materials and methods
8	Describe the methods of handling heterogeneity/between-study variance.	Para 4 of Materials and methods
9	Describe how the Hardy–Weinberg equilibrium and linkage disequilibrium were assessed.	Para 4 of Materials and methods
10	Describe and justify the choice of model for the analyses (per-allele vs per-genotype vs genetic model-free, random effects vs fixed effects).	Para 4 of Materials and methods
11	Describe whether a sensitivity analysis has been completed.	Para 4 of Materials and methods
12	Describe whether an assessment of the effects of population stratification has been conducted.	Para 3 of Materials and methods
13	Describe whether study-specific results have been assessed and if so the reasons for this (eg, forest plot).	Para 4 of Materials and methods
Results		
14	Include flow diagram for the studies included in the meta-analysis as the first figure for the manuscript	Para 1 of Results
15	Report details on allele/genotype prevalence.	Para 2 of Results
16	Report the effect size estimates and <i>P</i> -values for each analysis.	Para 2 of Results
Discussion		
17	Discuss the limitations of the meta-analysis, including genotyping errors/bias and publication bias.	Para 4 of Discussion
18	If the meta-analysis identifies an association within a subgroup of the population studied but not another, discuss the implications of these results, and if applicable the possibility of subgroup-specific publication bias.	Para 3 of Discussion
19	Discuss the suitability of the sample size employed to the research question and the power of the study.	Para 3 and Para 4 of Discussion

Figure S2 Meta-analysis on genetic association studies checklist

Abbreviations: Para, paragraph; SNP, single nucleotide polymorphisms.




Figure S3 Forest plot of MIF -173G/C polymorphism and cancer risk in heterozygote comparison.

Abbreviations: MIF, migration inhibitory factor; CI, confidence interval.




Figure S4 Forest plot of MIF -173G/C polymorphism and cancer risk in recessive model.

Abbreviations: MIF, migration inhibitory factor; CI, confidence interval.




Figure S5 Forest plot of MIF -173G/C polymorphism and cancer risk in homozygote comparison.

Abbreviations: MIF, migration inhibitory factor; CI, confidence interval.




Figure S6 Forest plot of MIF -173G/C polymorphism and cancer risk in allelic model.
Abbreviations: MIF, migration inhibitory factor; CI, confidence interval.

References

- Moher D, Liberati A, Tetzlaff J, Altman DG. The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med.* 6(6):e1000097.
- Ramireddy L, Lin CY, Liu SC, et al. Association study between macrophage migration inhibitory factor-173 polymorphism and acute myeloid leukemia in Taiwan. *Cell Biochem Biophys.* 2014;70(2):1159–1165.
- Wu S, Lian J, Tao H, Shang H, Zhang L. Correlation of macrophage migration inhibitory factor gene polymorphism with the risk of early-stage cervical cancer and lymphatic metastasis. *Oncol Lett.* 2011;2(6):1261–1267.
- Ziino O, D'Urbano LE, De Benedetti F, et al. The MIF-173G/C polymorphism does not contribute to prednisone poor response in vivo in childhood acute lymphoblastic leukemia. *Leukemia.* 2005;19(12):2346–2347.
- Razzaghi M, Mazloomfard M, Lotfi B. Macrophage migration inhibitory factor gene-173*C polymorphism and risk of prostate cancer. *Eur Urol Suppl.* 2012;19(12):142.
- Ramireddy L, Chen WT, Peng CT, et al. Association between genetic polymorphism of the MIF gene and colorectal cancer in Taiwan. *J Clin Lab Anal.* 2014.
- Meyer-Siegler KL, Vera PL, Iczkowski KA, et al. Macrophage migration inhibitory factor (MIF) gene polymorphisms are associated with increased prostate cancer incidence. *Genes Immun.* 2007;8(8):646–652.
- Yuan Q, Wang M, Wang M, Zhang Z, Zhang W. Macrophage migration inhibitory factor gene -173G>C polymorphism and risk of bladder cancer in southeast China: a case-control analysis. *Mol Biol Rep.* 2012;39(3):3109–3115.
- Arisawa T, Tahara T, Shibata T, et al. Functional promoter polymorphisms of the macrophage migration inhibitory factor gene in gastric carcinogenesis. *Oncol Rep.* 2008;19(1):223–228.
- Xue Y, Xu H, Rong L, et al. The MIF-173G/C polymorphism and risk of childhood acute lymphoblastic leukemia in a Chinese population. *Leuk Res.* 2010;34(10):1282–1286.
- Ding GX, Zhou SQ, Xu Z, et al. The association between MIF-173 G>C polymorphism and prostate cancer in southern Chinese. *J Surg Oncol.* 2009;100(2):106–110.

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic agents and protocols on

Submit your manuscript here: <http://www.dovepress.com/oncotargets-and-therapy-journal>

patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Dovepress