

Procalcitonin implication in renal cell apoptosis induced by acute pyelonephritis in children

Hafid Belhadj-Tahar^{1,2}

Yvon Coulais²

Mathieu Tafani²

François Bouissou³

¹Groupe Santé Recherche, Toulouse, France; ²EA 3033 Université Paul Sabatier III, Toulouse, France; ³Service de Néphrologie Pédiatrique, CHU Purpan, Toulouse, France

Abstract: The aim of this biomedical trial was to clarify the physiological role of procalcitonin (PCT) in renal parenchyma apoptosis and fibrosis caused by acute childhood pyelonephritis. This prospective study enrolled 183 children. All children were treated with bi-therapy according to the French consensus on acute pyelonephritis treatment dated November 16, 1990: intra-vascular administration of ceftriaxone 50 mg/kg/day and netromicine 7 mg/kg/day during the first 48 hours, followed by specific antibiotic therapy suited to antibiogram. On admission, PCT, C-reactive protein, and phospholipase A2 were quantified in serum. Scintigraphy monitoring with ^{99m}Tc-DMSA was performed on day 4 and 9 months later, in the presence of persistent abnormalities. On day 4, 78% presented renal parenchyma alterations and 30% renal fibrosis 9 months after admission. Paradoxically, PCT level was significantly lower in the presence of renal fibrosis due to cell apoptosis (4.19 vs 7.59 µgL⁻¹). A significant increase in PCT indicated favorable progress (recovery 7.55 vs aggravation 3.34) and no difference between recovery and improvement. This result suggests the protective effect of PCT against apoptosis by nitric oxide down-regulation.

Keywords: acute pyelonephritis, procalcitonin, apoptosis, fibrosis, technetium-DMSA

Background

Researchers generally agree that increased procalcitonin (PCT) synthesis is detrimental to the host because a high concentration of this calcitonine precursor reflects the severity of sepsis and is predictive of mortality (Bossink et al 1999). Procalcitonin, a 116 amino acid peptide, is produced physiologically by thyroid C-cells and cleaved by specific protease. This prohormone is found at low concentration (<0.1 ng/mL) in healthy subjects and at high concentration in critically ill patients with a particular infection or shock. Although the role and production site of the PCT are unclear, this peptide induced by tumor necrosis factor alpha (TNF- α) is currently considered to be an effective inflammatory marker used to monitor septic shock, pancreatic necrosis, or burn injuries (Venet et al 2002).

Recently, our research has focused on how PCT is implicated in acute childhood pyelonephritis (AP), followed in 30% of cases by renal fibrosis (RF) leading to chronic sequelae (such arterial hypertension gravidic toxemy). Renal fibrosis results from functional parenchyma disorder consecutive to renal cell apoptosis, usually mediated by lipopolysaccharide (LPS)-induced TNF- α production. During the infection of the renal parenchyma, the bacterial LPS of the membranes induces the release of TNF, interleukin-1 (IL1), and IL6 (Oberhoffer et al 1999).

Pro-inflammatory TNF- α activates nitric oxide (NO) release, which is catalyzed by nitric oxide synthase (NOS) in the presence of Ca²⁺/calmodulin. Thus intracellular calcium concentration is crucial for Ca²⁺/calmodulin complex formation and NOS activation-inducing apoptosis (Arena et al 2002).

Correspondence: Hafid Belhadj-Tahar
Groupe Santé Recherche, 35 Bernard de Ventadour, 31300 Toulouse (France)
Tel +33 5 61 49 20 30
Fax +33 5 61 49 20 40
Email belhadj.h@chu-toulouse.fr

In fact, cell viability and apoptosis depends on the spatiotemporal organization of cytosolic Ca⁺⁺ leading to transmembrane polarization magnitude (Lang et al 2005). In this context, the renal scintigraphy using the polar ^{99m}Tc-DMSA complex allows parenchyma exploration and then monitoring of acute and chronic lesions due to membrane adsorption attenuation (Belhadj-Tahar et al 2000).

The aim of this work is to clarify the physiological role of PCT in AP by monitoring the early and later renal lesions (edema/fibrosis) using ^{99m}Tc-DMSA parenchyma exploration.

Material and methods

This prospective study enrolled 183 children. All subjects exhibited clinical and biological infectious syndrome, with abdominal or lumbar pains and positive urine culture before antibiotic therapy administration. All children were treated with bi-therapy according to the French consensus on acute pyelonephritis treatment dated November 16, 1990: intravascular administration of ceftriaxone 50 mg/kg/day and netromicine 7 mg/kg/day during the first 48 hours, followed by specific antibiotic therapy suited to antibiogram.

On admission C-reactive protein (CRP), phospholipase A₂, and procalcitonin were measured by the following methods: CRP turbidimetric immunoassay (Olympus diagnostica, GmbH) PLA₂: fluometric assay (SFM 25 Kontron Instruments) and PCT: luminescent immunoassay (Brahms diagnostica GmbH). Procalcitonin in serum were quantified as well as CRP and phospholipase A₂. Scintigraphy monitoring with intravascular ^{99m}Tc-DMSA (CIS Bio International, France) was performed according to MacKenzie protocol (MacKenzie 1996) on day 4 and nine months after admission in the presence of persistent abnormalities.

The scintigraphic parenchyma lesions were classified into 4 categories: absence (stage 0), unique and focal (stage II), multiple but unilateral (stage II), and bilateral (stage III).

Results

At day 4, 78% (107/136) had scintigraphic parenchyma renal alterations and 30% (17/57) presented renal fibrosis at 9th month. On admission, except for CRP, both PLA₂ and PCT were significantly higher in patients with early scintigraphic alteration (PCT: 7.85 vs 2.36 µg/L and PLA₂: 72.1 vs 29.9 IU/L) (Table1).

Discussion

During the infection of the renal parenchyma, the bacterial LPS of the membranes induces the release of TNF, IL1, and IL6.

These cytokines are capable of evoking fever, inflammation syndrome, and cell apoptosis followed by fibrosis (Figure 1).

In the acute phase, IL1 and IL6 induce hepatic production of CRP and PLA₂ leukocyte release. PLA₂ is directly implied in the inflammatory process including the prostaglandins production from arachidonic acid in the presence of cyclooxygenase (Diaz 1996). Whereas, procalcitonin CRP and PLA₂ have weak discriminative power for predictive renal fibrosis regarding to their AUC values below 0.65 (see Figure 1).

We hypothesized that PCT down-regulates nitric oxide (NO) release produced by NO-Synthase activated by TNF-α pro-inflammatory cytokine (Figure 2). Indeed, PCT was paradoxically significantly lower in the presence of chronic renal fibrosis (4.19 ± 1.03 vs 7.59 ± 3.45 µgL⁻¹). We observe a significant PCT increase in the favorable progress (recovery 7.55 ± 4.02 vs aggravation 3.34 ± 1.09) and no difference between recovery and improvement evolution. These results suggest the protective effect of PCT against apoptosis resulting from down-regulation of nitric oxide release catalyzed by NO-synthase (Figure 1) and corroborate the previous *In vitro* observations (Hoffmann et al 2003).

Table I Characteristics of 107 patients with acute childhood pyelonephritis

	lesion	age	Sex ratio	CRP (mgL⁻¹)	PLA₂ (IU L⁻¹)	PCT(µgL⁻¹)
4th day	Absence	43.48 ± 7.39 n = 29	M: 7%; F: 15%	77.16 ± 10.21 n = 25	29.90 ± 7.83 n = 27	2.36 ± 1.36 n = 23
	Stage > 0	44.07 ± 4.26 n = 107	M: 29%; F: 71%	114.60 ± 9.74 n = 100	72.03 ± 7.94 n = 87	7.85 ± 1.93 n = 65
	Stage I	37.91 ± 4.47 n = 72	M: 15%; F: 38%	114.60 ± 9.0 n = 66	76.32 ± 13.08 n = 57	6.40 ± 1.99 n = 47
	Stage II	48.70 ± 7.30 n = 21	M: 5%; F: 10%	130.14 ± 23.36 n = 21	32.70 ± 6.72 n = 17	7.34 ± 4.66 n = 11
	Stage III	33.47 ± 8.79 n = 14	M: 1%; F: 9%	121.07 ± 16.46 n = 14	104.65 ± 28.47 n = 13	18.37 ± 9.46 n = 7
9th month	Absence	39.86 ± 6.78 n = 40	M: 17%; F: 53%	101.16 ± 12.20 n = 37	61.80 ± 12.22 n = 33	7.59 ± 3.45 n = 24
	Fibrosis (Stage I)	27.19 ± 5.54 n = 17	M: 7%; F: 23%	123.35 ± 21.35 n = 17	81.65 ± 20.38 n = 13	4.19 ± 1.03 n = 11




Figure 1 Receiver operating characteristics curves for C-reactive protein (CRP), phospholipase A₂ (PLa₂), procalcitonin (PCT), and renal fibrosis observed by ^{99m}Tc-DMSA scintigraphy.

Note: Areas under curve (AUC) of the 3 markers were respectively (CI: 0.26163–0.72184) for CRP; 0.5475 (CI: 0.33059–0.76445) for PLa₂ and 0.6240 (CI: 0.42373–0.82421) for PCT.




Figure 2 Physiopathology of pyelonephritis.

Abbreviations: IL, interleukin; NO, nitric oxide; NOS, NO synthase; PCT, procalcitonin; PLa₂, phospholipase A₂; CRP, C-reactive protein; FGF, fibroblast growth factor; TNF, tumor necrosis factor.

Table 2 Pyelonephritis progress according to procalcitonin concentration

	No sequelae	Fibrosis sequelae	F. test
Early lesion stage I-II	$7.59 \pm 3.45 \text{ mgL}^{-1}$ n = 24 Recovery stage(0)	$4.19 \pm 1.03 \text{ mgL}^{-1}$ n = 11 Aggravation stage (→I)	p < 0.001
Early lesion stage: I	$7.55 \pm 4.02 \text{ mgL}^{-1}$ n = 20 recovery stage (0)	$3.34 \pm 1.09 \text{ mgL}^{-1}$ n = 6 Improvement stage (\geq I→I)	p < 0.001
Early lesion stage: \geq II	$7.78 \pm 5.98 \text{ mgL}^{-1}$ n = 4	$4.60 \pm 2.36 \text{ mgL}^{-1}$ n = 5	ns

Note: The procalcitonin levels are listed according to pyelonephritis progress.

Conclusion

Our study shows that PCT plays key role in the renal parenchyma protection against cell apoptosis and fibrosis caused by acute pyelonephritis. this finding constitutes a original approach in the comprehension of apoptosis paradigm that causes parenchyma fibrosis.

Acknowledgments

The authors thank AFPreMed (Association Française de Promotion de la Recherche Médicale) for invaluable help. The authors report no conflicts of interest in this work.

References

- Arena S, Pattarozzi A, Thellung V, et al. 2002. Nitric oxide production stimulated by the basic fibroblast growth factor requires the synthesis of ceramide. *Ann NY Acad Sci*, 973:94–104.
- Belhadj-Tahar H, Darbieu MH, Boumahdi R. 2000. Conceptualisation of diagnostic agents : From empirical *in vivo* screening to rational *in vitro* predictive parameters. *ATLA*, 28:303–14.
- Bossink AW, Groeneveld AB, Thijss LG. 1999. Prediction of microbial infection and mortality in medical patients with fever: plasma procalcitonin, neutrophilic elastase-alpha1-antitrypsin, and lactoferrin compared with clinical variables. *Clin Infect Dis*, 29:398–407.
- Diaz BL, Arm JP. 2003. Phospholipase A(2). *Prostaglandins Leukot Essent Fatty Acids*, 69:87–97.
- Hoffmann G, Totzke G, Seibel M, et al. 2001. In vitro modulation of inducible nitric oxide synthase gene expression and nitric oxide synthesis by procalcitonin. *Crit Care Med*, 29:112–6.
- Lang F, Foller M, Lang KS, et al. 2005. Ion channels in cell proliferation and apoptotic cell death. *J Membr Biol*, 205:147–57.
- MacKenzie JR. 1996. A review of renal scarring in children. *Nucl Med Commun*, 17:176–90.
- Oberhoffer M, Karzai W, Meier-Hellmann A, et al. 1999. Sensitivity and specificity of various markers of inflammation for the prediction of tumor necrosis factor-alpha and interleukin-6 in patients with sepsis. *Crit Care Med*, 27:1814–8.
- Venet C, Tardy B, Zeni F. 2002. [Marqueurs biologiques de l'infection en réanimation chez l'adulte: place de la procalcitonine.] *Réanimation*, 11:156–71.