

Co-Existence of *mcr-1* and *bla*_{NDM-5} in an *Escherichia coli* Strain Isolated from the Pharmaceutical Industry, WWTP

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Huiming Han^{1,2}
Wenhong Liu³
Xinjie Cui³
Xu Cheng^{1,2}
Xiawei Jiang³

¹Medical College, Beihua University, Jilin, People's Republic of China; ²The Center for Infection and Immunity, Beihua University, Jilin, People's Republic of China; ³College of Basic Medical Sciences, Zhejiang Chinese Medical University, Hangzhou, Zhejiang, People's Republic of China

Abstract: The emergence of the plasmid-borne colistin-resistant gene (*mcr-1*) poses a great threat to human health. What is worse, the recent observations of the co-existence of *mcr-1* with other antimicrobial resistance genes in some bacteria cause further concern. Here, we present the first report of a wild *Escherichia coli* strain that co-carries an *mcr-1* encoding phage-like IncY plasmid (pR15_MCR-1) and a *bla*_{NDM-5} encoding IncX3 plasmid (pR15_NDM-5) from a pharmaceutical industry, wastewater treatment plant, in China. This study highlights the spreading of *E. coli* carrying both *mcr-1* and *bla*_{NDM-5} genes in the pharmaceutical industry.

Importance: *Escherichia coli* strains that carry both *mcr-1* and *bla*_{NDM-5} genes are of great health concern and are already found in humans and animals worldwide, yet there is a paucity of observations of this resistant strain in the environment. Here we present the first isolation of an *E. coli* strain (R15) that co-carries *mcr-1* and *bla*_{NDM-5} genes from a wastewater treatment plant in China. Whole-genome sequencing indicated that R15 harbored two plasmids, pR15_MCR-1 and pR15_NDM-5, that carry *mcr-1* and *bla*_{NDM-5}, respectively. The observation of this wild-derived *E. coli* strain that carries *mcr-1* and *bla*_{NDM-5} genes simultaneously calls for the urgency to improve monitoring and reducing its further spreading.

Keywords: *mcr-1*, *bla*_{NDM-5}, wastewater treatment plant, *Escherichia coli*

The transferable colistin resistance gene, *mcr-1*, has been a growing concern worldwide because colistin is considered as the last resort in the treatment of multidrug-resistant pathogens.¹ Recently, reports of bacteria that carry the *mcr-1* gene along with other types of antimicrobial resistance genes (ARGs), especially beta-lactam resistance genes, have highlighted the threat of these pan-drug resistant pathogens to public health. Several cases of infection caused by a pathogen which harbored *mcr-1* and beta-lactam resistance genes were reported worldwide.²⁻⁴ In addition, bacteria that carried *mcr-1* and beta-lactam resistance genes were also found in the natural environment. The occurrence of the *mcr-1* gene in extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* from well water was reported in rural China.⁵ In our previous study, we isolated six *mcr-1*-positive ESBL-producing *E. coli* strains from farm soils.⁶ Wastewater treatment plants (WWTPs), which have been taken as deep reservoirs of ARGs and antimicrobial-resistant bacteria (ARB), are believed to offer the opportunity for ARGs to flow into susceptible disease-causing bacteria.^{7,8} Thus, the surveillance of these environment-derived bacteria that carry the *mcr-1*

Correspondence: Xiawei Jiang
College of Basic Medical Sciences,
Zhejiang Chinese Medical University, No.
548 Binwen Road, Hangzhou 310053,
Zhejiang, People's Republic of China
Tel/Fax +86-571-86613613
Email xiawei.jiang@zcmu.edu.cn

gene with other types of ARGs is of great importance and the transmission patterns of these bacteria between different habitats should be elucidated.

In our previous study, we obtained several ESBL-producing *E. coli* strains from a WWTP located in Taizhou, Zhejiang province, China.⁹ This WWTP treats the wastewater from a pharmaceutical company which produces antimicrobial agents including beta-lactam antibiotics such as cefaclor, cefprozil, cefixime, cefepime, ceftriaxone, cefuroxime sodium, cefazolin, cefepime, cefepime (MIC ≥ 64 $\mu\text{g}/\text{mL}$), ertapenem (MIC ≥ 8 $\mu\text{g}/\text{mL}$), imipenem (MIC ≥ 16 $\mu\text{g}/\text{mL}$), amikacin (MIC ≥ 64 $\mu\text{g}/\text{mL}$), gentamicin (MIC ≥ 16 $\mu\text{g}/\text{mL}$), tobramycin (MIC ≥ 16 $\mu\text{g}/\text{mL}$), ciprofloxacin (MIC ≥ 4 $\mu\text{g}/\text{mL}$), levofloxacin (MIC ≥ 8 $\mu\text{g}/\text{mL}$), nitrofurantoin (MIC = 128 $\mu\text{g}/\text{mL}$) and trimethoprim (MIC ≥ 320 $\mu\text{g}/\text{mL}$). Resistance to colistin (MIC = 8 $\mu\text{g}/\text{mL}$) was examined using broth microdilution test as described before.⁴

The isolation and identification of R15 was performed as described previously.⁹ Antimicrobial susceptibility testing by VITEK 2 Compact (bioMérieux, France) showed that R15 was resistant to various types of antimicrobial agents, including ampicillin (MIC ≥ 32 $\mu\text{g}/\text{mL}$), amoxicillin (MIC ≥ 32 $\mu\text{g}/\text{mL}$), piperacillin (MIC ≥ 128 $\mu\text{g}/\text{mL}$), cefazolin (MIC ≥ 64 $\mu\text{g}/\text{mL}$), cefoxitin (MIC ≥ 64 $\mu\text{g}/\text{mL}$), ceftriaxone (MIC ≥ 64 $\mu\text{g}/\text{mL}$), cefepime (MIC ≥ 64 $\mu\text{g}/\text{mL}$), ertapenem (MIC ≥ 8 $\mu\text{g}/\text{mL}$), imipenem (MIC ≥ 16 $\mu\text{g}/\text{mL}$), amikacin (MIC ≥ 64 $\mu\text{g}/\text{mL}$), gentamicin (MIC ≥ 16 $\mu\text{g}/\text{mL}$), tobramycin (MIC ≥ 16 $\mu\text{g}/\text{mL}$), ciprofloxacin (MIC ≥ 4 $\mu\text{g}/\text{mL}$), levofloxacin (MIC ≥ 8 $\mu\text{g}/\text{mL}$), nitrofurantoin (MIC = 128 $\mu\text{g}/\text{mL}$) and trimethoprim (MIC ≥ 320 $\mu\text{g}/\text{mL}$). Resistance to colistin (MIC = 8 $\mu\text{g}/\text{mL}$) was examined using broth microdilution test as described before.⁴

To investigate the genotype in correspondence to its antimicrobial phenotype, whole-genome sequencing of R15 was performed using Illumina Hiseq 2000 sequencer (Illumina, USA) as described previously¹⁰ and reads were assembled into 140 contigs (PSSH00000000.1) using SPAdes software.¹¹ Analyzing with the draft genome sequence of R15 by MLST 2.0¹² showed that it belonged to ST-744. Further, we used ResFinder 3.2¹³ to predict the resistance genes encoded by strain R15 and 20 resistance genes were identified, including one copy of *mcr-1*, *bla*_{CTX-M-14}, *bla*_{NDM-5}, *bla*_{TEM-1B}, *fosA3*, *floR*, *aac(3)-IV*, *aph(3'')-Ib*, *aph(3'')-IIa*, *aph(4)-Ia*, *aph(6)-Id*, *rmtB*, *mdf(A)*, *mph(A)*, *oqxA*, *oqxB*, *tet(B)*, and *dfrA17*, and two copies of *sul2* genes. Some of these resistance genes might be responsible for the resistant phenotype of strain R15. For example, three beta-lactam resistance genes, *bla*_{CTX-M-14}, *bla*_{NDM-5}, *bla*_{TEM-1B}, which were harbored by strain R15, might be related to its resistance to several beta-lactams antimicrobial agents such as ampicillin, amoxicillin, piperacillin, and cefazolin, etc. In addition, one *mcr-1* gene was also identified which might be responsible for the resistance of colistin of strain R15. However, here we only represented

required resistance genes, so there might be some other mechanisms that exist in strain R15 which may also be related to its resistance phenotype. To a further extent, the co-existence of *mcr-1* and *bla*_{CTX-M} in *E. coli* strains has been observed in different sources of samples.¹⁴ Meanwhile, *E. coli* strains with both *mcr-1* and *bla*_{NDM-5} were also found in humans and animals.^{2-4,15,16} However, R15 was isolated from a WWTP which treats wastewater generated by a pharmaceutical factory. To the best of our knowledge, R15 is the first reported wild-derived *E. coli* strain that harbored both *mcr-1* and *bla*_{NDM-5} genes in China.

To gain insight into the locations and the gene environments of the *mcr-1* and *bla*_{NDM-5} genes in R15, whole-genome sequencing using Pacific Biosciences (PacBio) RSII sequencing system with single-molecule real-time (SMRT) analysis was performed. Two complete plasmids that carry *mcr-1* (pR15_MCR-1) and *bla*_{NDM-5} (pR15_NDM-5) were assembled from the sequencing reads. The complete sequences of these two plasmids were deposited in GenBank with the accession numbers, MK256965 (pR15_MCR-1) and MK256964 (pR15_NDM-5).

Plasmid pR15_MCR-1 is 109,908 bp in length with a G+C content of 47.0%. Plasmid typing revealed that it belongs to the IncY group. Nucleotide sequence alignment indicated that pR15_MCR-1 possesses 98% homology and 74% coverage with plasmid pMCR-1-P3 (KX880944). Further comparison of the gene arrangement of these two plasmids revealed the absence of the second IS*Apl1* element downstream of *mcr-1* gene in pR15_MCR-1 (Figure 1A). The absence of this second IS*Apl1* element was also reported in other studies and was considered as the indication of the dynamic changes of Tn*Apl* during the transposition process.^{3,17} Furthermore, an insertion of approximately 11 kb in pR15_MCR-1 was found downstream of the *mcr-1* gene. This insertion showed 99% nucleotide homology to the chromosome of a porcine extraintestinal pathogenic *E. coli* strain PCN033 (CP006632) isolated in China.¹⁸ An insertion element IS609 was found adjacent to this insertion that may be responsible for its transfer (Figure 1A). Since its first discovery, the *mcr-1* gene has been found in various incompatibility groups of plasmids, including IncX4, IncI2, IncP, IncFI, IncFII, IncHI1, IncHI2, IncX1-X2, IncX3-X4, and IncI2-IncFIB.¹⁷ Plasmid pR15_MCR-1, identified in the present study, belongs to the IncY incompatibility group. This group of plasmids has commonly been reported to be carrying the *bla*_{CTX-M-15} gene.¹⁹ Moreover, IncY plasmids contain a portion of phage-related sequences, which imply that phages may play an important role in disseminating of the *mcr-1* gene.²⁰ To date, there have only been two studies




Figure 1 Lineal comparison of antimicrobial-resistant regions. **(A)** Comparison of the *mcr-1* coding region of plasmid pR15_MCR-1 (MK256965) with plasmid pMCR-1-P3 (KX880944) and the chromosome of *E. coli* PCN033 (CP006632). **(B)** Comparison of the *bla*_{NDM-5} coding region of plasmids pR15_NDM-5 (MK256964) and pNDM_MGR194 (KF220657). Genes are portrayed by arrows and colored according to their functions.

reporting the carrying of the *mcr-1* gene in IncY group plasmids. One was pMCR-1-P3, identified in *E. coli* strain P3 isolated from pig anal swabs in Shandong province in China.²⁰ The other was pHYEC7-*mcr1*, which was harbored by *E. coli* strain HYEC7 isolated from a fecal sample collected from a pig farm in Guangdong province in China.²¹ Thus, both plasmids were harbored by *E. coli* strains of animal origin. However, pR15_MCR-1 in the present study was identified from an *E. coli* isolated from a WWTP in Zhejiang province in China, and this WWTP has no contact with animal waste. Thus, our report is the first discovery of an IncY plasmid that carries the *mcr-1* gene in an *E. coli* strain of environmental origin. Our work highlights the dissemination of this kind of plasmid in the environment.

Plasmid pR15_NDM-5, which carries the *bla*_{NDM-5} gene, is 46,161 bp in length. This plasmid belongs to the IncX3 group which was reported to be the most common Inc type plasmid that harbored *bla*_{NDM} gene in China.²² Nucleotide BLAST revealed that pR15_NDM-5 was highly similar to pNDM-MGR194 (99% identity and 100% coverage).²³ Furthermore, genetic environment characterization revealed a structure of IS26-*dsbC*-*trpF*-*ble*_{MBL}-*bla*_{NDM-5}-IS5 in both plasmids (Figure 1B). Thus, our discovery of this plasmid supplements previous studies and further highlights the dissemination of *bla*_{NDM-5} gene-carrying IncX3 group plasmids in WWTP.

In conclusion, we report the complete sequences of an IncY type *mcr-1* carrying plasmid and an IncX3 type *bla*_{NDM-5} carrying plasmid in an *E. coli* strain isolated from WWTP.

There is a growing body of literature that reports the co-existence of *mcr-1* and *bla*_{NDM-5} genes in *E. coli* strains of clinic-origin and animal-origin. However, our study represents the first report of a wild-derived *E. coli* strain that harbors *mcr-1* and *bla*_{NDM-5} genes simultaneously. To lower the risk of the dissemination of this multidrug-resistant strain in the environment, more surveillance is needed in future.

Acknowledgments

This work was supported by grants from: the National Key Research and Development Program of China (No.2016YFD0501105), National Natural Science Foundation of China (41406140 and 81741098), and Zhejiang Provincial Natural Science Foundation of China (No. LY17H190003); Innovation funding project for postgraduate students of Beihua University (Research and Innovation [2018] 002).

Disclosure

The authors report no conflicts of interest in this work.

References

- Liu -Y-Y, Wang Y, Walsh TR, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. *Lancet Infect Dis.* 2016;16(2):161–168. doi:10.1016/S1473-3099(15)00424-7
- Mediavilla JR, Patrawalla A, Chen L, et al. Colistin-and carbapenem-resistant *Escherichia coli* harboring *mcr-1* and *bla*_{NDM-5}, causing a complicated urinary tract infection in a patient from the United States. *MBio.* 2016;7(4):e01191–16. doi:10.1128/mBio.01191-16

3. Yu H, Qu F, Shan B, et al. Detection of the *mcr-1* colistin resistance gene in carbapenem-resistant Enterobacteriaceae from different hospitals in China. *Antimicrob Agents Chemother.* 2016;60(8):5033–5035. doi:10.1128/AAC.00440-16
4. Zheng B, Lv T, Xu H, et al. Discovery and characterisation of an *Escherichia coli* ST206 strain producing NDM-5 and MCR-1 from a patient with acute diarrhoea in China. *Int J Antimicrob Agents.* 2018;51(2):273–5. PubMed PMID: 28919194.
5. Sun P, Bi Z, Nilsson M, et al. Occurrence of blaKPC-2, blaCTX-M, and *mcr-1* in Enterobacteriaceae from well water in rural China. *Antimicrob Agents Chemother.* 2017;61(4):PubMed PMID: 28115344. PMCID: PMC5365680.
6. Zheng B, Huang C, Xu H, et al. Occurrence and genomic characterization of ESBL-producing, MCR-1-harboring *Escherichia coli* in farming soil. *Front Microbiol.* 2017;8:2510. doi:10.3389/fmicb.2017.02510
7. Guo J, Li J, Chen H, Bond PL, Yuan Z. Metagenomic analysis reveals wastewater treatment plants as hotspots of antibiotic resistance genes and mobile genetic elements. *Water Res.* 2017;123:468–478. doi:10.1016/j.watres.2017.07.002
8. Wright GD. Environmental and clinical antibiotic resistomes, same only different. *Curr Opin Microbiol.* 2019;51:57–63. doi:10.1016/j.mib.2019.06.005
9. Jiang X, Cui X, Xu H, et al. Whole genome sequencing of extended-spectrum beta-lactamase (ESBL)-producing *Escherichia coli* isolated from a wastewater treatment plant in China. *Front Microbiol.* 2019;10. doi:10.3389/fmicb.2019.01797(1797)
10. Zheng B, Jiang X, Cheng H, et al. Genome characterization of two bile-isolated *Vibrio fluvialis* strains: an insight into pathogenicity and bile salt adaptation. *Sci Rep.* 2017;7(1):11827. doi:10.1038/s41598-017-12304-8
11. Bankevich A, Nurk S, Antipov D, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J comput biol.* 2012;19(5):455–477. doi:10.1089/cmb.2012.0021
12. Larsen MV, Cosentino S, Rasmussen S, et al. Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol.* 2012;50(4):1355–1361. doi:10.1128/JCM.06094-11
13. Zankari E, Hasman H, Cosentino S, et al. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother.* 2012;67(11):2640–2644. doi:10.1093/jac/dks261
14. Wu C, Wang Y, Shi X, et al. Rapid rise of the ESBL and *mcr-1* genes in *Escherichia coli* of chicken origin in China, 2008–2014. *J Glob Antimicrob Resist.* 2018;7(1):30. doi:10.1038/s41426-018-0033-1
15. Yang R-S, Feng Y, Lv X-Y, et al. Emergence of NDM-5-and MCR-1-producing *Escherichia coli* clones ST648 and ST156 from a single muscovy duck (*Cairina moschata*). *Antimicrob Agents Chemother.* 2016;60(11):6899–6902. doi:10.1128/AAC.01365-16
16. Zhang Y, Liao K, Gao H, et al. Decreased fitness and virulence in ST10 *Escherichia coli* Harboring blaNDM-5 and *mcr-1* against a ST4981 strain with blaNDM-5. *Front Cell Infect Microbiol.* 2017;7:242. doi:10.3389/fcimb.2017.00242
17. Wang Q, Sun J, Li J, et al. Expanding landscapes of the diversified *mcr-1*-bearing plasmid reservoirs. *Microbiome.* 2017;5(1):70. doi:10.1186/s40168-017-0288-0
18. Liu C, Zheng H, Yang M, et al. Genome analysis and in vivo virulence of porcine extraintestinal pathogenic *Escherichia coli* strain PCN033. *BMC Genomics.* 2015;16(1):717. doi:10.1186/s12864-015-1890-9
19. Saidani M, Messadi L, Mefteh J, et al. Various Inc-types plasmids and lineages of *Escherichia coli* and *Klebsiella pneumoniae* spreading bla_{CTX-M-15}, bla_{CTX-M-1} and *mcr-1* genes in camels in Tunisia. *J Glob Antimicrob Resist.* 2019;19:280–283. doi:10.1016/j.jgar.2019.05.007.
20. Zhang C, Feng Y, Liu F, et al. A phage-like IncY plasmid carrying the *mcr-1* gene in *Escherichia coli* from a pig farm in China. *Antimicrob Agents Chemother.* 2017;61(3):e02035–16. doi:10.1128/AAC.02035-16
21. Li R, Xie M, Lv J, Wai-Chi Chan E, Chen S. Complete genetic analysis of plasmids carrying *mcr-1* and other resistance genes in an *Escherichia coli* isolate of animal origin. *J Antimicrob Chemother.* 2017;72(3):696–699. doi:10.1093/jac/dkw509
22. Yamamoto M, Matsumura Y, Gomi R, et al. Interspecies dissemination of a mobilizable plasmid harboring bla_{IMP-19} and the possibility of horizontal gene transfer in a single patient. *Antimicrob Agents Chemother.* 2016;60(9):5412–5419. doi:10.1128/AAC.00933-16
23. Krishnaraju M, Kamatchi C, Jha A, et al. Complete sequencing of an IncX3 plasmid carrying bla_{NDM-5} allele reveals an early stage in the dissemination of the bla_{NDM} gene. *Indian J Med Microbiol.* 2015;33(1):30. doi:10.4103/0255-0857.148373

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

Dovepress