

PTBPI Targets ILK to Regulate the Hypoxia-Induced Phenotypic Transformation of Pulmonary Artery Smooth Muscle Cells

Gaoliang Yan
Renhua Sun
Zhongpu Chen
Xiaodong Pan
Zulong Sheng
Chengchun Tang

Department of Cardiology, Zhongda
Hospital of Southeast University Medical
School, Nanjing, 210009, People's
Republic of China

Purpose: Pulmonary hypertension (PH) is a pathological process mainly characterized by the progressive increase in pulmonary vascular resistance. The degradation of pulmonary artery smooth muscle cells (PASMCs) from contractile/differentiated phenotype to synthetic/dedifferentiated phenotype is a key factor for hypoxic pulmonary hypertension.

Materials and Methods: In this study, qPCR was performed to evaluate the gene expression of mRNAs. Western blot, immunofluorescence and RNA pull down were used to detect gene expression levels.

Results: We found that the gene expression of polypyrimidine tract-binding protein1 (PTBPI) was increased significantly in a time-dependent manner in rats PA tissues and PASMCs after hypoxia. PTBPI knockdown can inhibit the phenotypic transition of PASMCs. PTBPI inhibits the phenotypic transition of PASMCs. In addition, PTBPI inhibits the integrin-linked kinase (ILK) expression under hypoxic conditions, thereby down-regulating the expression of downstream proteins. It inhibits the phenotypic transition of PASMCs and alleviates pulmonary hypertension.

Conclusion: In conclusion, PTBPI/ILK axis promotes the development of PH via inducing phenotypic transition of PASMCs. This may provide a novel therapy for PH.

Keywords: pulmonary hypertension, polypyrimidine tract-binding protein 1, integrin-linked kinase, pulmonary artery smooth muscle cells

Introduction

Pulmonary hypertension (PH) is a pathological process mainly characterized by the progressive increase in pulmonary vascular resistance and progressive failure of right heart function.^{1,2} PH is a common complication of many clinical diseases. Although extensive researches have been carried out to study the initiation and progression of pulmonary hypertension, the potential mechanisms are still unclear. Studies suggested PH may be related to several factors such as hypoxia, neurohumoral, congenital, genetic and others.^{3,4} Its pathogenesis has not been fully understood.

Polypyrimidine tract-binding protein1 (PTBPI) is a member of heterogeneous nuclear ribonucleoproteins (hnRNPs) (also named hnRNPI),⁵ which is involved in the transcriptional regulation of many genes. PTBPI modulates tumor metastasis by regulating metastasis-related genes. In tumor cells, PTBPI promotes the expression of PKM2 isoforms by regulating the alternative splicing of the pyruvate kinase PKM gene, thereby promoting the glycolysis process of tumor cells and

Correspondence: Gaoliang Yan
Department of Cardiology, Zhongda
Hospital of Southeast University Medical
School, Nanjing, 210009, People's
Republic of China
Tel +86 18761890380
Email yangaol@hotmail.com

thus inducing tumorigenesis.⁶ Integrin-linked kinase (ILK) is a serine/threonine kinase that regulates cell survival, proliferation, and apoptosis through glycogen synthesis kinase 3 β (GSK-3 β).⁷ Hypoxia suppresses the expression of myocardin in smooth muscle cells and may be an essential mechanism involved in the formation of hypoxic PH.^{8,9} ILK is upstream to many intracellular molecules involved in hypoxic stress. Investigating the role of the ILK signaling pathway in the molecular mechanism of PH may provide theoretical and experimental evidence for clinical drug treatment of pulmonary hypertension.

In this study, we explored the role of PTBP1 and ILK in the hypoxia-induced PASMCM phenotypic transition. We speculate that PTBP1 regulates PASMCM by inhibiting the expression of ILK. To test this hypothesis, we identified the role of PTBP1 in the phenotypic transformation of PASMCMs through silencing and overexpressing the PTBP1 gene *in vivo* and *in vitro*. Our results may provide a new therapeutic strategy for the prevention and treatment of PH.

Materials and Methods

Pulmonary Hypertension Animal Model

Thirty male SD rats, weighing 280–320g, were provided by the Animal Center of Zhongda Hospital of Southeast University Medical School. The rats were divided into control group and hypoxia treatment group. To establish the PH model, the rats were housed in a hypobaric hypoxia chamber depressurized to 380 mmHg (PO₂ was reduced to about 79.6 mmHg accordingly) for 4 weeks. All the rats were housed in a condition of 12:12 hour light-dark cycle with free access to food and water. The Ethics Committee in Zhongda Hospital of Southeast University Medical School approved the animal study. All animal procedures were performed following the institutional guidelines and approved by the Zhongda Hospital of Southeast University Medical School.

Relative Ventricular Weight

The relative ventricular weight was assessed after sacrificing of the rats and clearing the blood from isolated hearts. The moisture over the hearts was absorbed with filter paper, and we determined the hearts' weight. Then we calculated the right ventricular (RV)/left ventricle + septum (LV + S), RV weight-to-body weight (RV/BW), and (LV + S)/BW.

Hemodynamics Measurements

After hypoxia, the rats were injected with pentobarbital sodium (60 mg/kg). A polyethylene microcatheter was inserted into the right ventricle and pulmonary artery and the mean pulmonary arterial pressure (mPAP).

The Detection of Wall Thickness (WT) and Wall Area (WA)

The right lower lung from the sacrificed animals were immersed in 10% formalin fixation. Then, specimens were dissected and stained with Hematoxylin. Ten pulmonary arterioles, each having a tube diameter of 50 to 100 μ m, were randomly selected. The wall thickness (WT) and external diameter (ED) were determined to calculate the percentage of pulmonary arteriolar wall thickness (WT%) to the diameter of the tube using this equation ($WT\% = 2 \times WT/ED \times 100\%$). The wall area (WA) % of the blood vessel was calculated using this equation, $WA\% = (TA-LA)/TA \times 100\%$.

Primary Culture of Pulmonary Artery Smooth Muscle Cells

Wister rats were anaesthetized intraperitoneally with 1% sodium pentobarbital (40mg/kg). Rat heart and lung tissues were removed under aseptic conditions, pulmonary arteries were separated, and washed with Hanks' solution to remove blood stains on the tissue surface. Then, the connective tissue and blood vessels were removed and the vascular endothelial cells were scraped. The tissues were divided into 0.5–1 mm³ size pieces and rinsed with Hanks solution until it became clear. Next, Hanks solution was removed, and digestion was performed using 0.2 mL type II collagenase solution at 37°C for 15 min. The digested samples were filtered and centrifuged at 1000 r/min for 5 min, and the supernatant was discarded. The pellet was washed with Hanks solution, followed by centrifugation; then the cells were resuspended in DMEM culture medium containing 10% fetal bovine serum and seeded into a 24-well culture plate.

Inducing Cellular Hypoxia

PASMCMs were cultured medium containing 0.1% FBS for 24 hours. Then, PASMCMs were treated with CoCl₂ (100 μ M) under the hypoxic condition for 1 h, 6 h, 24 h, 48 h, 72 h. The cells were placed in a hypoxia-specific incubator (XBS-08, Aipu, Hangzhou, China) containing 5% CO₂ at 37°C. The control PASMCMs were maintained under normoxic conditions.

Plasmid Transfection

siRNA for PTBP1 were synthesized by Genewiz (Beijing, China). The overexpressing vector of PTBP1 and ILK were purchased from Hanbio (Shanghai, China). Plasmids were transfected using Lipofectamine 3000 (Invitrogen, USA) kit following the manufacturer's instructions. Adeno virus silencing PTBP1 was purchased from Hanbio (Beijing, China). Ad-sh-PTBP1 (0.5 mL of 4×10^8 plaque-forming units) was injected into the rats via tail vein.

Immunoprecipitation (IP)

Total protein was extracted from PSMCs with RIPA buffer. The protein sample was incubated with 25 μ L protein A/G agarose and 25 μ L immunoglobulin for 1 h at 4°C followed by centrifugation at $2000 \times g$ at 4°C for 5 min. Primary antibody (4 μ g) was added and incubated overnight at 4°C. Immunoglobulin (4 μ g) was added to the protein sample as a negative control. Then, 25 μ L of protein/G agarose was added, followed by centrifuging at 4°C for 2 h. After washing, the immunoprecipitation complex was boiled in $2 \times$ protein loading buffer for 10 min for further Western blot analysis.¹⁰

Immunofluorescence

The changes in phenotypic markers of PSMCs were detected by immunofluorescence staining. According to the referenced method, cells grown over coverslips were washed with PBS buffer, fixed with formaldehyde for 10min, and further washed with PBS. The primary antibody was added (1:1000 dilution) and incubated for 1 h at 37°C on a shaker. Coverslips were washed three times with PBS and incubated with a FITC-labeled secondary antibody diluted 1:80 at 37°C for 30 min on a shaker. We used an imaging analysis system at 400x magnification power for each specimen. Images were taken to observe the intensity of F-actin staining and cell morphology. PTPB1 and ILK co-localization was determined by double staining.¹¹

Western Blot

Proteins from tissue samples were isolated and quantified using BCA kit according to the instructions. Then, standard Western blotting was performed as following: SDS-PAGE electrophoresis, membrane transfer, primary antibody, and secondary antibody addition, development, and fixation. GAPDH was used as an internal reference. Band analysis

was performed using image analysis software. The ratio of the grayscale of the target protein to the grayscale of the GAPDH color represents the relative expression of the target protein.¹²

Statistical Analysis

SPSS 19.0 software package was used for statistical data analysis Each index is expressed as a mean value \pm standard deviation. Differences between different groups were analyzed using the Mann–Whitney non-parametric tests. $P < 0.05$ was considered as statistically significant.¹³

Results

Identification PH Animal Model

In order to study the pathogenesis of PH, we constructed the PH animal model. After establishing a PH model, we measured the RV/(LV + S), mPAP, WT, and WA (n=6). Compared to the control group, these indexes started to increase 2 weeks after establishing the model and reached their maximum at 4 weeks. These findings indicated the successful establishment of the PH model (Figure 1A). We examined the expression of PTBP1 in pulmonary arterial tissue of rats with PH by Western blot and immunofluorescence assays. The results showed that the expression of PTBP1 was significantly increased 3 weeks after hypoxia (Figure 1B). Then, we evaluated the expression of PTBP1 of PSMCs using Western blot and immunofluorescence. The results indicated that hypoxia significantly promoted the expression PTBP1 after 24h (Figure 1C and D).

PTBP1 Silencing Inhibits Phenotype Transition of PSMCs

Since the expression of PTBP1 was significantly increased in rats with HP, we speculated that it might play a role in the transformation of PSMCs, so we established a hypoxia model of PSMCs. We found that knockdown of PTBP1 inhibited the phenotype switching of PSMCs induced by hypoxia (Figure 2A), and the PSMCs length and area were significantly reduced (Figure 2B and C). The experimental results show that PTBP1 plays an essential role in the phenotypic transition of PSMCs.

PTBP1 Inhibits Expression of ILK and Downstream Signaling Pathway

In order to further confirm the regulation of PTBP1 on the deformation and transformation of PSMCs, we examined




Figure 1 Establishment of an animal model of pulmonary hypertension. **(A)** Detection of RV/(LV + S)%, mPAP, WT% and WA% in rats. The results showed that the model was successfully prepared. **(B)** The western method was used to detect the expression of PTBP1 in the pulmonary artery tissue of rats with pulmonary hypertension. **(C and D)** The Western blot and immunofluorescence method were used to detect the expression of PTBP1 in the PASMCs. n=6, *p<0.05, **p<0.01.




Figure 2 PTBP1 interference inhibits phenotypic switching of pulmonary artery smooth muscle cells. **(A)** F-actin staining was used to detect phenotypic changes in smooth muscle cells. **(B)** The cell length of PASMCs was calculated. **(C)** Results of the area of PASMCs were detected. n=6, **p<0.01.

the expression of PTBP1-related proteins. The results showed that knockdown of PTBP1 significantly decreased the protein level of ILK, meantime, it can inhibit the expression of Myocardin, SM α -actin, Calponin, and increase the

expression of osteopontin (Figure 3A). ILK overexpression can reverse the PTBP1 regulatory effect. This finding indicates that PTBP1 can regulate ILK and affect the expression of several downstream signaling proteins (Figure 3B).




Figure 3 PTBP1 inhibits ILK and downstream signaling pathway protein expression. (A) The expressions of ILK, Myocardin, SM α -actin, Calponin, and osteopontin were detected by Western blot. (B) Statistical results of expression changes of ILK, Myocardin, SM α -actin, Calponin, and osteopontin. $n=6$, $**p<0.01$.




Figure 4 PTBP1 interact with ILK and regulates its RNA stability. (A) RNA pulled down + silver staining were performed to identify that ILK mRNA interact with PTBP1 protein. (B) RIP experiments along with the PCR was used to confirm the interaction between PTBP1 and ILK. (C) RNA pull down was performed to detect the interaction between PTBP1 and ILK. (D) Bioinformatics analysis proves that PTBP1 and ILK can be combined. (E) qPCR was used to evaluate the mRNA expression of ILK at different time point in the PSMCs after actinomycin D treatment. (F) Western blot was performed to confirm that overexpression of PTBP1 could inhibit ILK protein expression while PTBP1 knockdown promoted ILK expression. (G) Fish experiment was conducted to detect the co-localization of PTBP1 and ILK. $n=6$, $**p<0.01$.




Figure 5 PTBPI silencing modulates the expression of phenotypic transition-related genes. **(A)** Western blot was used to detect the expression of phenotypic transition-related genes in PSMCs including PTBPI, ILK, Myocardin, SM α -actin, Calponin, and osteopontin. **(B)** Statistical results of protein expression was calculated. $n=6$, $**p<0.01$.

PTBPI Binds to ILK's RNA and Maintains the RNA Stability of ILK

To clarify the precise mechanism of action of PTBPI, we performed RNA pulled down and mass spectrometry experiments (Figure 4A) also, RIP experiments (Figure 4B). The results showed that PTBPI and ILK had direct interaction (Figure 4C). Bioinformatics analysis proves that these two molecules can interact with each other (Figure 4D). After the addition of transcription suppressor actinomycin D to the PSMCs, we found that PTBPI overexpression inhibited the stability of ILK mRNA, and PTBPI silencing can stabilize ILK's mRNA (Figure 4E). These results were further confirmed by Western blotting and immunofluorescence (Figure 4F). The FISH analysis demonstrated the colocalization of PTBPI and ILK, which verified the interaction of PTBPI and ILK (Figure 4G).

PTBPI Silencing Modulates the Expression of Phenotypic Transition Related Genes

Next, based on the above results, we detected the expression of phenotypic transition-related proteins. The results showed that ILK and osteopontin expression was significantly increased in PSMCs after hypoxia while ILK, myocardin, SM α -actin and Calponin were downregulated. However, this was reversed by PTBPI knockdown (Figure 5A and B).

PTBPI Silencing Alleviated the Pulmonary Hypertension of Rats

In order to further confirm the effect of PTBPI in pulmonary hypertension, we carried out in vivo study ($n=6$). Adeno virus silencing PTBPI was injected into rats via tail vein. We found that compared to the control group,

RV/(LV + S)%, mPAP, WT%, and WA% increased 4 weeks after establishing the model, and PTBPI silencing notably reduced these index compared to the Ad-sh-nc treatment group (Figure 6A–D).

Discussion

Pulmonary hypertension (PH) is a group of diseases characterized by a progressive increase in the circulation resistance of the pulmonary artery system.^{14–17} Its pathological changes include pulmonary vasoconstriction and remodeling, abnormal proliferation of pulmonary vascular smooth muscle and endothelial cells. However, the cause of hypertension causes is still unclear. Several reasons could be involved in pulmonary hypertension diagnosis difficulty. Therefore, it is crucial to improve the understanding of pulmonary hypertension, effective early screening, and early treatment.^{18–20}

PASMCs exist in two forms: synthetic phenotype and contractile phenotype. The contractile phenotype of PASMCs is rich in Myosin and α -actin, which are the main components of contractile proteins.^{21–23} Therefore, observing the expression of α -actin in PASMCs is an important method to determine the phenotypic transformation of cells. Previous reports showed that after 4 weeks of hypoxia, the content of α -actin in the cytoplasm of PASMCs is significantly reduced, and the degree of cell proliferation is increased. The previous results are consistent with the results of pulmonary hypertension modeling in the literature, providing a basis for better follow-up research.

PASMCs are the main components of the pulmonary artery vascular media, and play a significant role in pulmonary vascular remodeling. The abnormal proliferation of PASMCs manifested as the contraction, proliferation, differentiation, and matrix secretion of smooth muscle cells, which caused the thickening of the pulmonary artery




Figure 6 PTBP1 silencing alleviated pulmonary hypertension in rats. (A–D) After administration of Ad-sh-PTBP1 or Ad-sh-nc, RV/(LV + S)%, mPAP, WT% and WA% of the pulmonary hypertension or control rats were detected. $n=6$, $**p<0.01$.

vascular medial membrane. A variety of factors (such as hypoxia, inflammatory response, abnormal mechanical forces in blood vessels) induce PASMCs proliferation. It has been reported that PTBP1 knockout in glioma cells attenuates cell proliferation, and migration, and enhances cell adhesion via regulating the alternative splicing of the transmembrane factor RNT4.²⁴ PTBP1 has been indicated to mediate the miR-124's effect on regulating the endothelial cell glycolysis in pulmonary arterial hypertension. In PH, miR-124, through the alternative splicing factor PTBP1, regulates the PKM2/PKM1 ratio to modulate the proliferative, and inflammatory state of cells.

In the present study, we confirmed that PTBP1 was increased significantly in PASMC after hypoxia which

indicate the critical role of PTBP1 in PH progression. PTBP1 knockdown inhibited the phenotypic transition of PASMC. As PTBP1 is a RNA binding protein, to elucidate the mechanism of PTBP1, we performed RIP and mass spectrum assay. ILK was found to be bound with PTBP1. ILK was first discovered by Hannigan et al⁷. ILK is a central regulatory protein of cell signaling; it also depends on phosphatidylinositol-3 kinase (PI3K) activation, which in turn acts on the downstream GSK-3 β . Shen et al²⁵ reveal that ILK is one of the necessary genes to maintain the aortic vasoconstriction phenotype. ILK has a wide range of regulatory roles in various pathophysiological processes, such as cell differentiation, development, and tumor growth. In this study, PTBP1 bound with ILK

and inhibited the mRNA stability of the ILK, thus lead to the dysregulation of ILK. Moreover, ILK was involved in the modulation of phenotypic transition-related proteins including myocardin, SM α -actin, Calponin and osteopontin. However, more investigations such as rescue experiments will make this conclusion more credible.

Conclusion

This study found that PTBP1 was overexpressed in PSMCs after hypoxia. Silencing of PTBP1 inhibited the transformation of PSMCs by regulating ILK and its downstream signaling. This study suggests that PTBP1 is involved in regulating the occurrence and development of PH and may become a new biomarker for the diagnosis and prognosis of pulmonary hypertension.

Acknowledgment

This study is supported by National Natural Science Foundation of China (Research Grant no. 81600227).

Author Contributions

All authors made substantial contributions to the design and conception of the study, and acquisition, analysis and interpretation of data, and took part in either drafting or revising the manuscript. All authors gave final approval of the version to be published, have agreed on the journal to which the article has been submitted, and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Disclosure

There is no interest conflict of all the authors.

References

- Zhao L, Mason NA, Morrell NW, et al. Sildenafil inhibits hypoxia-induced pulmonary hypertension. *Circulation*. 2001;104(4):424–428. doi:10.1161/hc2901.093117
- Olschewski H, Simonneau G, Galie N, et al. Inhaled iloprost for severe pulmonary hypertension. *N Engl J Med*. 2002;11(6):19.
- Walsh-Sukys MC, Tyson JE, Wright LL, Bauer CR. Persistent pulmonary hypertension of the newborn in the era before nitric oxide: practice variation and outcomes. *Pediatrics*. 2000;105(1 Pt 1):14–20. doi:10.1542/peds.105.1.14
- Olschewski H, Ghofrani HA, Walmrath D, et al. Inhaled prostacyclin and iloprost in severe pulmonary hypertension secondary to lung fibrosis. *Am J Respir Crit Care Med*. 1999;160(2):600–607. doi:10.1164/ajrccm.160.2.9810008
- Kornbliht AR, Schor IE, Alló M, Dujardin G, Petrillo E, Muñoz MJ. Alternative splicing: a pivotal step between eukaryotic transcription and translation. *Nat Rev Mol Cell Biol*. 2013;14(3):153–165. doi:10.1038/nrm3525

- Christofk HR, Vander Heiden MG, Harris MH, et al. The M2 splice isoform of pyruvate kinase is important for cancer metabolism and tumour growth. *Nature*. 2008;452(7184):230–233. doi:10.1038/nature06734
- Hannigan GE, Leunghagesteijn C, Fitzgibbon L, et al. Regulation of cell adhesion and anchorage-dependent growth by a new β 1-integrin-linked protein kinase. *Nature*. 1996;379(6560):91–96. doi:10.1038/379091a0
- Jie W, Guo J, Shen Z, et al. Contribution of myocardin in the hypoxia-induced phenotypic switching of rat pulmonary arterial smooth muscle cells. *Exp Mol Pathol*. 2010;89(3):301–306. doi:10.1016/j.yexmp.2010.06.010
- Chettimada S, Gupte R, Rawat D, Gebb SA, Gupte SA, Physiology M. Hypoxia-induced glucose-6-phosphate dehydrogenase overexpression and activation in pulmonary artery smooth muscle cells: implication in pulmonary hypertension. *Am J Physiol Lung Cell Mol Physiol*. 2014;308(3):ajplung.00229.02014. doi:10.1152/ajplung.00229.2014
- Meng J, Ai X, Lei Y, et al. USP5 promotes epithelial-mesenchymal transition by stabilizing SLUG in hepatocellular carcinoma. *Theranostics*. 2019;9(2):573–587. doi:10.7150/thno.27654
- Xi X, Liu N, Wang Q, et al. ACT001, a novel PAI-1 inhibitor, exerts synergistic effects in combination with cisplatin by inhibiting PI3K/AKT pathway in glioma. *Cell Death Dis Oct*. 2019;10(10):757. doi:10.1038/s41419-019-1986-2
- Zhong W, Yang W, Qin Y, et al. 6-Gingerol stabilized the p-VEGFR2/VE-cadherin/beta-catenin/actin complex promotes microvessel normalization and suppresses tumor progression. *J Exp Clin Cancer Res*. 2019;38(1):285. doi:10.1186/s13046-019-1291-z
- Zhong W, Sun B, Gao W, et al. Salvianolic acid A targeting the transgelin-actin complex to enhance vasoconstriction. *E Bio Med*. 2018;37:246–258. doi:10.1016/j.ebiom.2018.10.041
- Rich S, Dantzker DR, Ayres SM, Bergofsky EH, Koerner SK. Primary pulmonary hypertension: a national prospective study. *Ann Intern Med*. 1987;107(2):216–223. doi:10.7326/0003-4819-107-2-216
- Park YM, Chung WJ, Lee SP, et al. Efficacy of inhaled iloprost in cor pulmonale and severe pulmonary hypertension associated with tuberculous destroyed lung. *J Cardiovasc Ultrasound*. 2014;22(2):95–97. doi:10.4250/jcu.2014.22.2.95
- McLaughlin VV, Shillington A, Rich S. Survival in primary pulmonary hypertension: the impact of epoprostenol therapy. *Circulation*. 2002;106(12):1477–1482. doi:10.1161/01.CIR.0000029100.82385.58
- Herbay AV, Illes A, Waldherr R, Otto HF. Pulmonary tumor thrombotic microangiopathy with pulmonary hypertension. *Cancer*. 1990;66(3):587–592. doi:10.1002/1097-0142(19900801)66:3<587::AID-CNCR2820660330>3.0.CO;2-J
- Galie N, Hoeper MM, Humbert M, et al. Guidelines for the diagnosis and treatment of pulmonary hypertension: the Task Force for the Diagnosis and Treatment of Pulmonary Hypertension of the European Society of Cardiology (ESC) and the European Respiratory Society (ERS), endorsed by the Internati. *Eur Heart J*. 2015;37(1):67–119. doi:10.1093/eurheartj/ehv317
- D'Alonzo GE, Barst RJ, Ayres SM, et al. Survival in patients with primary pulmonary hypertension: results from a national prospective registry. *Ann Intern Med*. 1991;115(5):343–349. doi:10.7326/0003-4819-115-5-343
- Fishman AP. Clinical classification of pulmonary hypertension. *Clin Chest Med*. 2001;22(3):385–391. doi:10.1016/S0272-5231(05)70278-1
- Abe K, Shimokawa H, Morikawa K, et al. Long-term treatment with a Rho-kinase inhibitor improves monocrotaline-induced fatal pulmonary hypertension in rats. *Circ Res*. 2004;94(3):385–393. doi:10.1161/01.RES.0000111804.34509.94
- Baber SR, Deng W, Master RG, et al. Intratracheal mesenchymal stem cell administration attenuates monocrotaline-induced pulmonary hypertension and endothelial dysfunction. *Am J Physiol Heart Circ Physiol*. 2007;292(2):H1120–H1128. doi:10.1152/ajpheart.00173.2006

23. Rosenberg HC, Rabinovitch M. Endothelial injury and vascular reactivity in monocrotaline pulmonary hypertension. *Am J Physiol.* 1988;255(6 Pt 2):H1484–H1491. doi:10.1152/ajpheart.1988.255.6.H1484
24. Cobbold LC, Wilson LA, Sawicka K, et al. Upregulated c-myc expression in multiple myeloma by internal ribosome entry results from increased interactions with and expression of PTB-1 and YB-1. *Oncogene.* 2010;29(19):2884–2891. doi:10.1038/onc.2010.31
25. Shen D, Li J, Lepore JJ, Anderson T. Aortic aneurysm generation in mice with targeted deletion of integrin-linked kinase in vascular smooth muscle cells. *Circ Res.* 2011;109(6):616–628. doi:10.1161/CIRCRESAHA.110.239343

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>