

Novel Anti-Tubulin Compounds from *Trigonella foenum-graecum* Seeds; Insights into In-vitro and Molecular Docking Studies

Sahar Saleh Alghamdi^{1,2}
Rasha Saad Suliman^{1,2}
Amjad Sulaiman Alsaeed¹
Khlood Khaled Almutairi¹
Norah Abdulaziz Aljammaz¹
Abdulelah Altolayyan²
Rizwan Ali²
Alshaimaa Alhallaj²

¹College of Pharmacy, King Saud bin Abdulaziz University for Health Sciences, Riyadh, Kingdom of Saudi Arabia;

²Medical Research Core Facility and Platforms, King Abdullah International Medical Research Center (KAIMRC), Ministry of National Guard Health Affairs, Riyadh, Kingdom of Saudi Arabia

Background: Fenugreek, also known as *Trigonella foenum-graecum* L, is a natural plant that belongs to the *Fabaceae* family and has been known as a promising source of bioactive compounds. It has been widely used as traditional medicine since it has shown to lower blood glucose, manage cholesterol levels and further aid in the prevention and treatment of cancer. Herein, we aim to evaluate the anticancer activity of methanolic fenugreek seed extract against several cancer cell lines.

Methods: We sought to investigate the phytochemical classes present in multiple fenugreek seeds extracts using HPLC-DAD followed by LC/MS, predict and investigate anticancer activity using PASS online webserver, the CellTiter-Glo assay, evaluate ADME properties, and perform molecular docking for all bioactive compounds via Maestro software.

Results: Multiple extracts exhibited distinct phytochemical classes that demonstrated different biological activities. Fenugreek methanolic extract contains flavonoid chemical class, which showed the highest anticancer activity against the HCT8 cell line of colorectal cancer (IC₅₀ of 8.83 µg/mL), followed by KAIMRC1 breast cancer cell line (IC₅₀ of 35.06 µg/mL), HL60 leukemia cell line (37.80 µg/mL), MDA-MB-231 breast cancer cell line (38.51 µg/mL), and lastly, HCT116 colorectal cancer cell line with IC₅₀ of 56.03 µg/mL. In contrast, the chloroform extract was inactive. The molecular docking study for all the bioactive compounds suggested that flavonoids F6 (−9.713 and −12.132), F7 (−10.166 and −12.411), and F11 (−10.084 and −13.516) possess the highest docking scores through SP and XP scores, respectively.

Conclusion: The obtained results confirm that the bioactive compounds present in fenugreek seeds exhibit anticancer activity against several cancer cells that can mediate via tubulin polymerization inhibition. Although our study has evaluated the anticancer potential of *Trigonella foenum-graecum* as a promising natural source for new anticancer agents, fenugreek biological activity needs further research and investigations on their mechanism of action and toxicity profile.

Keywords: fenugreek *Trigonella foenum-graecum*, tubulin inhibitors, computational pharmacokinetics ADME, QTOF-LC/Ms metabolites


Introduction

Cancer is recognized as a significant public health concern and is expected to rise as a result of changes in human lifestyles.¹ Despite the variety of anticancer agents available nowadays, these agents only provide limited therapeutic benefits due to undesirable side effects and the development of chemoresistance.² In the last few years, researchers have emphasized the need to discover novel chemotherapeutic drugs from natural resources, as the usage of herbs has provided a new therapeutic

Correspondence: Sahar Saleh Alghamdi
Tel +966 114299999 ext 99516
Email ghamdisa@ksau-hs.edu.sa



Graphical Abstract



approach in a variety of diseases. Moreover, it has been reported that the general community is using herbal products to complement or replace modern therapeutic medicine due to their potential safety profile and promising activity.^{3–6}

Fenugreek, known as *Trigonella foenum-graecum*, or Hilbeh in the Middle East, belongs to the *Fabaceae* family.⁷ The extract of fenugreek contains numerous phytochemical classes, including flavonoids, which are recognized as antioxidants that decrease oxidative stress.⁸ In addition, phytosterol diosgenin, a steroidal saponin with potential biological activities such as glucose preservation and lipid homeostasis, has a cytotoxic effect on lung cancer cells and an inhibitory effect of the hTERT gene. It also exhibits a potential anticarcinogenic activity toward breast cancer tissues due to its strong antioxidant effect.^{9–12} Moreover, the extract includes the soluble fiber galactomannan, which enhances blood glucose and lowers serum lipid in diabetic animal models, besides 4-Hydroxyisoleucine that stimulates insulin secretion and that has insulin-mimetic properties.^{13,14}

Hence, a wide variety of fenugreek seeds medical applications are well known, including dysmenorrhea, alleviate painful uterine contractions during menstruation, diabetes, wound healing, and stimulation of milk

production.¹⁵ Furthermore, several in-vivo and in-vitro studies have reported the promising anticancer activity of fenugreek extracts with the advantage of selective toxicity to cancer cells.¹⁶

Chatterjee et al studied the *Trigonella foenum-graecum* seeds extract on skin papilloma tumors. The authors found a remarkable reduction in tumor incidence compared to a control group by inhibiting the tumour necrosis factor (TNF) gene expression.¹⁷ Likewise, *Trigonella foenum-graecum* crude methanol extracts caused apoptosis induction in hepatocellular carcinoma (HepG2) using MTT assay.¹⁸ Another study examined the anticancer activities of fenugreek seed extract on MCF-7 breast cancer, HCAM liver cancer, and non-cancerous Vero cell lines and revealed that these seeds diminished cancer cell proliferation within the MCF-7 cell line. Unfortunately, no apoptosis or necrosis was observed.¹⁹

In a recent study, *T. foenum-graecum* extracts were tested on cancer cell lines, including Hep2, MCF7, and normal cell line HCEC. The methanolic extract showed considerable cytotoxicity to Hep2 and MCF7 cell lines, with IC₅₀ values ranging from 2.85 to 3.14 g/L, which was higher than the cytotoxicity of reference drugs Paclitaxel and Tamoxifen. Importantly, these extracts demonstrated

no cytotoxicity against normal cells, indicating that they were selectively cytotoxic to cancer cells.²⁰

Microtubule, a protein involved in the critical functions of several cells in the human body, has been proven to be a target in the development of novel anticancer agents.²¹ Microtubules are formed when α - and β -tubulin heterodimers polymerize to build a tube-shaped structure, which is then depolymerized in a dynamically balanced cycle.²² Since various isotypes of tubulin are expressed in cancerous cells, inhibiting these microtubules can interfere with neoplastic cell cycles and induce apoptosis. Indeed, multiple anti-tubulin agents are naturally derived from plants, such as Taxanes and Vinca Alkaloids. These agents impair cell division by disrupting the equilibrium state of microtubules via inhibiting polymerization or the depolymerization process.²³ Similarly, Nagireddy et al found that Noscopine analogs, derived from the *Papaver somniferum*, can inhibit tubulin protein and were considered a potential chemotherapeutic candidate for the treatment of pancreatic cancer. Furthermore, the presence of bromine in compounds can suppress cell growth at the G2/M phase of MCF-7 cells and induce apoptosis.²⁴

Thus, the main aim of this study is to obtain an appropriate extraction method of fenugreek seeds and to evaluate the anticancer activity against various cancer cell lines. We also predict the pharmacokinetic properties and potential targets for the bioactive constituents that mediate the anticancer activity of fenugreek seed extract.

Materials and Methods

Fenugreek Seed Extraction Protocols

Seeds Collection

Seeds were purchased from a certified company called Mountain Rose Herbs. The purchased seeds are carefully selected to be organic, without additives, and free of preservatives. Moreover, the seeds undergo more than 10 quality control tests. The seeds Lot number is 24330-R. The seeds have an authentication certificate and certificate of analysis. The local herbs shops were avoided because of the high chance of adulterants.

Seeds Extraction

First, we followed the standard method of fenugreek extraction using the Soxhlet apparatus. 250 g of the seeds was grounded by a blender to semi-grounded seeds. An Erlenmeyer of 500 mL was used to weigh 75 g of the grounded seeds. The seeds were placed in 3-filter tube each contains 75 g of grounded seeds to utilize it in

the Soxhlet apparatus. Methanol was used as an extraction solvent, the volume was used as recommended by the manufacturer, and a 50-mL volume of solvent was introduced into the extractor thimble from the top to facilitate flushing of the extractor chamber. After the extraction process has been done, the extraction was dried by rotary evaporator apparatus to get a powder form of the extraction. Further extraction has been done using other extract solvents by the maceration method. A 100 g of powder fenugreek was soaked with 1 liter of each solvent as follows: chloroform, water, diethyl ether, and methanol. The mixture was kept soaked under a magnetic stirrer hot plate at 40°C for 48 h. Then, the extract was collected and filtered using filter paper and bacterial filter. A rotary evaporator was used for the drying process, and all the extracts were stored in a 4°C refrigerator to be further used.

Chemical Identification

The extract obtained above was subjected to HPLC-DAD followed by LC/MS using the following protocol.

Chemicals

Formic acid and HPLC grade methanol were purchased from Sigma-Aldrich (St. Louis, MO, USA) and Honeywell (France), respectively.

RP-HPLC Method

All extracts were first injected into the Agilent 1260 Infinity HPLC system (Agilent, Germany) with Diode-Array Detection (DAD) detector. The separation carried out in a reversed-phase mode using Phenomenex Kinetex-C18 column (4.6 mm \times 250 mm, 5 μ m) with the following elution gradient methanol to ultrapure water; 0–1 min, 5% B; 1–11 min, 5–100% B; 11–13 min, 95% B; 13–15 min, 5% B; 15–16 min, 5% B using mobile phase A (0.1% HCOOH in water) and mobile phase B (0.1% HCOOH in Methanol). Samples were injected with 10 μ L, with 16 min run time, and the flow rate was set as 1 mL/min. The DAD collected UV spectrum at 200, 225, 250, 275, 300, 325, and 350. The data were processed using ChemStation software.

LC-QTOF-MS Method

The analysis of all extracts was performed on Agilent 1260 Infinity HPLC system (Agilent, Germany) coupled to Agilent 6530 Quadrupole Time of Flight (Agilent, Singapore). Separation was performed using Agilent Extend-C18 column (2.1 mm \times 50 mm, 1.8 μ m) with the following elution

gradient; 0–1 min, 5% B; 1–11 min, 5–100% B; 11–13 min, 95% B; 13–15 min, 5% B; 15–16 min, 5% B using mobile phase A (0.1% HCOOH in water) and mobile phase B (0.1% HCOOH in Methanol). The injection volume was 10 μ L, and the flow rate was set as 300 μ L/min. The MS1 acquisition method was achieved in positive mode with mass ranged from 100 to 600 m/z. The mass spectrometer parameters were set as follows: Gas Temperature = 300°C; Gas flow = 8 l/min; Nebulizer = 35 psig; SheathGas Temperature = 350 and SheathGas flow was 11. MS1 data was generated using the Agilent MassHunter qualitative analysis software.

In-vitro Anticancer Activity

The anticancer biological experiment was conducted at King Abdullah International Medical Research Center (KAIMRC), Riyadh, Saudi Arabia. To determine the effect of the fenugreek extracts on cell proliferation, MCF-7, MDA MB-231, KAIMRC1 (breast cancer cell lines), HCT8, and HCT116 (colorectal cancer cell lines) were used. All the cell lines were purchased from ATCC, USA, except KAIMRC1, which was isolated and established in the core laboratory facility KAIMRC, Riyadh KSA.²⁵ All the cell lines were maintained in advanced DMEM containing 10% FBS, 1% L-glutamine, and 1% antibiotics (Pen-Strep). Cells were plated on flat-bottom white 96-well plates at a density of 5×10^3 cells/well in a 100 μ L growth medium. Serial dilutions of the extracts, ranging from 100 to 0.01 μ g/100 μ L cell culture media were made in triplicates and transferred to the cell culture plates containing the cells. Additional rows with only cells were added to account for the compounds and cells' effect. Cells were incubated for 48 h at 37°C with 5% CO₂. Cell viability was determined using the CellTiter-Glo assay (Promega) according to the manufacturer's recommendations. Luminescence was measured using the EnVision plate reader (Perkin Elmer), and readings were normalized to averaged DMSO controls and expressed as a relative percentage. Mitoxantrone was used as a positive control, data were analyzed with GraphPad Prism 8.1 software, and the half-maximal inhibitory concentration (IC₅₀) was determined.

ADME Using SwissADME Webserver

The calculations of the absorption, distribution, metabolism, and elimination (ADME) properties of the fenugreek active compounds were performed using the SwissADME web server.²⁶ The SMILES of each active compound were used as input to generate the ADME data, and the results

were then compared and analyzed according to the reported literature values.

Prediction of Biological Activity Using PASS Online

The flavonoid-derived compounds from fenugreek methanolic extract were analyzed using the online service Prediction of Activity Spectra for Substances (PASS) to determine the biological probability of a substance being active (Pa) or inactive (Pi).²⁷

Molecular Docking Study on Tubulin Protein and Ligands Preparation

The 3D crystal structure of Tubulin (PDB: 4O2B) was obtained from the RCSB protein data bank, and the structure was prepared using the Protein Preparation Wizard tool (PrepWizard, Schrödinger Release 2021-1). The 3D Tubulin structure was refined, minimized, and optimized with the OPLS4 force field. Unnecessary water molecules, substrates, ions, and other subunits were removed and the β -Tubulin subunit that contains the Colchicine binding site was maintained for the docking study. The 2D structures for ligands (F1-11, S1-3, D1-4, and Colchicine) were prepared using Schrödinger's LigPrep tool, and several conformations were generated, optimized, and minimized for their lowest energy conformation.

Ligand-Protein Molecular Docking

A one-step docking tool was utilized to validate the docking modes, and the root-mean-square deviation (RMSD) values were below 0.366 Å for both standard precision (SP) and extra precision (XP) scoring functions of GLIDE. All the compounds were docking using SP and XP scoring functions, and post-docking analysis for the docked poses was performed. Moreover, the energies or the relative binding affinity for the best docking (complex) structures were carried out using Prime MM-GBSA in which VSGB was used as a solvation model and OPLS4 force field.

QikProp for Absorption, Distribution, Metabolism, and Excretion (ADME) Predictions

For all the ligands, the ADME parameters were calculated and computed using the QikProp tool, Schrödinger, LLC, New York, NY, 2021. More than twenty physical descriptors were generated, and the most important parameters for drug development were chosen for further analysis.

Results

Chemical Analysis and Compound Identifications in the LC/MS Based on the Solvent Used

Soxhlet Methanolic Extract

The extracted yield obtained was as follows: Soxhlet methanolic extract 68% of the started weight, chloroform extract 45%, water extract 73%, diethyl ether extract 67%, and methanol extract 84%. The obtained extracts were injected into LC/MS, and the peaks total ion chromatogram, which shows high abundance, was examined as shown in Figure 1. QTOF-LCMS is one of the recent methods that could be used to identify metabolites, since we are handling the same seed that was analysed previously and identified, and their chemical structure was illustrated using NMR and chromatography techniques. It is valid to tentatively map the specific mass value with the previously reported compounds isolated from the same seed.

Chloroform Extract

The obtained chloroform extract was injected into LC/MS, and the peaks total ion chromatogram, which shows high abundance, was examined as shown in Figure 2.

Water Extract

The obtained water extract was injected into LC/MS, and the peaks total ion chromatogram, which shows high abundance, was examined as shown in Figure 3.

Diethyl Ether Extract

The obtained water extract was injected into LC/MS, and the peaks total ion chromatogram, which shows high abundance, was examined as shown in (Figure 4).

Predicted Biological Activity of Bioactive Compounds Using PASS Online

The metabolic profiling results showed that the fenugreek seed extracts exhibited different phytochemical classes depending on the solvent used. Our results showed that




Figure 1 The LC/MS spectrum for soxhlet methanolic extract.




Figure 2 The LC/MS spectrum for chloroform extract.

methanol extract was rich in flavonoids, while the chloroform extract was rich with Phytosterol Diosgenin (Spirostan) derivatives. The Prediction of Activity Spectra for Substances (PASS) online service was used to assess the potential pharmacological activities of all the three different phytochemical classes that are present in different extracts.²⁷ As a result, compounds F6, F7, and F11 exhibited the highest probability of being active as anticarcinogenic and antineoplastic relative to the other compounds present in fenugreek extracts, as summarized in Table 1.

In-vitro AntiCancer Activity of Fenugreek Extracts Against Several Cancer Cell Lines

To confirm the predicted anticancer activity of fenugreek phytochemical compounds, the methanol and chloroform fenugreek extracts were examined against Leukemia cancer cell lines (HL60 and K562), breast cancer cell lines

(KAIMRC1 and MDA-MB-231), colorectal cancer cell lines (HCT8 and HCT116), and also against normal primary fibroblasts P1 and normal blood cells N1 to determine the selectivity of these chemical classes. Methanol extract exhibited a promising anticancer activity on all the cell lines except the K562 Leukemia cell, as plotted in Figure 5. The highest inhibitory activity of the methanol extract was against the HCT8 cell line of colorectal cancer with an IC_{50} of 8.83 $\mu\text{g/mL}$, followed by KAIMRC1 breast cancer cell line, HL60 leukemia cell line, MDA-MB-231 breast cancer cell line, and lastly HCT116 colorectal cancer cell line, with IC_{50} of 35.06, 37.80, 38.51, and 56.03 $\mu\text{g/mL}$, respectively (presented in Table 2). On the contrary, the chloroform extract was inactive on all cell lines, as seen in Figure 6.

ADME Properties of Fenugreek Extracts Bioactive Compounds

The selection of the ADME properties that were assessed in this study was based on Lipinski's rule of five (ROF)




Figure 3 The LC/MS spectrum for water extract.




Figure 4 The LC/MS spectrum for diethyl ether extract.

that estimates the drug-likeness properties of tested compounds.^{28,29} The pharmacokinetic properties of flavonoid derivatives (F1-F11) were within the acceptable range <500 Daltons in the molecular weight, except for F9 and

F10 that were out of range, suggesting the bulkiness of these two compounds (Table 3, Figure 7).²⁸ Flavonoid derivatives F6 to F11 showed greater than 5 in the number of H-bond donors, whereas F9 to F11 were out of range in

Table 1 Biological Activity Predictions Using PASS Online Webservice

Compounds Codes	Type of Activity	Pa	Pi
F6	Anticarcinogenic	0.926	0.002
	Antioxidant	0.831	0.003
	Antineoplastic	0.828	0.009
F7	Anticarcinogenic	0.856	0.004
	Antioxidant	0.802	0.003
	Antineoplastic	0.800	0.012
F11	Anticarcinogenic	0.943	0.002
	Antioxidant	0.915	0.003
	Antineoplastic	0.854	0.007

the number of H-bond acceptors, which consequently would affect the blood-brain barrier (BBB) permeability and GI absorption. Moreover, all the flavonoid derivatives (F1-F11) showed low lipophilicity and high solubility profile that affected the BBB penetration and oral activity. Furthermore, it was predicted that F1 to F8 could potentially have oral activities; however, F9 to F11 are orally inactive due to violation of two or more of Lipinski's rules.

The pharmacokinetic properties of diosgenin derivatives were computed (Table 3, Figure 7), and all the four derivatives were within the acceptable range <500 Daltons in the molecular weight and hydrogen bond donors and acceptors requirements. The computed lipophilicity (Log $P_{o/w}$) and solubility (LogS) parameters of diosgenin derivatives were higher

compared to the optimal range except for D4, suggesting these compounds D1 to D3 are highly lipophilic and thus poorly soluble.^{30,31} Additionally, most of the compounds were permeable to the BBB except for D2, and all the four derivatives demonstrated high GI absorption suggesting these diosgenin derivatives could be orally active, with no violation of two or more of Lipinski's rules.

Moreover, the ADME predictions (Table 3, Figure 7) of polyphenol stilbenes derivatives were within the acceptable range <500 Daltons in the molecular weight and hydrogen bond donors and acceptors except for S1 that demonstrated H-bond donors of more than 5. All three derivatives showed good lipophilicity (Log $P_{o/w}$) and solubility (LogS) values that were within the recommended range. Also, ADME results showed that S3 could penetrate the BBB, and this might be due to the balance between the solubility and lipophilicity values; however, the other two stilbenes S1 and S2 derivatives could not penetrate BBB due to their low lipophilicity and high solubility.³² Additionally, S2 and S3 showed high GI absorption, whereas S1 showed low GI absorption that correlates with H-bond donor ability that affects the absorption. Overall, all the Stilbenes derivatives were predicted to be orally active due to no violation of two or more of Lipinski's rules.

Docking of Fenugreek Bioactive Chemical Classes into Tubulin Crystal Structure

In this study, we sought to determine the potential mechanism that mediates the anticancer activity of the fenugreek


**Figure 5** The inhibitory activity of fenugreek methanol extract against six cancer cell lines.

Table 2 The IC₅₀ Values of Fenugreek Extracts Against Six Cancer Cell Lines

Cancer		Leukemia		Normal Primary Fibroblast	Normal Blood Sample	Breast Cancer		Colorectal Cancer	
Cell Lines		HL60	K562	PI	NI	KAIMRC1	MDA-MB-231	HCT8	HCT116
IC ₅₀ (µg/mL)	Fenugreek Methanol Extract	37.80	702.4	1888	17.38	35.06	38.51	8.831	56.03
	Fenugreek Chloroform Extract	165.3	557.7	1516	155.0	411.3	-	424.5	410.4

bioactive constituents. The colchicine (reference ligand) docking score with SP (-11.572) method was better compared to the XP scoring (-7.707) function and a similar observation was found with MM/GBSA binding energy in which SP (-88.10 Kcal mol⁻¹) was significantly higher compared to XP (-20.81 Kcal mol⁻¹) docking (Table 4). From the docked compounds, the best ligand was F7 that demonstrated a value of -10.166 and -12.411 through SP and XP docking methods, respectively. It is worth mentioning that ligands demonstrated higher docking scores with XP scoring function; however, this was not the case with the reference ligand (colchicine). Moreover, ligand F11 exhibited the third-best docking scores with values of -10.084 for SP and -13.516 for XP docking methods. Ligand F6 exhibited the fourth-best docking values with -9.713 and -12.132 , respectively, through SP and XP scoring functions. In the case of MM/GBSA binding

energy, the order of the ligands is different, as ligand F6 exhibits the highest binding energy, while F11 demonstrates the lowest with both SP and XP scoring methods. Of note, all the ligands were docked into β -Tubulin using SP and XP; however, our results showed that the flavonoid chemical class exhibit the highest docking scores with tubulin confirming the above-discussed results.

The post-docking analysis of the ligands with SP docking function exhibited several valuable molecular interactions. For instance, colchicine exhibits multiple hydrogen bond (H-bond) interactions with residues Cys241, Val238, Val181, and Thr353 (Figure 8). While XP docking interaction demonstrates less interaction while maintaining only one H-bond with Cys241 (Figure 9) that could explain the lower docking score compared to SP, and a similar trend was observed with the docked ligands. Moreover, multiple H-bonds interactions were demonstrated with the SP best


**Figure 6** The inhibitory activity of fenugreek chloroform extract against six cancer cell lines.

Table 3 The Pharmacokinetic ADME Properties of Fenugreek Bioactive Constituents

Compounds Codes	Molecular Weight	HB Donor	HB Acceptor	Log Po/w (WLOGP)	Log S (SILICOS-IT)	BBB Permeant	GI Absorption	Rule of Five (ROF)
Diosgenin I (D1)	414.62 g/mol	1	3	5.71 High lipophilic	-4.49 Poorly soluble	Yes	High	Yes; 1 violation: MLOGP>4.15
D2	414.62 g/mol	0	3	6.00 Highly lipophilic	-5.20 Moderately soluble	No	High	Yes; 1 violation: MLOGP>4.15
D3	412.60 g/mol	0	3	5.78 Highly lipophilic	-4.74 Moderately soluble	Yes	High	Yes; 1 violation: MLOGP>4.15
D4	446.62 g/mol	2	5	3.94	-3.56 Soluble	Yes	High	Yes; 0 violation
Stilbene I (S1)	420.41 g/mol	6	9	0.24	-1.71 Soluble	No	Low	Yes; 1 violation: NHorOH>5
S2	404.41 g/mol	5	8	0.53	-2.30 Soluble	No	High	Yes; 0 violation
S3	258.27 g/mol	3	4	2.76	-3.41 Soluble	Yes	High	Yes; 0 violation
Flavonoid I (F1)	286.24 g/mol	4	6	2.28	-3.82 Soluble	No	High	Yes; 0 violation
F2	270.24 g/mol	3	5	2.58	-4.40 Moderately soluble	No	High	Yes; 0 violation
F3	302.24 g/mol	5	7	1.99	-3.24 Soluble	No	High	Yes; 0 violation
F4	330.29 g/mol	3	7	2.59	-4.63 Moderately soluble	No	High	Yes; 0 violation
F5	286.24 g/mol	4	6	2.28	-3.82 Soluble	No	High	Yes; 0 violation
F6	432.38 g/mol	6	10	0.05	-2.69	No	Low	Yes; 1 violation: NHorOH>5
F7	432.38 g/mol	7	10	-0.23	-2.38 soluble	No	Low	Yes; 1 violation: NHorOH>5
F8	432.38 g/mol	7	10	-0.23	-2.38 soluble	No	Low	Yes; 1 violation: NHorOH>5
F9	564.49 g/mol	10	14	-2.40	-0.71 soluble	No	Low	No; 3 violations: MW>500, NorO>10, NHorOH>5
F10	594.52 g/mol	11	15	-3.04	-0.27 soluble	No	Low	No; 3 violations: MW>500, NorO>10, NHorOH>5
F11	448.38 g/mol	7	11	0.49	-2.08 Soluble	No	Low	No; 2 violations: NorO>10, NHorOH>5

docking pose of ligand F6 with Asn249, Gln11, Glu183, Asn101, and Thr179. Using XP docking, ligand F6 demonstrated additional two H-bonds interactions with Gln247 and Tyr224 residues. The highest SP docking score was attained by ligand F7 that showed H-bonds interactions with Asn101, Ser178, Lys254, Gln11, Gln247, and Val315. Using the XP scoring function, ligand F7 exhibited additional H-bonds with residues Cys241, Val238, Asn101, Asn249, Gln11, Gln247, and Glu183. Additionally, the H-bonds interactions

exhibited by ligand F11 within the binding site involve Cys241, Asn101, Ser178, Val181, and Asn350 through the SP docking method, while XP docking exhibited six H-bonds interactions with Cys241, Asn101, Lys254, Ser178, Val181, and Val315.

Discussion

In Soxhlet methanolic extract, the first spectrum screening at Rt (0.691–1.294 min): The peaks, which appear at Rt range




Figure 7 Log $P_{o/w}$ and LogS of fenugreek phytochemical classes.

Table 4 SP and XP Glide Scores for the Best Docked Compounds/Protein Complexes with MM/GBSA Binding Energy Calculations

Ligand	SP Docking			XP Docking		
	Glide Score	Glide Energy (Kcal mol ⁻¹)	MM/GBSA Binding Energy (Kcal mol ⁻¹)	Glide Score	Glide Energy (Kcal mol ⁻¹)	MM/GBSA Binding Energy (Kcal mol ⁻¹)
Colchicine	-11.572	-64.602	-88.10	-7.707	-40.594	-20.81
F6	-9.713	-65.402	-53.00	-12.132	-61.975	-52.68
F7	-10.166	-54.596	-43.56	-12.411	-50.704	-49.73
F11	-10.084	-54.034	-18.79	-13.516	-54.718	-12.25

(0.691–1.294 min), are correlated with the compounds Quercitrin (F11) and Vicenin-1 (F9), respectively,³³ F11 with [M+H]⁺ 448.101 and molecular formula of [C₂₁H₂₀O₁₁]⁺, and compound F9 with [M+H]⁺ 564.148 and molecular formula of [C₂₆H₂₈O₁₄]⁺. Second spectrum screening at Rt (4.712–5.110 min): The peaks, which appear at Rt range (4.712–5.110 min), are correlated with the compounds Afroside (F6),³⁴ F6 with [M+H]⁺ 432.106 and molecular formula of [C₂₁H₂₀O₁₀]⁺. Third spectrum screening at Rt (10.816–12.825 min): The peaks, which appear at Rt range (10.816–12.825 min), are correlated with the compounds Naringenin (F2) and Vicenin-2 (F10), respectively,³⁴ F2 with [M+H]⁺ 270.053 and molecular formula of [C₁₅H₁₀O₅]⁺, and compound F10 with [M+H]⁺ 594.158 and molecular formula of [C₂₇H₃₀O₁₅]⁺. Fourth spectrum screening at Rt (25.146–25.411 min): The peaks, which appear at Rt range (25.146–25.411 min), are associated with the compounds Quercetin (F3) and Tricin-2 (F4), respectively,³⁴ F2 with [M+H]⁺ 270.053 and molecular formula of [C₁₅H₁₀O₅]⁺, and compound F10 with [M+H]⁺ 594.158 and molecular formula of [C₂₇H₃₀O₁₅]⁺.


In chloroform extract, the first spectrum screening at Rt (10.348–12.645 min): The peaks that appear at Rt range (10.348–12.645 min) are correlated with the compounds Diosgenin-1 (D1),³³ D1 with [M+H]⁺ 414.621 and molecular formula of [C₂₇H₄₂O₃]⁺. Second spectrum screening at Rt (25.835–26.934 min): The peaks that appear at Rt range (25.835–26.934 min) are related with the compounds (25R)-5x-Spirost-2-en-6-one (D3) and (25R)-2a,3az-Dihydroxy-5a-spirostan-6-one (D4), respectively,³³ Compound D3 with [M+H]⁺ 412.298 and a molecular formula of [C₂₇H₄₀O₃]⁺, and compound D4 with [M+H]⁺ 446.303 and a molecular formula of [C₂₇H₄₂O₅]⁺. Third spectrum screening at Rt (27.805–28.692 min): The peaks that appear at Rt range (27.805–28.692 min) are correlated with the compounds (25R)-3cx,5-Cyclo-5a-spirostan-6-one

(D2) Compound D2 with [M+H]⁺ 432.106 and a molecular formula of [C₂₁H₂₀O₁₀]⁺.³³


In water extract, the first spectrum screening at Rt (0.752–1.266 min): The peaks that appear at Rt range (0.752–1.266 min) are associated with the compounds Rhapontigenin (S3) and Afroside (F6), respectively,^{34,35} compound S3 with [M+H]⁺ 258.089 and molecular formula of [C₁₅H₁₄O₄]⁺, and compound F6 with [M+H]⁺ 432.106 and molecular formula of [C₂₁H₂₀O₁₀]⁺. Second spectrum screening at Rt (25.295–25.884 min): The peaks that appear at Rt range (25.295–25.884 min) are correlated with the compounds Desoxyrhaponticin (S2) and Vicenin-1 (F9), respectively,^{34,35} compound S2 with [M+H]⁺ 402.147 and a molecular formula of [C₂₁H₂₄O₈]⁺, and compound F9 with [M+H]⁺ 564.148 and a molecular formula of [C₂₆H₂₈O₁₄]⁺. Third spectrum screening at Rt (28.046–28.510 min): The peak that appears at Rt range (28.046–28.510 min) is correlated with the compound Rhaponticin (S1).³⁵ Compound S1 with [M+H]⁺ 420.3362 and a molecular formula of [C₂₁H₂₄O₉]⁺.

In diethyl ether, the first spectrum screening at Rt (25.084–25.766 min): The peaks that appear at Rt range (25.084–25.766 min) are related with the compounds Desoxyrhaponticin (S2) and Vicenin-1 (F9), respectively,^{34,35} compound S2 with [M+H]⁺ 402.147 and a molecular formula of [C₂₁H₂₄O₈]⁺, and compound F9 with [M+H]⁺ 564.148 and a molecular formula of [C₂₆H₂₈O₁₄]⁺. Second spectrum screening at Rt (25.146–25.411 min): The peaks that appear at Rt range (25.146–25.411 min) are correlated with the compounds Desoxyrhaponticin (S2) and Rhaponticin (S1), respectively,³⁵ compound S2 with [M+H]⁺ 402.147 and a molecular formula of [C₂₁H₂₄O₈]⁺, and compound S1 with [M+H]⁺ 420.3362 and a molecular formula of [C₂₁H₂₄O₉]⁺.


Using the PASS online webserver, the flavonoid class of compounds demonstrated the most promising anticancer and




Colchicine



F6



F7



F11

Figure 8 Ligand and protein molecular interactions for colchicine, F6, F7, and F11 with the β -Tubulin binding site using glide SP scoring tool.




Figure 9 Ligand and protein molecular interactions for colchicine, F6, F7, and F11 with the β -Tubulin binding site using glide XP scoring tool.

anti-neoplastic activities among others, while the polyphenol stilbenes and phytosterol diosgenin derivatives showed lower activity compared to flavonoids, suggesting the flavonoid class of compounds could be responsible for the anticancer activity of the fenugreek plant. Since flavonoids predominate in the methanolic extract, our in-vitro cytotoxicity findings appear to be consistent with previously reported studies that support its anticancer properties. For example, Torres et al found that THDF, a natural flavonoid, is a potent inhibitor of cytotoxic activity on human myeloid leukemia HL-60 and U937 cells.³⁶ Moreover, Centaureidin, a flavone derived from *Polymnia fruticosa*, has been shown to inhibit tubulin polymerization, making it one of the first known flavonoids with an antimetabolic effect.³⁷ Furthermore, Liu et al proposed that Casticin, a polymethoxyflavone derived from *Vitex rotundifolia*, reduced the viability of glioma cells U251, U87, and U373, through a variety of mechanisms, one of which is attenuating the polymerization of tubulin.³⁸ Similarly, Isoliquiritigenin, a flavonoid isolated from *liquorice*, has been shown to suppress proliferation and promote differentiation in glioma stem cells isolated from SHG44 human glioma cells.³⁹ These studies confirm our results and suggest that fenugreek methanolic extract possesses a promising anticancer activity that can mediate through inhibition of tubulin polymerization as reported in the literature. Meanwhile, chloroform extract demonstrated less anticancer activity, confirming the predicted activities using PASS online webserver.

Furthermore, we assessed and evaluated the absorption, distribution, metabolism, and elimination (ADME) properties of natural active compounds present in the fenugreek seeds extract. Understanding the ADME parameters for the tested compounds allows the design and development of highly effective therapeutic agents with good pharmaceutical properties.⁴⁰ Therefore, our results show that the most important ADME parameters that possess a crucial role in the drug discovery process are acceptable among the three different classes. Additionally, since several reported studies have suggested β -Tubulin could mediate beneficial flavonoid effects, a molecular docking study with β -Tubulin was carried out, and our docking results showed that flavonoids exhibit the highest docking scores with tubulin crystal structure, suggesting the cytotoxic activity of flavonoids could be mediated through inhibition of tubulin polymerization, which warrants further investigation.^{41–44}

Conclusion

This study demonstrates that the use of solvents has a significant impact on the obtained phytochemical class

present in the fenugreek seeds. Among the four various solvents used, the phytochemical class obtained with maceration using chloroform is steroidal skeleton diosgenin, while methanol solvent contained the flavonoids. Furthermore, the computational activity predictions and in-vitro anticancer results confirmed the promising anticancer activity of flavonoids against several cancer cell lines that can potentially mediate through inhibition of tubulin polymerization. Thus, fenugreek methanolic extract obtained from *Trigonella foenum-graecum* seeds exhibits anti-tubulin activity, and further research is needed to identify new chemotherapeutic agents with less harmful side effects.

Data Sharing Statement

The data associated with the paper are available, and under what conditions the data can be accessed.

Ethical Statement

Written consent was obtained from the donors under project # RC16/096 according to the Institutional Review Board (IRB) of King Abdullah International Medical Research Center (KAIMRC), Riyadh, Saudi Arabia. All the used cancer cell lines were purchased from authorized accredited commercial suppliers. The cell lines are patented and previously reported and published.

Acknowledgments

The authors want to express their sincerest gratitude to the College of Pharmacy (COP) at King Saud bin Abdulaziz University for Health Sciences (KSAU-HS) for their continued support and Dr. Mohamed Boudjelal for the generous help during the experimental work.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis, and interpretation, or in all these areas, have drafted or written, or substantially revised or critically reviewed the article, have agreed on the journal to which the article will be submitted, reviewed and agreed on all versions of the article before submission, during revision, the final version accepted for publication, and any significant changes introduced at the proofing stage, agreed to take responsibility and be accountable for the contents of the article.

Funding

The authors acknowledge financial support from King Abdullah International Medical Research Center (KAIMRC), Ministry of National Guard Health Affairs, Riyadh, Kingdom of Saudi Arabia; Grant # (SP19.271.R).

Disclosure

The authors declare no conflicts of interest in this work.

References

- Siegel RL, Miller KD, Jemal A. Cancer statistics, 2020. *CA Cancer J Clin*. 2020;70(1):7–30. doi:10.3322/caac.21590
- Pearce A, Haas M, Viney R, et al. Incidence and severity of self-reported chemotherapy side effects in routine care: a prospective cohort study. *PLoS One*. 2017;12(10):e0184360. doi:10.1371/journal.pone.0184360
- Azaizeh H, Saad B, Cooper E, Said O. Traditional Arabic and Islamic medicine, a re-emerging health aid. *Evid Based Complement Alternat Med*. 2010;7(4):419–424. doi:10.1093/ecam/nen039
- Saad B, Azaizeh H, Said O. Tradition and perspectives of Arab herbal medicine: a review. *Evid Based Complement Alternat Med*. 2005;2(4):475–479. doi:10.1093/ecam/neh133
- Omeish AF, Abbadi W, Ghanma IM, et al. Hospital-based study on the use of herbal medicine in patients with coronary artery disease in Jordan. *J Pak Med Assoc*. 2011;61(7):683–687.
- Tang J, Song X, Zhu M, Zhang J. Study on the pharmacokinetics drug-drug interaction potential of Glycyrrhiza uralensis, a traditional Chinese medicine, with lidocaine in rats. *Phytother Res*. 2009;23(5):603–607. doi:10.1002/ptr.2450
- Nagulapalli Venkata KC, Swaroop A, Bagchi D, Bishayee A. A small plant with big benefits: fenugreek (*Trigonella foenum-graecum* Linn.) for disease prevention and health promotion. *Mol Nutr Food Res*. 2017;61(6):1600950. doi:10.1002/mnfr.201600950
- Pietta PG. Flavonoids as antioxidants. *J Nat Prod*. 2000;63(7):1035–1042. doi:10.1021/np9904509
- Kaviarasan S, Vijayalakshmi K, Anuradha CV. Polyphenol-rich extract of fenugreek seeds protect erythrocytes from oxidative damage. *Plant Foods Hum Nutr*. 2004;59(4):143–147. doi:10.1007/s11130-004-0025-2
- Nair S, Nagar R, Gupta R. Antioxidant phenolics and flavonoids in common Indian foods. *J Assoc Physicians India*. 1998;46(8):708–710.
- Feki A, Jaballi I, Cherif B, et al. Therapeutic potential of polysaccharide extracted from fenugreek seeds against thiamethoxam-induced hepatotoxicity and genotoxicity in Wistar adult rats. *Toxicol Mech Methods*. 2019;29(5):355–367. doi:10.1080/15376516.2018.1564949
- Hannan JM, Ali L, Rokeya B, et al. Soluble dietary fibre fraction of *Trigonella foenum-graecum* (fenugreek) seed improves glucose homeostasis in animal models of type 1 and type 2 diabetes by delaying carbohydrate digestion and absorption, and enhancing insulin action. *Br J Nutr*. 2007;97(3):514–521. doi:10.1017/S0007114507657869
- Sauvaire Y, Petit P, Broca C, et al. 4-Hydroxyisoleucine: a novel amino acid potentiator of insulin secretion. *Diabetes*. 1998;47(2):206–210. doi:10.2337/diab.47.2.206
- Fuller S, Stephens JM. Diosgenin, 4-hydroxyisoleucine, and fiber from fenugreek: mechanisms of actions and potential effects on metabolic syndrome. *Adv Nutr*. 2015;6(2):189–197. doi:10.3945/an.114.007807
- Basch E, Ulbricht C, Kuo G, Szapary P, Smith M. Therapeutic applications of fenugreek. *Altern Med Rev*. 2003;8(1):20–27.
- Alsemari A, Alkhodairy F, Aldakan A, et al. The selective cytotoxic anti-cancer properties and proteomic analysis of *Trigonella foenum-graecum*. *BMC Complement Altern Med*. 2014;14:114. doi:10.1186/1472-6882-14-114
- Chatterjee S, Kumar M, Kumar A. Chemomodulatory effect of *Trigonella foenum graecum* (L.) seed extract on two stage mouse skin carcinogenesis. *Toxicol Int*. 2012;19(3):287–294. doi:10.4103/0971-6580.103670
- Khalil MI, Ibrahim MM, El-Gaaly GA, Sultan AS. *Trigonella foenum* (Fenugreek) induced apoptosis in hepatocellular carcinoma cell line, HepG2, mediated by upregulation of p53 and proliferating cell nuclear antigen. *Biomed Res Int*. 2015;2015:914645. doi:10.1155/2015/914645
- Al-Timimi LAN. Antibacterial and anticancer activities of fenugreek seed extract. *Asian Pac J Cancer Prev*. 2019;20(12):3771–3776. doi:10.31557/APJCP.2019.20.12.3771
- Ahmed SI, Hayat MQ, Zahid S, et al. Isolation and identification of flavonoids from anticancer and neuroprotective extracts of *Trigonella foenum-graecum*. *Tropical J Pharm Res*. 2017;16(6):1391–1398. doi:10.4314/tjpr.v16i6.25
- Binarová P, Tuszynski J. Tubulin: structure, functions and roles in disease. *Cells*. 2019;8(10):1294. doi:10.3390/cells8101294
- Field JJ, Díaz JF, Miller JH. The binding sites of microtubule-stabilizing agents. *Chem Biol*. 2013;20(3):301–315. doi:10.1016/j.chembiol.2013.01.014
- Naaz F, Haider MR, Shafi S, et al. Anti-tubulin agents of natural origin: targeting taxol, vinca, and colchicine binding domains. *Eur J Med Chem*. 2019;171:310–331. doi:10.1016/j.ejmech.2019.03.025
- Nagireddy PKR, Kommalapati VK, Siva Krishna V, Sriram D, Tangatur AD, Kantevari S. Imidazo[2,1-*b*]thiazole-coupled natural nospapine derivatives as anticancer agents. *ACS Omega*. 2019;4(21):19382–19398. doi:10.1021/acsomega.9b02789
- Ali R, Samman N, Al Zahrani H, et al. Isolation and characterization of a new naturally immortalized human breast carcinoma cell line, KAIMRC1. *BMC Cancer*. 2017;17(1):803. doi:10.1186/s12885-017-3812-5
- Daina A, Michielin O, Zoete V. SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. *Sci Rep*. 2017;7:42717. doi:10.1038/srep42717
- Filimonov D, Lagunin A, Glorizova T, et al. Prediction of the biological activity spectra of organic compounds using the PASS online web resource. *Chem Heterocycl Compd*. 2014;50(3):444–457.
- Lipinski CA. Lead- and drug-like compounds: the rule-of-five revolution. *Drug Discov Today Technol*. 2004;1(4):337–341. doi:10.1016/j.ddtec.2004.11.007
- Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. *Adv Drug Deliv Rev*. 2001;46(1–3):3–26. doi:10.1016/s0169-409x(00)00129-0
- Leo A, Hansch C, Elkins D. Partition coefficients and their uses. *Chem Rev*. 1971;71(6):525–616.
- Jorgensen WL, Duffy EM. Prediction of drug solubility from Monte Carlo simulations. *Bioorg Med Chem Lett*. 2000;10(11):1155–1158. doi:10.1016/s0960-894x(00)00172-4
- Ajay BGW, Murcko MA. Designing libraries with CNS activity. *J Med Chem*. 1999;42(24):4942–4951. doi:10.1021/jm990017w
- Melo PS, De Azevedo MB, Zullo MA, Fabrin-Neto JB, Haun M. Cytotoxicity of the phytosterol diosgenin and its derivatives in rat cultured hepatocytes and V79 fibroblasts. *Hum Exp Toxicol*. 2004;23(10):487–493. doi:10.1191/0960327104ht4740a

34. Skalsta H. Chemical constituents. In: Georgios AP, editor. *Fenugreek, The Genus Trigonella*. New York: Taylor & Francis Inc; 2002:132.
35. Li G, Luan G, He Y, et al. Polyphenol stilbenes from fenugreek (*Trigonella foenum-graecum*L.) seeds improve insulin sensitivity and mitochondrial function in 3T3-L1 adipocytes. *Oxid Med Cell Longev*. 2018;2018:7634362. doi:10.1155/2018/7634362
36. Torres F, Quintana J, Estévez F. 5,7,3'-trihydroxy-3,4'-dimethoxyflavone-induced cell death in human leukemia cells is dependent on caspases and activates the MAPK pathway. *Mol Carcinog*. 2010;49(5):464–475. doi:10.1002/mc.20619
37. Beutler JA, Cardellina JH II, Lin CM, et al. Centaureidin, a cytotoxic flavone from *Polymnia fruticosa*, inhibits tubulin polymerization. *Bioorg Med Chem Letters*. 1993;3(4):581–584.
38. Liu E, Kuang Y, He W, Xing X, Gu J. Casticin induces human glioma cell death through apoptosis and mitotic arrest. *Cell Physiol Biochem*. 2013;31(6):805–814.
39. Lin Y, Sun H, Dang Y, Li Z. Isoliquiritigenin inhibits the proliferation and induces the differentiation of human glioma stem cells. *Oncology Rep*. 2018;39(2):687–694.
40. Butina D, Segall MD, Frankcombe K. Predicting ADME properties in silico: methods and models. *Drug Discovery Today*. 2002;7(11):S83–S88.
41. Mukhtar E, Adhami VM, Sechi M, Mukhtar H. Dietary flavonoid fisetin binds to β -tubulin and disrupts microtubule dynamics in prostate cancer cells. *Cancer Letters*. 2015;367(2):173–183.
42. Torres F, Quintana J, Estévez F. 5, 7, 3'-trihydroxy-3, 4'-dimethoxyflavone inhibits the tubulin polymerization and activates the sphingomyelin pathway. *Mol Carcinog*. 2011;50(2):113–122.
43. Gupta K, Panda DJB. Perturbation of microtubule polymerization by quercetin through tubulin binding: a novel mechanism of its antiproliferative activity. *Biochemistry*. 2002;41(43):13029–13038.
44. Beutler JA, Hamel E, Vlietinck AJ, et al. Structure– activity requirements for flavone cytotoxicity and binding to tubulin. *J Med Chem*. 1998;41(13):2333–2338.

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also

been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>