



# Comparing Efficacy of Different Analgesic Modalities in Patients Undergoing Total Knee Arthroplasty [Response To Letter]

Rui Xiao <sup>1,\*</sup>, Li-Fang Liu<sup>1,\*</sup>, Ya-Ru Luo<sup>1,\*</sup>, Chang Liu<sup>1</sup>, Xiao-Bin Jin<sup>1</sup>, Wei Zhou <sup>1</sup>, Guang-Hong Xu<sup>1,2</sup>

<sup>1</sup>Department of Anesthesiology, First Affiliated Hospital, Anhui Medical University, Hefei, Anhui, 230022, People's Republic of China; <sup>2</sup>Department of Neurology, First Affiliated Hospital, Anhui Medical University, Hefei, Anhui, 230022, People's Republic of China

\*These authors contributed equally to this work

Correspondence: Guang-Hong Xu, Department of Anesthesiology, First Affiliated Hospital, Anhui Medical University, Hefei, Anhui, 230022, People's Republic of China, Tel +86-551-62922344, Fax +86 551 62923704, Email xuguanghong2004@163.com

## Dear editor

In the letter to editor named “comparing efficacy of different analgesic modalities in patients undergoing total knee arthroplasty”, Gao et al noted some issues to discuss and offer some recommendations for our study. Thank you for their suggestions, they are conducive for further improving scientific research, and we'd be happy to answer their queries.

First, in our study, there were ten patients in each group in the preliminary study, and the results of pre-experiment were not included in the final analysis. Previous studies showed that reductions in pain scores of around 30–40% are needed to reflect clinically useful improvements in pain,<sup>1</sup> and a 30% difference in pain scores on the Visual Analog Scale score was selected to calculate sample size.<sup>2</sup> Although the non-inferiority margin of our study is set based on the results of the pre-experiment, the results of the pre-experiment were only used to calculate the sample size. Next, we used the results of a previous similar study<sup>3</sup> for sample size estimation. When the 0.5 times standard deviation of the pain score at postoperative 24 hours during motion in the control group was selected as the non-inferiority margin, the estimated sample size was 42 patients in each group. The total sample size in our study was 87 patients, which meets the requirements of the study. Thus, the sample size in our study is adequate to achieve an effective inferiority test.

Second, it is certain that it is more appropriate to convert all postoperative analgesics used to equivalent opioid consumption. But we can not find an equivalent morphine dose in other previous studies. So, the dosages of different rescue analgesics were expressed separately as performed in other studies,<sup>4–6</sup> which did not perform equal analgesia conversion of all analgesics.

Finally, the quality of early postoperative recovery and patients' satisfaction with postoperative analgesia may be helpful for the evaluation of postoperative pain control effect, and it does help to improve the quality of research. But they are not necessary for the assessment of postoperative pain, as shown in previous studies.<sup>5,6</sup>

## Disclosure

The authors report no conflicts of interest in this communication.

## References

1. Myles PS, Myles DB, Gallagher W, et al. Measuring acute postoperative pain using the visual analog scale: the minimal clinically important difference and patient acceptable symptom state. *Br J Anaesth*. 2017;118(3):424–429. doi:10.1093/bja/aew466
2. Patterson ME, Bland KS, Thomas LC, et al. The adductor canal block provides effective analgesia similar to a femoral nerve block in patients undergoing total knee arthroplasty—a retrospective study. *J Clin Anesth*. 2015;27(1):39–44. doi:10.1016/j.jclinane.2014.08.005

3. Wang Q, Hu J, Zeng Y, Li D, Yang J, Kang P. Efficacy of two unique combinations of nerve blocks on postoperative pain and functional outcome after total knee arthroplasty: a prospective, double-blind, randomized controlled study. *J Arthroplasty*. 2021;36(10):3421–3431. doi:10.1016/j.arth.2021.05.014
4. Shin H, Do S, Lee J, Kim T, Na H. Comparison of intraoperative sedation with dexmedetomidine versus propofol on acute postoperative pain in total knee arthroplasty under spinal anesthesia. *Anesth Analg*. 2019;129(6):1512–1518. doi:10.1213/ANE.0000000000003315
5. Palanne R, Rantasalo M, Vakkuri A, et al. Effects of anaesthesia method and tourniquet use on recovery following total knee arthroplasty: a randomised controlled study. *Br J Anaesth*. 2020;125(5):762–772. doi:10.1016/j.bja.2020.03.036
6. Wiesmann T, Hüttemann I, Schilke N, et al. Ultrasound-guided single injection versus continuous sciatic nerve blockade on pain management and mobilisation after total knee arthroplasty (CoSinUS trial). *Eur J Anaesth*. 2018;35(10):782–791. doi:10.1097/EJA.0000000000000830

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Drug Design, Development and Therapy 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Drug Design, Development and Therapy editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

## Drug Design, Development and Therapy

Dovepress

### Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>

<https://doi.org/10.2147/DDDT.S362387>