

Molecular Pathways of Diabetic Kidney Disease Inferred from Proteomics

Lan Wei ¹, Yuanyuan Han², Chao Tu ¹

¹Department of Internal Medicine, The Third Affiliated Hospital of Soochow University, Changzhou, People's Republic of China; ²Institute of Medical Biology, Chinese Academy of Medical Sciences and Peking Union Medical College, Yunnan Key Laboratory of Vaccine Research and Development on Severe Infectious Diseases, Kunming, People's Republic of China

Correspondence: Chao Tu, Department of Internal Medicine, The Third Affiliated Hospital of Soochow University, 185 Juqian Road, Changzhou, 213000, People's Republic of China, Email tcmedical21@czfph.com

Abstract: Diabetic kidney disease (DKD) affects an estimated 20–40% of type 2 diabetes patients and is among the most prevalent microvascular complications in this patient population, contributing to high morbidity and mortality rates. Currently, changes in albuminuria status are thought to be a primary indicator of the onset or progression of DKD, yet progressive nephropathy and renal impairment can occur in certain diabetic individuals who exhibit normal urinary albumin levels, emphasizing the lack of sensitivity and specificity associated with the use of albuminuria as a biomarker for detecting diabetic kidney disease and predicting DKD risk. According to the study, a non-invasive method for early detection or prediction of DKD may involve combining proteomic analytical techniques such as second generation sequencing, mass spectrometry, two-dimensional gel electrophoresis, and other advanced system biology algorithms. Another category of proteins of relevance may now be provided by renal tissue biomarkers. The establishment of reliable proteomic biomarkers of DKD represents a novel approach to improving the diagnosis, prognostic evaluation, and treatment of affected patients. In the present review, a series of protein biomarkers that have been characterized to date are discussed, offering a theoretical foundation for future efforts to aid patients suffering from this debilitating microvascular complication.

Keywords: diabetic kidney disease, proteomics, diagnostic biomarkers, therapeutic targets

Introduction

Diabetic kidney disease (DKD) is a common microvascular complication that impacts 20–40% of individuals diagnosed with type 2 diabetes, resulting in high rates of morbidity and mortality. DKD is diagnosed based on the detection of persistent proteinuria, an albumin creatinine ratio greater than 30 mg/g creatinine, a reduction in the estimated glomerular filtration rate (eGFR), and the progressive deterioration of kidney function.^{1,2} Persistent hyperglycemia in individuals with diabetes can drive the increased synthesis of extracellular matrix (ECM) proteins, enhanced cell proliferation, and the dysfunction of endothelial cells together with tubular atrophy, interstitial fibrosis, and the thickening of the glomerular and tubular basement membrane as a consequent of inflammation, oxidative stress, and advanced glycation end product (AGE) production.³ DKD development is further driven by the increased accumulation of protein within the renal extracellular matrix, the expansion of the mesangial matrix, glomerular hyperfiltration, tubulointerstitial fibrosis, and the aggravated thickening of the glomerular basement membrane. Injury to glomerular podocytes can also contribute to DKD incidence.¹

Both a reduction in eGFR and albuminuria are considered relatively non-specific biomarkers of DKD given that they are also altered in the context of many other chronic glomerular diseases, with 60% of nephron function already being absent with eGFR values falling to the threshold of 60 mL/min/1.73 m².⁴ Given that DKD is a prevalent cause of end-stage renal disease (ESRD), it is critical that reliable biomarkers that can guide the diagnosis and treatment of this condition be defined. Both urinary albumin (UALB) and β 2-microglobulin (U β 2m) have been leveraged as biomarkers of glomerular and tubular injury,⁵ yet these levels and eGFR changes are only evident after substantial glomerular damage has taken place. However, many patients exhibit histopathological alterations in renal structures prior to the onset of microalbuminuria.⁶ While these changes can be detected through renal biopsy, enabling the more robust diagnosis and

monitoring of affected patients, the utility of this approach is limited by the inherent invasivity of biopsy procedures. As such, there is a pressing need for the establishment of new, non-invasive biomarkers that can gauge the risk of future DKD onset or detect this disease while in its earliest stages.

Certain urinary and plasma protein biomarkers have been linked to the pathophysiology of many forms of renal disease, and some of these biomarkers have even been successfully implemented in clinical settings. The kidneys directly produce urine. It is relatively simple and non-invasive to collect urine protein because it is so stable. It can be a useful biomarker and is difficult to degrade.^{7,8} Blood samples are also easy to collect, serving as an optimal sample type for efforts to diagnose early-stage DKD.⁸⁻¹⁰ Proteins found in renal tissue allow for the monitoring of biomolecules' effects on tissues. As proteomic indicators of DKD, they are currently becoming more and more well-known.¹¹ Indeed, the proteome has long been regarded as an ideal focus for efforts to understand the molecular etiology of disease and to define prognostic or diagnostic biomarkers associated with particular disease states. The use of blood, urine, and renal tissue proteomic analyses to non-invasively detect and monitor DKD has emerged as an increasingly important field that can clarify the pathophysiological mechanisms underlying DKD while identifying key targets for future efforts to prevent subsequent disease progression.¹² The systematic examination of these proteomic targets offers more comprehensive insight into DKD and other complex metabolic diseases while also providing a foundation to design prognostic tools and targeted therapies. DKD biomarkers have been the subject of numerous transcriptomics and metabonomics investigations in the past. Lei et al conducted work on transcriptomics,¹³ and Mu et al conducted experiments on metabonomics.¹⁴ Proteomics is the topic of this essay. This article primarily examines and discusses the most recent developments in the proteome analysis of urine, blood, and kidney tissue samples for the purpose of identifying potential specific biomarkers for the diagnosis of DKD.

Proteins and Proteomics

The proteome is a catch-all term that refers to all proteins produced by a given tissue, cell, or organism. A range of proteomic techniques have been leveraged to date to characterize the proteome or the expression of particular protein subsets, including low molecular mass polypeptides. Protein-based biomarkers of DKD have been explored through enzyme-linked immunosorbent assay (ELISAs), 2-way gel electrophoresis (2-DE),¹⁵ equivalent labeling for relative and absolute quantification (iTRAQ),^{16,17} tandem mass labeling (TMT) and mass spectrometry, unlabeled approaches, and biochip analyses.¹⁸ Proteomic analyses can enable the establishment of DKD-specific biomarkers through analyses of patient urine, blood, or renal tissue samples,^{19,20} with advances in the underlying techniques having made the robust and routine screening of biofluid samples increasingly feasible, potentially enabling the detection of novel targets for the treatment of DKD.²¹ These biomarkers also have the potential to advance current understanding of the mechanistic basis for DKD-related injury and to provide new diagnostic, monitoring, and classification options to support the individualized treatment of affected patients.²² Labeled and unlabeled mass spectrometry methods, which may simultaneously screen thousands of target proteins, are currently the most widely used proteomic tools for locating biomarkers relevant to DKD diagnosis and prognosis.

Urinary Protein Biomarkers of DKD

In addition to being easy to collect in a non-invasive manner, urine can remain stable for years in storage such that even archived urine samples can be used to screen for DKD-related biomarkers.²³ Urinary proteomics analyses thus offer an opportunity to define small proteins and peptides characteristic of pathophysiological shifts associated with early-stage DKD. In one study, a proteomic biosignature consisting of 273 different urinary protein fragments known as the CKD273 classifier was shown to be capable of differentiating between diabetic kidney disease patients and individuals with other forms of chronic renal disease.²⁴ The CKD273 classifier is a more accurate predictor of the incidence rate of DKD and can aid in the diagnosis, monitoring, and prognosis assessment of DKD patients. As it can detect individuals at risk of developing DKD, it can also help protect these patients from adverse renal outcomes.²⁵ Limonte et al determined that high urinary levels of the lysosomal proteolytic enzyme cathepsin D are linked to intracellular protein degradation, autophagy, and apoptotic cell death.²⁶ Elevated urinary cathepsin D levels are closely associated with rapid eGFR declines, tubular atrophy, tubulointerstitial inflammation, and fibrosis.²⁷ Ahn et al determined that urinary proteomic analyses are more reliable than measurement of albuminuria alone as a tool to predict DKD patient prognosis, with a proteomic biosignature comprised of ACP2, CTSA, GM2A, MUC1, and SPARCL1 offering substantial clinical

utility.²⁸ Chen et al determined that β 2-microglobulin and Clara-cell protein are protein biomarkers associated with DKD that can aid in predicting the incidence of this condition and associated risks.

ITRAQ, TMT, etc. are technologies that can detect proteins in a specific way.²⁹ Through the use of a 2D-DIGE-MALDIQ-TOF strategy and related techniques, Patel et al were able to detect novel DKD-related protein biomarkers in at-risk individuals corresponding to the processes of renal tubulointerstitial fibrosis and tubulointerstitial fibrosis, including apolipoprotein A1, α -1microglobulin, and zinc α -2 glycoprotein.^{30,31} Guo et al discovered that - 1-antitrypsin - 1-acid glycoprotein, ceruloplasmin, prostate stem cell antigen, APOA4 and other apolipoprotein AIV are prospective protein biomarkers of DKD in 23 diabetic patients and 16 healthy volunteers.³² Liao et al also observed a significant increase in urinary haptoglobin (Hp) levels in DKD patients, allowing for the more accurate diagnosis and prognostic assessment of individuals with this disease.³³ Urinary retinol-binding protein 4 (uRBP4) and SH3 domain-binding glutamic acid-rich-like protein 3 (uSH3BGRL3) protein levels are closely associated with eGFR.³⁴ Through an SDS-PAGE approach, Araumi et al determined that urinary afamin (uAFM), uRBP4, and uSH3BGRL3 levels were elevated in DKD patients relative to individuals without renal damage, underscoring their potential utility as diagnostic biomarkers in this context.^{20,35,36} Fan et al have also demonstrated that haptoglobin (HPT) and α -1-microglobulin / bicunin precursor (AMBP) are two protein biomarkers that can effectively differentiate between healthy individuals and diabetic patients who are or are not affected by DKD. Urinary proteomic techniques can aid in the non-invasive diagnosis of DKD and the identification of patients considered at high risk.³⁷

The low molecular weight protein RBP is filtered by the glomeruli and ultimately reabsorbed via the proximal tubules, and has been found to offer substantial diagnostic utility in DKD patients who exhibit pronounced albuminuria. Al-Rubeaan et al reported the increased excretion of urinary transferrin, RBP, and serum osteopontin with diabetic kidney disease development, emphasizing the value of these biomarkers when diagnosing type 2 diabetic kidney disease.³⁸ Urinary L-FABP is a recently detected urinary protein biomarker used for early DKD detection, and Panduru et al reported higher urinary L-FABP levels in DKD patients relative to controls in a manner associated with disease severity. Urinary L-FABP thus represents a specific, sensitive protein biomarker capable of guiding early DKD diagnosis.³⁹ Nucleated cells produce the low molecular mass protein cystatin C (CST3), and cysteine aminotransferase activity can be used as a metric for kidney function in diabetic individuals with higher CST3 levels in urine being indicative of renal tubular damage. Through a CE-MS approach, Garg et al determined that urinary CST3 levels were significantly elevated in DKD patients relative to controls ($P < 0.05$), supporting the diagnostic utility of this protein ($P < 0.05$), indicating that urinary CST3 has diagnostic value.⁴⁰ A DKD-related proteomic classifier consisting of collagen fragments, α -microglobulin, β 2-microglobulin (β 2MG), A1AT, uromodulin, and other peptides has also shown promise as a tool for identifying type 2 diabetes patients at a high risk of DKD.⁴¹ For further details regarding these urinary protein biomarkers, see [Table 1](#).

Blood-Based Protein Biomarkers of DKD

Tubular cells are activated in response to high glucose levels, resulting in the secretion of inflammatory mediators and ECM components that contribute to tubular thinning and cell damage. Glomerular injury-associated proteinuria can act as a stressor that adversely impacts proximal renal tubular cells, driving inflammatory mediator secretion and renal damage. Blood can be kept stable in storage for a number of years and can be used to check for DKD-related biomarkers. Circulating biomarkers can offer insight into kidney damage in patients diagnosed with diabetes,⁴² offering new potential means of guiding patient diagnosis, treatment, and prognostic assessment.⁴³ Plasma u/PC3 levels in the serum of DKD patients have been shown to be elevated as compared to diabetic individuals free of renal damage, emphasizing the potential diagnostic utility of this serum biomarker.³⁵ Chiang et al determined that SH3BGRL3 can function as an epidermal growth factor receptor conjugate, and that there is a significant positive correlation between uSH3BGRL3 levels and eGFR such that it can serve as a DKD-related biomarker.³⁴ Caseiro et al further detected increases in MMP-9, neutrophil gelatinase-associated lipocalin (NGAL), aminopeptidase N, azurocidin, and kallikrein 1 activity in type 1 DKD patients as compared to healthy individuals, suggesting that this protein biosignature may offer value in this type of DKD.⁴⁴ Moresco et al also confirmed that circulating NGAL levels correspond to renal tubular function, and a positive correlation has been reported between serum NGAL concentrations and proteinuria whereas these levels are negatively correlated with eGFR.⁴⁵ The lipocalin protein family member β -trace protein (β -TP), also known as lipoprotein

Table 1 Biomarkers of Urine Proteomics in DKD

PMID	Types (The Main Biomarkers)	Population	Authors	Methods	Disease	Main Findings
35377940	Urinary cathepsin D	Human and mouse models	Limonte et al ²⁶	LC-MS	Type 1 DKD	Elevated urinary cathepsin D was associated with rapid decline in EGFR, tubulointerstitial inflammation, fibrosis, and tubular atrophy.
32545899	ACP2, CTSA, GM2A, MUC1 和 SPARCL1	Human	Ahn et al ²⁸	Nano-LC-ESI-MS/MS Analysis	Type 2 DKD	A biomarker composed of five proteins, ACP2, CTSA, GM2A, MUC1 and SPARCL1, is important in the diagnosis of DKD
35327145	Glutathione S-transferase Mu 1, Zinc finger C2HC domain-containing protein 1A, Haptoglobin, Glutaredoxin-3, Zinc finger protein 2	Animal (dog)	Winiarczyk et al ⁷⁰	MALDI-TOF MS	DKD	All five proteins were up-regulated in diabetic canine urine, and the only protein up-regulated in microalbuminuric dogs was zinc finger protein 2.
29729954	U/PC3	Human	Li et al ²²	Sodium dodecyl sulfate (SDS)- polyacrylamide gel electrophoresis	DKD	The level of urinary C3 in diabetic nephropathy group was significantly higher than that in diabetic group without renal damage.
27293888	uRBP4	Human	Fiseha et al ³⁶	Sodium dodecyl sulfate (SDS)- polyacrylamide gel electrophoresis	DKD	uRBP4 is a urinary marker of early tubular damage in DKD, and the glomerular filtration rate of DKD is negatively correlated with uRBP4.
30486327	Haptoglobin and α -1-microglobulin/bikunin precursor	Human	Liao et al ³³	iTRAQ	Type 2 DKD	Haptoglobin and α -1-microglobulin/bikunin precursor are early indicators of DKD and markers associated with ERFD in diabetic patients.
30024955	β 2-microglobulin and clara-cell protein	Human	Chen et al ²⁹	C18 plate-MALDI-TOF	Type 2 DKD	MALDI-TOF MS has high specificity in detecting protein markers by measuring protein quality, β 2-microglobulin and clara-cell protein are DKD-related biomarkers.
32856461	408 N-linked glycoproteins, A1AT, and ceruloplasmin	Human	Jin et al ⁷¹	iTRAQ	DKD	408 N-linked glycoproteins, A1AT, and ceruloplasmin as common or specific biomarkers for DKD.
31131043	Apolipoprotein A1, α -1 microglobulin, zinc α -2 glycoprotein	Human	Patel et al ³⁰	2D-DIGE-MALDI Q-TOF	DKD	Proteomic analysis may be a useful tool to enable early detection of new markers in patients susceptible to diabetic nephropathy, such as apolipoprotein A1, α -1 microglobulin, zinc α -2 glycoprotein, etc.
26608305	Alpha-1-antitrypsin, alpha-1-acid glycoprotein 1, ceruloplasmin, prostate stem cell antigen and apolipoprotein AIV	Human	Guo et al ³²	2D-LC-MS/MS and iTRAQ quantification	Type 2 DKD	Using 2D-LC-MS/MS and iTRAQ quantification technology, it was found that α -1-antitrypsin, α -1-acid glycoprotein, ceruloplasmin, prostate stem cell antigen and apolipoprotein AIV in urine are candidate protein biomarkers for DKD.
34458592	uAFM, uSH3BGRL3, uRBP4	Human	Araumi et al ³⁵	Sodium dodecyl sulfate (SDS)- polyacrylamide gel electrophoresis	DKD	uAFM, uRBP4 and uSH3BGRL3 are promising diagnostic markers for DKD.
34963468	HPT, AMBP	Human	Fan et al ³⁷	Nano HPLC-MS	DKD	HPT and AMBP are the two protein biomarkers with the highest capacity for diagnosing DKD patients.

30574040	Transthyretin precursor, Ig k chain C region, carbonicanhydrase 1, retinol binding protein, beta-2-microglobulin precursor and beta-2-glycoprotein 1.	Human	Bellei et al ⁷²	2-DE, Q-ToF-LC/MS, Agilent-Technologies, Western blot, ELISA	Type 2 DKD	In a proteome-related study, significantly differentially expressed proteins were detected in DKD patients compared with healthy control subjects.
29854824	Protein S100A9, Kallikrein-I, Pepsin A-3, Uteroglobin, Prostatic acid phosphatase, Ganglioside GM2, VCAM-1.	Human	Guillén et al ⁷³	TMT labeling, LC-MS, ELISA.	Type 2 DKD	Detection of these proteins in urine could serve as a potential new tool for biomarkers of DKD progression.
28577020	Urinary transferrin, urinary Retinol binding protein (RBP) and serum osteopontin	Human	Al-Rubeaan et al ³⁸	Biochip assay, ELISA	Type 2 DKD	Urinary transferrin, RBP and serum osteopontin have great diagnostic value in different stages of type 2 diabetic nephropathy.
23378622	L-FABP	Human	Panduru et al ³⁹	CE-MS, CKD273 classifier analysis	Type 2 DKD	The concentration of L-FABP in the urine of DKD patients was significantly increased, and it was closely related to the development and severity of the disease.
25634253	CST3	Human	Garg et al ⁴⁰	CE-MS analysis	DKD	Cystatin C (CST3) is a low-molecular-weight protein produced by nucleated cells in vivo. CST3 can be used to assess renal function. Urinary CST3 indicates renal tubular damage.
22872235	Collagen fragment, α -microglobulin, β 2-microglobulin (β 2MG), A1AT, uromodulin and other peptides	Human	Zürbig et al ⁴¹	iTRAQ labeling, Nano-LC-MS/MS, Label-Free Quantitative Proteomics, ELISA	DKD	Collagen fragments, α -microglobulin, β 2-microglobulin (β 2MG), A1AT, uromodulin and other peptides have been shown to be promising protein biomarkers.
28396115	TNFR1, TNFR2	Human	Yamanouchi et al ⁵⁰	LC-MS	T2D, T1D	The median serum concentrations of TNFR1 and TNFR2 at baseline were almost twice that of T1D patients with normal proteinuria and normal renal function.
28851702	apoA4, CD5L and IBP3	Human	Peters et al ⁵²	Mass spectrometry	T2D, T1D	apoA4, CD5L, and CIQB independently predicted rapid decline (odds ratio 2.40 [95% CI 1.24–4.61], 0.52 [0.29–0.93], and 2.41 [1.14–5.11], respectively) and improved model performance and fit in 35 participants ($P < 0.001$).
23780945	AGEs, OPs	Human	Beisswenger et al ⁵⁴	LC-MS/MS	DKD	Blood samples from 103 normal albuminuria subjects demonstrated that the increase of advanced glycation end products (AGEs) and oxidation products (OPs) was considered to lead to the progress of DKD.

prostaglandin D2 synthase (L-PGDS), is reportedly an independent and reliable serum protein biomarker that responds to eGFR levels more reliably than do anhydride or cystatin.⁴⁶ Lu et al employed Q-Exactive mass spectrometry and proteomic iTRAQ techniques which ultimately revealed that DKD patient serum exhibited higher concentrations of gelatin, collectin-11, protein tyrosine phosphatase type J receptor (PTPRJ), and protein kinase anchoring protein-7 (AKAP-7), supporting their possible diagnostic value.⁴³ The circulating glycoprotein Hp is primarily produced in the liver, and plasma Hp levels are significantly higher in individuals with DKD. Plasma gelatin protein (PGSN) is primarily produced by skeletal muscle, smooth muscle, and the myocardium. Pikel et al determined that PGSN can promote PKC activation, thereby contributing to renal reactive oxygen species (ROS) production, inflammation, and DKD development.⁴⁷ DKD patients exhibit higher PGSN levels than do controls, emphasizing the potential value of this protein as a diagnostic biomarker in this pathological context.⁴⁸ Yang et al reported that serum levels of the inflammatory cytokine tumor necrosis factor- α (TNF α) are significantly correlated with early glomerular dysfunction and early-stage DKD.⁴⁹ Yamanouchi et al discovered that 221 individuals with type 2 diabetes and 279 patients with type 1 diabetes had higher serum levels of the tumor necrosis factor receptors (TNFR) 1 and 2. End-stage renal disease (ESRD), T2D, and T1D were all highly correlated with TNFR. According to the study, this indicator had an 85% predictive value and a 72% sensitivity.^{50,51} In Phase II of the diabetes study, Peters et al used mass spectroscopy on 35 community patients. They discovered that the participants' plasma levels of apolipoprotein (apo) A-IV (apoA4), CD5 antigen like (CD5L), and insulin-like growth factor binding protein 3 (IBP3) independently predicted rapid decline (odds ratio 2.40 [95% CI 1.24–4.61], 0.52 [0.29–0.93], and 2.41 [1.14–5 (P < 0.001)]. New plasma biomarkers (apoA4, CD5L, and IBP3) have been discovered by recent study that may help predict the rapid loss in renal function in type 2 diabetic patients. Among these, the increase in ApoA4 level is a precursor to mild to moderate DKD, the increase in IBP3 level is connected to low baseline EGFR, and the inhibitor of apoptosis of CD5L or macrophage protein is related to immunological and inflammatory response.^{52,53} Beisswenger et al performed an LC-MS/MS investigation on blood samples from 103 people with normal albuminuria diabetes and discovered that the progression of DKD was thought to be accompanied by an increase in advanced glycation end products (AGEs) and oxidation products (OPs).⁵⁴ El-Asrar et al found soluble CD40 ligand (CD40L) to represent a promising plasma protein biomarker associated with progressive DKD that can participate in endothelial injury and inflammatory activity.⁵⁵ In a separate analysis, Kobayashi et al detected an independent association between the LAYN, DLL1, MAPK11, MATN2, Endostatin, and ROR1 proteins and the risk of chronic renal failure owing to their role in the progression of DKD, emphasizing their promise as targets for renoprotective therapeutic interventions.²⁷ Niewczas et al additionally determined that plasma TNF-R1 and TNF-R2 play roles in renal inflammation and can predict the decline in renal function that is associated with DKD incidence.⁵⁶ For further details regarding blood-based protein biomarkers of DKD characterized to date, see [Table 2](#).

Renal Tissue Protein Biomarkers of DKD

Recent advances in proteomic techniques have enabled the routine assessment of proteins in urine, plasma, and kidney tissue samples, allowing for the more in-depth interrogation of the mechanistic basis for DKD progression.⁵⁷ Renal protein levels may reflect the actual pathogenesis of tubular atrophy, glomerular injury, and tubulointerstitial fibrosis. To aid in the characterization of renal proteomic profiles, high-throughput matrix-assisted laser desorption/ionization imaging MS has been leveraged to analyze the proteins present in tissue sections.⁵⁸ Renal tissue proteomic analyses can detect biomarkers of glomerular injury such as albumin, as well as renal tubular injury-related proteins including α 1MG, β 2MG, RBP4, N acetyl β D glucosaminidase (NAG), alkaline phosphatase (ALP), gamma-glutamyltransferase (GGT), liver type fatty acid binding protein (LFABP), NGAL, and renal injury molecule 1 (KIM1).⁵⁹ Moresco et al determined that the transmembrane glycoprotein Kim-1 is specifically expressed on proximal tubular cells on the apical membrane, serving as a reliable biomarker of proximal tubule injury that is closely linked to the extent of renal tubulointerstitial injury and fibrosis, offering a tool to diagnose early-stage DKD.⁴⁵ Vaidy et al observed elevated NAG and Kim-1 levels in type 1 diabetes patients with microalbuminuria, with these levels being even higher in patients with albuminuria, emphasizing the link between these protein biomarkers and type 1 diabetes-related DKD.⁶⁰ The 25 kDa lipocalin family member NGAL is often studied as a sensitive, specific biomarker for renal tubular injury that can predict DKD development.⁶¹ Regucalcin is a mediator of cellular Ca²⁺ homeostasis, oxidative stress regulation, and ascorbic acid biosynthesis. Using a 2D-DGE and MS-based

Table 2 Biomarkers of Blood Proteomics in DKD

PMID	Types (The Main Biomarkers)	Population	Authors	Methods	Disease	Main Findings
34458592	Plasma U/PC3	Human	Araumi et al ³⁵	Q-Exactive, Thermo Scientific	DKD	Plasma U/PC3 was highest in diabetic nephropathy group, which was significantly higher than that in non-diabetic nephropathy group (P<0.05).
26286913	SH3BGRL3	Human	Chiang CY et al ³⁴	Q-Exactive, Thermo Scientific	DKD	SH3BGRL3 is a conjugate of epidermal growth factor receptor, and uSH3BGRL3 in DKD patients was significantly positively correlated with estimated glomerular filtration rate (P>0.05).
22975642	MMP-9/neutrophil gelatinase-associated lipocalin, aminopeptidase N, azurocidin and kal-likrein I	Human	Caseiro et al ⁴⁴	LC-MS/MS	Type I DKD	Fifteen patients with type I DKD had higher MMP-9/neutrophil gelatinase-associated lipoprotein, aminopeptidase N, azurocidin, and kallikrein I activities compared with healthy controls.
23485645	NGAL	Human	Moresco et al ⁴⁵	LC-MS/MS	DKD	Serum NGAL is positively correlated with proteinuria and negatively correlated with glomerular filtration rate, and is a promising biomarker for the diagnosis of DKD.
26124640	βTP	Human	Bacci et al ⁴⁶	LC-MS/MS	Type I DKD	β-Trace protein (βTP), a member of the Lipocalin protein family, is a new independent and reliable serum protein marker, a better GFR marker than creatinine and cystatin.
31384238	Gelsolin, collectin-I I, PTPRJ, and AKAP-7 proteins	Human	Lu et al ⁴³	iTRAQ, LC-MS/MS	Type 2 DKD	The contents of gelatin, collectin-I I, protein tyrosine phosphatase J-type receptor (PTPRJ) and protein kinase-anchored protein-7 (AKAP-7) were higher in DKD patients.
30149613	PGSN	Human	Piktel et al ⁴⁷	iTRAQ, LC-MS/MS	DKD	PGSN content is higher in DKD patients than in controls, so PGSN is also a promising protein diagnostic marker for DKD.
25986995	TNFRs	Human	Yang et al ⁴⁹	Surface-enhanced Laser desorption/ionization time-of-flight Massenspektrometrie (SELDI-TOF-MS)	Type 2 DKD	The levels of TNFRs were significantly correlated with early glomerular dysfunction and were protein biomarkers associated with early stages of DKD.
26975878	Orosomuroid	Human	El-Beblawy et al ⁷⁴	TMT labeling, LC-MS, ELISA	Type I DKD	Elevated Orosomuroid in the blood of DKD patients is an independent risk factor for kidney damage.
20205888	Transthyretin, Apolipoprotein A1, Apolipoprotein C1 and Cystatin C	Human	Overgaard et al ⁷⁵	CE-ESI-MS LC-MS, MALDI-MS	Type I DKD	Transthyretin, apolipoprotein A1, apolipoprotein C1 and cystatin C are potential plasma protein biomarkers.
22702645	sCD40L	Human	El-Asrar et al ⁵⁵	LC-MS	Type I DKD	Soluble CD40 ligands involved in inflammatory processes and markers of endothelial damage have emerged as potential markers of progressive DKD.
35618095	LAYN, ESAM, DLL1, MAPK1 I and endostatin	Human	Kobayashi et al ²⁷	LC-MS, MALDI-MS	DKD	LAYN, DLL1, MAPK1 I, MATN2, Endostatin and ROR1 proteins are involved in the progression of DKD disease.
31011203	TNF-R1, TNF-R2, KRIS proteins	Human	Niewczas et al ⁵⁶		DKD	Plasma levels of tumor necrosis factor-R1 or tumor necrosis factor-R2 are involved in renal inflammatory responses and are predictors of DKD.

Table 3 Biomarkers of Renal Proteomics in DKD

PMID	Types (The Main Biomarkers)	Population	Authors	Methods	Disease	Main Findings
23485645	KIM I	Human	Moresco et al ⁴⁵	2-DE, CE-ESI-MS	Type 1 DKD	Kim-I is closely related to the degree of tubulointerstitial injury and fibrosis, which provides the possibility of a new marker for early diagnosis of DKD.
20980978	NAG	Human	Vaidya et al ⁶⁰	Spectrophotometrically	Type 1 DKD	NAG and Kim-I levels were elevated in type 1 diabetic patients with microalbuminuria, and NAG and Kim-I levels were highest in diabetic patients with proteinuria.
21779943	NGAL	Human	Fu et al ⁶¹	Sandwich ELISA kits	Type 2 DKD	NGAL is a sensitive and specific protein biomarker that reflects renal tubular injury, and has become a sensitive and specific early biomarker for predicting DKD.
26072307	L-FABP Regucalin	Animal.	Zubiri et al ⁶²	2-dimensional differential gel electrophoresis, mass spectrometry, Matrix Assisted Laser Desorption/Ionization MS	DKD	Differential analysis was performed on the renal tissue proteome of the early DKD rat model, and it was found that the Regucalin content in the renal tissue of DKD rats was significantly reduced compared with the control tissue.
29134286	Metalloproteinase TIMP3	Animal.	Rossi et al ⁶⁵	Nano-liquid chromatography–electrospray ionization–tandem mass spectrometry (nLC–ESI–MSE) analyses	DKD	
25884625	Periostin	Animal.	Satirapoj et al ⁶⁶	ELISA	Type 2 DKD	Glomerular periostin levels are elevated in patients with type 2 diabetes before overt albuminuria in diabetic patients.
22172726	Nephronectin, apolipo-protein E	Human	Nakatani et al ⁶⁸	QSTAR Elite liquid chromatogram-phy with tandem mass spectrometry, LC–MS/MS and iTRAQ	DKD	Nephronectin and apolipo-protein E play key roles in the development of renal extracellular matrix and are related to the development of diabetic glomerulosclerosis.
27457269	Ribosomal proteins, cytoskeletal proteins	Pig	Malagrino et al ⁷⁶	LC-ESI-MS/MS	DKD	Kidney tissue proteins can be used as non-invasive biomarkers for DKD.
21544065	Vanin-1, Ennp6 protein, alpha-1-acid glycoprotein, uromodulin	Rat	Fugmann et al ⁶⁹	MALDI-TOF, MALDI-TOF/TOF MS	Type 1 DKD	The up-regulated proteins in DKD were Vanin-1 and cystatin C, while the expression of uromodulin was down-regulated in diabetic rat samples.
28938946	PKM2	Rat	Chun et al ⁷⁷	LC–MS/MS	DKD	Pyruvate kinase M2 (PKM2) is one of the highly upregulated proteins in DKD glomeruli.
27356525	(MDR1/P-gp, BCRP/Bcrp and MRPs/Mrps 2 and 3	Human	Fallon et al ⁷⁸	NanoLC-MS/MS	DKD	(MDR1/P-gp, BCRP/Bcrp and MRPs/Mrps 2 and 3 are elevated in DKD.
23846697	Heme oxygenase-1 and other AngII-regulated proteins	Mice	Konvalinka et al ⁷⁹	LC-MS/MS	DKD	Heme oxygenase-1 and other AngII- regulated proteins represented novel markers of AngII activity.

approach, Zubiri et al compared the proteomic profiles of renal tissue samples from rats with early-stage DKD and observed significant decreases in regucalin levels in tissues from DKD model rats relative to controls.^{62,63} The matrix protein TIMP3 (tissue inhibitor of metalloproteinases) can inhibit proteolytic enzyme activity and receptor activation in the ECM. DKD has been shown to be associated with TIMP3 downregulation.⁶⁴ Rossi et al employed differential proteomic analyses of renal tissues from 6-week-old wild-type (WT) and *Timp3*^{-/-} mice using a nano-liquid chromatography-electrospray ionization-tandem mass spectrometry (nLC-ESI-MSE) approach, observing significant TIMP3 downregulation in DKD mice consistent with its possible relevance as a therapeutic target in this disease.⁶⁵ The cell adhesion molecule periostin (POSTN) has also been posited to serve as a protein-based marker of diabetes-related renal damage, with elevated glomerular POSTN concentrations in patients with type 2 diabetes even before the onset of significant proteinuria such that it may hold value as a protein marker of early-stage renal injury.⁶⁶ Increased glomerular ECM protein deposition is a primary mechanism governing the pathogenesis of DKD, contributing to higher glomerular membrane ECM accumulation that ultimately contributes to glomerular membrane dilatation and glomerulosclerosis.⁶⁷ LC-MS/MS and iTRAQ techniques have both been used to compare renal tissues from DKD and nondiabetic patients. Nakatani et al determined that both nephronectin and apolipoprotein E are important mediators of renal extracellular matrix development, while renal connexin is an integrin $\alpha 8\beta 1$ ligand associated with diabetic glomerulosclerosis onset and progression. Further studies of these and other targets in patient renal tissues may aid in further understanding of DKD.⁶⁸ Fugmann et al further determined that significant renal Vanin-1 and cystatin C upregulation were evident in their model system, corresponding to proinflammatory and cytoprotective effects, while uromodulin was downregulated in diabetic model rates such that these proteins may offer value as therapeutic targets in DKD.⁶⁹ For further details regarding known renal tissue protein biomarkers of DKD, see [Table 3](#).

Conclusions and Outlook

As the studies discussed herein emphasize, there is growing interest in the use of alternative biomarkers other than urinary albumin and creatinine when detecting early-stage DKD, including protein-based biomarkers present in urine, blood, and renal tissue samples. Proteomic technologies offer an increasingly robust means of detecting low-abundance proteins related to the progressive pathogenesis of kidney injury.

Numerous biomarkers discovered thus far have demonstrated strong diagnostic potential, including urine-based CKD273 proteomic biomarkers, which have demonstrated excellent potential. In the clinic, more and more diagnostics are now available, allowing for the independent and accurate prediction of microalbuminuria associated with DKD. These proteomic biomarker analyses are often simpler and less invasive than standard diagnostic techniques, potentially reducing the need for renal biopsy procedures as the primary means of diagnosing DKD. These strategies also offer a theoretical foundation for more reliably treating this disease and preventing its progression. Despite such promise, however, proteomics-based biomarker panels are not routinely applied in most clinical chemistry laboratories. Further research focused on these proteomic biomarkers will guide the early and noninvasive diagnosis of DKD, allow for more reliable patient monitoring, and support better patient outcomes at many levels. We anticipate that further research efforts centered on the role of targeted proteomics in diabetic kidney disease will yield novel breakthroughs in this field with clear clinical value.

Ethics Statement

This review does not involve animal or human trials, therefore it does not require ethical review.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This review is supported by Yunnan Fundamental Research Projects (grant NO. 202001AT070145, 202101AT070288) and Changzhou Health Commission Youth Project (grant NO. QN202206).

Disclosure

The authors report no conflicts of interest in this review.

References

1. Liu S, Gui Y, Wang MS, et al. Serum integrative omics reveals the landscape of human diabetic kidney disease. *Mol Metab.* 2021;54:101367. doi:10.1016/j.molmet.2021.101367
2. Tofte N, Lindhardt M, Adamova K, et al. Early detection of diabetic kidney disease by urinary proteomics and subsequent intervention with spironolactone to delay progression (PRIORITY): a prospective observational study and embedded randomised placebo-controlled trial. *Lancet Diabetes Endocrinol.* 2020;8:301–312. doi:10.1016/S2213-8587(20)30026-7
3. Gordin D, Shah H, Shinjo T, et al. Characterization of glycolytic enzymes and pyruvate kinase M2 in type 1 and 2 diabetic nephropathy. *Diabetes Care.* 2019;42:1263–1273. doi:10.2337/dc18-2585
4. Uyy E, Suica VI, Boteanu RM, et al. Diabetic nephropathy associates with deregulation of enzymes involved in kidney sulphur metabolism. *J Cell Mol Med.* 2020;24:12131–12140. doi:10.1111/jcmm.15855
5. Shah IM, Mackay SP, McKay GA. Therapeutic strategies in the treatment of diabetic nephropathy - a translational medicine approach. *Curr Med Chem.* 2009;16:997–1016. doi:10.2174/092986709787581897
6. Nicholas SB. Use of urinary proteomics in diagnosis and monitoring of diabetic kidney disease. *Lancet Diabetes Endocrinol.* 2020;8:261–262. doi:10.1016/S2213-8587(20)30067-X
7. Klein J, Ramirez-Torres A, Ericsson A, et al. Urinary peptidomics provides a noninvasive humanized readout of diabetic nephropathy in mice. *Kidney Int.* 2016;90:1045–1055. doi:10.1016/j.kint.2016.06.023
8. Shao C, Wang Y, Gao Y. Applications of urinary proteomics in biomarker discovery. *Sci China Life Sci.* 2011;54:409–417. doi:10.1007/s11427-011-4162-1
9. Greco V, Piras C, Pieroni L, et al. Direct assessment of plasma/serum sample quality for proteomics biomarker investigation. *Methods Mol Biol.* 2017;1619:3–21.
10. Overgaard AJ, Thingholm TE, Larsen MR, et al. Quantitative iTRAQ-based proteomic identification of candidate biomarkers for diabetic nephropathy in plasma of type 1 diabetic patients. *Clin Proteomics.* 2010;6:105–114. doi:10.1007/s12014-010-9053-0
11. Cefalu WT, Bakris G, Blonde L, Boulton A, DAlessio D, Groot M. Standards of medical care in diabetes-2017: summary of Revisions. *Diabetes Care.* 2017;40:S4–S5. doi:10.2337/dc17-S003
12. Aluksanasuwan S, Plumworasawat S, Malaitad T, et al. High glucose induces phosphorylation and oxidation of mitochondrial proteins in renal tubular cells: a proteomics approach. *Sci Rep.* 2020;10:5843. doi:10.1038/s41598-020-62665-w
13. Lei L, Bai Y, Fan Y, et al. Comprehensive diagnostics of diabetic nephropathy by transcriptome RNA sequencing. *Diabetes Metab Syndr Obes.* 2022;15:3069–3080. doi:10.2147/DMSO.S371026
14. Mu X, Yang M, Ling P, et al. Acylcarnitines: can they be biomarkers of diabetic nephropathy? *Diabetes Metab Syndr Obes.* 2022;15:247–256. doi:10.2147/DMSO.S350233
15. Shen L, Zhao D, Chen Y, et al. Comparative proteomics analysis of serum proteins in gestational diabetes during early and middle stages of pregnancy. *Proteomics Clin Appl.* 2019;13:e1800060. doi:10.1002/prca.201800060
16. Moulder R, Bhosale SD, Goodlett DR, et al. Analysis of the plasma proteome using iTRAQ and TMT-based isobaric labeling. *Mass Spectrom Rev.* 2018;37(5):583–606. doi:10.1002/mas.21550
17. Bringans SD, Ito J, Stoll T, et al. Comprehensive mass spectrometry based biomarker discovery and validation platform as applied to diabetic kidney disease. *EuPA Open Proteom.* 2017;14:1–10. doi:10.1016/j.euprot.2016.12.001
18. Frantzi M, Latosinska A, Fluhe L, et al. Developing proteomic biomarkers for bladder cancer: towards clinical application. *Nat Rev Urol.* 2015;12:317–330. doi:10.1038/nrurol.2015.100
19. Geyer PE, Holdt LM, Teupser D, et al. Revisiting biomarker discovery by plasma proteomics. *Mol Syst Biol.* 2017;13:942. doi:10.15252/msb.20156297
20. Siwy J, Zurbig P, Argiles A, et al. Noninvasive diagnosis of chronic kidney diseases using urinary proteome analysis. *Nephrol Dial Transplant.* 2017;32:2079–2089. doi:10.1093/ndt/gfw337
21. Beasley-Green A. Urine proteomics in the era of mass spectrometry. *Int Neurourol J.* 2016;20:S70–75. doi:10.5213/inj.1612720.360
22. Li XQ, Chang DY, Chen M, et al. Complement activation in patients with diabetic nephropathy. *Diabetes Metab.* 2019;45:248–253. doi:10.1016/j.diabet.2018.04.001
23. Yang J, Liu D, Liu Z. Integration of metabolomics and proteomics in exploring the endothelial dysfunction mechanism induced by serum exosomes from diabetic retinopathy and diabetic nephropathy patients. *Front Endocrinol.* 2022;13:830466. doi:10.3389/fendo.2022.830466
24. Lindhardt M, Persson F, Zurbig P, et al. Urinary proteomics predict onset of microalbuminuria in normoalbuminuric type 2 diabetic patients, a sub-study of the DIRECT-Protect 2 study. *Nephrol Dial Transplant.* 2017;32(11):1866–1873. doi:10.1093/ndt/gfw292
25. Khan NU, Lin J, Liu X, et al. Insights into predicting diabetic nephropathy using urinary biomarkers. *Biochim Biophys Acta Proteins Proteom.* 2020;1868:140475. doi:10.1016/j.bbapap.2020.140475
26. Limonte CP, Valo E, Drel V, et al. Urinary proteomics identifies cathepsin D as a biomarker of rapid eGFR decline in type 1 diabetes. *Diabetes Care.* 2022;45:1416–1427. doi:10.2337/dc21-2204
27. Kobayashi H, Looker HC, Satake E, et al. Results of untargeted analysis using the SOMAscan proteomics platform indicates novel associations of circulating proteins with risk of progression to kidney failure in diabetes. *Kidney Int.* 2022;102:370–381. doi:10.1016/j.kint.2022.04.022

28. Ahn HS, Kim JH, Jeong H, et al. Differential urinary proteome analysis for predicting prognosis in type 2 diabetes patients with and without renal dysfunction. *Int J Mol Sci.* 2020;21. doi:10.3390/ijms22010021
29. Chen CJ, Liao WL, Chang CT, et al. Urine proteome analysis by C18 plate-matrix-assisted laser desorption/ionization time-of-flight mass spectrometry allows noninvasive differential diagnosis and prediction of diabetic nephropathy. *PLoS One.* 2018;13:e0200945. doi:10.1371/journal.pone.0200945
30. Patel DN, Kalia K. Characterization of low molecular weight urinary proteins at varying time intervals in type 2 diabetes mellitus and diabetic nephropathy patients. *Diabetol Metab Syndr.* 2019;11:39. doi:10.1186/s13098-019-0430-1
31. Chebotareva N, Vinogradov A, McDonnell V, et al. Urinary protein and peptide markers in chronic kidney disease. *Int J Mol Sci.* 2021;23(1):22. doi:10.3390/ijms23010022
32. Guo Z, Liu X, Li M, et al. Differential urinary glycoproteome analysis of type 2 diabetic nephropathy using 2D-LC-MS/MS and iTRAQ quantification. *J Transl Med.* 2015;13:371. doi:10.1186/s12967-015-0712-9
33. Liao WL, Chang CT, Chen CC, et al. Urinary proteomics for the early diagnosis of diabetic nephropathy in Taiwanese patients. *J Clin Med.* 2018;7:483.
34. Chiang CY, Pan CC, Chang HY, et al. SH3BGR1 protein as a potential prognostic biomarker for urothelial carcinoma: a novel binding partner of epidermal growth factor receptor. *Clin Cancer Res.* 2015;21:5601–5611. doi:10.1158/1078-0432.CCR-14-3308
35. Araumi A, Osaki T, Ichikawa K, et al. Urinary and plasma proteomics to discover biomarkers for diagnosing between diabetic nephropathy and minimal change nephrotic syndrome or membranous nephropathy. *Biochem Biophys Res.* 2021;27:101102. doi:10.1016/j.bbrep.2021.101102
36. Fiseha T, Tamir Z. Urinary markers of tubular injury in early diabetic nephropathy. *Int J Nephrol.* 2016;2016:4647685. doi:10.1155/2016/4647685
37. Fan G, Gong T, Lin Y, et al. Urine proteomics identifies biomarkers for diabetic kidney disease at different stages. *Clin Proteomics.* 2021;18:32. doi:10.1186/s12014-021-09338-6
38. Al-Rubeaan K, Siddiqui K, Al-Ghonaim MA, et al. Assessment of the diagnostic value of different biomarkers in relation to various stages of diabetic nephropathy in type 2 diabetic patients. *Sci Rep.* 2017;7(1):2684. doi:10.1038/s41598-017-02421-9
39. Panduru NM, Forsblom C, Saraheimo M, et al. Urinary liver-type fatty acid-binding protein and progression of diabetic nephropathy in type 1 diabetes. *Diabetes Care.* 2013;36:2077–2083. doi:10.2337/dc12-1868
40. Garg V, Kumar M, Mahapatra HS, et al. Novel urinary biomarkers in pre-diabetic nephropathy. *Clin Exp Nephrol.* 2015;19(5):895–900. doi:10.1007/s10157-015-1085-3
41. Zurbig P, Jerums G, Hovind P, et al. Urinary proteomics for early diagnosis in diabetic nephropathy. *Diabetes.* 2012;61:3304–3313. doi:10.2337/db12-0348
42. Canadas-Garre M, Anderson K, McGoldrick J, et al. Proteomic and metabolomic approaches in the search for biomarkers in chronic kidney disease. *J Proteomics.* 2019;193:93–122. doi:10.1016/j.jprot.2018.09.020
43. Lu H, Deng S, Zheng M, et al. iTRAQ plasma proteomics analysis for candidate biomarkers of type 2 incipient diabetic nephropathy. *Clin Proteomics.* 2019;16(1):33. doi:10.1186/s12014-019-9253-1
44. Caseiro A, Ferreira R, Quintaneiro C, et al. Protease profiling of different biofluids in type 1 diabetes mellitus. *Clin Biochem.* 2012;45:1613–1619. doi:10.1016/j.clinbiochem.2012.08.027
45. Moresco RN, Sangoi MB, De Carvalho JA, et al. Diabetic nephropathy: traditional to proteomic markers. *Clin Chim Acta.* 2013;421:17–30. doi:10.1016/j.cca.2013.02.019
46. Bacci MR, Cavallari MR, de Rozier-Alves RM, et al. The impact of lipocalin-type-prostaglandin-D-synthase as a predictor of kidney disease in patients with type 2 diabetes. *Drug Des Devel Ther.* 2015;9:3179–3182. doi:10.2147/DDDT.S82100
47. Piktel E, Levental I, Durnas B, et al. Plasma gelsolin: indicator of inflammation and its potential as a diagnostic tool and therapeutic target. *Int J Mol Sci.* 2018;20(1):19. doi:10.3390/ijms20010019
48. Cheng Y, Hu X, Liu C, et al. Gelsolin inhibits the inflammatory process induced by LPS. *Cell Physiol Biochem.* 2017;41:205–212. doi:10.1159/000456043
49. Yang Y, Zhang S, Lu B, et al. Predicting diabetic nephropathy by serum proteomic profiling in patients with type 2 diabetes. *Wien Klin Wochenschr.* 2015;127:669–674. doi:10.1007/s00508-014-0679-1
50. Yamanouchi M, Skupien J, Niewczas MA, et al. Improved clinical trial enrollment criterion to identify patients with diabetes at risk of end-stage renal disease. *Kidney Int.* 2017;92:258–266. doi:10.1016/j.kint.2017.02.010
51. Saulnier PJ, Gand E, Velho G, et al. Association of circulating biomarkers (Adrenomedullin, TNFR1, and NT-proBNP) with renal function decline in patients with type 2 diabetes: a French prospective cohort. *Diabetes Care.* 2017;40:367–374. doi:10.2337/dc16-1571
52. Peters KE, Davis WA, Ito J, et al. Identification of novel circulating biomarkers predicting rapid decline in renal function in type 2 diabetes: the Fremantle diabetes study Phase II. *Diabetes Care.* 2017;40:1548–1555. doi:10.2337/dc17-0911
53. Peters KE, Xu J, Bringans SD, et al. PromarkerD predicts renal function decline in type 2 diabetes in the Canagliflozin Cardiovascular Assessment Study (CANVAS). *J Clin Med.* 2020;9(10):3212. doi:10.3390/jcm9103212
54. Beisswenger PJ, Howell SK, Russell GB, et al. Early progression of diabetic nephropathy correlates with methylglyoxal-derived advanced glycation end products. *Diabetes Care.* 2013;36(10):3234–3239. doi:10.2337/dc12-2689
55. El-Asrar MA, Adly AA, Ismail EA. Soluble CD40L in children and adolescents with type 1 diabetes: relation to microvascular complications and glycemic control. *Pediatr Diabetes.* 2012;13:616–624. doi:10.1111/j.1399-5448.2012.00881.x
56. Niewczas MA, Pavkov ME, Skupien J, et al. A signature of circulating inflammatory proteins and development of end-stage renal disease in diabetes. *Nat Med.* 2019;25:805–813. doi:10.1038/s41591-019-0415-5
57. Barutta F, Bellini S, Canepa S, et al. Novel biomarkers of diabetic kidney disease: current status and potential clinical application. *Acta Diabetol.* 2021;58(7):819–830. doi:10.1007/s00592-020-01656-9
58. Blutke A. Opening a treasure chest: glomerular proteome analyses of formalin-fixed paraffin-embedded kidney tissue in the investigation of diabetic nephropathy. *Nephrol Dial Transplant.* 2012;27(5):1695–1698. doi:10.1093/ndt/gfs082
59. Hoyer KJR, Dittrich S, Bartram MP, et al. Quantification of molecular heterogeneity in kidney tissue by targeted proteomics. *J Proteomics.* 2019;193:85–92. doi:10.1016/j.jprot.2018.03.001
60. Vaidya VS, Niewczas MA, Ficociello LH, et al. Regression of microalbuminuria in type 1 diabetes is associated with lower levels of urinary tubular injury biomarkers, kidney injury molecule-1, and N-acetyl-beta-D-glucosaminidase. *Kidney Int.* 2011;79:464–470. doi:10.1038/ki.2010.404

61. Fu WJ, Xiong SL, Fang YG, et al. Urinary tubular biomarkers in short-term type 2 diabetes mellitus patients: a cross-sectional study. *Endocrine*. 2012;41:82–88. doi:10.1007/s12020-011-9509-7
62. Zubiri I, Posada-Ayala M, Benito-Martin A, et al. Kidney tissue proteomics reveals regucalcin downregulation in response to diabetic nephropathy with reflection in urinary exosomes. *Transl Res*. 2015;166:474–484 e474. doi:10.1016/j.trsl.2015.05.007
63. de la Cuesta F, Barderas MG, Calvo E, et al. Secretome analysis of atherosclerotic and non-atherosclerotic arteries reveals dynamic extracellular remodeling during pathogenesis. *J Proteomics*. 2012;75:2960–2971. doi:10.1016/j.jprot.2011.12.005
64. Mi H, Poudel S, Muruganujan A, et al. PANTHER version 10: expanded protein families and functions, and analysis tools. *Nucleic Acids Res*. 2016;44(D1):D336–D342. doi:10.1093/nar/gkv1194
65. Rossi C, Marzano V, Consalvo A, et al. Proteomic and metabolomic characterization of streptozotocin-induced diabetic nephropathy in TIMP3-deficient mice. *Acta Diabetol*. 2018;55:121–129. doi:10.1007/s00592-017-1074-y
66. Satirapoj B, Tassanasorn S, Charoenpitakchai M, et al. Periostin as a tissue and urinary biomarker of renal injury in type 2 diabetes mellitus. *PLoS One*. 2015;10:e0124055. doi:10.1371/journal.pone.0124055
67. Jung CY, Yoo TH. Pathophysiologic mechanisms and potential biomarkers in diabetic kidney disease. *Diabetes Metab J*. 2022;46:181–197. doi:10.4093/dmj.2021.0329
68. Nakatani S, Wei M, Ishimura E, et al. Proteome analysis of laser microdissected glomeruli from formalin-fixed paraffin-embedded kidneys of autopsies of diabetic patients: nephronectin is associated with the development of diabetic glomerulosclerosis. *Nephrol Dial Transplant*. 2012;27:1889–1897. doi:10.1093/ndt/gfr682
69. Fugmann T, Borgia B, Revesz C, et al. Proteomic identification of vanin-1 as a marker of kidney damage in a rat model of type 1 diabetic nephropathy. *Kidney Int*. 2011;80:272–281. doi:10.1038/ki.2011.116
70. Winiarczyk D, Winiarczyk M, Michalak K, et al. Urinary proteome differences in canine diabetes with and without the presence of microalbuminuria. *Animals*. 2022;12(6):748. doi:10.3390/ani12060748
71. Jin J, Gong J, Zhao L, et al. iTRAQ-based comparative proteomics analysis reveals specific urinary biomarkers for various kidney diseases. *Biomark Med*. 2020;14(10):839–854. doi:10.2217/bmm-2019-0556
72. Bellei E, Monari E, Bergamini S, et al. Urinary proteomics in biomarker discovery of kidney-related disorders: diabetic nephropathy and drug-induced nephrotoxicity in chronic headache. *EJIFCC*. 2018;29:290–297.
73. Guillen-Gomez E, Bardaji-de-quixano B, Ferrer S, et al. Urinary proteome analysis identified neprilysin and VCAM as proteins involved in diabetic nephropathy. *J Diabetes Res*. 2018;2018:6165303. doi:10.1155/2018/6165303
74. El-Beblawy NM, Andrawes NG, Ismail EA, et al. Serum and urinary orosomucoid in young patients with type 1 diabetes: a link between inflammation, microvascular complications, and subclinical atherosclerosis. *Clin Appl Thromb Hemost*. 2016;22:718–726. doi:10.1177/1076029616637185
75. Overgaard AJ, Hansen HG, Lajer M, et al. Plasma proteome analysis of patients with type 1 diabetes with diabetic nephropathy. *Proteome Sci*. 2010;8:4. doi:10.1186/1477-5956-8-4
76. Malagrino PA, Venturini G, Yogi PS, et al. Proteome analysis of acute kidney injury - Discovery of new predominantly renal candidates for biomarker of kidney disease. *J Proteomics*. 2017;151:66–73. doi:10.1016/j.jprot.2016.07.019
77. Chun N, Wyatt CM, He JC. Identification of a protective proteomic signature and a potential therapeutic target in diabetic nephropathy. *Kidney Int*. 2017;92:780–781. doi:10.1016/j.kint.2017.08.002
78. Fallon JK, Smith PC, Xia CQ, et al. Quantification of four efflux drug transporters in liver and kidney across species using targeted quantitative proteomics by isotope dilution NanoLC-MS/MS. *Pharm Res*. 2016;33:2280–2288. doi:10.1007/s11095-016-1966-5
79. Konvalinka A, Zhou J, Dimitromanolakis A, et al. Determination of an angiotensin II-regulated proteome in primary human kidney cells by stable isotope labeling of amino acids in cell culture (SILAC). *J Biol Chem*. 2013;288:24834–24847. doi:10.1074/jbc.M113.485326

Diabetes, Metabolic Syndrome and Obesity

Dovepress

Publish your work in this journal

Diabetes, Metabolic Syndrome and Obesity is an international, peer-reviewed open-access journal committed to the rapid publication of the latest laboratory and clinical findings in the fields of diabetes, metabolic syndrome and obesity research. Original research, review, case reports, hypothesis formation, expert opinion and commentaries are all considered for publication. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/diabetes-metabolic-syndrome-and-obesity-journal>