


Use of Cryopreserved Amniotic Membrane During Pterygium Excision: Health Economic Analysis

Neel R Desai, Bryan Adams 

Ophthalmology, The Eye Institute of West Florida, Largo, FL, USA

Correspondence: Neel R Desai, The Eye Institute of West Florida, Eye Care Partners, 1225 W Bay Dr, Largo, FL, 33770, USA, Email desaivision2020@gmail.com

Purpose: To determine the health economic opportunity cost or gain associated with performing pterygium excision surgery using the TissueTuck technique with cryopreserved amniotic membrane (AM) instead of conjunctival autograft (CAU).

Methods: We performed a literature review to determine the average surgical duration of pterygium surgery using CAU with fibrin glue or sutures to calculate the average time saved with the TissueTuck technique. Such time savings was then used to determine the opportunity revenue gain per national average Medicare reimbursement if adjusted to the average surgical duration of cataract surgery.

Results: The time savings achieved using the TissueTuck technique over CAU with fibrin glue is 8.9 min per procedure, which can be applied to additional MSICS or phacoemulsification procedures to generate an opportunity revenue gain of \$1167 or \$762 per 2022 National Average Medicare reimbursement, respectively. After subtracting the current list cost of AmnioGraft (ie, \$645), the opportunity gain is \$522 or \$117 if the time saving is applied to the above procedures, respectively. Alternatively, the time savings achieved by using the TissueTuck technique over CAU with sutures is 23.4 min per procedure, which can be applied to additional MSICS or phacoemulsification procedures to generate an opportunity revenue gain of \$3068 and \$2004 per TissueTuck procedure or \$2423 or \$1359 when accounting for the list cost of AmnioGraft, respectively.

Conclusion: The TissueTuck surgical technique using cryopreserved AM for pterygium takes less time, has lower recurrence rates, and provides an opportunity gain compared to pterygium excision with CAU.

Keywords: amniotic membrane, conjunctival autograft, health economics, pterygium

Introduction

Pterygium is a relatively common ocular disorder in which abnormal fibrovascular tissue of the conjunctiva encroaches onto the cornea, which can impair vision and quality of life.¹ Currently, surgical excision is the only effective treatment option for pterygia that involves meticulous excision of the pterygium followed generally by graft placement with either conjunctival autograft (CAU) or cryopreserved amniotic membrane (AM).² Compared to the bare sclera technique, use of a CAU or AM graft has been shown to improve post-surgical outcomes and reduce recurrence rates to less than 10%.^{3–6}

Ambulatory and outpatient facilities have been increasingly utilized to perform the majority of ophthalmic procedures worldwide. Between 1996 and 2006, the volume of outpatient surgeries at freestanding ambulatory surgery centers (ASCs) increased by 300%.^{7,8} As the demand for outpatient and ambulatory surgery increases, there is also increasing importance to minimize the supply costs or minimize the time to perform ophthalmic surgeries in order to allocate more time for additional revenue generating surgeries. This is particularly relevant for pterygium excision as there are multiple surgical approaches, and the reimbursement to ASCs is the same when using either AM or CAU graft for pterygium excision (CPT code 65426). However, the supply cost of AM grafts is not separately identified or billed to Medicare, which may impel ASCs and their physicians to use CAU to mitigate additional costs. Nevertheless, cryopreserved AM is readily available and avoids the need to harvest a graft, thus, potentially shortening operative time. A recent retrospective study of 582 eyes demonstrated that the TissueTuck procedure takes 14.7 minutes on average.⁹ A shorter surgical duration of using AM with the TissueTuck technique may enable other revenue generating procedures such as cataract

surgery to be performed to offset the supply costs of AM. Thus, the purpose of this health economics study is to determine whether the TissueTuck procedure with AM is a more economically efficient procedure for pterygium excision compared to CAU in the ASC environment.

Materials and Methods

We performed a literature review to determine the average surgical duration for pterygium surgery using CAU with fibrin glue or sutures as well as the average surgical duration for phacoemulsification and manual small incision cataract surgery (MSICS). The difference in surgical times between the TissueTuck and CAU techniques (fibrin glue or sutures) was then calculated to determine the time savings with the TissueTuck technique. The time saved with the TissueTuck technique was then extrapolated to calculate the potential revenue gain if used to perform additional cataract surgeries (MSICS or phacoemulsification). Because pterygium recurrence is a clinically meaningful outcome that is widely reported following surgical excision, a literature review was also conducted to compare the average recurrence rate of the TissueTuck technique to that of CAU with sutures or fibrin glue.

Surgical Time: Cataract Surgery

A literature search of the PubMed electronic database and Google Scholar was performed in July 2022 to retrieve all relevant studies that reported the surgical times of phacoemulsification and MSICS. All studies included in this review were retrieved either from the above databases or from references cited therein. The search results were restricted to articles that were published within the last twenty years (ie, 2002 and onward). No language restrictions were applied. The search strategy combined terms related to disease (“cataract”) with terms related to the procedure (“phacoemulsification”) OR (“manual small incision” OR “MSICS”) and surgical time (“minutes” or “min” or “surgical time”). The titles and abstracts of studies were screened to identify those that fulfilled the inclusion criteria: prospective studies; patients undergoing cataract surgery with phacoemulsification or MSICS; studies that reported the average surgical time. The outcome measure assessed was the surgical time, which was generally defined as total time spent to complete surgery. The mean surgical duration was calculated for both phacoemulsification and MSICS by taking the weighted average of all relevant studies.

Surgical Time: Pterygium Excision with Conjunctival Autograft

A literature search in the PubMed electronic database and Google Scholar was similarly conducted in July 2022 for studies assessing the reported surgical times for pterygium excision with CAU secured with sutures or fibrin glue. Only publications within the last five years (ie, 2017 and onward) were assessed. There were no language restrictions. The keywords used in the search were “fibrin glue”, “suture”, “pterygium”, “conjunctival autograft”, and “surgical time”, or “operative time”, or “minutes”, or “min”. Abstracts were reviewed to determine eligibility, and when the abstract alone could not provide the necessary information, the full text of the article was reviewed. References of relevant articles were also reviewed for eligibility. Only studies that fulfilled the following inclusion criteria were included in the analysis: prospective studies; patients undergoing pterygium surgery with CAU; use of fibrin glue and/or sutures to secure the graft; and studies that reported the average surgical time, which was generally defined as total time spent to complete surgery. The mean surgical duration was calculated for both CAU with fibrin glue and CAU with sutures by taking the weighted average of all relevant studies.

HEOR Analysis

The opportunity gain of performing the TissueTuck procedure with AM (AmnioGraft; BioTissue, Miami, FL) was determined by applying the time savings with the TissueTuck technique to perform additional cataract surgeries. The surgical duration of pterygium excision using CAU with either fibrin glue or sutures was calculated by taking the weighted average of the reported surgical times over the last five years, which were found using the methods listed previously. The average surgical duration of the TissueTuck technique was recently reported to be 14.7 minutes in a large retrospective case series of 582 eyes.⁹ The resultant time savings was measured by calculating the difference in the average operative time between the TissueTuck procedure and the CAU techniques (fibrin glue or sutures).

The time saved with the TissueTuck technique was then extrapolated to calculate the potential revenue gain if used to perform additional cataract surgeries (MSICS or phacoemulsification). This was calculated by dividing the times saving with the TissueTuck technique by the average surgical time of the given cataract surgery (phacoemulsification or MSICS). To determine the potential revenue gain, the above value was then multiplied by the national average Medicare reimbursement for cataract surgery to ASCs (CPT: 66984), which is \$1062 per Medicare.gov. The current list price of AmnioGraft, \$645, was then subtracted to yield the opportunity gain per TissueTuck procedure. All costs were expressed in US dollars.

$$\left(\frac{(CAU - TissueTuck)_{surgicaltime}}{Surgical\ time\ of\ cataract\ surgery} \times \$1062 \right) - \$645$$

Recurrence of Pterygium Excision

A literature search of the PubMed electronic database and Google Scholar was performed in July 2022 to retrieve all prospective, comparative studies that assessed and compared the recurrence rates of primary pterygium excision with CAU using fibrin glue versus sutures. All studies included in this review were either searched directly from the above databases or from relevant references cited therein. The keywords used in the search were “pterygium”, “conjunctival autograft”, “fibrin glue”, “suture”, and “recurrence”. The titles and abstracts of studies were screened to identify those that fulfilled the inclusion criteria: prospective comparative studies or randomized controlled trials; patients with primary pterygium; pterygium surgery with CAU; comparison on the efficacy of fibrin glue versus sutures; recurrence rate at six months. Only studies published within the last twenty years were included, and no language restrictions were applied. The outcome measure assessed was the rate of pterygium recurrence at six months, which was generally defined as the presence of postoperative fibrovascular tissue extending beyond the limbus onto the cornea. The number of cases with recurrence was calculated for each study, and the total recurrence rate was calculated by dividing the total number of recurrences by the total number of subjects of the included studies to yield a cumulative average. The recurrence rate of each procedure was then compared to that of the TissueTuck technique.⁹ To ensure comparability, the recurrence rate using the TissueTuck technique was calculated only for those subjects with primary pterygium and those who completed at least six months of follow-up.

Results

Operative Time: Pterygium Excision with CAU versus TissueTuck Technique

A total of 18 studies assessing the surgical duration of CAU with fibrin glue or sutures met the eligibility criteria and were included for analysis. All studies were randomized controlled trials aside from four prospective comparative studies.^{10–13} Eight studies compared the use of fibrin glue versus sutures,^{12,14–20} six studies compared CAU with fibrin glue to other treatment arms,^{11,21–25} and four studies compared CAU with sutures to other treatment arms.^{10,13,26,27} In brief, the majority of studies harvested the CAU from the superotemporal quadrant of the bulbar conjunctiva, seven of which reported the use of conjunctival limbal autograft to include part of the superficial limbus.^{10,12,13,17,18,24,27} Aside from one study that reported the use of autologous fibrin glue,²⁰ the fibrin glues used in the studies were commercially available products such as Tisseel® (Baxter Healthcare Corporation, Glendale, CA) and FIBINGLURAAS® (Shanghai RAAS Blood Products Co., Ltd, China). In the majority of the studies assessing CAU with sutures, the graft was anchored to the episclera and attached to the adjacent conjunctiva with interrupted 8–0 Vicryl sutures or 10–0 nylon sutures. Cakmak et al¹⁰ and Wang et al¹⁹ reported the use of 7–0 Vicryl and 10–0 Vicryl, respectively. Use of mitomycin c (MMC) was reported in only one study.²⁰ Studies varied with respect to the populations evaluated and surgical technique used; however, these variations were not taken into consideration in this analysis.

Of these 18 studies, the surgical duration was reported for CAU with fibrin glue in 14 prospective studies and CAU with sutures in 12 prospective studies. Of 796 eyes/cases that underwent pterygium excision and CAU with fibrin glue, the weighted average duration of surgery was 23.6 minutes (Table 1).^{11,12,14–25} Twelve studies reported the surgical duration for CAU with sutures, with an average operative time of 38.1 minutes among 640 eyes (Table 2).^{10,12–20,26,27} As

Table 1 Average Operative Time for Pterygium Excision with CAU Secured with Fibrin Glue

Study (Author, Year)	Eyes (N)	Time (Min)
Jha and Simba, 2022 ²²	118	12
Sune and Sune, 2021 ²⁴	31	23.2
Sutyawan et al, 2021 ¹⁷	15	21.8
Huang et al, 2020 ²⁵	30	20.1
Patil and Dasar, 2020 ¹⁴	30	24.2
Singh et al, 2019 ¹⁵	312	33.3
Sati et al, 2019 ²¹	42	15.1
Alamdari et al, 2018 ²⁰	60	10.4
Kumar and Singh, 2018 ¹²	20	36.2
Gong et al, 2018 ¹¹	40	18.8
Wang et al, 2017 ¹⁹	28	20.2
Suryaa et al, 2017 ¹⁶	25	19.5
Wadgaonkar et al, 2017 ¹⁸	25	15.8
Shrivastava et al, 2017 ²³	20	23
TOTAL	N=796	23.6 min

Abbreviations: CAU, conjunctival autograft; Min, Minutes; N, Number.

Table 2 Average Operative Time for Pterygium Excision with CAU Secured with Sutures

Study (Author, Year)	Eyes (N)	Time (Min)
Yang et al, 2021 ¹³	56	36.5
Sutyawan et al, 2021 ¹⁷	15	41.7
Patil and Dasar, 2020 ¹⁴	30	36.1
Singh et al, 2019 ¹⁵	312	43.3
Yan et al, 2019 ²⁷	34	24.3
Alamdari et al, 2018 ²⁰	60	13.3
Kumar and Singh, 2018 ¹²	20	53.3
Celik, 2018 ²⁶	15	48
Cakmak et al, 2017 ¹⁰	20	33.8
Suryaa et al, 2017 ¹⁶	25	50.4
Wadgaonkar et al, 2017 ¹⁸	25	33.6
Wang et al, 2017 ¹⁹	28	32.4
TOTAL	N=640	38.1 min

Abbreviations: CAU, conjunctival autograft; Min, Minutes; N, Number.

previously reported, the average operative time for pterygium excision using the TissueTuck technique is 14.7 minutes.⁹ Thus, the time saved in performing the TissueTuck technique in lieu of CAU with fibrin glue or sutures is 8.9 and 23.4 minutes, respectively.

Operative Time of Cataract Surgery

We performed a similar literature review to determine the average operative time of phacoemulsification and MSICS in order to calculate the opportunity gain if the time saved with the TissueTuck technique was then used to perform cataract surgery. A total of 23 prospective studies reported surgical times for MSICS and phacoemulsification. The average operative time for MSICS and phacoemulsification was reported in 11 and 17 prospective studies, respectively. Among 1721 eyes that underwent MSICS,^{28–38} the average operative time was 8.1 minutes (Table 3). Among 2748 eyes that underwent phacoemulsification, the average operative time was 12.4 minutes in 17 prospective studies (Table 4).^{28,30,35,37–50}

Table 3 Average Operative Time for MSICS

Study (Author, Year)	Eyes (N)	Time (Min)
Parajuli et al, 2020 ³⁴	68	8
Kongsap, 2019 ³³	21	13.2
Bhargava et al, 2016 ²⁸	76	11.2
Singh et al, 2012 ³⁶	110	7.7
Venkatesh et al, 2010 ³⁸	117	8.8
Singh et al, 2009 ³⁷	89	5.3
Ruit et al, 2007 ³⁵	54	9
Gogate et al, 2007 ³⁰	200	8.5
Degenring, 2004 ²⁹	128	11.6
Gogate et al, 2003 ³¹	358	12
Hennig et al, 2002 ³²	500	4
TOTAL	N=1721	8.1 min

Table 4 Average Procedure Times for Phacoemulsification

Study (Author, Year)	Eyes (N)	Time (Min)
Day et al, 2020 ⁴⁰	393	17.8
Roberts et al, 2019 ⁴⁷	200	14.7
Perone et al, 2018 ⁴⁴	85	8
Titiyal et al, 2018 ⁴⁸	77	8.3
Bhargava et al, 2016 ²⁸	70	14.2
Joshi, 2016 ⁴²	32	18
Ugurbas et al, 2012 ⁵⁰	128	12
Baillif et al, 2012 ³⁹	101	19.2
Venkatesh et al, 2010 ³⁸	113	12.2
Raskin et al, 2010 ⁴⁵	50	9.9
Liyanage et al, 2009 ⁴³	105	15
Singh et al, 2009 ³⁷	93	7
Ruit et al, 2007 ³⁵	54	15.5
Gogate et al, 2007 ³⁰	200	15.5
Rengaraj et al, 2004 ⁴⁶	306	6.8
Tsuneoka et al, 2002 ⁴⁹	637	8.7
Funatsu et al, 2002 ⁴¹	104	19.2
TOTAL	N = 2748	12.4 min

HEOR Analysis

The National Average Medicare reimbursement to ASCs for any cataract surgery is \$1062 (CPT code: 66984). Thus, the opportunity gain of performing TissueTuck over CAU with fibrin glue is 109.9% (8.9 min/8.1 min) of the time/revenue of one MSICS (\$1167). When accounting for the current list price of AmnioGraft (\$645), the opportunity gain is \$522 per TissueTuck procedure. If performing phacoemulsification, the opportunity gain is 71.8% (8.9 min/12.4 min) of the time/revenue of one cataract procedure (\$762), yielding an additional \$117 per TissueTuck procedure. Alternatively, if TissueTuck is performed instead of CAU with sutures, the opportunity gain is 288.9% (23.4 min/8.1 min) of one MSICS (\$3068) or an additional \$2423 per TissueTuck procedure. If performing phacoemulsification, the opportunity gain is 188.7% (23.4 min/12.4 min) of one cataract procedure (\$2004), resulting in an additional \$1359 per TissueTuck procedure (Table 5). Based on this analysis, the TissueTuck procedure is no longer economically feasible if the price of AmnioGraft exceeds \$762.

Table 5 Opportunity Gain of the TissueTuck Technique

	Opportunity Gain with MSICS	Opportunity Gain with Phaco
CAU with fibrin glue	\$522 /TT procedure	\$117 /TT procedure
CAU with sutures	\$2423 /TT procedure	\$1359 /TT procedure

Abbreviations: CAU, conjunctival autograft; MSICS, manual small incision cataract surgery; phaco, phacoemulsification; TT, TissueTuck.

Table 6 Recurrence Rates at 6 Months Following Pterygium Excision and Conjunctival Autografting with Sutures or Fibrin Glue

Study (Author, Year)	Recurrence Rate, % (N)	
	Fibrin Glue	Sutures
Cioba et al, 2022 ⁵¹	12.1% (4/33)	0% (0/28)
Sarkar et al, 2020 ⁶⁰	7.7% (2/26)	23.1% (6/26)
Singh et al, 2019 ¹⁵	1.3% (4/312)	2.2% (7/312)
Kumar and Singh, 2018 ¹²	0% (0/20)	5% (1/20)
Parekh et al, 2017 ⁵⁵	0% (0/30)	0% (0/30)
Patkar and Muley, 2017 ⁵⁶	0% (0/30)	6.6% (2/30)
Wang et al, 2017 ¹⁹	3.6% (1/28)	7.1% (2/28)
Qi-Feng and Wei, 2015 ⁵⁸	3.3% (1/30)	10% (3/30)
Priya and Hedge, 2015 ⁵⁷	5% (2/40)	5% (2/40)
Nishant et al, 2014 ⁵³	3.3% (1/30)	20% (6/30)
Vichare et al, 2013 ⁶¹	3.3% (1/30)	10% (3/30)
Rubin et al, 2011 ⁵⁹	4.8% (1/21)	7.7% (2/26)
Yüksel et al, 2010 ⁶²	6.9% (2/29)	13.8% (4/29)
Ozdamar et al, 2008 ⁵⁴	0% (0/12)	0% (0/12)
Koranyi et al, 2004 ⁵²	10% (2/20)	17.4% (4/23)
TOTAL	3.0% (21/691)	6.1% (42/694)

Rate of Recurrence: Pterygium Excision with CAU versus TissueTuck Technique

A total of 15 prospective comparative studies^{12,15,19,51–62} assessed the recurrence rate at six months for eyes receiving CAU with sutures or fibrin glue following excision of primary pterygium. Twelve studies were randomized controlled trials, and three studies were prospective comparative studies, in which randomization was not reported.^{12,55,61} Eight studies reported the use of conjunctival limbal autograft, in which the autograft was dissected towards the cornea to include part of the superficial limbus.^{12,52,54–56,60–62} Aside from one study that reported the use of autologous fibrin glue,⁵¹ the fibrin glue used in the studies were commercially available products such as Tisseel® (Baxter Healthcare Corporation, Glendale, CA). No studies reported use of intra-operative MMC.

From the 15 studies, the overall recurrence rate at six months following excision of primary pterygium with CAU is 3.0% (21/691 eyes) when securing the graft with fibrin glue and 6.1% (42/694 eyes) when using sutures (Table 6).^{12,15,19,51–62} In contrast, the recurrence rate using the TissueTuck technique is 0.4% (2/486 eyes) at six months in eyes presenting with primary pterygium.⁹

Discussion

Compared to hospital and inpatient settings, ASC facilities have allowed for greater productivity and lower healthcare costs without sacrificing the quality of care, which has impacted all relevant healthcare stakeholders including patients, payers, and physicians.^{63,64} ASCs have been particularly beneficial for ophthalmological care, as more than 95% of

ocular surgeries are performed in outpatient settings.⁶⁵ Nonetheless, ASC facilities are confronted with increasing costs of supplies, administration, and labor and decreasing payer care contracts. Hence, importance has been placed on procedural efficiency to identify surgical techniques that can lower input costs while maximizing outputs, such as revenue and of quality of care.⁶⁶

In this health economic analysis, we focused on the procedural efficiency of pterygium surgery which is a commonly performed procedure in ophthalmic ASCs.⁶⁷ Meta-analysis suggests that the overall prevalence of pterygium is 10.2%, though not all cases proceed with surgical excision.⁶⁸ The most common techniques of performing pterygium surgery involve the use of a CAU or AM graft that is applied to the ocular surface with fibrin glue or sutures. Of these techniques, our analysis demonstrated that the average surgical time of using AM with fibrin glue via the TissueTuck procedure was substantially lower than the time needed to perform pterygium excision with CAU with either fibrin glue or sutures, saving approximately 9 to 23 minutes, respectively. Furthermore, using AM with fibrin glue was the most cost-effective intervention, even though the initial supply cost of AM was greater than CAU. This cost effectiveness is primarily a result of the opportunity to perform additional cataract surgeries with the time savings, which is one of the most commonly performed surgical procedures.^{69,70} The overall opportunity gain of using AM with fibrin glue over CAU techniques ranged from \$117 to \$2423 per pterygium surgery. In order for CAU with fibrin glue or suture to be the lower cost alternative procedure, the overall supply cost of AM would need to exceed \$762 or \$2004, respectively, which is far greater than the current list price of \$645. As a result, the procedural efficiency of AM greatly reduces the operating time and overall total operation cost despite an obvious supply cost difference between AM and CUA.

Clinically, our analysis also demonstrated that the pterygium recurrence rate is lower with AM and fibrin glue when compared to CAU with fibrin glue or suture (0.4% vs 3.0–6.1%). Although the clinical superiority of AM over CAU is still an ongoing debate, and the recurrence rates after pterygium surgery are variable in the literature, we have reported that the TissueTuck technique using AM and fibrin glue is a highly reproducible and formulaic technique that produces similar outcomes regardless of pterygium size.⁹ Aside from a low rate of recurrence, we have also previously shown that the TissueTuck technique yields a great aesthetic outcome with low rates of other complications attributed to pterygium excision such as granuloma formation and diplopia in extreme lateral gaze. Furthermore, cryopreserved AM is readily available and avoids the need to harvest a graft in case of future glaucoma surgery. Hence, procedural efficiency of performing surgery with AM and fibrin glue is substantiated by increasing productivity without adversely affecting the quality of care.

There are several limitations with the present study. First, the cost-effectiveness of using AM may vary depending on the negotiated supply cost at different ASCs across different regions of the country. Furthermore, the current analysis did not consider the supply cost of suture versus fibrin glue, hospital versus freestanding ASC surgical times, potential realized and opportunity cost of revision surgeries and postoperative care due to pterygium recurrence or other complications, overtime for staff to complete the day's list of cases, and surgical duration from other surgeons using the TissueTuck technique. However, our analysis is quite thorough as we compared the outcomes of 582 cases using the TissueTuck technique to 640 to 796 cases of CAU technique using only prospective studies. Furthermore, there may be additional value from the patients' perspective, as there could be shorter waiting times and better quality of life as prolonged operative duration has been associated with complications.⁷¹ Those patients may also be more likely to return for other surgeries and refer their colleagues to the surgeon that performed the pterygium excision. This is especially relevant to many physicians performing the surgeries, as 90% of ASCs are owned in-part by physicians and 65% are solely owned by physicians.⁷² Additionally, this will benefit the overall healthcare burden as it accommodates the increased demand for outpatient services, as there is an increasing incidence of surgical procedures performed and improved patient access to better insurance coverage. Another limitation of this analysis is that the rate of recurrence was assessed at six months, which has been reported by several prospective comparative studies as demonstrated by our literature review. Nevertheless, we previously reported that the recurrence rate using the TissueTuck technique was 1.7% (7/420) at one year for both primary and recurrent pterygium,⁹ which is still lower than the cumulative average rate of recurrence at six months for CAU with sutures and fibrin glue (6.1% and 3.0%, respectively). Overall, this study should aid those ASC facilities interested in optimizing procedural efficiencies and builds off prior findings that showed CAU with fibrin glue is more cost-effective than CAU with sutures.⁷³

Acknowledgments

The authors would like to thank TissueTech for awarding an investigator-initiated grant for this project and assisting in the drafting of the manuscript.

Disclosure

Dr. Desai is a speaker, consultant, and stockholder for BioTissue, Inc. and received a grant to support the analysis and publication cost of this study.

References

1. Rezvan F, Khabazkhoob M, Hooshmand E, Yekta A, Saatchi M, Hashemi H. Prevalence and risk factors of pterygium: a systematic review and meta-analysis. *Surv Ophthalmol*. 2018;63:719–735. doi:10.1016/j.survophthal.2018.03.001
2. Hacıoglu D, Erdol H. Developments and current approaches in the treatment of pterygium. *Int Ophthalmol*. 2017;37:1073–1081. doi:10.1007/s10792-016-0358-5
3. Prajna NV, Devi L, Seeniraj SK, Keenan JD. Conjunctival autograft versus amniotic membrane transplantation after double pterygium excision: a randomized trial. *Cornea*. 2016;35:823–826. doi:10.1097/ICO.0000000000000812
4. Kucukerdonmez C, Akova YA, Altinors DD. Comparison of conjunctival autograft with amniotic membrane transplantation for pterygium surgery: surgical and cosmetic outcome. *Cornea*. 2007;26:407–413. doi:10.1097/ICO.0b013e318033b3d4
5. Akbari M, Soltani-Moghadam R, Elmi R, Kazemnejad E. Comparison of free conjunctival autograft versus amniotic membrane transplantation for pterygium surgery. *J Curr Ophthalmol*. 2017;29:282–286. doi:10.1016/j.joco.2017.08.003
6. Ma DH, See LC, Liao SB, Tsai RJ. Amniotic membrane graft for primary pterygium: comparison with conjunctival autograft and topical mitomycin C treatment. *Br J Ophthalmol*. 2000;84:973–978. doi:10.1136/bjo.84.9.973
7. Munnich EL, Parente ST. Procedures take less time at ambulatory surgery centers, keeping costs down and ability to meet demand up. *Health Aff*. 2014;33:764–769. doi:10.1377/hlthaff.2013.1281
8. Cullen KA, Hall MJ, Golosinskiy A. Ambulatory surgery in the United States, 2006. *Natl Health Stat Report*. 2009;2009:1–25.
9. Desai NR, Adams B. Cryopreserved amniotic membrane using the tissuetuck technique: a sutureless approach for pterygium surgery. *Cornea*. 2022;42(2):181–185. doi:10.1097/ICO.00000000000003111
10. Cakmak HB, Dereli Can G, Can ME, Cagil N. A novel graft option after pterygium excision: platelet-rich fibrin for conjunctivoplasty. *Eye*. 2017;31:1606–1612. doi:10.1038/eye.2017.109
11. Gong J, Fan J, Shen T, Jiang J. Comparison of self-made cryopreservative fibrin glue and commercial fibrin glue kit in pterygium surgery: 1-year follow-up. *Acta Ophthalmol*. 2018;96:e152–e155. doi:10.1111/aos.13478
12. Kumar S, Singh R. Pterygium excision and conjunctival autograft: a comparative study of techniques. *Oman J Ophthalmol*. 2018;11:124–128. doi:10.4103/ojo.OJO_6_2017
13. Yang N, Xing Y, Zhao Q, Zeng S, Yang J, Du L. Application of platelet-rich fibrin grafts following pterygium excision. *Int J Clin Pract*. 2021;75:e14560. doi:10.1111/ijcp.14560
14. Patil M, Dasar LV. Study of efficacy of fibrin glue versus sutures in pterygium. *IJBAMR*. 2020;9:226–232.
15. Singh TA, Mamota N, Uday V, et al. Comparative study of conjunctival autograft with suture and fibrin glue in primary progressive pterygium. *IJCB*. 2019;34(2):230–233. doi:10.1007/s12291-017-0728-4
16. Suryaa MA, Prasanth HR, Nirmale SD, Hansdak A. A randomized control trial comparing the efficacy of fibrin glue with sutures in primary pterygium surgery. *Int J Ocul Oncol Oculoplasty*. 2017;3:36–39.
17. Sutayawan E, Niti S, Widiana R. Effect of conjunctiva-limbus transplantation with fibrin glue compare to suture technique on stability of the graft attachment in pterygium surgery. *Bali Med J*. 2021;2:27–32.
18. Wadgaonkar S, Tiwari R, Patil P, Kamble B. Fibrin glue versus suture technique for pterygium excision: a prospective study in tertiary-based rural hospital. *J Clin Ophthalmol Res*. 2017;5:23–27. doi:10.4103/2320-3897.195305
19. Wang X, Zhang Y, Zhou L, Wei R, Dong L. Comparison of fibrin glue and Vicryl sutures in conjunctival autografting for pterygium surgery. *Mol Vis*. 2017;23:275–285.
20. Alamdari DH, Sedaghat M-R, Alizadeh R, Zarei-Ghanavati S, Naseri H, Sharifi F. Comparison of autologous fibrin glue versus nylon sutures for securing conjunctival autografting in pterygium surgery. *Int Ophthalmol*. 2018;38:1219–1224. doi:10.1007/s10792-017-0585-4
21. Sati A, Banerjee S, Kumar P, Kaushik J, Khera A. Mini-simple limbal epithelial transplantation versus conjunctival autograft fixation with fibrin glue after pterygium excision: a randomized controlled trial. *Cornea*. 2019;38:1345–1350. doi:10.1097/ICO.0000000000002007
22. Jha A, Simba A. Conjunctival autograft versus combined amniotic membrane and mini-simple limbal epithelial transplant for primary pterygium excision. *J Ophthalmic Vis Res*. 2022;17:4.
23. Shrivastava S, Patkar P, Ramakrishnan R, Kanhere M, Riaz Z. Comparison of fibrin glue and autologous serum for conjunctival autograft fixation in pterygium. *Delta J Ophthalmol*. 2017;18:133–137. doi:10.4103/DJO.DJO_18_17
24. Sune M, Sune P. Comparison between autologous blood and fibrin glue for adhering conjunctival autografts after pterygium excision- a randomised clinical trial. *J Clin Diagn Res*. 2021;15:NC1–NC4.
25. Huang H, Li S, Zhong J, et al. Evaluation of the safety and efficacy of a low-temperature plasma surgical system for pterygium. *Cornea*. 2020;39:1581–1587. doi:10.1097/ICO.0000000000002487
26. Celik T. In situ blood coagulum versus sutures for autograft fixation after pterygium excision. *Curr Eye Res*. 2018;43:977–980. doi:10.1080/02713683.2018.1470247
27. Yan B, Peng L, Peng H, Zhou S, Chen B. Modified sutureless and glue-free method versus conventional sutures for conjunctival autograft fixation in primary pterygium surgery: a randomized controlled trial. *Cornea*. 2019;38:1351–1357. doi:10.1097/ICO.0000000000002137

28. Bhargava R, Kumar P, Sharma SK, Arora Y. Phacoemulsification versus manual small incision cataract surgery in patients with fuchs heterochromic iridocyclitis. *Asia Pac J Ophthalmol*. 2016;5:330–334. doi:10.1097/APO.0000000000000191
29. Degenring RF, Vey S, Kampmpeter B, Budde W, Sauder G, Jonas JB. Central retinal thickness after uncomplicated small-incision cataract surgery. *Invest Ophthalmol Vis Sci*. 2004;45:2981.
30. Gogate P, Deshpande M, Nirmalan PK. Why do phacoemulsification? Manual small-incision cataract surgery is almost as effective, but less expensive. *Ophthalmology*. 2007;114:965–968. doi:10.1016/j.ophtha.2006.08.057
31. Gogate PM, Deshpande M, Wormald RP. Is manual small incision cataract surgery affordable in the developing countries? A cost comparison with extracapsular cataract extraction. *Br J Ophthalmol*. 2003;87:843–846. doi:10.1136/bjo.87.7.843
32. Hennig A, Kumar J, Yorston D, Foster A. Sutureless cataract surgery with nucleus extraction: outcome of a prospective study in Nepal. *Br J Ophthalmol*. 2003;87:266–270. doi:10.1136/bjo.87.3.266
33. Kongsap P. Central corneal thickness changes following manual small incision cataract surgery versus phacoemulsification for white cataract. *Rom J Ophthalmol*. 2019;63:61. doi:10.22336/rjo.2019.10
34. Parajuli S, Shrestha R, Chapagain S, Singh P, Shrestha R. Changes in central corneal thickness (CCT) and central macular thickness (CMT) following uncomplicated small-incision cataract surgery (SICS). *medRxiv*. 2020;2020(11):20228932.
35. Ruit S, Tabin G, Chang D, et al. A prospective randomized clinical trial of phacoemulsification vs manual sutureless small-incision extracapsular cataract surgery in Nepal. *Am J Ophthalmol*. 2007;143(32–38):e2. doi:10.1016/j.ajo.2006.07.023
36. Singh P, Singh S, Bhargav G, Singh M. Conjunctival flap in manual sutureless small-incision cataract surgery: a necessity or dogmatic. *Int Ophthalmol*. 2012;32:349–355. doi:10.1007/s10792-012-9569-6
37. Singh S, Winter I, Surin L. Phacoemulsification versus small incision cataract surgery (SICS): which one is a better surgical option for immature cataract in developing countries? *Nepal J Ophthalmol*. 2009;1:95–100. doi:10.3126/nepjoph.v1i2.3682
38. Venkatesh R, Tan CSH, Sengupta S, Ravindran RD, Krishnan KT, Chang DF. Phacoemulsification versus manual small-incision cataract surgery for white cataract. *J Cataract Refract Surg*. 2010;36:1849–1854. doi:10.1016/j.jcrs.2010.05.025
39. Baillif S, Roure-Sobas C, Le-Duff F, Kodjikian L. Aqueous humor contamination during phacoemulsification in a university teaching hospital. *J Fr Ophtalmol*. 2012;35:153–156. doi:10.1016/j.jfo.2011.05.007
40. Day AC, Burr JM, Bennett K, et al. Femtosecond laser-assisted cataract surgery compared with phacoemulsification cataract surgery: randomized noninferiority trial with 1-year outcomes. *J Cataract Refract Surg*. 2020;46:1360–1367. doi:10.1097/j.jcrs.0000000000000257
41. Funatsu H, Yamashita H, Noma H, Shimizu E, Mimura T, Hori S. Prediction of macular edema exacerbation after phacoemulsification in patients with nonproliferative diabetic retinopathy. *J Cataract Refract Surg*. 2002;28:1355–1363. doi:10.1016/S0886-3350(02)01243-9
42. Joshi RS. Phacoemulsification without preoperative mydriasis in patients with age-related cataract associated with type 2 diabetes. *Clin Ophthalmol*. 2016;10:2427–2432. doi:10.2147/OPTH.S122107
43. Liyanage SE, Angunawela RI, Wong SC, Little BC. Anterior chamber instability caused by incisional leakage in coaxial phacoemulsification. *J Cataract Refract Surg*. 2009;35:1003–1005. doi:10.1016/j.jcrs.2009.02.015
44. Perone JM, Boiche M, Lhuillier L, et al. Correlation between postoperative central corneal thickness and endothelial damage after cataract surgery by phacoemulsification. *Cornea*. 2018;37:587–590. doi:10.1097/ICO.0000000000001502
45. Raskin E, Paula JS, Cruz AAV, Coelho RP. Effect of bevel position on the corneal endothelium after phacoemulsification. *Arq Bras Oftalmol*. 2010;73:508–510. doi:10.1590/S0004-27492010000600008
46. Rengaraj V, Radhakrishnan M, Eong K-GA, et al. Visual experience during phacoemulsification under topical versus retrobulbar anesthesia: results of a prospective, randomized, controlled trial. *Am J Ophthalmol*. 2004;138:782–787. doi:10.1016/j.ajo.2004.06.023
47. Roberts HW, Wagh VK, Sullivan DL, et al. A randomized controlled trial comparing femtosecond laser-assisted cataract surgery versus conventional phacoemulsification surgery. *J Cataract Refract Surg*. 2019;45:11–20. doi:10.1016/j.jcrs.2018.08.033
48. Titiyal JS, Kaur M, Ramesh P, et al. Impact of clear corneal incision morphology on incision-site Descemet membrane detachment in conventional and femtosecond laser-assisted phacoemulsification. *Curr Eye Res*. 2018;43:293–299. doi:10.1080/02713683.2017.1396616
49. Tsuneoka H, Shiba T, Takahashi Y. Ultrasonic phacoemulsification using a 1.4 mm incision: clinical results. *J Cataract Refract Surg*. 2002;28:81–86. doi:10.1016/S0886-3350(01)01235-4
50. Ugurbas SC, Caliskan S, Alpay A, Ugurbas SH. Impact of intelligent phacoemulsification software on torsional phacoemulsification surgery. *Clin Ophthalmol*. 2012;6:1493–1498. doi:10.2147/OPTH.S35283
51. Cioba C, Marafon SB, Fortes BGB, et al. Autologous fibrin glue versus suture for conjunctival autograft in primary pterygium: a randomized clinical trial. *Res Square*. 2022;2022:1.
52. Koranyi G, Seregard S, Kopp ED. Cut and paste: a no suture, small incision approach to pterygium surgery. *Br J Ophthalmol*. 2004;88:911–914. doi:10.1136/bjo.2003.032854
53. Nishant K, Prasad V, Shah Nawaz A, Akbar MA. Comparison of ‘cut and paste (using fibrin glue)’ vs ‘cut and suture (using 8-0 vicryl sutures)’ techniques of pterygium surgery. *Int J Curr Res Rev*. 2014;6:64–76.
54. Ozdamar Y, Mutevelli S, Han U, et al. A comparative study of tissue glue and vicryl suture for closing limbal-conjunctival autografts and histologic evaluation after pterygium excision. *Cornea*. 2008;27:552–558. doi:10.1097/ICO.0b013e318165b16d
55. Parekh K, Patil MS, Balwir D. Fibrin glue vs vicryl (10-0) suture in limbo conjunctival autograft in the management of primary pterygium: a prospective comparative study. *MVP J Med Sci*. 2017;2017:4.
56. Patkar P, Muley S. A comparative study of fibrin glue vs vicryl sutures for conjunctival autografting in pterygium surgery. *Indian J Res*. 2019;8:36–39.
57. Priya VR, Hegde SS. A prospective comparative study to evaluate the outcome of fibrin glue versus sutures for conjunctival autografting after primary pterygium excision. *Int J Health Sci Res*. 2015;5:93–99.
58. Qi-Feng L, Wei C. Clinical observation on fibrin glue application during pterygium surgery. *Int Eye Sci*. 2015;2015:364–366.
59. Rubin MR, Dantas PE, Nishiwaki-Dantas MC, Felberg S. Eficácia do adesivo tecidual de fibrina na fixação de enxerto conjuntival autógeno em cirurgias de pterígio primário [Efficacy of fibrin tissue adhesive in the attachment of autogenous conjunctival graft on primary pterygium surgery]. *Arq Bras Oftalmol*. 2011;74:123–126. Portuguese. doi:10.1590/S0004-27492011000200011
60. Sarkar KC, Sarkar P, Mondal S, Saha RN. Comparison between fibrin glue and polyglactin sutures in limbal based conjunctival autograft technique in primary pterygium surgery in adults- A randomised controlled trial. *IJCEO*. 2020;6:35–40. doi:10.18231/j.ijceo.2020.009

61. Vichare N, Choudhary T, Arora P. A comparison between fibrin sealant and sutures for attaching conjunctival autograft after pterygium excision. *Med J Armed Forces India*. 2013;69:151–155. doi:10.1016/j.mjafi.2012.09.002
62. Yüksel B, Unsall SK, Onat S. Comparison of fibrin glue and suture technique in pterygium surgery performed with limbal autograft. *Int J Ophthalmol*. 2010;3:316–320. doi:10.3980/j.issn.2222-3959.2010.04.09
63. Barron E, Knoble JK. Ambulatory surgery offers quality, savings. *Hospitals*. 1980;54:74–76.
64. Fan YP, Boldy D, Bowen D. Comparing patient satisfaction, outcomes and costs between cataract day surgery and inpatient surgery for elderly people. *Aust Health Rev*. 1997;20:27–39. doi:10.1071/AH970027
65. Stagg BC, Talwar N, Mattox C, Lee PP, Stein JD. Trends in use of ambulatory surgery centers for cataract surgery in the United States, 2001–2014. *JAMA Ophthalmol*. 2018;136:53–60. doi:10.1001/jamaophthalmol.2017.5101
66. Marcinko DE, Hetico HR. Economic outcomes analysis from an ambulatory surgical center. *J Foot Ankle Surg*. 1996;35:544–549. doi:10.1016/S1067-2516(96)80128-1
67. Usmani B, Iftikhar M, Latif A, Shah SMA. Epidemiology of primary ophthalmic procedures performed in the United States. *Can J Ophthalmol*. 2019;54:727–734. doi:10.1016/j.jcjo.2019.03.006
68. Liu L, Wu J, Geng J, Yuan Z, Huang D. Geographical prevalence and risk factors for pterygium: a systematic review and meta-analysis. *BMJ open*. 2013;3:e003787. doi:10.1136/bmjopen-2013-003787
69. Brunin G, Khan K, Biggerstaff KS, Wang L, Koch DD, Khandelwal SS. Outcomes of femtosecond laser-assisted cataract surgery performed by surgeons-in-training. *Graefes Arch Clin Exp Ophthalmol*. 2017;255:805–809. doi:10.1007/s00417-016-3581-x
70. Hansen MS, Hardten DR. Financially efficient cataract surgery in today's healthcare environment. *Curr Opin Ophthalmol*. 2015;26:61–65. doi:10.1097/ICU.0000000000000120
71. Cheng H, Clymer JW, Po-Han Chen B, et al. Prolonged operative duration is associated with complications: a systematic review and meta-analysis. *J Surg Res*. 2018;229:134–144. doi:10.1016/j.jss.2018.03.022
72. Badlani N. Ambulatory surgery center ownership models. *J Spine Surg*. 2019;5:S195–S203. doi:10.21037/jss.2019.04.20
73. Bouhout S, Kam J, Robert MC, Harissi-Dagher M. Cost-effectiveness analysis: fibrin glue versus sutures for conjunctival fixation during pterygion surgery. *Can J Ophthalmol*. 2022;57:41–46. doi:10.1016/j.jcjo.2021.02.016

Clinical Ophthalmology

Dovepress

Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-ophthalmology-journal>