

Impaired Sensitivity to Thyroid Hormones Is Associated with Mild Cognitive Impairment in Euthyroid Patients with Type 2 Diabetes [Letter]

Jingxia Sun, Wensheng Lu

Department of Endocrinology and Metabolism, Guangxi Academy of Medical Sciences and the People's Hospital of Guangxi Zhuang Autonomous Region, Nanning, Guangxi, 530021, People's Republic of China

Correspondence: Wensheng Lu, Email Lwsxqz@163.com

Dear editor

We extend our appreciation to Yu et al for their original publication entitled “Impaired Sensitivity to Thyroid Hormones is Associated with Mild Cognitive Impairment in Euthyroid Patients with Type 2 Diabetes”¹ in Journal Clinical Interventions in Ageing, which presents us with some novel ideas.

This study mainly explored the association between mild cognitive impairment (MCI) and thyroid hormone sensitivity in euthyroid type 2 diabetes mellitus (T2DM) patients. The study eventually concluded that impaired sensitivity to thyroid hormones is associated with MCI in euthyroid patients with T2DM.

Given their extensive and continuous research in this field over the years, we wholeheartedly agree with the conclusions the China Medical University endocrine metabolism team reached. However, the design of this study, particularly the exclusion criteria, still requires refining to provide precise data for routine clinical practice. The following points, in our opinion, require further clarification. First, there is a lack of Vitamin D data. In older people with T2DM, vitamin D level is tightly related to cognitive function.² Usually, the vitamin D levels of Chinese people are generally low. Second, there is a lack of bone mineral density (BMD) and bone turnover markers (BTMs) data. According to extensive evidence-based evidence, patients with osteoporosis and bone loss often suffer from mild cognitive impairment.³ Thirdly, lack of nutrition status data. Nutritional status, including anemia, hypoproteinemia, and hyponatremia, is strongly connected with cognitive impairment.⁴ Finally, there is a lack of an estimated glomerular filtration rate (eGFR). Cognitive impairment usually occurs in CKD.⁵

As a result, the essential way to get the correct findings from the study design is to include the four scenarios indicated above in the exclusion criteria. Despite a few flaws, we nevertheless value the authors' original viewpoint. Their findings will remind clinicians to pay closer attention to impaired cognition in euthyroid patients with T2DM as early as possible and adopt effective intervention strategies to prevent the occurrence of cognitive impairment to enhance patient quality of life in clinical practice.

Disclosure

The authors report no conflicts of interest in this communication.

References

1. Yu ZW, Pu SD, Sun XT, et al. Impaired sensitivity to thyroid hormones is associated with mild cognitive impairment in euthyroid patients with type 2 diabetes. *Clin Interv Aging*. 2023;18:1263–1274. doi:10.2147/CIA.S413584
2. Arosio B, Rossi PD, Ferri E, Cesari M, Vitale G. Characterization of Vitamin D status in older persons with cognitive impairment. *Nutrients*. 2022;14(6):1142. doi:10.3390/nu14061142
3. Zhou R, Zhou H, Rui L, Xu J. Bone loss and osteoporosis are associated with conversion from mild cognitive impairment to Alzheimer's disease. *Curr Alzheimer Res*. 2014;11(7):706–713. doi:10.2174/1567205011666140812115818

4. Scarmeas N, Anastasiou CA, Yannakoulia M. Nutrition and prevention of cognitive impairment. *Lancet Neurol.* 2018;17(11):1006–1015. doi:10.1016/S1474-4422(18)30338-7
5. Drew DA, Weiner DE, Sarnak MJ. Cognitive impairment in CKD: pathophysiology, management, and prevention. *Am J Kidney Dis.* 2019;74(6):782–790. doi:10.1053/j.ajkd.2019.05.017

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Interventions in Aging 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Interventions in Aging editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Clinical Interventions in Aging

Dovepress

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>

<https://doi.org/10.2147/CIA.S437073>